

## Secretion Defects That Activate the Phage Shock Response of *Escherichia coli*

Susan E. Jones, Louise J. Lloyd, Kum K. Tan, and Martin Buck\*

Department of Biological Sciences, Imperial College London, South Kensington Campus, London SW7 2AZ, United Kingdom

Received 23 April 2003/Accepted 20 August 2003

**The phage shock protein (*psp*) operon of *Escherichia coli* is induced by membrane-damaging cues. Earlier studies linked defects in secretion across the inner membrane to induction of the *psp* response. Here we show that defects in *yidC* and *sec* secretion induce *psp* but that defects in *tat* and *srp* have no effect. We have also determined the cellular location of PspB and PspD proteins.**

The *pspABCDE* operon of *Escherichia coli* is specifically associated with a wide variety of membrane stresses. Expression of the *pspABCDE* operon is strongly induced by the infection of *E. coli* with filamentous phage f1 and, more specifically, by expression of f1 gene IV, which encodes protein IV (pIV). The pIV protein is a member of the secretin family that forms an outer membrane (OM) pore through which phage is extruded. Other stresses, including heat shock, osmotic shock, and exposure to organic solvents, induce *psp* (for a review, see reference 24). Overexpression of PspA, an OM secretin, also mediated up-regulation of PspA; however, this effect was reduced when the periplasmic chaperone PulS was coexpressed (14). Translocation of mutant forms of the OM protein maltoporin, encoded by *lamB* (7), and PhoE, a pore-forming OM protein (16), resulted in up-regulation of PspA. Furthermore, chromosomal mutations in *secA*, *secD*, and *secF* up-regulate expression of PspA, whereas mutations in *secE* and *secY* do not. For the *secA51* mutation, it was speculated that proteins might be trapped inside the translocase due to interruption of SecA ATP hydrolysis functioning, but this was not confirmed experimentally (16). SecDF depletion leads to cells being unable to maintain the proton motive force (PMF) across the inner membrane (2). PspA is not part of the secretion apparatus but may act to maintain PMF, since in an *E. coli* mutant lacking *pspA*, PMF decreased specifically compared to the level in the wild type when a mutant form of PhoE was overproduced (17). Recent work has identified new participants in protein secretion in *E. coli*, which led us to reexamine the determinants for induction of the *psp* response. We wished to clarify whether *psp* specifically responds to Sec-related stress or to more general secretion stresses at the inner membrane.

**Secretion of proteins destined for the periplasm or OM is mediated by several different systems in *E. coli*.** The SecYEG translocase translocates proteins across the inner membrane. Secretion requires energy (from ATP hydrolysis) and the proton gradient and electrochemical potential components of the PMF. Since earlier studies linking secretion problems to *psp* induction were carried out, several other components of the

secretion apparatus, including the bacterial signal recognition particle (SRP), the Tat system, and YidC, have been identified. The bacterial SRP, which presents proteins to the Sec translocase, functions to target and integrate cytoplasmic membrane proteins. In *E. coli*, the SRP comprises *ffh* (48-kDa protein), *ffs* (4.5S RNA), and *ftsY* (the SRP receptor in the inner membrane) (3, 25, 27, 32). The Tat system (twin-arginine translocation pathway) transports a specific subset of fully folded *E. coli* proteins across the bacterial inner membrane (reviewed in reference 26). YidC copurifies with SecY and is part of the *secYEG/secDF yajC* complex. YidC interacts with the translocase through a bridging interaction with SecDF YajC. YidC can work with the Sec translocase but also functions independently to facilitate the insertion of Sec-dependent and Sec-independent proteins into the membrane bilayer (9). Some evidence suggests that the SecDF complex is involved in the release of proteins into the periplasmic space (21) and can prevent the backsliding of partly translocated proteins from the translocase to the cytoplasm (11). YidC is hypothesized to contribute to the clearing of the translocon channel. YidC depletion does not affect protein export unless a Sec-dependent protein is overexpressed. Its primary role is in membrane protein topogenesis (reviewed in reference 9). Yi et al. (35) have recently shown a role for YidC in the function of the F<sub>1</sub>F<sub>0</sub> ATP synthase.

One hypothesis for *psp* induction is that slowed translocation, or slowed folding of an OM protein into its correct conformation, may be a trigger for PspA synthesis. Therefore, we hypothesized that defects in the SRP, Tat, or YidC may trigger the induction of *psp*. We decided to further probe the nature of the inducing signal for *psp* through a combination of depletion experiments and defined mutations. Induction signals for *psp* are poorly defined, placing an emphasis on the need to establish which translocation systems might be sensed by the *psp* system. Crucially, several systems previously used to study the induction of *psp* utilized an overexpression of membrane proteins which might themselves interact with Psp proteins. All the mutants used in this study have been well characterized and are chromosomally located, and the assays that were applied rely on conventional growth conditions without overexpression of membrane proteins, except key positive control assays. Pairs of strains used were isogenic or very closely related.

\* Corresponding author. Mailing address: Department of Biological Sciences, Sir Alexander Fleming Building, Imperial College London, South Kensington Campus, London SW7 2AZ, United Kingdom. Phone: 44 020 7594 5442. Fax: 44 020 7594 5419. E-mail: m.buck@imperial.ac.uk.

**Mutations in all the components of the bacterial SRP do not up-regulate *pspA* expression.** A reporter gene fusion construct (pSJ1) was made by PCR amplification of the *pspA* promoter with primers SE5 and SE6 (13) and then cloning into vector pMR25 to fuse the *pspA* promoter to *lacZ*. Verification was done by sequence analysis. The clone, pSJ1, allowed a specific assay for levels of *pspA* transcription, since stop codons are present in all three reading frames and a translational start site is provided for the promoterless *lacZ* gene (Table 1). The vector was present at one to five copies per cell.  $\beta$ -Galactosidase assays of the appropriate plasmid-containing strain, with tetracycline present at 10  $\mu$ g/ml, were carried out with a 100-fold dilution of overnight culture into Luria-Bertani broth (LB) with the appropriate antibiotics (28). Cultures were grown at 37°C to an optical density at 600 nm (OD<sub>600</sub>) of 0.6 to 0.8, and the activity was assayed (23) and expressed as the number of units at an OD<sub>600</sub> (Miller units). By classical induction with pIV harbored on pPMR129 (10), *pspA* up-regulation was confirmed (data not shown). pMR25 (control vector) or pSJ1 (*pspA-lacZ*) was transformed into HPT strains (Table 1), each of which harbors defined point mutations in various components of the bacterial SRP (33).  $\beta$ -Galactosidase activity was determined after growth overnight (data not shown) or after growth to an OD<sub>600</sub> of between 0.6 and 0.8 (Table 1). In all cases, the levels of  $\beta$ -galactosidase observed were comparable to those of the appropriate parental strain. Hence, mutations that cause defects in the presentation of inner membrane proteins to the translocon (and hence, accumulation of proteins in the cytoplasm) do not confer any significant up-regulation of *psp*. pMR25 and pSJ1 were also introduced into MC4100 (wild type) and HDB45 (*ffh::kan-1*; resulting in pHDB4, where pHDB4 harbors *ffh* under *trc* promoter control) (4). HDB45 can be used to deplete levels of the SRP. Strains were all grown in the presence of 10 mM IPTG (isopropyl- $\beta$ -D-thiogalactopyranoside) to maintain Ffh levels. Where necessary, cultures were washed with LB to remove IPTG and grown in LB without IPTG to determine the effect of depleting Ffh. Table 1 shows that depletion of Ffh in HDB45 did not up-regulate *pspA* expression, confirming that defects in the SRP fail to induce *pspA*. In contrast, both HPT73 and HPT93 up-regulated *psp* at least sixfold (compare the results for the HPT57 wild type harboring pSJ1 to those for HPT73 and HPT93 harboring pSJ1) (Table 1). Both strains harbor mutations in *secM*, which is not part of the SRP. Western blotting was used to determine whether PspA was detectably up-regulated at the protein level (Fig. 1). For generation of antibodies, the *pspA* gene from pET28b<sup>+</sup> harboring a six-His-tagged *pspA* gene (13) was subcloned into pET29a<sup>+</sup> with *Bam*HI and *Nde*I to allow overexpression of native nontagged PspA. PspA was overexpressed by using *E. coli* B834(DE3) and partially purified by extraction with 1.1% CHAPS {3-[(3-cholamidopropyl)-dimethylammonio]-1-propanesulfonate}, essentially as described by Elderkin et al. (13). PspA was excised from a sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) gel (15% polyacrylamide), eluted with 0.1 M sodium acetate and 0.1% (wt/vol) SDS, and then dried under a vacuum to lyophilize the protein. Naïve BALB/c mice were injected with 10  $\mu$ g of PspA protein mixed 1:1 (vol/vol) with Freund's adjuvant (Sigma). Mice received a 10- $\mu$ g booster injection of protein, as before, on days 20 and 40 after the primary injection.

TABLE 1. Survey of induction of *pspA* by chromosomal mutations in secretion loci of *E. coli*

Strain <sup>a</sup>	$\beta$ -Galactosidase assay result (Miller units) <sup>b</sup>	Reference
JS7131(pMR25) ara	6 $\pm$ 0	29
JS7131(pSJ1) ara	59 $\pm$ 0.4	29
JS7131(pMR25) glu	8 $\pm$ 0.29	29
JS7131(pSJ1) glu	277 $\pm$ 1.9	29
MC4100(pMR25)	15.4 $\pm$ 0.7	8
MC4100(pSJ1)	183 $\pm$ 3.5	8
BILKO (MC4100 $\Delta$ <i>tarC</i> )(pMR25)	17 $\pm$ 0.3	6
BILKO (MC4100 $\Delta$ <i>tarC</i> )(pSJ1)	180 $\pm$ 0.17	6
ELV15 (MC4100 $\Delta$ <i>tatA</i> )(pMR25)	33 $\pm$ 0.9	30
ELV15 (MC4100 $\Delta$ <i>tatA</i> )(pSJ1)	176 $\pm$ 2.6	30
J1M1 (MC4100 $\Delta$ <i>tatE</i> )(pMR25)	31 $\pm$ 0.8	30
J1M1 (MC4100 $\Delta$ <i>tatE</i> )(pSJ1)	160 $\pm$ 0.9	30
JARV15 (MC4100 $\Delta$ <i>tatA</i> $\Delta$ <i>tatE</i> )(pMR25)	34 $\pm$ 0.4	30
JARV15 (MC4100 $\Delta$ <i>tatA</i> $\Delta$ <i>tatE</i> )(pSJ1)	172 $\pm$ 1.7	30
HPT57 (MC1000 <i>phoA</i> <sup>+</sup> <i>phoR</i> <i>leu</i> <sup>+</sup> <i>102::malF-lacZ102</i> )(pMR25)	3.6 $\pm$ 0.7	33
HPT57 (MC1000 <i>phoA</i> <sup>+</sup> <i>phoR</i> <i>leu</i> <sup>+</sup> <i>102::malF-lacZ102</i> )(pSJ1)	39 $\pm$ 0.8	33
HPT-UV46 ( <i>ffs-46</i> ) (pMR25)	4.6 $\pm$ 0.8	33
HPT-UV46 ( <i>ffs-46</i> ) (pSJ1)	45.6 $\pm$ 2.9	33
HPT-UV70 ( <i>ftsY70</i> ) (pMR25)	5 $\pm$ 2.3	33
HPT-UV70 ( <i>ftsY70</i> ) (pSJ1)	26 $\pm$ 0.8	33
HPT-UV73 ( <i>secM73</i> ) (pMR25)	5.2 $\pm$ 0.2	33
HPT-UV73 ( <i>secM73</i> ) (pSJ1)	418 $\pm$ 0.8	33
HPT-UV77 ( <i>ffh-77</i> ) (pMR25)	2.2 $\pm$ 1.3	33
HPT-UV77 ( <i>ffh-77</i> ) (pSJ1)	34 $\pm$ 0.65	33
HPT-UV87 ( <i>ffh-87</i> ) (pMR25)	4.8 $\pm$ 0.3	33
HPT-UV87 ( <i>ffh-87</i> ) (pSJ1)	23.1 $\pm$ 0.84	33
HPT-UV89 ( <i>ffh-89</i> ) (pMR25)	2.9 $\pm$ 0.9	33
HPT-UV89 ( <i>ffh-89</i> ) (pSJ1)	49 $\pm$ 2.2	33
HPT-UV93 ( <i>secM93</i> ) (pMR25)	7.4 $\pm$ 2.3	33
HPT-UV93 ( <i>secM93</i> ) (pSJ1)	420 $\pm$ 9.9	33
HPT-UV103 ( <i>ffh-103</i> ) (pMR25)	4.9 $\pm$ 0.49	33
HPT-UV103 ( <i>ffh-103</i> ) (pSJ1)	37.7 $\pm$ 0.27	33
MC4100(pMR25) <sup>c</sup>	1.36 $\pm$ 0.79	4
MC4100(pMR25) <sup>d</sup>	2.57 $\pm$ 0.8	4
MC4100(pSJ1) <sup>c</sup>	37.3 $\pm$ 0.8	4
MC4100(pSJ1) <sup>d</sup>	38.4 $\pm$ 0.6	4
HDB45(pMR25) <sup>c</sup>	3.0 $\pm$ 0.58	4
HDB45(pMR25) <sup>d</sup>	3.3 $\pm$ 0.86	4
HDB45(pSJ1) <sup>c</sup>	37.1 $\pm$ 0.35	4
HDB45(pSJ1) <sup>d</sup>	36.4 $\pm$ 1.09	4

<sup>a</sup> Where present, the pMR25 vector numbered one to five copies per cell (C. Mohr and R. Roberts, unpublished data) (a gift from Dickon Alley). ara or glu, cultures were grown with LB plus 0.2% arabinose or glucose, respectively. pSJ1 is pMR25 with a *pspA-lacZ* transcriptional fusion. JS7131 strains were supplied by T. Palmer. The BILKO, ELV15, J1M1, and JARV15 strains, as well as the MC4100(pMR25) and MC4100(pSJ1) strains described in reference 8, were supplied by R. Dalbey. The HPT strains were supplied by J. Beckwith. The HBD strains and the MC4100(pMR25) and MC4100(pSJ1) strains described in reference 4 were supplied by Harris Bernstein.

<sup>b</sup> Assays were in triplicate using bacteria grown in LB supplemented with the appropriate antibiotic (tetracycline at 10  $\mu$ g/ml). Results are expressed as means  $\pm$  standard errors of the means.

<sup>c</sup> Grown with IPTG.

<sup>d</sup> Grown without IPTG.

Serum was taken from mice on day 60 and was judged by Western blotting exactly as described previously (13) with a 1:10,000 dilution of PspA and a 1:8,000 dilution of horseradish peroxidase-conjugated rabbit anti-mouse antibody (Sigma) to detect bound PspA antibody from samples of whole-cell extract of *E. coli* B834(DE3) harboring pET29a<sup>+</sup> (*pspA*). The wild type (HPT57), HPT73, and HPT93 were grown to satu-




FIG. 1. (A) JS3171(*paraBAD-yidC*) samples were grown to saturation in LB with 0.2% arabinose and then washed twice in 5 ml of LB to remove arabinose. Cells were diluted to an  $OD_{600}$  of 0.05 and grown in LB with 0.2% arabinose (lanes 1 to 3) or LB with 0.2% glucose (lanes 4 to 6). Samples were taken at 2 h (lanes 1 and 3), 4 h (lanes 2 and 4), or 6 h (lanes 3 and 6) postdilution. Western blot analysis used a mouse PspA antibody and horseradish peroxidase detection. PspA is clearly induced and is indicated. (B) Plasmid pMR25 or pSJ1 was introduced into HPT57 (wild type), HPT73 (*secM* point mutant), and HPT93 (*secM* point mutant). Following growth to saturation, cells were diluted to an  $OD_{600}$  of 0.05 and grown for 2 h (lanes 1, 4, and 7), 4 h (lanes 2, 5, and 8), or 6 h (lanes 3, 6, and 9). Samples were prepared for Western blotting using mouse PspA antibody and horseradish peroxidase detection. Lanes 1 to 3 contain HPT57 (wild type), lanes 4 to 6 contain HPT73 (*secM73*), and lanes 7 to 9 contain HPT93 (*secM93*).

ration and then diluted to an  $OD_{600}$  of 0.05 in LB and grown at 37°C. Samples were taken at 2, 4, and 6 h postdilution and were analyzed by SDS-PAGE. Equal amounts of cells were loaded in each lane, and no significant differences in growth rate were noticed. Western blotting of samples using the PspA antibody clearly revealed significant up-regulation of PspA protein (Fig. 1B) in the *secM* mutants.

**Depletion of YidC dramatically up-regulates *pspA* expression.** Depletion of YidC was used in these experiments, since *yidC* is essential. JS7131 contains a single copy of *yidC* integrated into the *attB* locus of MC1061 under the control of the *araBAD* promoter (29). JS7131 harboring pMR25 (vector control) or pSJ1 was grown to saturation in LB (plus 0.2% arabinose) and then washed twice with 5 ml of LB to remove arabinose. Cells were diluted to an  $OD_{600}$  of 0.05 in either LB plus 0.2% arabinose or LB plus 0.2% glucose and grown for 4 h. Triplicate samples were assayed for  $\beta$ -galactosidase activity as described above for Table 1. An approximately 4.7-fold up-regulation of *pspA* transcription was observed following YidC depletion. Following depletion of YidC, exactly as before, cell samples were analyzed by SDS-PAGE, and Western blotting clearly showed up-regulation of PspA protein at 2 and 4 h postdilution to remove arabinose (Fig. 1A).

**Mutations in *tat* loci do not affect *pspA* expression.** *E. coli* strain MC4100 harboring in-frame deletions in *tatA*, *tatE*, *tatAE*, and *tatC* were all assessed for *pspA* expression. Mutations in *tatC* and *tatAE* completely block translocation of *tat* substrates, while deletions in either *tatA* or *tatE* lead to partial blocking of *tat* substrate transport since their functions overlap (30; T. Palmer, personal communication). pMR25 (vector control) and pSJ1 were introduced into *tat* mutants and a wild-type reference strain (MC4100). All strains were grown to an  $OD_{600}$  of 0.6 to 0.8 or saturation. PspA was not up-regulated under any of the conditions tested (Table 1). Western blotting was employed to confirm these results; again no up-regulation of PspA protein was observed (data not shown). We did not test *psp* induction under anaerobic conditions, since *tat* genes are constitutively expressed, indicating a requirement for the Tat export apparatus under all growth conditions (15).

**Which Psp protein(s) could monitor stress at the inner membrane?** Recent evidence indicated that regulation of the *psp* operon proceeds through protein-protein interactions (1). PspA is a cytoplasmic protein associated with the inner mem-

brane (12). PspE is a periplasmic protein (22), while PspC is known to span the inner membrane once with an N-terminus-in, C-terminus-out topology (17). PspB is an inner membrane protein with one membrane-spanning domain, but the topological arrangement has not been proven. PspD was identified as a peripherally bound inner membrane protein by Western blotting and cell fractionation studies (1). To confirm the cellular location of PspB and PspD, we used a gene fusion approach, a well-characterized method of determining the topology of proteins (20). We fused *lacZ* or *phoA* to the C termini of PspD and PspB and used enzyme assays to determine the activities of the fusion proteins produced (Table 2). Since both PspB and PspD fused to *lacZ* gave reproducible  $\beta$ -galactosidase activity, whereas fusion of either to *phoA* gave no detectable alkaline phosphatase activity, the C termini of these proteins must be located in the cytoplasm. Western blotting with

TABLE 2. Roles of PspB and PspD in the phage shock response<sup>a</sup>

Strain	$\beta$ -Galactosidase assay result (Miller units)	Alkaline phosphatase assay result (U/h)
DH5 $\alpha$ (vector)	57.2 $\pm$ 0.05	6 $\pm$ 0.05
DH5 $\alpha$ (vector + <i>pspD</i> )	1968 $\pm$ 26	7 $\pm$ 0.02
DH5 $\alpha$ (vector + <i>pspB</i> )	686 $\pm$ 8	7 $\pm$ 0.02
MG1655(pSJ1, pGZ119EH, NI)	183.4 $\pm$ 6.4	
MG1655(pSJ1, pGZ119EH, I)	242.9 $\pm$ 6.4	
MG1655(pSJ1, pPMR129, NI)	196 $\pm$ 6.6	
MG1655(pSJ1, pPMR129, I)	657.2 $\pm$ 9.3	
MG1655 $\Delta$ <i>pspD</i> (pSJ1, pGZ119EH, NI)	191.0 $\pm$ 3.3	
MG1655 $\Delta$ <i>pspD</i> (pSJ1, pGZ119EH, I)	197.24 $\pm$ 2.2	
MG1655 $\Delta$ <i>pspD</i> (pSJ1, pPMR129, NI)	199.6 $\pm$ 4.0	
MG1655 $\Delta$ <i>pspD</i> (pSJ1, pPMR129, I)	700.6 $\pm$ 18.6	

<sup>a</sup> *E. coli* strain DH5 $\alpha$  was the host for gene fusion constructs. This strain lacks endogenous  $\beta$ -galactosidase activity and has minimal alkaline phosphatase activity. The vector was pLKC480 for fusions made to *lacZ* (34), and *pspB* and *pspD* were amplified with primers and clones containing appropriate restriction sites (verified by sequencing; primers are available on request). The vector was pBAF for fusions made to *phoA* (13a), and cloning sites and primers were as described above for fusions to *lacZ*. Assays were in triplicate using bacteria grown in LB supplemented with the appropriate antibiotic (ampicillin at 100  $\mu$ g/ml), and results are means  $\pm$  standard errors. The pGZ119EH vector is described in the work of Lessel et al. (18). Strains were grown with no induction (NI) in LB supplemented with the appropriate antibiotics. For strains grown with induction (I), IPTG at a concentration of 1 mM was used throughout growth to induce the expression of pIV (encoding the secretin) from pPMR129. pPMR129 is vector pGZ119EH containing the pIV gene from phage f1 under *lac* promoter control (10).

an antibody to LacZ (Cp Laboratories) demonstrated the production of stable fusion proteins of the expected size (data not shown). Hence, alterations in PMF (or improperly inserted membrane proteins) might be sensed by PspA, PspB, PspC, or PspD. A previous study showed that both *pspB* and *pspC* are required for induction of the *psp* response but that *pspE* is dispensable for induction (24). Eviction-exchange mutagenesis was used to delete *pspD* from the chromosome of *E. coli* MG1655. A PCR product containing an in-frame deletion of *pspD* was constructed as described by Link et al. (19) with primers listed on the website <http://arep.med.harvard.edu/labgc/adnan/projects/EcoliKOprimers/EcoliKOprimers.html> in which the *SaI* restriction site is replaced with a *Bam*HI site. The PCR product was digested with *Bam*HI and *Not*I and ligated into the same sites in plasmid pSR47s (modified *sacB* vector supplied by Paul Casaz). Eviction-exchange mutagenesis was carried out as described by Link et al. (19) by using *E. coli* MG1655 CGSC 7740 (5) as the parent strain for the gene deletion. *pspD* deletion mutants were detected by colony PCR with the same primers used to make the original PCR product. Amplified DNA from putative mutants was sequenced to verify that the deletion was in frame. Induction of *pspA* with pPMR129 (which harbors pIV under *lac* promoter control) was not affected in a strain with *pspD* deleted (Table 2). Hence, although it remains a possibility that PspD may influence the kinetics of induction by pIV or other inducers, it seems likely that either PspA, PspB, or PspC (or a combination) is the primary sensor of membrane stresses.

**Implications.** Transport of proteins across the bacterial inner membrane may impose stress on the cell, perhaps through modulation of the PMF. Using a simple reporter system for *pspA* expression and a specific antibody to detect PspA levels, we have shown that while stress(es) imposed by depletion of YidC and impaired functioning of SecM leads to an increased synthesis of PspA (and, in concert, *pspBCDE*, since these genes are arranged as an operon), defects in the bacterial SRP and Tat systems have no effect on *psp*. Since the *tat* system can open a large pore while maintaining the PMF intact (31) and mutations in the SRP affect only the presentation of substrates to the Sec translocon, this study confirms that the elimination of PMF is a good candidate for *psp* up-regulation. PMF is, however, required for translocation through the Tat pore; perhaps the mutations in the Tat apparatus analyzed in this study did not lead to the dissipation of PMF. Alternatively, perhaps even though export in a *tatA* background is severely affected (T. Palmer, personal communication), the number of exported proteins affected is significantly lower than when the *sec* machinery is corrupted. The Psp system may sense problems specifically at the Sec translocon. The effects of *secA*, *secD*, *secF*, and *secM* upon the induction of the *psp* operon are clear, yet *secY* and *secE* mutations do not up-regulate *psp*. The basis for this differential effect is not yet clear, but it might be related to some redundancy in the roles that *secY* and *secE* can play in the cell. It has been suggested that YidC acts with the Sec translocon, and we hypothesize that one or more of the Psp proteins may sense alterations in the PMF directly, or perhaps they interact with members of the Sec translocon or YidC itself. SecM has been suggested to participate directly in the insertion of membrane proteins into the inner membrane. The *secM* mutants used in this study affected membrane protein insertion

into the inner membrane but had no effect on SecA levels (33). YidC can also independently insert membrane proteins into the inner membrane. So the possibility also exists that a Psp protein(s) can sense incorrectly inserted proteins directly, perhaps by close association with SecM, SecD, or SecF and YidC. The differences in *psp* expression between *secM* and *srp* mutants is striking. An involvement of SecM in the regulation of SecA may be one basis for the clear effects of SecM upon *psp* expression. Although *secM* and *srp* mutations have the property of having weak effects upon membrane protein insertion, the *secM* mutant has the additive effect of interfering with protein translocation. By utilizing chromosomal mutations in several components of the secretion apparatus, as opposed to determining the effects of overexpressed proteins on the *psp* system, we have established a firm link between secretion-imposed stress and the *psp* system of *E. coli*. Recent reports that regulation of the phage shock response proceeds through protein-protein interactions speculated that PspC was the most likely candidate for the prime sensor of membrane integrity (1). PspD is dispensable for induction of *psp* (this study), leaving PspA, PspB, or PspC as likely sensors of membrane stress. The challenge now is to identify how the Psp protein(s) senses stress and to work out how PspA can act to restore PMF and hence aid the secretion of proteins across the bacterial inner membrane.

We thank Jon Beckwith, Harris Bernstein, Tracy Palmer, Ross Dalbey, and Marjorie Russel, who kindly provided strains and plasmids used in this work. We are grateful to Paul Casaz for the gift of plasmid pSR47 and tips for the gene knockout methodology.

This work was supported by funding from the Wellcome Trust to M.B. L.J.L. is the recipient of a Wellcome Trust studentship.

#### REFERENCES

- Adams, H., W. Teertstra, J. Demmers, R. Boesten, and J. Tommassen. 2003. Interactions between phage-shock proteins in *Escherichia coli*. *J. Bacteriol.* **185**:1174–1180.
- Arkowitz, R. A., and W. Wickner. 1994. SecD and SecF are required for the proton electrochemical gradient stimulation of preprotein translocation. *EMBO J.* **13**:954–963.
- Bernstein, H. D., M. A. Poritz, K. Strub, P. J. Hoben, S. Brenner, and P. Walter. 1989. Model for signal sequence recognition from amino-acid sequence of 54K subunit of signal recognition particle. *Nature* **340**:482–486.
- Bernstein, H. D., and J. B. Hyndman. 2001. Physiological basis for conservation of the signal recognition particle targeting pathway in *Escherichia coli*. *J. Bacteriol.* **183**:2187–2197.
- Blattner, F. R., G. Plunkett III, C. A. Bloch, N. T. Perna, V. Burland, M. Riley, J. Collado-Vides, J. D. Glasner, C. K. Rode, G. F. Mayhew, J. Gregor, N. W. Davis, H. A. Kirkpatrick, M. A. Goeden, D. J. Rose, B. Mau, and Y. Shao. 1997. The complete genome sequence of *Escherichia coli* K-12. *Science* **277**:1453–1474.
- Bogsch, E. G., F. Sargent, N. R. Stanley, B. C. Berks, C. Robinson, and T. Palmer. 1998. An essential component of a novel bacterial protein export system with homologues in plastids and mitochondria. *J. Biol. Chem.* **273**:18003–18006.
- Carlson, J. H., and T. J. Silhavy. 1993. Signal sequence processing is required for the assembly of LamB trimers in the outer membrane of *Escherichia coli*. *J. Bacteriol.* **175**:3327–3334.
- Casadaban, M. J., and S. N. Cohen. 1979. Lactose genes fused to exogenous promoters in one step using a Mu-lac bacteriophage: in vivo probe for transcriptional control sequences. *Proc. Natl. Acad. Sci. USA* **76**:4530–4533.
- Chen, M., K. Xie, F. Jiang, L. Yi, and R. E. Dalbey. 2002. YidC, a newly defined evolutionarily conserved protein, mediates membrane protein assembly in bacteria. *Biol. Chem.* **383**:1565–1572.
- Daefler, S., M. Russel, and P. Model. 1997. Module swaps between related translocator proteins pIV(f1), pIV(IKe) and PulD: identification of a specificity domain. *J. Mol. Biol.* **266**:978–992.
- Duong, F., and W. Wickner. 1997. The SecDFyajC domain of preprotein translocase controls preprotein movement by regulating SecA membrane cycling. *EMBO J.* **16**:4871–4879.
- Dworkin, J., G. Jovanovic, and P. Model. 2000. The PspA protein of *Esch-*

- erichia coli* is a negative regulator of  $\sigma^{54}$ -dependent transcription. J. Bacteriol. **182**:311–319.
13. **Elderkin, S., S. Jones, J. Schumacher, D. Studholme, and M. Buck.** 2002. Mechanism of action of the *Escherichia coli* phage shock protein PspA in repression of the AAA family transcription factor PspF. J. Mol. Biol. **320**: 23–37.
  - 13a. **Fulkerson J. F., and Mobley H. C.** 2000. Membrane topology of the NixA nickel transporter of *Helicobacter pylori*: two nickel transport-specific motifs within transmembrane helices II and III. J. Bacteriol. **182**:1722–1730.
  14. **Hardie, K. R., S. Lory, and A. P. Pugsley.** 1996. Insertion of an outer membrane protein in *Escherichia coli* requires a chaperone-like protein. EMBO J. **15**:978–988.
  15. **Jack, R. L., F. Sargent, B. C. Berks, G. Sawers, and T. Palmer.** 2001. Constitutive expression of *Escherichia coli* *tat* genes indicates an important role for the twin-arginine translocase during aerobic and anaerobic growth. J. Bacteriol. **183**:1801–1804.
  16. **Kleerebezem, M., and J. Tommassen.** 1993. Expression of the *pspA* gene stimulates efficient protein export in *Escherichia coli*. Mol. Microbiol. **7**:947–956.
  17. **Kleerebezem, M., W. Crielaard, and J. Tommassen.** 1996. Involvement of stress protein PspA (phage shock protein A) of *Escherichia coli* in maintenance of the protonmotive force under stress conditions. EMBO J. **15**:162–171.
  18. **Lessl, M., D. Balzer, R. Lurz, V. L. Waters, D. G. Guiney, and E. Lanka.** 1992. Dissection of IncP conjugative plasmid transfer: definition of the transfer region Tra2 by mobilization of the Tra1 region in *trans*. J. Bacteriol. **174**:2493–2500.
  19. **Link, A. J., D. Phillips, and G. M. Church.** 1997. Methods for generating precise deletions and insertions in the genome of wild-type *Escherichia coli*: application to open reading frame characterization. J. Bacteriol. **179**:6228–6237.
  20. **Manoil, C.** 1991. Analysis of membrane protein topology using alkaline phosphatase and beta-galactosidase gene fusions. Methods Cell Biol. **34**:61–75.
  21. **Matsuyama, S., Y. Fujita, and S. Mizushima.** 1993. SecD is involved in the release of translocated secretory proteins from the cytoplasmic membrane of *Escherichia coli*. EMBO J. **12**:265–270.
  22. **Mielke, D. L., and M. Russel.** 1992. A modified *TnphoA* useful for single-strand sequencing. Gene **118**:93–95.
  23. **Miller, J. H.** 1972. Experiments in molecular genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
  24. **Model, P., G. Jovanovic, and J. Dworkin.** 1997. The *Escherichia coli* phage-shock-protein (*psp*) operon. Mol. Microbiol. **24**:255–261.
  25. **Poritz, M. A., H. D. Bernstein, K. Strub, D. Zopf, H. Wilhelm, and P. Walter.** 1990. An *E. coli* ribonucleoprotein containing 4.5S RNA resembles mammalian signal recognition particle. Science **250**:1111–1117.
  26. **Robinson, C., and A. Bolhuis.** 2001. Protein targeting by the twin-arginine translocation pathway. Nat. Rev. Mol. Cell Biol. **2**:350–356.
  27. **Romisch, K., J. Webb, J. Herz, S. Prehn, R. Frank, M. Vingron, and B. Dobberstein.** 1989. Homology of 54K protein of signal-recognition particle, docking protein and two *E. coli* proteins with putative GTP-binding domains. Nature **340**:478–482.
  28. **Sambrook, J., E. F. Fritsch, and T. Maniatis.** 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
  29. **Samuelson, J. C., M. Chen, F. Jiang, I. Moller, M. Wiedmann, A. Kuhn, G. J. Phillips, and R. E. Dalbey.** 2000. YidC mediates membrane protein insertion in bacteria. Nature **406**:637–641.
  30. **Sargent, F., E. G. Bogsch, N. R. Stanley, M. Wexler, C. Robinson, B. C. Berks, and T. Palmer.** 1998. Overlapping functions of components of a bacterial Sec-independent protein export pathway. EMBO J. **17**:3640–3650.
  31. **Sargent, F., B. C. Berks, and T. Palmer.** 2002. Assembly of membrane-bound respiratory complexes by the Tat protein-transport system. Arch. Microbiol. **178**:77–84.
  32. **Struck, J. C., H. Y. Toschka, and V. A. Erdmann.** 1988. Nucleotide sequence of the 4.5S RNA gene from *Thermus thermophilus* HB8. Nucleic Acids Res. **16**:9042.
  33. **Tian, H., and J. Beckwith.** 2002. Genetic screen yields mutations in genes encoding all known components of the *Escherichia coli* signal recognition particle pathway. J. Bacteriol. **184**:111–118.
  34. **Tiedeman, A. A., and J. M. Smith.** 1988. Lac24 gene fusion cassettes with kanR resistance. Nucleic Acids Res. **16**:3587.
  35. **L. Yi, F. Jiang, M. Chen, B. Cain, A. Bolhuis, and R. E. Dalbey.** 2003. YidC is strictly required for membrane insertion of subunits a and c of the F-1f-OATP synthase and SecE of the SecUEG translocase. Biochemistry **42**: 10537–10544.