

Short report

Open Access

Complete genomic sequence analysis of infectious bronchitis virus Ark DPI strain and its evolution by recombination

Arun Ammayappan^{1,2}, Chitra Upadhyay¹, Jack Gelb Jr³ and Vikram N Vakharia*¹

Address: ¹Center of Marine Biotechnology, University of Maryland Biotechnology Institute, Baltimore, 701 East Pratt Street, Baltimore, Maryland 21202-3101, USA, ²Department of Veterinary Medicine, University of Maryland, College Park, MD 20742, USA and ³Avian Biosciences Center, University of Delaware, 531 South College Avenue, Newark, DE 19716-2150, USA

Email: Arun Ammayappan - ammayapp@umbi.umd.edu; Chitra Upadhyay - upadhyay@umbi.umd.edu; Jack Gelb - jgelb@udel.edu; Vikram N Vakharia* - vakharia@umbi.umd.edu

* Corresponding author

Published: 22 December 2008

Received: 22 September 2008

Virology Journal 2008, 5:157 doi:10.1186/1743-422X-5-157

Accepted: 22 December 2008

This article is available from: <http://www.virologyj.com/content/5/1/157>

© 2008 Ammayappan et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

An infectious bronchitis virus Arkansas DPI (Ark DPI) virulent strain was sequenced, analyzed and compared with many different IBV strains and coronaviruses. The genome of Ark DPI consists of 27,620 nucleotides, excluding poly (A) tail, and comprises ten open reading frames. Comparative sequence analysis of Ark DPI with other IBV strains shows striking similarity to the Conn, Gray, JMK, and Ark 99, which were circulating during that time period. Furthermore, comparison of the Ark genome with other coronaviruses demonstrates a close relationship to turkey coronavirus. Among non-structural genes, the 5'untranslated region (UTR), 3C-like proteinase (3CL^{pro}) and the polymerase (RdRp) sequences are 100% identical to the Gray strain. Among structural genes, S1 has 97% identity with Ark 99; S2 has 100% identity with JMK and 96% to Conn; 3b 99%, and 3C to N is 100% identical to Conn strain. Possible recombination sites were found at the intergenic region of spike gene, 3'end of S1 and 3a gene. Independent recombination events may have occurred in the entire genome of Ark DPI, involving four different IBV strains, suggesting that genomic RNA recombination may occur in any part of the genome at number of sites. Hence, we speculate that the Ark DPI strain originated from the Conn strain, but diverged and evolved independently by point mutations and recombination between field strains.

Findings

Avian infectious bronchitis virus (IBV) is a pathogen of domestic chickens that causes acute, highly contagious respiratory disease [1]. IBV is a member of the *Coronaviridae*, order *Nidovirales* [2] and its genome consists of a 27.6 kb single stranded positive-sense RNA molecule that encodes for four structural proteins; the spike (S) glycoprotein, the small envelope (E) protein, the membrane (M) glycoprotein, and the nucleocapsid (N) protein [3,4].

Six subgenomic mRNAs are transcribed from the IBV genome in virus-infected cells. The mRNA 1 contains two large overlapping open reading frames, encoding two polyproteins 1a and 1b [5], among which 1b is produced as 1ab polyprotein by ribosomal frame-shifting mechanism [6].

Many serotypes have been described for IBV, probably due to the frequent point mutations that occur in RNA

viruses and also due to recombination events demonstrated for IBV [7-9]. For this reason, the characterization of virus isolates existing in the field is very important. The Ark DPI strain was first isolated from Delmarva Peninsula broiler flock [10,11] and it is currently being used as a vaccine in the USA and Europe. In this study, we characterized the entire genome of virulent Ark DPI strain (embryo passage 11) and compared it with other IBV strains and coronaviruses from all over the world.

The Ark DPI virus was inoculated into 9-day-old SPF chicken eggs and allantoic fluid was collected 72 h post inoculation. The fluid was clarified by low speed centrifugation and clear supernatant was stored at -80°C. Genomic RNA was extracted from virus-infected allantoic fluid with Qiagen RNeasy kit, following the manufacturer's instructions, and stored at -80°C until further use. Oligonucleotides were designed based on consensus sequence of the following IBV strains: Cal 99 [GenBank:AY514485], Mass 41 [GenBank:AY851295] and BJ [GenBank:AY319651]. Overlapping primers were designed in a manner such that each pair of primer covered approximately two kb of genome. The RT-PCR was carried out as described earlier [12] and the RT-PCR products were cloned into pCR2.1 TOPO TA vector (Invitrogen, CA). Plasmid DNA from various clones was sequenced by dideoxy chain termination method, using an automated DNA sequencer (Applied Biosystems, CA). Three independent clones were sequenced for each amplicon to exclude errors that can occur from RT and PCR reactions. The assembly of contiguous sequences and multiple sequence alignments were performed with the GeneDoc software [13]. The pair-wise nucleotide identity and comparative sequence analyses were conducted using Vector NTI Advance 10 software (Invitrogen, CA) and BLAST search, NCBI. Phylogenetic analyses were conducted using the MEGA4 program [14].

The GenBank accession number for the Ark DPI sequence is EU418976. The complete genomes of following strains are obtained from GenBank: TCoV10, [NC_010800](#); Beaudette, [NC_001451](#); M41, [AY851295](#); CK/CH/LSD/05I, [EU637854](#); A2, [EU526388](#); LX4, [AY338732](#); SAIBK, [DQ288927](#); The accession numbers of IBV gene sequences which are used in this study are as follows: For replicase gene sequences: (a) 5'UTR: Conn, [AY392049](#); Florida, [AY392050](#); CU-T2, [AY561724](#); Ark 99, [AY392051](#); DE072, [AY392054](#); GA98, [AY392053](#); Gray, [AY392056](#); (b) PL^{pro}: Conn, [AY392059](#); Florida, [AY392060](#); CU-T2, [AY561734](#); Ark 99, [AY392061](#); DE072, [AY392064](#); GA98, [AY392063](#); Gray, [AY392066](#); (c) M^{pro}: Conn, [AY392069](#); Florida, [AY392070](#); CU-T2, [AY561744](#); Ark 99, [AY392071](#); DE072, [AY392074](#); GA98, [AY392073](#); Gray, [AY392076](#); (d) RdRp: Conn, [AY392079](#); Florida, [AY392080](#); CU-T2, [AY561754](#); Ark99, [AY392081](#);

DE072, [AY392084](#); GA98, [AY392083](#); Gray, [AY392086](#). For Structural genes (a) Complete structural genes: HK, [AY761141](#); Vic, [DQ490221](#); KB8523, [M21515](#); TW2296/95, [DQ646404](#) (b) S1: Jilin, [AY839144](#); Gray, [L18989](#); Conn, [EU526403](#); Holte, [L18988](#); UK/2/91, [Z83976](#); Qu16, [AF349620](#); JMK, [L14070](#); H120, [M21970](#); GAV-92, [AF094817](#); DE072, [AF274435](#); IS/1366, [EU350550](#); (c) S2: JMK, [AF239982](#); Jilin, [AY839146](#); Holte, [AF334685](#); DE072, [AY024337](#); Conn, [AF094818](#); Gray, [AF394180](#); H120, [AF239982](#); (d) S: Ark 99, [L10384](#); CU-T2, [U04739](#); (e) Gene 3: Jilin, [AY846833](#); Conn, [AY942752](#); CU-T2, [U46036](#); Ark 99, [AY942751](#); Gray, [AF318282](#) (f) M: Jilin, [AY846833](#); JMK, [AF363608](#); Conn, [AY942741](#); H120, [AY028295](#); Gray, [AF363607](#); (g) Gene 5: Jilin, [AY839142](#); Gray, [AF469011](#); Conn, [AF469013](#); DE072, [AF203000](#); (h) N: Jilin, [AY839145](#); Ind/TN/92/03, [EF185916](#); Conn, [AY942746](#); H120, [AY028296](#); Gray, [M85245](#); (i) 3'UTR: H120, [AI278336](#).

The genome of Ark DPI consists of 27,620 nucleotides (nts), excluding poly (A) tail, and comprises ten open reading frames (ORFs) flanked by 5' (528 nts) and 3' (507 nts) untranslated regions (UTRs). The genome organization is ORF1ab (529–20360), ORF2 (20311–23820), ORF3abc [3a, (23820–23993), 3b (23993–24187), 3c (24168–24491)], ORF4 (24469–25140), ORF5ab [5a (25500–25697), 5b (25694–25942)] and ORF6 (25885–27114).

The details of genome organization of Ark DPI are shown in Fig. 1. IBV polyprotein is cleaved into 15 cleavage products, among which first two N-terminal products are cleaved by PL^{pro} and rest of the C-terminal products are cleaved by 3CL^{pro} [15]. The putative domains and their cleavage sites (Fig. 1) are predicted by comparison of amino acid sequences of each non-structural protein (nsp) of Ark DPI with those of IBV-Beaudette which is available in Coronavirus Database (CoVDB) [16]. The nucleotide and the amino acid identity of Ark DPI with other IBVs and coronaviruses are listed in Tables 1, 2, 3. The whole structural gene of Jilin is 100% identical to Ark DPI, which suggests that Jilin strain is actually Ark DPI, which is currently used as a vaccine in China [17]. The whole genome comparison of IBV strains reveals a close relationship of Ark DPI with Cal 99 (96% identity), as shown in Fig. 2. Earlier studies have shown that Cal 99 probably evolved from Ark DPI [18].

The complete genome sequence analysis of Ark DPI strain shows striking similarity to the Conn, Gray, JMK, and Ark 99 IBV strains, which were circulating during that time period [1,19-21]. The 5'UTR, PL^{pro}, M^{pro} and RdRp sequence analysis demonstrates that Ark DPI is 100% identical to Gray strain, except for PL^{pro} which has 87% identity, as shown in Table 1. It was suggested that PL^{pro}




Figure 1
Classical Genome Organization of IBV-Ark DPI. The genome of Ark DPI is 27,620 nt long, excluding poly (A) tract. Middle: ten genes and its ORFs. Ribosomal frameshift and position of transcriptional regulatory sequences (TRS) of each gene is indicated. Top: putative domains of ORF1a/1b polyprotein: nsp-non-structural protein, Ac-acidic domain, X-unknown domain X, PL1- papain like proteinase I, PL2-papain like proteinase 2; Y-unknown domain Y; HD-hydrophobic domain, 3CL-3C-like proteinase, G-Growth factor like protein, RdRp-RNA dependent RNA polymerase, Hel-helicase, ExoN-exoribonuclease, Ne-nidoviral uridylyate-specific endoribonuclease, MT- 2'-O-ribose methyltransferase. Bottom: details of spike protein; SP-signal peptide, RRSRR/S- spike protein cleavage site between 544 and 545aa, TM-transmembrane domain of spike protein.

gene has high genetic variation because of selection pressure [22]. From this analysis, it appears that genetic mutation may have occurred at PL^{pro} gene level. The modern strain GA98 maintains 100% identity with Ark DPI in replicase proteins and because of unavailability of sequence information for rest of the genome; we speculate that GA98 may be a derivative of the Ark DPI strain.

Analysis of the structural region of Ark DPI clearly demonstrates that it is a chimera of three strains. The S1 gene of Ark DPI is probably derived from Ark 99 (97% identical) and because of genetic mutations in the S1 region, Ark DPI may have evolved independently. There is an A-T rich sequence TGTGTTGATTATAAT (Fig. 3) at the 3' terminus of S1 gene (~300 nts upstream from the end of S1 gene) which is conserved among most of the IBV strains. The S1 gene of Ark 99 maintains its identity with Ark DPI up to this conserved region, but from this point onwards to the end of S2, the nucleotide sequence is 100% identical to JMK strain. The recombination between JMK and Ark 99 had taken place presumably between above mentioned conserved region and intergenic (IG) region of S gene,

which is located 49 nts upstream of start codon of S gene. It is speculated that IG sequences serve as "hot spots" for recombination because of its consensus nature [23]. Gray and JMK strains share 99% homology both in the S1 and S2 genes of Ark DPI, but JMK shows greater identity than Gray strain, as shown in Table 2. It is interesting that very few residues in the S1 gene make the Gray strain nephrotropic, whereas JMK is pneumotropic [24].

Out of 174 nts of gene 3a of Ark DPI, last 74 nts are 100% identical to Conn, whereas first 100 nts are only 86% identical. Even though it is not clear whether the 5'-end of 3a was derived from Conn or JMK, but it is evident that the recombination event may have occurred between JMK and Conn at gene 3a. The 3b gene of Ark DPI and Conn differed only by two nucleotides and both share 99% identity, suggesting that 3b belongs to Conn strain. From gene 3c to N gene, Ark DPI shares 100% identity with Conn. It is obvious that the entire structural genome, except spike, belongs to Conn strain. Cross protection studies carried out by Gelb and coworkers [11] demonstrated that the birds immunized with Ark DPI showed

Table 1: Percent (%) nucleotide identity of Ark DPI non-structural genes and ORF1ab, ORF2-6 and complete genome with other IBV strains^{a, b, c}

IBV Strains	5'UTR	PL ^{pro}	M ^{pro}	RdRp	ORF1	ORF2-6	Complete genome
Ark99	98	87	99	99	NA	96	NA
A2	95	83	85	99	86	85	86
Beaudette	96	85	93	93	91	91	91
BJ	96	83	88	93	86	85	86
Cal99*	100	99	97	99	96	94	96
CK/CH/LSD/051	97	84	88	90	89	90	89
Connecticut	97	87	100	99	NA	96	NA
CU-T2*	100	94	99	99	NA	94	NA
DE072	100	100	90	90	NA	NA	NA
Florida	97	87	100	99	NA	NA	NA
GA98*	100	100	100	100	NA	NA	NA
Gray	100	87	100	100	NA	NA	NA
Jilin	NA	NA	NA	NA	NA	100	NA
LX4	94	83	85	89	87	84	86
M41	97	87	91	94	91	91	91
SAIBK	92	85	90	90	90	86	89

^a Sequences with > 95% identity are in bold letters

^b NA-not analyzed

^c Parental strains of Ark DPI are shown in bold letters and immediate derivative of Ark DPI is indicated by asterisk (*).

95%, 90% and 63% protection against Conn, Ark 99 and JMK strains, respectively. Indeed, these results suggest that major part of Ark DPI genome was derived from Conn. The level of protection for JMK is 80%, when Ark 99 was used as immunogen [25]. On the other hand, Conn and JMK immunization induces inadequate immunity against Ark-type IBV challenge, suggesting that Ark cross-immunity to JMK and Conn is a one-way relationship [10,11,26].

Recombination hot spots have been demonstrated for IBV isolates by many researchers. These hot spots have been detected in the IG region [23], S1 gene [27], 3' terminus of S2, N and between N gene and 3'UTR [8,28]. Some earlier

sequencing studies had provided circumstantial evidence of recombination events in field isolates of IBV [7,29,30]. More or less recombination sites were detected over the entire genome of coronavirus [31]. Based on these results, we speculate that the Ark DPI strain originated from the Conn strain, but diverged and evolved independently by point mutations and recombination between field strains. These findings suggest that there is high level of genetic diversity among currently circulating IBV serotypes. Most of them come from genetic changes which already exist in the IBV field strains and from IBV live vaccines. So frequent monitoring is highly essential to track the emergence of new variants and is mandatory to develop efficient vaccination strategies to control and prevent IB.

Table 2: Percent (%) nucleotide identity of Ark DPI structural genes with other IBV strains ^{a, b, c, d}

BV Strains	S1	S2	3a	3b	3c	M	5a	5b	N	3'UTR
Ark99	97(96)	96(95)	92	99	96(92)	97(97)	93	97	98(98)	97
Beaudette	81(79)	95(94)	91	84	88(83)	91(93)	85	93	93(95)	97
BJ	77(75)	85(89)	88	76	87(79)	90(93)	NA	NA	89(93)	87
Cal99*	87(84)	94(93)	92	99	94(90)	97(96)	100	98	95(98)	89
CK/CH/LSD/051	78(75)	88(91)	89	84	(87)	96(96)	99	100	100(99)	90
Conn	81(77)	96(96)	92	99	100(100)	100(100)	100	100	100(100)	NA
CU-T2*	96(94)	93(93)	88	98	92(87)	88(87)	97	99	95(96)	97
DE072	62(50)	75(76)	NA	NA	NA	NA	98	98	NA	98
Gray	83(80)	99(99)	NA	NA	96(92)	98(98)	97	99	97(97)	98
GAV-92	94(92)	NA	NA	NA	NA	NA	NA	NA	NA	NA
H120	81(78)	NA	NA	NA	NA	92(94)	NA	NA	93(96)	83
HK*	81(78)	96(96)	99	100	100(100)	100(100)	100	100	100(100)	NA
Holte	83(80)	95(95)	NA	NA	NA	NA	NA	NA	NA	NA
Ind/TN/92/03	NA	NA	NA	NA	NA	NA	NA	NA	92(94)	NA
IS/1366	78(75)	NA	NA	NA	NA	NA	NA	NA	92(95)	NA
Jilin*	100(99)	100(100)	100	100	100(100)	100(100)	100	100	100(100)	100
JMK	84(82)	100(100)	NA	NA	NA	97(98)	NA	NA	NA	NA
KB8523	81(78)	91(92)	NA	NA	NA	93(95)	NA	NA	95(96)	NA
LX4	77(76)	85(88)	86	76	88(80)	91(91)	82	90	89(93)	NA
M41	81(78)	95(94)	91	85	88(83)	91(95)	90	97	94(95)	97
Qu16	84(81)	NA	NA	NA	NA	NA	NA	NA	NA	NA
SAIBK	79(77)	87(91)	86	83	85(80)	89(93)	84	96	87(92)	88
TW2296/95	79(77)	86(90)	86	85	(83)	91(92)	82	96	89(92)	NA
UK/2/91	78(76)	NA	NA	NA	NA	NA	NA	NA	NA	NA
Vic	81(79)	89(92)	88	88	88(88)	89(95)	87	94	90(94)	NA

^a Sequences with > 95% identity are indicated in bold letters

^b Amino acid sequences within the parenthesis

^c NA-Not Analyzed

^d Parental strains of Ark DPI are shown in bold letters and immediate derivative of Ark DPI is indicated by asterisk (*).

Table 3: Percent (%) amino acid identity of Ark DPI replicase and structural proteins to other coronaviruses c, d

Coronaviruses ^a	3CL ^{pro}	RdRp	S	E	M	N	Complete genome ^b
BatCoV	40	70	22	11	29	25	46
BCoV	41	67	21	13	30	24	47
ECoV	41	67	22	13	30	25	47
FCoV	45	62	22	16	23	23	48
HCoV 229E	40	63	23	12	26	23	49
TGEV	46	61	22	20	25	26	49
MHV A59	40	69	21	14	31	26	46
SARS CoV	46	68	21	17	29	22	45
SWI	56	79	25	28	36	35	50
TCoV	96	98	34	91	97	95	88

^a BatCoV, Bat coronavirus; FCoV, feline coronavirus; HCoV, human coronavirus; BCoV, bovine coronavirus; MHV, mouse hepatitis virus; SARS-CoV, severe acute respiratory syndrome coronavirus; SWI, beluga whale coronavirus; TCoV, turkey coronavirus.

^b Percent nucleotide identity of entire genome

^c Sequences with > 50% identity are in bold letters

^d Gene in bold letters (RdRp) is highly conserved; TCoV exhibits significant identity with IBV-Ark DPI (marked in bold letters).




Figure 2
Phylogenetic tree analysis of complete Ark DPI genome sequence with other IBV strains. Phylogenetic tree analysis was conducted by neighbor-joining method using bootstrap analysis (100 replications). The scale at the bottom indicates the number of substitution events.




Figure 3
Schematic presentation of the structural region of Ark DPI genome. Entire genome of Ark DPI was analyzed for its similarity with other IBV strains. Top panel: 5'UTR & ORF1. Shadowed regions were used for comparative analysis. 5'UTR-5'-untranslated region; PLI-papain like proteinase I; M^{pro}-main or 3C-like proteinase; RdRp-RNA-dependent-RNA polymerase. Bottom panel: ORF2 to 3'UTR. Structural genes and their ORFs are marked by (●). Conserved sequence TGTGTTGATTATAAT in S1 gene is shown; ◆ denotes plausible recombination site in Ark DPI.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

VNV and JG conceived the study. AA planned the experimental design, AA and CU carried out cloning and sequencing. AA drafted the manuscript. All authors critically reviewed and approved the final manuscript.

Acknowledgements

The project was supported by the National Research Initiative of the USDA Cooperative State Research, Education and Extension Service; grant number 2004-35204-14814 to V.N.V. and J.G.

References

- Chomiak TW, Luginbuhl RE, Jungherr EL: **The propagation and cytopathogenic effect of an egg-adapted strain of infectious bronchitis virus in tissue culture.** *Avian Dis* 1958, **2**:456-465.
- Cavanagh D, Naqi S: **Infectious Bronchitis.** In *Diseases of Poultry* 11th edition. Edited by: Saif YM, Barnes HJ, Glisson JR, Fadly AM, McDougald LR, Swayne DE. Ames: Iowa State University Press; 2003:101-120.
- Spaan W, Cavanagh D, Horzinek MC: **Coronaviruses: structure and genome expression.** *J Gen Virol* 1988, **69**:2939-2952.
- Sutou S, Sato S, Okabe T, Nakai M, Sasaki N: **Cloning and sequencing of genes encoding structural proteins of avian infectious bronchitis virus.** *Virology* 1988, **165**:589-595.
- Bournnell MEG, Brown TDK, Foulds IJ, Green PF, Tomely FM, Binns MM: **Completion of the sequence of the genome of the coronavirus avian infectious bronchitis virus.** *J Gen Virol* 1987, **68**:57-77.
- Brierley I, Digard P, Inglis SC: **Characterization of an efficient coronavirus ribosomal frameshifting signal: Requirement for an RNA pseudoknot.** *Cell* 1989, **57**:537-547.
- Cavanagh D, Davis PJ, Cook J: **Infectious bronchitis virus: evidence for recombination within the Massachusetts serotype.** *Avian Path* 1992, **21**:401-408.
- Kottier SA, Cavanagh D, Britton P: **Experimental evidence of recombination in coronavirus infectious bronchitis virus.** *Virology* 1995, **213**:569-580.
- Gelb J, Weisman Y, Ladman BS, Meir R: **S1 gene characteristics and efficacy of vaccination against infectious bronchitis virus field isolates from the United States and Israel (1996 to 2000).** *Avian Pathol* 2005, **34**:194-203.
- Gelb J, Perkins BE, Rosenberger JK, Allen PH: **Serologic and cross-protection studies with several infectious bronchitis virus isolates from Delmarva-reared broiler chickens.** *Avian Dis* 1981, **25**:655-666.
- Gelb J, Leary JH, Rosenberger JK: **Prevalence of Arkansas-type infectious bronchitis virus in Delmarva Peninsula chickens.** *Avian Dis* 1983, **27**:667-678.
- Brandt M, Yao K, Liu M, Heckert RA, Vakharia VN: **Molecular determinants of virulence, cell tropism, and pathogenic phenotype of infectious bursal disease virus.** *J Virol* 2001, **75**:11974-11982.
- Nicholas KB, Nicholas HBJ, Deerfield DW: **GeneDoc: analysis and visualization of genetic variation.** *EMBNEW NEWS* 1997, **4**:14.
- Tamura K, Dudley J, Nei M, Kumar S: **MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0.** *Mol Biol Evol* 2007, **24**:1596-1599.
- Ziebuhr J, Snijder EJ, Gorbalenya AE: **Virus-encoded proteinases and proteolytic processing in the Nidovirales.** *J Gen Virol* 2000, **81**:853-879.
- Huang H, Lau SKP, Woo PCY, Yuen K: **CoVDB: a comprehensive database for comparative analysis of coronavirus genes and genomes.** *Nucleic Acids Res* 2007, **36**:1-8.
- Liu S, Chen J, Han Z, Zhang Q, Shao Y, Kong X, Tong G: **Infectious bronchitis virus: S1 gene characteristics of vaccines used in China and efficacy of vaccination against heterologous strains from China.** *Avian Pathol* 2006, **35**:394-399.
- Mondal SP, Cardona CJ: **Genotypic and phenotypic characterization of the California 99 (Cal99) variant of infectious bronchitis virus.** *Virus Genes* 2007, **34**:327-341.
- Winterfield RW, Hitchner SB, Appleton GS: **Immunological characteristics of a variant of infectious bronchitis virus isolated from chickens.** *Avian Dis* 1964, **8**:40-47.
- Winterfield RW, Hitchner SB: **Etiology of an infectious nephritis-nephrosis syndrome of chickens.** *Am J Vet Res* 1962, **23**:1273-1279.
- Johnson RB, Marquardt WW, Newman JA: **A new serotype of infectious bronchitis virus responsible for respiratory disease in Arkansas broiler flocks.** *Avian Dis* 1973, **17**:518-523.
- Mondal SP, Cardona CJ: **Comparison of four regions in the replicase gene of heterologous infectious bronchitis virus strains.** *Virology* 2004, **324**:238-248.
- Lee CW, Jackwood MW: **Evidence of genetic diversity generated by recombination among avian coronavirus IBV.** *Arch Virol* 2000, **145**:2135-2148.
- Kwon HM, Jackwood MW: **Molecular cloning and sequence comparison of the S1 glycoprotein of the Gray and JMK strains of avian infectious bronchitis virus.** *Virus Genes* 1995, **9**:219-229.
- Hofstad MS: **Cross-immunity in chickens using seven isolates of avian infectious bronchitis virus.** *Avian Dis* 1981, **25**:650-654.
- Johnson RB, Marquardt WW: **The neutralizing characteristics of strains of infectious bronchitis virus as measured by the constant-virus variable-serum method in chicken tracheal cultures.** *Avian Dis* 1975, **19**:82-90.
- Callison S, Hilt D, Jackwood M: **Using DNA shuffling to create novel infectious bronchitis virus S1 genes: Implications for S1 gene recombination.** *Virus Genes* 2005, **31**:5-11.
- Jia W, Karaca K, Parrish DR, Naqi SA: **A novel variant of avian infectious bronchitis virus resulting from recombination among three different strains.** *Arch Virol* 1995, **140**:259-271.
- Cavanagh D, Davis PJ: **Evolution of avian coronavirus IBV: Sequence of the matrix glycoprotein gene and intergenic region of several serotypes.** *J Gen Virol* 1988, **69**:621-629.
- Wang L, Junker D, Collisson EW: **Evidence of natural recombination within the S1 gene of infectious bronchitis virus.** *Virology* 1993, **192**:710-716.
- Lai MMC: **RNA recombination in animal and plant viruses.** *Microbiol Rev* 2000, **56**(1):61-79.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

