

Hemolysin of *Streptococcus faecalis* Subspecies *zymogenes* Contributes to Virulence in Mice

YASUYOSHI IKE,^{1*} HAJIME HASHIMOTO,¹ AND DON B. CLEWELL²

Department of Microbiology, School of Medicine, Gunma University, Maebashi, Japan,¹ and Departments of Oral Biology and Microbiology/Immunology, and The Dental Research Institute, Schools of Dentistry and Medicine, The University of Michigan, Ann Arbor, Michigan 48109²

Received 13 February 1984/Accepted 8 May 1984

The conjugative plasmid pAD1 (56.7 kilobases) in *Streptococcus faecalis* confers hemolysin-bacteriocin (Hly-Bcn) expression and a mating response to the sex pheromone cAD1 excreted by recipient cells. We examined the contribution of hemolysin to pathogenicity in intraperitoneally infected mice by using Tn916 and Tn917 insertion mutants altered in hemolysin expression. Strains exhibiting the normal hemolysin phenotype were significantly more virulent than the nonhemolytic insertion mutants. A mutant plasmid with an increased copy number which gave rise to a larger-than-normal zone of hemolysis on blood agar rendered host strains more virulent than the wild-type streptococci in mice.

Hemolysin production is common among gram-positive and gram-negative bacteria; however, the extent to which this property contributes to virulence is, in most cases, not clear. There have been reports that urinary tract infections caused by *Escherichia coli* are more likely to involve hemolytic strains (7, 8, 18), and a cloned *E. coli* hemolysin determinant has been shown to enhance virulence in an animal model (21). *Streptococcus faecalis*, a normal inhabitant of the gut, can also cause urinary tract infection. In addition, it can be involved in endocarditis and root canal infection (4). *S. faecalis* subsp. *zymogenes* is distinguished from other *S. faecalis* strains by its production of a cytotoxin which is active in the lysis of human, rabbit, and horse erythrocytes. Strains producing this beta-hemolysin also produce a bacteriocin which is active against a wide variety of gram-positive bacteria, including most *Streptococcus* spp. (3). It is believed that hemolysin and bacteriocin are mediated by the same genetic determinant (1, 2, 4, 14). To date, the hemolysin-bacteriocin (Hly-Bcn) activity of *S. faecalis* strains has been shown to reside in a conjugative plasmid (4). The plasmids usually confer a sex pheromone response (4) and transfer to recipient cells at a relatively high frequency (10^{-3} to 10^{-1} per donor cell) in broth culture (10, 17). It has not been shown that the hemolytic property contributes to pathogenicity in human infection or to virulence in experimental animal infection.

Cloned fragments of DNA carrying portions of the Hly-Bcn determinants of plasmid pAD1 (56.7 kilobases) originally identified on *S. faecalis* var. *zymogenes* DS16 (5, 19), have recently been shown to share extensive homology with DNAs of other Hly-Bcn plasmids isolated from *S. faecalis* var. *zymogenes* strains of diverse geographical origin (17). These results suggest that the Hly-Bcn trait has been disseminated among *S. faecalis* strains in natural environments. In this communication, we report that the Hly-Bcn determinant of pAD1 enhanced virulence in intraperitoneal infections in mice.

The expression of pAD1-mediated Hly-Bcn activity in certain *S. faecalis* derivatives can be altered by the insertion of the tetracycline (Tc) resistance transposon Tn916 (11-13)

and the erythromycin (Em) resistance transposon Tn917 (5, 20) into specific regions of the plasmid. A physical map of the plasmid and the location of the Hly-Bcn determinant, revealed by insertion mutagenesis, have been reported (5). Insertion mutants altered in hemolysin expression were used to examine effects on virulence. The derivatives of pAD1 and the *EcoRI* fragments on which Tn916 and Tn917 are located are shown in Table 1. Plasmid pAM714 expresses normal hemolysin in its host strain (6, 16), whereas derivatives pAM307 and pAM211 do not express hemolysin (5, 12). Derivative pAM710 is a plasmid copy number mutant which gives rise to larger-than-normal zones of hemolysis for pAM714 on blood agar (15). The copy number was increased to 4.8 times that of pAM714 (one to two copies per chromosome) in a strain OG1RF1 background (15). We used *S. faecalis* OG1RF1 (9), carrying the various pAD1 derivatives, to test for virulence.

Figure 1 shows representative results on the lethality of the various strains harboring the pAD1 derivatives after intraperitoneal injection. Table 1 shows the 50% lethal dose and the time to death of the experimental mice. These data show a significant increase in mouse mortality when pAM714 was present in OG1RF1, suggesting that the plasmid has a determinant contributing to virulence. The strains harboring pAM307 or pAM211, which do not express hemolysin, did not exhibit the effect seen with pAM714. This implies that the Hly-Bcn expression encoded by pAD1 is responsible for the increased lethal behavior. The copy number mutant derivative pAM710 conferred a higher degree of virulence than did pAM714, and all of the OG1RF1-(pAM710)-infected mice died 1 to 2 h after injection. An increase in the gene dosage of hemolysin is presumed to be the cause.

These results indicate that hemolysin is toxic to mice. We examined the question of whether the OG1RF1(pAM710) culture supernatant alone was responsible for the lethality. The OG1RF1(pAM710) suspensions of 2×10^9 or 2×10^{10} cells per ml were centrifuged for 10 min at 10,000 rpm at 4°C in a Kubota RA-3 rotor, and 0.5 ml of these supernatants was intraperitoneally injected into mice by the method described in the legend to Fig. 1. None of the mice died within 7 days of the injection. These results imply that either (i) the hemolysin may be unstable and become inactivated soon after the

* Corresponding author.

TABLE 1. 50% Lethal dose of *S. faecalis* harboring pAD1 derivatives and time to death of mice after infection

Strain ^a	Phenotype	50% Lethal dose ^b	Time to death (h) of mice after infection of 10 ⁹ cells
OG1RF1(pAM710)	Hyperhemolytic	1.6 × 10 ⁸	1–2 ^c
OG1RF1(pAM714)	Normal hemolytic	2.6 × 10 ⁸	4–5 ^c
OG1RF1(pAM307)	Nonhemolytic	>3.0 × 10 ⁹	— ^d
OG1RF1(pAM211)	Nonhemolytic	>3.0 × 10 ⁹	— ^d
OG1RF1	Nonhemolytic	>3.0 × 10 ⁹	— ^d

^a pAM710, pAM714, pAM307, and pAM211 are representative derivatives of pAD1 which were altered in hemolysin expression by Tn917 or Tn916 transposon insertion. Tn917 was inserted into *EcoRI* fragment B in pAM710 and pAM714 and into *EcoRI* fragment H in pAM307. Tn916 was inserted into *EcoRI* fragment F in pAM211.

^b 50% lethal dose was derived from the data of Fig. 1.

^c All of the mice died within this time.

^d None of the mice died within 7 days of injection of 10⁹ cells. All of the mice died between 18 and 72 h after injection of 10¹⁰ cells.

excretion, and the continuous excretion by cells is necessary to provide enough toxin to be lethal; or (ii) the hemolysin may be stable, but it requires some additional substance of bacterial cell origin to become lethal.

We examined the question of whether the death of the mice was caused by hemolysin lysis of erythrocytes. Just before the death of mice inoculated with 10⁹ cells of OG1RF1(pAM710), blood samples were obtained from the orbital venous plexus with hematocrit capillary tubes and centrifuged for 5 min at 12,000 rpm with Kubota hematocrit KH-120A. The separated sera showed no reddish discoloration from hemolysis. These results indicate that the death of mice was not simply caused by hemolysin lysis of erythrocytes.




FIG. 1. Virulence of *S. faecalis* strains harboring different pAD1 derivatives. A 0.05-ml sample of an overnight culture of bacteria in Penassay broth (Difco) was plated out onto Penassay broth agar plates (Difco). The plates were incubated for 18 h at 37°C. The cells grown on the agar plates were suspended in 2 ml of phosphate-buffered saline, dilutions into phosphate-buffered saline were made, the optical density at 530 nm was checked, and the viable cells were counted on AB3 broth agar. Samples (0.5 ml) of appropriate dilutions containing the number of cells indicated in the figure were intraperitoneally injected into each mouse: Ten female ICR mice (weight, 25 to 27 g) were used for each inoculum size of bacteria. The mice were observed for 7 days. Symbols: ○, OG1RF1(pAM710); △, OG1RF1(pAM714); ●, OG1RF1(pAM307); ▲, OG1RF1(pAM211); □, OG1RF1.

Although the mode of action of *S. faecalis* hemolysin in pathogenicity is still not known, our data provide the most definitive evidence that hemolysin contributes to virulence in *S. faecalis* infection in mice. The role of hemolysin in human *S. faecalis* infection remains obscure; to our knowledge, no epidemiological data showing a correlation of the hemolytic trait with a specific type of infection have been reported.

LITERATURE CITED

- Bassinger, S. F., and R. W. Jackson. 1968. Bacteriocin (hemolysin) of *Streptococcus zymogenes*. J. Bacteriol. **96**:1895–1902.
- Brock, T. D., and J. M. Davie. 1963. Probable identity of a group D hemolysin with a bacteriocine. J. Bacteriol. **86**:708–712.
- Brock, T. D., B. Peacher, and D. Pierson. 1963. Survey of the bacteriocines of enterococci. J. Bacteriol. **86**:702–707.
- Clewell, D. B. 1981. Plasmids, drug resistance, and gene transfer in the genus *Streptococcus*. Microbiol. Rev. **45**:409–436.
- Clewell, D. B., P. K. Tomich, M. C. Gawron-Burke, A. E. Franke, Y. Yagi, and F. Y. An. 1982. Mapping of *Streptococcus faecalis* plasmids pAD1 and pAD2 and studies relating to transposition of Tn917. J. Bacteriol. **152**:1220–1230.
- Clewell, D. B., Y. Yagi, Y. Ike, R. A. Craig, B. L. Brown, and F. An. 1982. Sex pheromones in *Streptococcus faecalis*: multiple pheromone systems in strain DS5, similarities of pAD1 and pAMγ1, and mutants of pAD1 altered in conjugative properties, p. 97–100. In D. Schlessinger (ed.), *Microbiology—1982*. American Society for Microbiology, Washington, D.C.
- Cook, E. M., and S. P. Ewins. 1975. Properties of strains of *Escherichia coli* isolated from a variety of sources. J. Med. Microbiol. **8**:107–111.
- Dudgeon, L. S., E. Wordley, and F. Bawtree. 1921. On bacillus coli infections of the urinary tract, especially in relation to haemolytic organisms. J. Hyg. **20**:137–164.
- Dunny, G. M., B. L. Brown, and D. B. Clewell. 1978. Induced cell aggregation and mating in *Streptococcus faecalis*: evidence for a bacterial sex pheromone. Proc. Natl. Acad. Sci. U.S.A. **75**:3479–3483.
- Dunny, G. M., and D. B. Clewell. 1975. Transmissible toxin (hemolysin) plasmid in *Streptococcus faecalis* and its mobilization of a noninfectious drug resistance plasmid. J. Bacteriol. **124**:784–790.
- Franke, A. E., and D. B. Clewell. 1980. Evidence for conjugal transfer of a *Streptococcus faecalis* transposon (Tn916) from a chromosomal site in the absence of plasmid DNA. Cold Spring Harbor Symp. Quant. Biol. **45**:77–80.
- Franke, A. E., and D. B. Clewell. 1981. Evidence for a chromosomeborne resistance transposon (Tn916) in *Streptococcus faecalis* that is capable of "conjugal" transfer in the absence of a conjugative plasmid. J. Bacteriol. **145**:494–502.
- Gawron-Burke, M. C. and D. B. Clewell. 1982. A transposon in *Streptococcus faecalis* with fertility properties. Nature (London) **300**:865–868.
- Granato, P. A., and R. W. Jackson. 1969. Bicomponent nature of lysin from *Streptococcus zymogenes*. J. Bacteriol. **100**:865–868.
- Ike, Y., and D. B. Clewell. 1984. Genetic analysis of the pAD1 pheromone response in *Streptococcus faecalis*, using transposon Tn917 as an insertional mutagen. J. Bacteriol. **158**:777–783.
- Ike, Y., R. A. Craig, B. A. White, Y. Yagi, and D. B. Clewell. 1983. Modification of *Streptococcus faecalis* sex pheromones after acquisition of plasmid DNA. Proc. Natl. Acad. Sci. U.S.A. **80**:5369–5373.
- LeBlanc, D. J., L. N. Lee, D. B. Clewell, and D. Behnke. 1983. Broad geographical distribution of a cytotoxin gene mediating beta-hemolysis and bacteriocin activity among *Streptococcus faecalis* strains. Infect. Immun. **40**:1015–1022.
- Minshew, B. H., J. Jorgensen, G. W. Counts, and S. Falkow. 1978. Association of hemolysin production, hemagglutination of human erythrocytes, and virulence for chicken embryos of extraintestinal *Escherichia coli* isolates. Infect. Immun. **20**:50–54.

19. Tomich, P., F. An, S. Damale, and D. B. Clewell. 1979. Plasmid related transmissibility and multiple drug resistance in *Streptococcus faecalis* subsp. *zymogenes* strain DS16. *Antimicrob. Agents Chemother.* **15**:828–830.
20. Tomich, P. K., F. Y. An, and D. B. Clewell. 1980. Properties of erythromycin-inducible transposon Tn917 in *Streptococcus faecalis*. *J. Bacteriol.* **141**:1366–1374.
21. Welch, R. A., E. P. Dellinger, B. Minshew, and S. Falkow. 1981. Haemolysin contributes to virulence of extra-intestinal *E. coli* infections. *Nature (London)* **294**:665–667.