

Research

Open Access

Pore dilation occurs in TRPA1 but not in TRPM8 channels

Jun Chen^{*1}, Donghee Kim^{*2}, Bruce R Bianchi¹, Eric J Cavanaugh²,
Connie R Faltynek¹, Philip R Kym¹ and Regina M Reilly¹

Address: ¹Neuroscience, Global Pharmaceutical Research and Development, Abbott Laboratories, Abbott Park, IL 60064-6125, USA and
²Department of Physiology, Rosalind Franklin University of Medicine and Science, The Chicago Medical School, 3333 Green Bay Road, North
Chicago, IL60064, USA

Email: Jun Chen^{*} - jun.x.chen@abbott.com; Donghee Kim^{*} - Donghee.Kim@rosalindfranklin.edu;
Bruce R Bianchi - bruce.bianchi@abbott.com; Eric J Cavanaugh - Eric.Cavanaugh@rosalindfranklin.edu;
Connie R Faltynek - connie.r.faltynek@abbott.com; Philip R Kym - Phil.R.Kym@abbott.com; Regina M Reilly - Regina.Reilly@abbott.com

^{*} Corresponding authors

Published: 21 January 2009

Received: 20 October 2008

Molecular Pain 2009, **5**:3 doi:10.1186/1744-8069-5-3

Accepted: 21 January 2009

This article is available from: <http://www.molecularpain.com/content/5/1/3>

© 2009 Chen et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Abundantly expressed in pain-sensing neurons, TRPV1, TRPA1 and TRPM8 are major cellular sensors of thermal, chemical and mechanical stimuli. The function of these ion channels has been attributed to their selective permeation of small cations (e.g., Ca²⁺, Na⁺ and K⁺), and the ion selectivity has been assumed to be an invariant fingerprint to a given channel. However, for TRPV1, the notion of invariant ion selectivity has been revised recently. When activated, TRPV1 undergoes time and agonist-dependent pore dilation, allowing permeation of large organic cations such as Yo-Pro and NMDG⁺. The pore dilation is of physiological importance, and has been exploited to specifically silence TRPV1-positive sensory neurons. It is unknown whether TRPA1 and TRPM8 undergo pore dilation. Here we show that TRPA1 activation by reactive or non-reactive agonists induces Yo-Pro uptake, which can be blocked by TRPA1 antagonists. In outside-out patch recordings using NMDG⁺ as the sole external cation and Na⁺ as the internal cation, TRPA1 activation results in dynamic changes in permeability to NMDG⁺. In contrast, TRPM8 activation does not produce either Yo-Pro uptake or significant change in ion selectivity. Hence, pore dilation occurs in TRPA1, but not in TRPM8 channels.

Background

Abundantly expressed in sensory neurons, TRPV1, TRPA1 and TRPM8 are involved in sensory function, pain and neurogenic inflammation [1]. The function of these ion channels has been attributed to their ability to pass certain ion species across the plasma membrane. Once activated, TRPV1, TRPA1 and TRPM8 are permeable to small cations such as Ca²⁺, K⁺, Na⁺; hence, channel activation simultaneously depolarizes the plasma membrane and raises intracellular Ca²⁺, which subsequently triggers a variety of physiological processes. By analogy to voltage-gated K⁺

channels, it is assumed that ion selectivity of TRP channels should be an invariant signature to the respective channel. However, this notion has been challenged recently. When activated, TRPV1 exhibits time and agonist-dependent changes in ion selectivity [2]. In fact, TRPV1 undergoes pore dilation and allows permeation of large organic cations, including spermine (202.3 Da), NMDG (195.2 Da), Yo-Pro (376 Da), gentamycin (477.6 Da) and QX-314 [3-7]. Here we explored whether TRPA1 and TRPM8 undergo pore dilation by examining Yo-Pro uptake and changes in ion selectivity upon channel activation.

Results and discussion

Yo-Pro is a divalent cation impermeable to the plasma membrane. However, under certain conditions, it can enter cells, bind nucleic acids and emit fluorescence. Hence the uptake of Yo-Pro has been used previously as an indicator of pore dilation [2,8,9]. In HEK293-F cells transiently expressing rat TRPA1, allyl isothiocyanate (AITC) evoked robust increases in intracellular Ca^{2+} (Fig. 1A). Concomitantly, AITC also induced Yo-Pro uptake in a concentration-dependent manner (Fig. 1B). At higher concentrations of AITC (100 or 300 μM), the increase in fluorescence was immediately noticeable and continued to increase for about 50 min. In addition, AITC also induced Ca^{2+} influx and Yo-Pro uptake in cells expressing human TRPA1 and mouse TRPA1, but not in untransfected cells (data not shown). In cells expressing human TRPM8, menthol activated TRPM8 as indicated by the concentration-dependent Ca^{2+} influx, but failed to induce Yo-Pro uptake (Fig. 1C and 1D). Other TRPM8 agonists (e.g., icilin) also evoked Ca^{2+} influx but failed to induce

Yo-Pro uptake (data not shown). Hence, Yo-Pro uptake occurs upon activation of TRPA1, but not TRPM8.

In addition to AITC, TRPA1 can be activated by many other electrophilic agonists (e.g., cinnamaldehyde or CA, 4-hydroxynonanal or 4-HNE), and non-reactive agonists (e.g., URB597, farnesyl thiosalicylic acid or FTS) [10-14]. We investigated whether the Yo-Pro uptake is limited to AITC. CA, 4-HNE, FTS and URB597 all evoked Ca^{2+} influx and Yo-Pro uptake in a concentration dependent-manner (Fig. 2A and 2B). In the Ca^{2+} assay, the EC_{50} was $6.5 \pm 0.35 \mu\text{M}$ for AITC, $6.8 \pm 1.5 \mu\text{M}$ for CA, $4.4 \pm 0.6 \mu\text{M}$ for 4-HNE, $33.2 \pm 8.1 \mu\text{M}$ for FTS and $85.6 \pm 10.4 \mu\text{M}$ for URB597 ($n = 4-8$). Compared to AITC, the maximal signals were 104% for CA, 88% for 4-HNE, 107% for FTS and 82% for URB597. In the Yo-Pro uptake assay, the EC_{50} was $16.0 \pm 3.8 \mu\text{M}$ for AITC, $5.9 \pm 0.7 \mu\text{M}$ for CA, $7.1 \pm 0.2 \mu\text{M}$ for 4-HNE, $41.8 \pm 10.7 \mu\text{M}$ for FTS and $85.4 \pm 19.8 \mu\text{M}$ for URB597 ($n = 4-8$). Compared to AITC, the maximal signals were 98% for CA, 82% for 4-HNE, 117% for FTS and




Figure 1
The activation of TRPA1, but not TRPM8, induced Yo-Pro uptake. A, in HEK-293F cells expressing rat TRPA1, AITC elevated intracellular Ca^{2+} , as represented by increases of fluorescence signals (RFU) in the FLIPR based Ca^{2+} assay. B, in cells expressing TRPA1, AITC evoked robust Yo-Pro uptake in a concentration-dependent manner from the FLIPR based Yo-Pro uptake assays. C, in cells expressing human TRPM8, menthol activated TRPM8 and elevated intracellular Ca^{2+} . D, in cells expressing TRPM8, menthol failed to induce Yo-Pro uptake. Compounds are in μM and additions are indicated by arrows.

Figure 2
Yo-Pro uptake was evoked by various TRPA1 agonists and blocked by TRPA1 antagonists. Concentration-effect relationships for agonist responses in the Ca^{2+} assay (A) and Yo-Pro uptake (B). Reactive agonists: AITC, CA and 4-HNE. Non-reactive agonists: FTS and URB597. Data are represented as percentage of maximal AITC responses. C, representative traces of Yo-Pro uptake in response to a first addition of AP18 (0 to 100 μM) and a second addition of AITC (30 μM). AP18 inhibited AITC evoked Yo-Pro uptake in a concentration-dependent manner. D, concentration-effect relationship of Yo-Pro uptake inhibition by AP18, HC-030031 and RR. ($n = 4 - 8$).

84% for URB597, respectively. Hence, TRPA1 activation by different agonists all induced Yo-Pro uptake.

Several small molecule inhibitors of TRPA1 have been described recently, including AP18, HC-030031 and ruthenium red (RR) [15,16]. We tested whether these antagonists blocked Yo-Pro uptake. AP18 attenuated 30 μM AITC-induced Yo-Pro uptake in a concentration-dependent manner, with an IC_{50} of $10.3 \pm 0.8 \mu\text{M}$ (Fig. 2C and 2D). Likewise, HC-030031 and RR also completely blocked Yo-Pro uptake (IC_{50} : $9.4 \pm 0.6 \mu\text{M}$ for HC-030031 and $10.0 \pm 1.6 \mu\text{M}$ for RR). Taken together, these data show that agonist-evoked Yo-Pro uptake is related to TRPA1 channel activities.

Next, we investigated whether TRPA1 undergoes changes in ion selectivity upon channel activation. Currents were recorded under the outside-out patch configuration using NMDG⁺ as the sole external cation and Na⁺ as the major internal cation. Patch membrane potential was held at -80 mV, and a ramp voltage from -140 mV to 0 mV (500 ms duration) was applied every 3 seconds. Before addition of AITC, a small basal current was present, consistent with previous reports [17,18]. The reversal potential (E_{rev}) of basal currents was $-95.3 \pm 4.8 \text{ mV}$ ($n = 5$). Compared to activation of TRPV1 by capsaicin, activation of TRPA1 by AITC was relatively slow, probably due to the covalent reaction that is needed to activate TRPA1. Addition of AITC (100 μM) elicited gradual activation of TRPA1 and rightward shift in reversal potential (Fig. 3A). The shift in E_{rev} occurred as early as 6 s following addition of AITC, and continued to increase with nearly maximum shift at ~ 15 s. Addition of 10 μM RR nearly completely blocked AITC-evoked NMDG⁺ and Na⁺ currents (Fig. 3A inset), indicating the observed currents were mediated by TRPA1 channels.

In contrast, TRPM8 showed no shift in E_{rev} following addition of 500 μM menthol, despite a clear increase in current (Fig. 3B). The time-dependent changes in E_{rev} for TRPA1 and TRPM8 following their activation are shown in Fig. 4A. The shift in E_{rev} for TRPA1 was not due to an increase in anion selectivity, as removal of Cl⁻ in the bath solution caused a similar shift in E_{rev} from -96 mV to -42 mV. From E_{rev} values, permeability ratios ($P_{\text{NMDG}}/P_{\text{Na}}$) before and 60 s after agonist addition were derived. As shown in Fig. 4B, $P_{\text{NMDG}}/P_{\text{Na}}$ increased ~ 4.4 -fold for TRPA1 from 0.05 ± 0.003 to 0.22 ± 0.013 ($n = 4$, $P < 0.05$, paired t-test), comparable to the ~ 5.5 fold increase reported for TRPV1 [2]. In contrast, $P_{\text{NMDG}}/P_{\text{Na}}$ did not change significantly for TRPM8. It is interesting that the shift in E_{rev} occurred much earlier than the increase in TRPA1 currents (Fig. 4C), indicating that pore dilation occurs well before maximal channel activation.




Figure 3
Ionic currents of TRPA1, but not TRPM8, exhibited shifts in reversal potential. Outside-out patches were formed from HeLa cells expressing rat TRPA1 or human TRPM8 plus GFP. NMDG⁺ was the sole external cation and Na⁺ was the major internal cation. Membrane potential was held at -80 mV, and a voltage ramp from -140 to 0 mV (500 ms duration) was applied every 3 s immediately. **A**, current traces from a representative TRPA1-containing patch during 60 s application of AITC (100 μM). To illustrate shifts in E_{rev} , only currents between -6 to 8 pA were plotted. Inset shows the AITC-evoked currents were almost completely blocked by 10 μM RR. **B**, currents from a representative TRPM8-containing patch during 60 s application of menthol (500 μM). Note the shifts in E_{rev} for TRPA1, but no shift for TRPM8.

The accuracy of E_{rev} measurement could be compromised by small current amplitudes, especially for basal currents and currents immediately following AITC application. However, the E_{rev} of basal currents was consistent across patches ($-95.3 \pm 4.8 \text{ mV}$, $n = 5$), and E_{rev} shifts consistently occurred in TRPA1, but not in TRPM8. In addition, even for relatively large TRPA1 currents, significant shifts in E_{rev} occurred. For example, the shift in E_{rev} was 31.6 mV between 6 s and 15 s pulses, and 14.2 mV between 9 s and




Figure 4
Time-dependent changes in ion permeability occurred in TRPA1 but not in TRPM8. A, E_{rev} values were determined (from Fig. 3 experiments) and plotted as a function of time after application of AITC or menthol. B, permeability ratios (P_{NMDG}/P_{Na}) before and 60 s after agonist addition. Student's t-test was used with $p < 0.05$ as the criterion for significance (indicated by *). C, changes in E_{rev} and relative currents at 0 mV (from representative recordings) were plotted as a function of time after application of AITC. Outward currents were measured at 0 mV and normalized against the current obtained after 60 s addition of AITC.

15 s pulses (Fig. 3A and 4A). Taken together, these data suggests that the dynamic change in E_{rev} results from TRPA1 channel activity. Another concern in extrapolating the change in E_{rev} to the change in ion selectivity is that ion accumulation can occur during prolonged activation, particularly when large currents are conducted under

whole cell configuration. However, the ion accumulation should not significantly compromise our TRPA1 experiments using outside-out patch configuration, in which extracellular and intracellular ionic conditions were well controlled. In addition, reversal potentials changed within seconds of AITC application when currents were small, but reached a steady state (from 15 to 60 s) when currents were relatively large (Fig. 3A and 4A). Furthermore, TRPM8 conducted currents with similar amplitudes but without significant shifts in reversal potential. Consistent with the electrophysiology data, the large divalent cation Yo-Pro did not cross the membrane when the channel was closed or blocked by antagonists, but permeated the membrane freely when the channel was open (Fig. 1B and 2C). Collectively, our data suggest that TRPA1, but not TRPM8, undergoes pore dilation.

Pore dilation has been previously described for the ATP-gated P2X and TRPV1 [2,8,9]. For P2X channels, the mechanism underlying pore dilation remains controversial. Several alternative mechanisms have been proposed including a direct change in ion selectivity, formation of channel multimers, and recruitment of a downstream, nonselective pore [19,20]. For TRPV1, pore dilation most likely arises from a change in ion selectivity, as indicated by the dynamic change in ion selectivity during agonist stimulation, and the effects of mutations and chemical modification of certain residues within the selectivity filter [2]. In the current study, we did not elucidate the biophysical mechanism underlying pore dilation of TRPA1. However, there were several notable observations. First, the permeability to NMDG⁺ increased almost immediately upon channel activation. Second, the outside-out patch configuration should largely disrupt cytoskeletal structures and washout cytosolic factors. Third, the AITC evoked- NMDG⁺ and Na⁺ conductance was sensitive to blockade by ruthenium red. Finally, under identical conditions, TRPM8 conducted large currents, but did not exhibit Yo-Pro uptake or a significant change in NMDG⁺ permeability. Thus, TRPA1 pore dilation most likely represents a direct change in ion selectivity. Nonetheless, our present study does not completely rule out the involvement of other proteins.

TRPV1, TRPA1 and TRPM8 are major TRP channels involved in somatosensation. Within dorsal root ganglia, TRPV1 and TRPA1 are co-expressed and interact functionally in one population of sensory neurons, while TRPM8 is expressed largely in a separate neuronal population. Interestingly, pore dilation occurs in TRPA1, TRPV1 but not TRPM8, suggesting that this property is not ubiquitous, but rather specific to subtypes of channels within a subpopulation of neurons. The change in cation permeability, in turn, may alter channel function, affect a host of downstream processes (e.g., neurotransmitter release, cel-

lular toxicity) and contribute to pain hypersensitivity [21]. Recently, it was reported that TRPV1-mediated pore dilation could be utilized to deliver QX-314 (a membrane-impermeant sodium channel blocker) specifically to TRPV1-positive sensory neurons, achieving analgesic effects without motor deficits associated with local anesthetics [5]. However, this strategy of targeting TRPV1-positive neurons could be compromised by several factors, including the broad expression pattern of TRPV1, its role in regulating body temperature, and its involvement in hippocampal synaptic plasticity [22,23]. By analogy to TRPV1, the pore dilation of TRPA1 could be exploited to mediate entry of QX-314 specifically into TRPA1-positive neurons. Given the restrictive expression of TRPA1 in sensory neurons, this strategy may offer analgesic efficacy without unwanted side effects.

In conclusion, the present study demonstrates that pore dilation occurs in TRPA1 but not in TRPM8 channels. This finding raises many interesting questions: What is the exact biophysical mechanism underlying pore dilation of TRPA1? What are the physiological, pathological and therapeutic implications? Why does pore dilation not occur in TRPM8? What are the pore behaviors of other TRP channels? Answers to these questions will certainly extend our understanding of this family of ion channels.

Methods

Transient expression of recombinant TRPs

Full length cDNAs for rat TRPA1 (GenBank Accession: [NM_207608](#)), human TRPA1 ([NM_007332](#)), mouse TRPA1 ([NM_177781](#)) and human TRPM8 ([NM_024080](#)) were cloned into pcDNA3.1/V5-His TOPO vector and transiently expressed in HEK293-F or HeLa cells [24]. For the Ca²⁺ influx or Yo-Pro uptake assay, HEK293-F cells were transfected with TRP cDNA, collected 48 hours post transfection, and used either fresh or following storage at -70°C. For electrophysiological experiments, HeLa cells were transfected with TRPA1 or TRPM8 plus GFP, and used 48 hours later.

Ca²⁺ influx and Yo-Pro uptake assays

Ca²⁺ influx assay was performed using the FLIPR™ and calcium assay kit R8033 (MDS Analytical Technology) as reported previously [25]. After incubation with 100 μl of 1 × Ca²⁺ dye for ~2 hours at room temperature, a two-addition protocol was used for evaluating agonist activities (i.e., activation of Ca²⁺ influx) and antagonist activities (i.e., inhibition of agonist responses): 10 s baseline readout, addition of 50 μl assay buffer or antagonist (4 × stock), 3–4 min readout, addition of 50 μl agonist (4 × stock), and readout for 2.5 min. Maximum minus minimum signals before the second addition and at the end of the experiment were obtained.

Yo-Pro uptake was determined using the FLIPR™ and Mg²⁺/Ca²⁺-free DPBS buffer as reported previously [26]. Briefly, immediately after loading with 100 μl Yo-Pro dye (2 μM), a two-addition protocol was used for evaluating agonist activity (i.e., Yo-Pro uptake) and antagonist activities (i.e., inhibition of agonist evoked Yo-Pro uptake): 10 s baseline readout, addition of 50 μl assay buffer or antagonist, 3 min readout, addition of 50 μl agonists, and readout for 60 min. Max-min fluorescence signals before the second addition and at the end of the experiment were obtained.

Outside-out patch recording

Before forming cell-attached patches from HeLa cells expressing TRPA1 or TRPM8, pipette offset was adjusted to give a zero current value. Outside-out patches with the least amount of leak were used, as judged by the very small DC shift (<5 mV) of the basal current at different membrane potentials. Currents were recorded using AxoPatch200B. The pipette solution contained (mM): 140 NaCl, 1 MgCl₂, 5 mM EGTA, and 10 HEPES (pH 7.3). For experiments with NMDG⁺, the bath solution contained (mM): 150 mM NMDG⁺, 115 mM Cl⁻, 5 mM EGTA, and 10 HEPES (pH 7.3). For Cl⁻ replacement experiments, the bath contained 150 mM NMDG⁺, 62 mM EGTA and 10 HEPES (pH 7.3). Patch membrane potential was held at -80 mV, and then a voltage ramp from -140 mV to 0 mV (500 ms duration) was applied every 3 seconds. Current was filtered at 1 kHz using 8-pole Bessel filter (-3 dB; Frequency Devices) and transferred directly to a computer using the Digidata 1320 interface (Axon Instruments) at a sampling rate of 10 kHz. Permeability ratio (P_X/P_{Na}) was calculated using the equation: $P_X/P_{Na} = ([X]_o/[Na]_o) \cdot \exp(\Delta E_{rev} \cdot F/RT)$; where ΔE_{rev} represents the shift in E_{rev} after addition of AITC in NMDG⁺ external/Na⁺ internal solution, and F/RT is 0.040 mV⁻¹. The activity coefficient of Na⁺ and NMDG⁺ was taken as 0.75 and 0.81, respectively. Student's t-test was used with $p < 0.05$ as the criterion for significance. Data are represented as mean ± S.E. unless specified otherwise.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

JC conceived, coordinated the study and made the initial finding. BRB and JC conducted FLIPR experiments. DK and EJC conducted the electrophysiology experiments. CRF, PRK and RMR supported the study. JC and DK wrote the manuscript. All authors read and approved the final manuscript.

References

1. Wang H, Woolf CJ: **Pain TRPs**. *Neuron* 2005, **46**(1):9-12.

2. Chung MK, Guler AD, Caterina MJ: **TRPV1 shows dynamic ionic selectivity during agonist stimulation.** *Nat Neurosci* 2008, **11(5)**:555-64.
3. Ahern GP, Wang X, Miyares RL: **Polyamines are potent ligands for the capsaicin receptor TRPV1.** *J Biol Chem* 2006, **281(13)**:8991-5.
4. Meyers JR, MacDonald RB, Duggan A, Lenzi D, Standaert DG, Corwin JT, Corey DP: **Lighting up the senses: FMI-43 loading of sensory cells through nonselective ion channels.** *J Neurosci* 2003, **23(10)**:4054-65.
5. Binshtok AM, Bean BP, Woolf CJ: **Inhibition of nociceptors by TRPV1-mediated entry of impermeant sodium channel blockers.** *Nature* 2007, **449(7162)**:607-10.
6. Myrdal SE, Steyger PS: **TRPV1 regulators mediate gentamicin penetration of cultured kidney cells.** *Hear Res* 2005, **204(1-2)**:170-82.
7. Hellwig N, Plant TD, Janson W, Schafer M, Schultz G, Schaefer M: **TRPV1 acts as proton channel to induce acidification in nociceptive neurons.** *J Biol Chem* 2004, **279(33)**:34553-61.
8. Virginio C, MacKenzie A, Rassendren FA, North RA, Surprenant A: **Pore dilation of neuronal P2X receptor channels.** *Nat Neurosci* 1999, **2(4)**:315-21.
9. Khakh BS, Bao XR, Labarca C, Lester HA: **Neuronal P2X transmitter-gated cation channels change their ion selectivity in seconds.** *Nat Neurosci* 1999, **2(4)**:322-30.
10. Story GM, Peier AM, Reeve AJ, Eid SR, Mosbacher J, Hricik TR, Earley TJ, Hergarden AC, Andersson DA, Hwang SW, McIntyre P, Jegla T, Bevan S, Patapoutian A: **ANKTM1, a TRP-like channel expressed in nociceptive neurons, is activated by cold temperatures.** *Cell* 2003, **112(6)**:819-29.
11. Niforatos W, Zhang XF, Lake MR, Walter KA, Neelands T, Holzman TF, Scott VE, Faltynek CR, Moreland RB, Chen J: **Activation of TRPA1 channels by the fatty acid amide hydrolase inhibitor 3'-carbamoylbiphenyl-3-yl cyclohexylcarbamate (URB597).** *Mol Pharmacol* 2007, **71(5)**:1209-16.
12. Cavanaugh EJ, Simkin D, Kim D: **Activation of transient receptor potential A1 channels by mustard oil, tetrahydrocannabinol and Ca(2+) reveals different functional channel states.** *Neuroscience* 2008, **154(4)**:1467-76.
13. Maher M, Ao H, Banke T, Nasser N, Wu NT, Breitenbucher JG, Chaplan SR, Wickenden AD: **Activation of TRPA1 by farnesyl thiosalicylic acid.** *Mol Pharmacol* 2008, **73(4)**:1225-34.
14. Trevisani M, Siemens J, Materazzi S, Bautista DM, Nassini R, Campi B, Imamachi N, Andre E, Patacchini R, Cottrell GS, Gatti R, Basbaum AI, Bunnett NW, Julius D, Geppetti P: **4-Hydroxynonenal, an endogenous aldehyde, causes pain and neurogenic inflammation through activation of the irritant receptor TRPA1.** *Proc Natl Acad Sci USA* 2007, **104(33)**:13519-24.
15. Petrus M, Peier AM, Bandell M, Hwang SW, Huynh T, Olney N, Jegla T, Patapoutian A: **A role of TRPA1 in mechanical hyperalgesia is revealed by pharmacological inhibition.** *Mol Pain* 2007, **3**:40.
16. McNamara CR, Mandel-Brehm J, Bautista DM, Siemens J, Deranian KL, Zhao M, Hayward NJ, Chong JA, Julius D, Moran MM, Fanger CM: **TRPA1 mediates formalin-induced pain.** *Proc Natl Acad Sci USA* 2007, **104(33)**:13525-30.
17. Nagata K, Duggan A, Kumar G, Garcia-Anoveros J: **Nociceptor and hair cell transducer properties of TRPA1, a channel for pain and hearing.** *J Neurosci* 2005, **25(16)**:4052-61.
18. Zhang XF, Chen J, Faltynek CR, Moreland RB, Neelands TR: **Transient receptor potential A1 mediates an osmotically activated ion channel.** *Eur J Neurosci* 2008, **27(3)**:605-11.
19. Fujiwara Y, Kubo Y: **Density-dependent changes of the pore properties of the P2X2 receptor channel.** *J Physiol* 2004, **558(Pt 1)**:31-43.
20. Locovei S, Scemes E, Qiu F, Spray DC, Dahl G: **Pannexin1 is part of the pore forming unit of the P2X(7) receptor death complex.** *FEBS Lett* 2007, **581(3)**:483-8.
21. Bautista D, Julius D: **Fire in the hole: pore dilation of the capsaicin receptor TRPV1.** *Nat Neurosci* 2008, **11(5)**:528-9.
22. Gavva NR, Bannan AW, Surapaneni S, Hovland DN Jr, Lehto SG, Gore A, Juan T, Deng H, Han B, Klionsky L, Kuang R, Le A, Tamir R, Wang J, Youngblood B, Zhu D, Norman MH, Magal E, Treanor JJ, Louis JC: **The vanilloid receptor TRPV1 is tonically activated in vivo and involved in body temperature regulation.** *J Neurosci* 2007, **27(13)**:3366-74.
23. Gibson HE, Edwards JG, Page RS, Van Hook MJ, Kauer JA: **TRPV1 channels mediate long-term depression at synapses on hippocampal interneurons.** *Neuron* 2008, **57(5)**:746-59.
24. Chen J, Zhang XF, Kort ME, Huth JR, Sun C, Miesbauer LJ, Cassar SC, Neelands T, Scott VE, Moreland RB, Reilly RM, Hajduk PJ, Kym PR, Hutchins CW, Faltynek CR: **Molecular determinants of species-specific activation or blockade of TRPA1 channels.** *J Neurosci* 2008, **28(19)**:5063-71.
25. Chen J, Lake MR, Sabet RS, Niforatos W, Pratt SD, Cassar SC, Xu J, Gopalakrishnan S, Pereda-Lopez A, Gopalakrishnan M, Holzman TF, Moreland RB, Walter KA, Faltynek CR, Warrior U, Scott VE: **Utility of large-scale transiently transfected cells for cell-based high-throughput screens to identify transient receptor potential channel A1 (TRPA1) antagonists.** *J Biomol Screen* 2007, **12(1)**:61-9.
26. Donnelly-Roberts DL, Namovic MT, Faltynek CR, Jarvis MF: **Mitogen-activated protein kinase and caspase signaling pathways are required for P2X7 receptor (P2X7R)-induced pore formation in human THP-1 cells.** *J Pharmacol Exp Ther* 2004, **308(3)**:1053-61.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

