

Review

Calcium Signaling Network in Plants

An Overview

Narendra Tuteja*

Shilpi Mahajan

Plant Molecular Biology Group; International Centre for Genetic Engineering and Biotechnology; New Delhi, India

*Correspondence to: Narendra Tuteja; Plant Molecular Biology Group; International Centre for Genetic Engineering and Biotechnology; New Delhi- 67, India; Tel.: +91.11.26189358; Fax: +91.11.26162316; Email: narendra@icgeb.res.in

Original manuscript submitted: 03/22/07

Manuscript accepted: 03/22/07

Previously published online as a *Plant Signaling & Behavior* E-publication: <http://www.landesbioscience.com/journals/psb/abstract.php?id=4176>

KEY WORDS

Calcium binding proteins, Ca²⁺ channel, Ca²⁺-dependent protein kinases, Ca²⁺/H⁺ antiport, calcium memory, calcium sensors, calcium signatures, Ca²⁺-transporters, EF hand motifs, plant signal transduction

ACKNOWLEDGEMENTS

I thank Dr. Renu Tuteja for critical reading and corrections on the article. This work was partially supported by the grants from the Department of Biotechnology (DBT), Department of Science and Technology (DST) and Defence Research and Development Organisation (DRDO), Government of India. I apologize if some references could not be cited due to space constraint.

ABSTRACT

Calcium ion (Ca²⁺) is one of the very important ubiquitous intracellular second messenger molecules involved in many signal transduction pathways in plants. The cytosolic free Ca²⁺ concentration ([Ca²⁺]_{cyt}) have been found to increased in response to many physiological stimuli such as light, touch, pathogenic elicitor, plant hormones and abiotic stresses including high salinity, cold and drought. This Ca²⁺ spikes normally result from two opposing reactions, Ca²⁺ influx through channels or Ca²⁺ efflux through pumps. The removal of Ca²⁺ from the cytosol against its electrochemical gradient to either the apoplast or to intracellular organelles requires energized 'active' transport. Ca²⁺-ATPases and H⁺/Ca²⁺ antiporters are the key proteins catalyzing this movement. The increased level of Ca²⁺ is recognised by some Ca²⁺-sensors or calcium-binding proteins, which can activate many calcium dependent protein kinases. These kinases regulate the function of many genes including stress responsive genes, resulted in the phenotypic response of stress tolerance. Calcium signaling is also involved in the regulation of cell cycle progression in response to abiotic stress. The regulation of gene expression by cellular calcium is also crucial for plant defense against various stresses. However, the number of genes known to respond to specific transient calcium signals is limited. This review article describes several aspects of calcium signaling such as Ca²⁺ requirement and its role in plants, Ca²⁺ transporters, Ca²⁺-ATPases, H⁺/Ca²⁺-antiporter, Ca²⁺-signature, Ca²⁺-memory and various Ca²⁺-binding proteins (with and without EF hand).

INTRODUCTION

Calcium plays a fundamental role in plant's growth and development. Many extracellular signals and environmental cues including light, abiotic and biotic stress factors, elicit change in the cellular calcium levels, termed as calcium signatures.¹ Ca²⁺ has been well established as a second messenger and the concentration of Ca²⁺ is delicately balanced by the presence of 'Ca²⁺ stores' like vacuoles, endoplasmic reticulum, mitochondria and cell wall. Ca²⁺ is present in millimolar concentrations in the cell wall and vacuoles and is released whenever required by the cell.^{1, 2} Recently, Xiong et al. (2006)³ have demonstrated that the organelles surrounded by a double membrane (e.g. mitochondria, chloroplasts and nuclei) are also equipped to generate calcium signal on their own, which is delimited by a double membrane.

Ca²⁺ ion represents an important signaling molecule and a convergence point of many disparate signaling pathways. Facing the environmental challenge, plant cells reprogram their cellular set up by triggering a network of signaling events that start with stress perception at the membrane level of the cell and ends with a cellular response (Fig. 1). A generic signal transduction pathway has following steps:

(a) Perception of the signal by the membrane receptors; (b) Generation of second messengers; (c) A cascade of protein phosphorylation/dephosphorylation events that may target transcription factors controlling specific set of stress regulated genes; (d) Stress tolerance, plant adaptation and other phenotypic responses.

The extracellular stress signal transduces inside the nucleus to induce multiple stress responsive genes, the products of which ultimately lead to plant adaptation to stress tolerance directly or indirectly (Fig. 1).¹ Overall, the stress response could be a coordinated action of many genes, which may cross-talk with each others. Signal transduction requires the proper spatial and temporal coordination of all signaling molecules. In plant cell, many molecules acting as second messenger in signaling pathways have been reported; these include Ca²⁺, lipids like IP₃ and cyclic GMP (cGMP). However no single messenger except cytosolic Ca²⁺ has been demonstrated to be involved in diverse




Figure 1. Generic pathway for plant response to stress. The extracellular stress signal is first perceived by the membrane receptors and then activate large and complex signaling cascade intracellularly including the generation of secondary signal molecules. The signal cascade results into the expression of multiple stress responsive genes, the products of which can provide the stress tolerance directly or indirectly. ABA, abscisic acid; LEA, late embryogenesis abundant; InsP, inositol phosphate; ROS, reactive oxygen species.




Figure 2. A generic pathway for calcium regulated gene expression and the stress response. The calcium level increases in response to the stimuli. The increased level of Ca^{2+} is recognised by some Ca^{2+} -sensors or calcium-binding proteins, which can activate many calcium dependent protein kinases. These kinases regulate the function of many genes including stress responsive genes, resulted in the phenotypic response of stress tolerance.

pathways and respond to numerous stimuli. A generic pathway for calcium regulated expression of stress responsive genes is shown in Figure 2. The elevated level of Ca^{2+} can be recognized by calcium sensors or Ca^{2+} -binding proteins, which together can activate protein kinases.^{1, 2} These activated protein kinases can phosphorylate many regulatory proteins including transcription factors, which

regulate the gene expression level of regulatory proteins resulting in the altered metabolism followed by phenotypic response of stress tolerance (Fig. 2). The response could also be growth inhibition or cell death, which will depend upon how many and what kinds of genes are getting upregulated or down-regulated in response to the elevated calcium.

Overall, in response to calcium several genes are reported to be upregulated. An analysis of transcriptome changes revealed 230 calcium-responsive genes, of which 162 were upregulated and 68 were downregulated. These include known early stress-responsive genes as well as genes of unknown function.⁴ Recently, a blue light receptor phototropins, which regulate growth and development of plants, has also been shown to involved in calcium signaling in higher plants.⁵ Ca^{2+} signaling pathway can also regulates a K^+ channel for low- K response in *Arabidopsis*. Calcium is also reported to be an essential component of the sucrose signaling pathway that leads to the induction of fructan synthesis.⁶ Calcium signaling is also involved in the regulation of cell cycle progression in response to abiotic stress. Understanding the immense significance of Ca^{2+} ions, this review has been solely dedicated to the salient features associated with calcium signaling. Various aspects regarding the Ca^{2+} requirements of plant, Ca^{2+} deficiency, Ca^{2+} transporters, efflux pumps, $\text{Ca}^{2+}/\text{H}^+$ antiporters, Ca^{2+} signatures, Ca^{2+} memory, Ca^{2+} sensor and transducer proteins have been briefly covered in this review.

Role of calcium in plants. (1) Calcium is an essential plant nutrient required for growth and development of plant, especially the root and shoot tip. The tips are meristematic and cell division occurs by mitosis. Ca^{2+} helps in the formation of microtubules and microtubules in turn are essential for the anaphasic movement of chromosomes.

(2) Ca^{2+} is an important divalent cation and is required for structural roles in the cell wall and membranes where it exists as Ca^{2+} pectate. Ca^{2+} accumulates as calcium pectate in the cell wall and binds the cells together.

(3) It is also required as a counter-cation for inorganic and organic anions in the vacuole and as an intracellular messenger in the cytosol.¹ Externally supplied Ca^{2+} reduces the toxic effects of NaCl and ameliorates stress.

(4) Ca^{2+} is required for pollen tube growth and elongation.⁸

The calcium requirement of plants and its uptake from soil. Calcium was first described as an essential macro nutrient element in plants more than a century ago.⁹ Ca^{2+} is taken up by roots from the soil solution and delivered to the shoot via the xylem. Ca^{2+} may traverse the root either through the cytoplasm of the cells, linked together by plasmodesmata (the symplast) or through the spaces between the cells (the apoplast). The cytosolic free Ca^{2+} concentration is maintained typically at 200 nM.¹⁰ However, the Ca^{2+} content of the cytosol is far higher than this because of the high affinity of Ca^{2+} to a range of Ca^{2+} binding proteins. Ca^{2+} mobility within the cell is very low as a consequence of its low free concentration and rapid chelation. It moves predominantly apoplastically rather than symplastically.

Plants vary markedly in their Ca^{2+} content and requirements. Ca^{2+} content can vary between 0.1 and >5.0% dry wt.¹¹ for different plants and organs. Soil Ca^{2+} also varies widely from <0.01% calcium in acid laterites to very high abundance in chalky soils.⁹ Ecologists have classified plant species into calcifuges and calcicoles. Calcifuges are plants occurring on acidic soils with low Ca^{2+} . The term calcicole is used to describe plants, which can grow on calcareous/ Ca^{2+} rich soils. These soils are generally Ca^{2+} and base-rich soils. However, plants vary in their Ca^{2+} requirements and the ability to extract Ca^{2+}

from complex soil environments. In particular, monocots require less Ca^{2+} for optimal growth than do the dicots.¹² Ca^{2+} enters plant cells through Ca^{2+} -permeable ion channels in their plasma membranes. Ca^{2+} competes with other cations both for these sites and for the uptake from the soil. The presence of high levels of Ca^{2+} is known to ameliorate the effects of the uptake of toxic cations (Al^{3+} and Na^+) from the soil while the presence of high levels of other cations (K^+ , Mg^{2+}) may reduce Ca^{2+} uptake. Calcium uptake and requirements for optimal growth are thus strongly dependent on the presence of other cations.⁹

Calcium deficiency is rare in nature, but may occur on soils with low base saturation and/or high levels of acidic deposition. Calcium deficiency may occur because of competition by other cations, or because of low transpiration, where the xylem flow is inadequate to supply the calcium requirements of rapidly growing tissue. Calcium deficiency results in stunted root growth and altered leaf appearances.¹¹ More severe symptoms generally result from the failure of cell membrane integrity and include bitter-pit in apple fruit, blossom end rot in tomato and tip burn in lettuce. Deficiency symptoms develop first in the regions of plant most distal to the region of Ca^{2+} uptake.

Ca^{2+} TRANSPORTERS/ EFFLUX PUMPS IN THE CELLULAR MEMBRANES

Plant cells, like animal cells, maintain a low regulated free Ca^{2+} concentration ranging from 30 nM to 400 nM in higher plants. Maintaining the low cytosolic Ca^{2+} concentration observed in plant cells requires active transport of Ca^{2+} from the cytosol. Active efflux pumping is a prerequisite for the restoration of low levels cytosolic calcium after the signaling event. The removal of Ca^{2+} from the cytosol against its electrochemical gradient to either the apoplast or to intracellular organelles requires energized 'active' transport. Ca^{2+} -ATPases and $\text{H}^+/\text{Ca}^{2+}$ antiporters are the key proteins catalyzing this movement.

By removing Ca^{2+} from the cytosol several important functions are performed by these enzymes.¹³

(1) They maintain a low $[\text{Ca}^{2+}]_{\text{cyt}}$ in the resting or the unstimulated cell which is appropriate for the cytoplasmic metabolism.

(2) They restore $[\text{Ca}^{2+}]_{\text{cyt}}$ levels to the resting normal levels following a $[\text{Ca}^{2+}]_{\text{cyt}}$ perturbation, thereby influencing the magnitude kinetic and subcellular location of $[\text{Ca}^{2+}]_{\text{cyt}}$ signals.

(3) They replenish intracellular and extracellular Ca^{2+} stores for subsequent $[\text{Ca}^{2+}]_{\text{cyt}}$ signals and permit the generation of localized $[\text{Ca}^{2+}]_{\text{cyt}}$ oscillations through their interplay with Ca^{2+} channels.¹⁴

(4) They provide Ca^{2+} in the ER for the secretory system to function.

(5) They remove some divalent cations, such as Ni^{2+} , Zn^{2+} , Mg^{2+} and Mn^{2+} from the cytosol to prevent mineral toxicity.¹³

Hirschi (2001)¹³ suggested that the Ca^{2+} -ATPases, which have high affinity ($K_m = 1\text{--}10 \mu\text{M}$) but low capacity for Ca^{2+} transport, are responsible for maintaining $[\text{Ca}^{2+}]_{\text{cyt}}$ homeostasis in the resting cells. Whereas the $\text{H}^+/\text{Ca}^{2+}$ -antiporters, which have lower affinities ($K_m = 10\text{--}15 \mu\text{M}$) but high capacities for Ca^{2+} transport, are likely to remove Ca^{2+} from the cytosol during $[\text{Ca}^{2+}]_{\text{cyt}}$ signals and thereby modulate $[\text{Ca}^{2+}]_{\text{cyt}}$ perturbations. This hypothesis is supported by the fact that $\text{H}^+/\text{Ca}^{2+}$ antiporter, but not the vacuolar Ca^{2+} -ATPase, resets $[\text{Ca}^{2+}]_{\text{cyt}}$ in yeast following hypertonic shock.¹⁵

Plant Ca^{2+} -ATPases belongs to two major families (a) The P-type ATPase II A family and (b) The P-type ATPase II B family.¹⁶

(a) **The P-type ATPase II A family.** Nucleotide specificity of these pumps is broad (30-60% activity achieved with GTP and ITP). The pumps are inhibited by erythrosine B ($\text{IC}_{50} \leq 1 \mu\text{M}$) and estimate Ca^{2+} affinity is in the range of 0.4-12 μM . The first family (The P-type ATPase II A family) lacks N-terminal auto regulatory domain. Four members of this family have been identified in the *Arabidopsis* genome (termed *AtECAs 1 to 4* by Axelsen and Palmgren, (2001).¹⁶ They are likely to be present in the plasma membrane, tonoplast ER and the Golgi apparatus.

(b) **The P-type ATPase II B family.** The second family of plant Ca^{2+} -ATPases (the P-type ATPase IIB family) is characterized by an auto inhibitory N-terminal domain that contains a binding site for Ca-CaM and in addition a serine-residue phosphorylation site. Their catalytic activity can be modulated by $[\text{Ca}^{2+}]_{\text{cyt}}$ either through activation upon binding CaM or by inhibition following phosphorylation by Ca^{2+} -dependent protein kinases (CDPK).¹⁷ Since CaM binding sites are generally quite diverse, each type-II B Ca^{2+} -ATPase may have different affinity for CaM or may bind a different CaM isoform. Ten members of the type-II B, Ca^{2+} -ATPase family have been identified in the *Arabidopsis* genome (termed as *ACAS1, 2 and 4* and *ALACAs 7 to 13* by Axelsen and Palmgren, (2001).¹⁶ These Ca^{2+} -ATPases reside on various cellular membranes including the plasma membrane (*AtACA8*), the tonoplast (*AtACA4*), and the plastid inner membrane (*AtACA1*). The relative molecular mass of type II B Ca^{2+} -ATPase pumps has been estimated to be between 115,000 Da and 135,000 Da.¹⁸

The abundance of Ca^{2+} -ATPase isoforms suggest that individual isoforms are functionally distinct and may respond differentially to distinct cellular processes involving specific Ca^{2+} signals. They also imply a requirement for CaM-independent and CaM-dependent regulation of Ca^{2+} -ATPase activities in the modulation of $[\text{Ca}^{2+}]_{\text{cyt}}$ perturbations during cell signaling. The expression of many Ca^{2+} -ATPases is increased upon exposure to high salinity or high $[\text{Ca}^{2+}]_{\text{cyt}}$ and some Ca^{2+} -ATPase genes are expressed only under stress conditions.¹⁹ This may reflect a role in maintaining $[\text{Ca}^{2+}]_{\text{cyt}}$ homeostasis or in reducing Na^+ influx to the cytosol in saline environments.

$\text{Ca}^{2+}/\text{H}^+$ ANTIPORTERS

In addition to a P-type pump, the presence of a low affinity $\text{Ca}^{2+}/\text{H}^+$ antiport mechanism at the plasma membrane level had been suggested. The first plant $\text{H}^+/\text{Ca}^{2+}$ antiporter to be cloned and functionally expressed was *CAX1* (Calcium exchanger 1).²⁰ The gene was identified by its ability to restore growth on a high Ca^{2+} medium to a yeast mutant defective in vacuolar Ca^{2+} transport.

The $\text{H}^+/\text{Ca}^{2+}$ - antiporters present in the plasma membrane and tonoplast have been characterized biochemically.⁸ These have a lower affinity for Ca^{2+} than Ca^{2+} -ATPases and may also transport Mg^{2+} . The stoichiometry of the dominant $\text{H}^+/\text{Ca}^{2+}$ -antiporter in the tonoplast is apparently $3\text{H}^+/1\text{Ca}^{2+}$. Eleven genes encoding putative $\text{H}^+/\text{Ca}^{2+}$ antiporters (*AtCAX*) have been identified in the genome of *Arabidopsis thaliana*.¹³ The transporters *AtCAX1*, *AtCAX2* and *AtCAX4* are located at the tonoplast.¹³ The *AtCAX1* antiporter exhibits both a high affinity and high specificity for Ca^{2+} . By contrast, the *AtCAX2* transporter is a high-affinity, high capacity $\text{H}^+/\text{heavy metal cation}$ antiporter. The *AtCAXs* have homologues in other plant species and their physiological roles have been investigated using transgenic plants.¹³ Transgenic tobacco overexpressing

AtCAX1 exhibits Ca^{2+} -deficiency disorders, which includes tip burn, metal-hypersensitivity and susceptibility to chilling that can be reversed by increasing Ca^{2+} supply. By raising Ca^{2+} supply the expression of *AtCAX1* and *AtCAX3* (but not *AtCAX2* or *AtCAX4*) genes was increased.¹³

$\text{Ca}^{2+}/\text{H}^+$ antiporters, utilize the H^+ gradient generated by the tonoplast V-type H^+ -pump and by a proton-pumping pyrophosphatase to sequester Ca^{2+} in the Vacuole.²¹ In many plant cells the vacuole occupy more than 50% of the cell volume, and it is evident that trans tonoplast Ca^{2+} transport makes a very significant contribution to the regulation of cytosolic Ca^{2+} concentrations. Ca^{2+} concentrations within the vacuole range from 0.1 to 10 mM.²²

Ca^{2+} -ATPases are estimated to represent only <0.1% of the membrane protein and are thus 30–100 fold less abundant than H^+ -ATPases in the PM (3%) and the endomembranes (5–10%). All Ca^{2+} pumps are inhibited by orthovanadate. $\text{H}^+/\text{Ca}^{2+}$ antiporters are efflux transporter and are different from Ca^{2+} -ATPases in that they do not require ATP and they are not sensitive to vanadate.

In contrast Ca^{2+} influx to the cytosol is mediated via Ca^{2+} channels. The principal roles of Ca^{2+} permeable channels in the plasma membrane appear to be in all signaling. Ca^{2+} permeable channels have been found in all plant membranes. They have been classified on the basis of their voltage-dependence into depolarization activated (DACC), hyperpolarization-activated (HACC) and voltage independent cation channels (VICC).⁸

THE Ca^{2+} SIGNATURES

The cytosolic Ca^{2+} in plant cells increases in response to various environmental challenges like abiotic and biotic stresses and developmental cues. This transient increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ is considered critical for the production of a physiological response. Elevation of $[\text{Ca}^{2+}]_{\text{cyt}}$ is considered a universal response to stress. The perturbations in cytosolic calcium levels (termed as $[\text{Ca}^{2+}]_{\text{cyt}}$ “signature”) elicited by each environmental challenge and developmental cue is unique and results in an appropriate physiological response to a particular stimulus. The uniqueness is manifested in the sub-cellular location and/or the kinetics of magnitude of the $[\text{Ca}^{2+}]_{\text{cyt}}$ perturbation.²³ An increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ is effected by Ca^{2+} influx to the cytosol either from the apoplast, across the plasma membrane, or from the intracellular organelles. The Ca^{2+} influx is mediated by Ca^{2+} permeable ion channels, and their type, cellular localization and the abundance influences the spatial characteristics of $[\text{Ca}^{2+}]_{\text{cyt}}$ perturbations. Since the diffusion of Ca^{2+} within the cytoplasm is low, and the buffering of Ca^{2+} in the cytoplasm is high (0.1 to 1 mM)²⁴ the opening of Ca^{2+} channel produces a local increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ that dissipates rapidly after the channel has been closed. The subcellular localization of Ca^{2+} channel is therefore critical for the targeting of different cellular processes. Cytosolic calcium “waves” are produced within the cytoplasm by the successive recruitment of particular Ca^{2+} channels to coordinate cellular responses. It was suggested that a local elevation of $[\text{Ca}^{2+}]_{\text{cyt}}$ might generate soluble second messengers, such as IP_3 or cADPR, that diffuse through the cytoplasm to activate a relay of spatially separated Ca^{2+} channels.²⁴ This theory was supported in the plant cells responding to salt stress.²⁵

Franklin-Tong et al. (2002)²⁶ suggested that repetitive Ca^{2+} influx across the plasma membrane contributed in the $[\text{Ca}^{2+}]_{\text{cyt}}$ that occur following the application of ABA to guard cells. This transient increase was first close to the plasma membrane and subsequently adjacent to the vacuole.²⁷ These waves are thought to reflect the

sequential opening of hyperpolarization-activated Ca^{2+} channels at the plasma membrane and then second-messenger activated Ca^{2+} channels in the tonoplast.²⁸

In addition to these sub cellular waves, “waves” of cells with high $[\text{Ca}^{2+}]_{\text{cyt}}$ may also propagate through the plant tissue. This can be induced in root tissues by mechanical stimulation²⁹ or saline shock³⁰, in cotyledons by cold shock³¹ and in leaves by chilling plant roots briefly.³² Electrical action potentials, osmotic perturbations or chemical signals may trigger these waves. Although an elevated $[\text{Ca}^{2+}]_{\text{cyt}}$ is necessary for signal transduction, a prolonged increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ is lethal. Sustained high $[\text{Ca}^{2+}]_{\text{cyt}}$ is implicated in apoptosis, both during the normal development and in hypersensitive responses to pathogens.³³

To effect adaptive responses, the $[\text{Ca}^{2+}]_{\text{cyt}}$ perturbations must be either of low amplitude or transient. Transient increases in $[\text{Ca}^{2+}]_{\text{cyt}}$ can be single (spike), double (biphasic) or multiple (oscillations). The perturbations generated may differ in their cellular location, role and extent of propagation and on their amplitude during propagation. Calcium signatures may also be tissue specific. For example, within the root, the $[\text{Ca}^{2+}]_{\text{cyt}}$ perturbations induced by mechanical perturbation, salinity, osmotic stress, cold shock or slow cooling differ markedly between cell types.³⁰

Several abiotic challenges result in an immediate, transient increase in $[\text{Ca}^{2+}]_{\text{cyt}}$ that is restored to basal levels within minutes. Such changes include mechanical perturbations and rapid cooling for brief periods, termed as ‘cold shock’. The duration, periodicity and amplitude of oscillations vary considerably, and their form is often dependent on the strength and combination of specific stimuli.²⁷

CALCIUM “MEMORY”

The term “memory” was put forward first by Knight et al. (1996).³⁴ There is considerable evidence that $[\text{Ca}^{2+}]_{\text{cyt}}$ signatures are modified by previous experience. A diminished $[\text{Ca}^{2+}]_{\text{cyt}}$ elevation upon repetitive stimulation by the same environmental challenge or a developmental cue is a common observation. Many examples support this. The magnitude of the $[\text{Ca}^{2+}]_{\text{cyt}}$ perturbation elicited by the wind-induced motion becomes progressively smaller upon repeated stimulation and a refractory period of several minutes is required before a full response is observed again. A second exposure to an elicitor does not influence $[\text{Ca}^{2+}]_{\text{cyt}}$ for several hours after its initial application.³⁵ For example, plant cells challenged with H_2O_2 fail to respond to H_2O_2 for several hours.³⁶ There is also evidence that the $[\text{Ca}^{2+}]_{\text{cyt}}$ signatures elicited by one environmental challenge can be modified by prior exposure to a contrasting one. For example, the magnitude of the $[\text{Ca}^{2+}]_{\text{cyt}}$ perturbations in response to oxidative stress was reduced by prior exposure to hyperosmotic stress and vice-versa was also found to be true. These observations also imply a cross talk between the signaling cascades. The attenuated response of $[\text{Ca}^{2+}]_{\text{cyt}}$ after repeated stimulation by various elicitors forms a part of cellular memory and the cells are able to retain the previous information. This ‘memory’ is significant and helps the cells to respond better to a particular stress without disturbing the delicate balance of Ca^{2+} levels and maintaining cellular $[\text{Ca}^{2+}]$ homeostasis. It is noteworthy that *Nicotiana plumbaginifolia*, which could not retain ‘cold memory’ during acclimation of the plant by pretreatment with nonfreezing cold temperature was cold-sensitive whereas *Arabidopsis* which could retain ‘cold memory’ was more resistant to cold stress.³⁴




Figure 3. (A) Helix-loop-helix pattern of an EF hand. The EF hand is a highly conserved 29 amino-acid motif consisting of a α helix E (residue 1-10), a loop (residue 10-21), which binds the Ca^{2+} ion and a second α -helix F (residue 19-29). (B) The canonical EF hand consensus sequence of the 12 amino acid Ca^{2+} binding loop. The loop consists of 12 residues with the pattern X (D), Y(D), Z (D), -Y (K), -X (D), -Z (E) which participate in binding Ca^{2+} and the intervening residues are represented by an asterisks (*).

CALCIUM BINDING PROTEINS

Transient increase in the cytoplasmic Ca^{2+} in response to signals is sensed by an array of Ca^{2+} sensors. Ca^{2+} sensors are small proteins that bind Ca^{2+} and change their conformation in a Ca^{2+} dependent manner. Specificity in the signaling pathway is provided by the uniqueness in calcium signatures and also by plethora of Ca^{2+} sensors, which can decode the Ca^{2+} perturbations quite precisely. Once Ca^{2+} sensors decode the elevated $[\text{Ca}^{2+}]_{\text{cyt}}$, Ca^{2+} efflux into the cell exterior and/or the sequestration into cellular organelles such as vacuoles, ER and mitochondria restores its levels to resting state. The properties of binding proteins are described below:

(a) Properties essential for Ca^{2+} -binding proteins (CaBPs) to function as intracellular- Ca^{2+} receptor.

(1) The Ca^{2+} receptor/sensor must have Ca^{2+} binding sites essentially unoccupied in the resting cell (free Ca^{2+} is 10^{-7}M) and occupied at levels, which are reached upon stimulation ($\sim 10^{-7}\text{M}$ – 10^{-6}M). This means that the affinity constants (K_a) for Ca^{2+} should be $\sim 10^{-6}\text{M}^{-1}$.

(2) The protein must show selectivity and preference for Ca^{2+} in the presence of other cations like Mg^{2+} and K^+ .

(3) After binding to Ca^{2+} , a Ca^{2+} sensor must undergo a conformational change that either alters its interaction with other molecules or changes its activity if it is an enzyme. Formation of Ca^{2+} protein complex is reversible.

(4) The kinetics of interaction should be very fast so as to correspond with the short lived Ca^{2+} signature.

(b) **EF Hand, Structure and Function.** Most of the Ca^{2+} sensors bind Ca^{2+} using a helix-loop-helix motif termed as the 'EF hand' or the elongation factor, which binds a single Ca^{2+} molecule with high affinity. The Ca^{2+} sensors utilize the side chain oxygen atoms of the EF hand motif for Ca^{2+} coordination. In 1973 Kretsinger and Nockolds³⁷ first discovered the EF hand structural motif in the crystal structure of parvalbumin. The properties, structure and function of EF hands are described below.

(1) The EF hand motifs mostly exist in pairs, which helps in the stabilization of the protein structure. Exceptions to this norm include proteins such as, parvalbumin and *Plasmodium falciparum* membrane surface protein. Frequently, EF hand pairs interact through antiparallel β -sheets, which allow cooperation in Ca^{2+} binding.

(2) The EF hand is a highly conserved 29 amino-acid motif consisting of a α helix E (residue 1-10), a loop (residue 10-21), which binds the Ca^{2+} ion and a second α -helix F (residue 19-29) (Moews and Kretsinger, 1975). The loop consists of 12 residues with the

pattern X (D), Y(D), Z (D), -Y (K), -X (D), -Z (E) which participate in binding Ca^{2+} and the intervening residues are represented by an asterisks (*). A representation of the EF hand and loop is shown in (Fig. 3A and B), respectively.

(3) The Ca^{2+} ion is coordinated by an oxygen atom or by a bridging water molecule (-x) of the side chains of residues 10 (X), 12 (Y), 14 (Z) and 18 (-X). The ligand at vertex (-Y) is the carbonyl oxygen at residue 16.³⁸ Position 21 (-Z) is usually Glutamate and is the sixth residue to coordinate Ca^{2+} .

(4) In most of the functional EF hand motifs, the first amino acid is Aspartate (Asp) and the twelfth is Glutamate (Glu), Glu contributes both its side chain oxygen atoms to the metal ion coordination. Since there are 7 oxygen ligands, the Ca^{2+} coordination exhibits pentagonal bipyramidal symmetry, whose axis is 10(X)–18(-X). Glycine (Gly), present at the apex of the loop is an invariant residue and allows the loop to take a sharp bend. Each loop is flanked on either side by 8-10 residues of alternating hydrophilic and hydrophobic character. This arrangement favors the formation of the outer and inner surface of an alpha helical cylinder respectively.

(5) Most of the EF hand proteins are characterized by the relatively high percentage of acidic residues (Troponin C, 29%, CaM, 25%, intestinal Ca^{2+} binding protein 23%, and parvalbumin, 18%). Several isoforms of an EF hand protein may exist in a single organism. The Ca^{2+} binding affinities of the EF hand protein vary substantially ($K_d = 10^{-4}$ – 10^{-9}M) and depend on the amino acid sequence of the protein, especially with regards to the 12 residue consensus loop that provide all the acids that directly ligate to Ca^{2+} ions. When Ca^{2+} binds to $[\text{Ca}^{2+}]_{\text{cyt}}$ sensors their structural and/or enzymatic properties change and their subsequent interactions with target proteins can alter enzymatic activities, cytoskeletal orientation, protein phosphorylation cascades and gene expression. It is believed that these changes result in stress tolerance or a developmental switch.

(c) **Important calcium sensors in plants.** The major family of Ca^{2+} sensors includes; (1) Calmodulin (CaMs); (2) Calmodulin like proteins; (3) Calcium dependent protein kinases (CDPKs); (4) Calcineurin B-like proteins.

(1) **Calmodulin: CaM.** Calmodulin (CaM) is a small (17 KDa), highly conserved, acidic protein with two globular domains each containing two EF hands connected by a flexible α -helical linker.³⁹ CaM is found in the apoplast and in the cytosol, ER and the nucleus of plant cells. Within the cytosol, the estimated CaM concentration is 5 to 40 μM .⁴⁰ Role of CaM has been implicated in many physiological processes like light, gravity, mechanical stress, phytohormones, pathogens, osmotic stress, heat shock and chilling.^{23,29,40}

The structure of CaM was first solved by Babu et al., 1985, and revealed that all 4 EF hands are saturated by Ca^{2+} ions (4 Ca^{2+}). CaM appears to be regulatory protein and induces large changes in inter-helical angles as Ca^{2+} is bound. The affinity of CaM for Ca^{2+} is influenced by the presence of particular target proteins.⁴⁰ CaM can also regulate gene expression by binding to specific transcription factors.⁴¹

(2) **CaM-like proteins.** Plants also possess CaM-like proteins, which differ from the CaM in containing more than 148 amino-acid residues and have between one to six EF hand motifs. They possess limited homology to CaM (75% identity with canonical CaM isoforms.³⁹ In *Arabidopsis*, they include: CaBP-22, TCH2 and TCH3, AtCP1, NADPH oxidases, and Ca^{2+} binding protein phosphatases such as ABI 1 and ABI 2. These proteins have been implicated in cellular responses to diverse environmental, developmental and pathological challenges.

(3) **Ca²⁺-dependent protein kinases.** 5 types of Ca²⁺ regulated protein kinases have been reported in plants: (i) Ca²⁺ dependent and CaM independent protein kinases (CDPKs); (ii) CDPK-related protein kinases (CRKs); (iii) CaM dependent protein kinases (CaMKs); (iv) Ca²⁺/CaM-dependent protein kinases (CCaMK); (v) SOS3/CBL interacting protein kinases (SIPKs/CIPKs).

(i) **Ca²⁺-dependent and CaM-independent protein kinases CDPKs.** The CDPKs are ubiquitous in plants. There are at least 34 genes encoding CDPKs in the *Arabidopsis* genome⁴² and similar numbers in other plant species. Apart from plants, CDPKs are also found in protozoans and algae. They generally have four EF hands at their C-terminus that bind Ca²⁺ and activate the serine/threonine kinase activity of the enzyme. These kinases require micromolar concentrations of Ca²⁺ for their activity and have no requirement of CaM or lipids. They have a unique structure as N-terminal protein kinase domain is fused with C-terminal auto-regulatory domain and a CaM like domain, which has Ca²⁺ binding EF hand or helix-loop-helix motif. The autoinhibitory domain of CDPKs is a 30 amino acid sequence, which acts as a pseudo-substrate.⁴³ The N-terminal domain of CDPKs is variable and provides specificity to different CDPK isoforms. These enzymes show several fold stimulation with Ca²⁺ and show autophosphorylation. The binding of Ca²⁺ to some of CDPKs, is modulated by lipids or phosphorylation.⁴² Ca²⁺ binding to CDPK effects conformation of the kinase and relieve the inhibition caused by the autoinhibitory region.

CDPKs are implicated in pollen development, control of cell cycle, phytohormone signaling, light-regulated gene expression, gravitropism, thigmotropism, cold acclimation, salinity tolerance, drought tolerance and responses to pathogens.⁴⁴

(ii) **CDPK-related protein kinases (CRKs).** CRKs are similar to CDPKs except that the CaM-like region is poorly conserved with degenerate or truncated EF hands that may not be able to bind Ca²⁺. There are at least seven CRKs in *Arabidopsis* genome, and orthologues of these are present in many plant species. However, the regulation and function of these kinases are not known.⁴⁵

(iii) **CaM kinases (CaMKs).** Several CaMKs have been cloned from *Arabidopsis* and other plants.⁴⁶ Their kinase activity is stimulated by CaM dependent autophosphorylation and their catalytic activity is also modulated by CaM. They are highly expressed in rapidly growing cells and tissues of the root and flower.⁴⁶

(iv) **Ca²⁺/CaM-dependent protein kinases (CCaMKs).** These are a group of Ca²⁺-dependent kinases, which in addition to Ca²⁺ also requires CaM for their activity. Thus CaM besides acting directly could also exert its effect by binding to protein kinases and modulating their activity. A Ca²⁺/CaM-dependent protein kinase (CCaMK) was characterized from lily and other plant species.⁴⁷ Sequence analysis revealed the presence of an N-terminal catalytic domain, a centrally located CaM-binding domain and a C-terminal visinin-like domain containing only three EF hands. Biochemical studies of CCaMK established that Ca²⁺ and CaM stimulates CCaMK activity. In the absence of CaM, Ca²⁺ promotes autophosphorylation of CCaMK. The phosphorylated form of CCaMK possesses more kinase activity than the non-phosphorylated form.

(v) **SOS3/CBL interacting protein kinases (SIPKs/CIPKs).** Calcineurin B-like proteins were found to interact specifically with a class of serine-threonine protein kinases known as CBL interacting protein kinases (CIPKs).^{1, 2} Recently, a novel CIPK from pea has been reported and found to interact and phosphorylate the pea.⁴⁸

(4) **Calcineurin B-like proteins.** Calcineurin B-like proteins (CBLs) are relatively a new class of calcium sensors discovered in

Arabidopsis originally, in search for the genes imparting salt tolerance and maintaining cellular ion homeostasis.^{49, 50} Molecular analysis of the sos mutants opened a new chapter in relation to salt stress signalling that led to the discovery of a pathway that transduce a salt stress induced Ca²⁺ signal to reinstate cellular ion homeostasis. Currently 10 CBL and 25 CBL-interacting protein kinases (CIPKs) have been reported from *Arabidopsis*.² The CBL-CIPK network is also widely distributed among higher plants but except *Arabidopsis*, the complexity and characterization of this pathway remains largely unrevealed. As different plants vary in their genome complexity, phenotype and physiology therefore, specie specific function or functional diversification can be expected. The essential role imparted by CBL-CIPK genes in stress tolerance, necessitates their detailed characterization from higher plants. In fact, there has been no report on experimental characterization of CBL from any higher plant except *Arabidopsis*. *AtCBL3*, in particular has been largely overlooked even in *Arabidopsis*. Recently, we (Mahajan et al., 2006)⁴⁸ have reported the cloning and characterization of a novel CIPK and its interacting partner CBL from pea. Pea CIPK showed autophosphorylation and could phosphorylate pea CBL. Both pea CBL and CIPK were found to be coordinately upregulated in response to various stresses such as cold, salinity but not to dehydration stress.⁴⁸

(d) **Ca²⁺-binding proteins without EF hands.** There are several proteins that bind Ca²⁺ but do not contain EF hand motifs. These include the phospholipase D (PLD), annexins, calreticulin and Pistil-expressed Ca²⁺ binding protein (PCP).

Phospholipase D. The activity of PLD, which cleaves membrane phospholipids into a soluble head group and PA, is regulated by [Ca²⁺]_{cyt} through a Ca²⁺/phospholipids binding-site termed as the 'C2 domain'.⁵¹ PLD activity is implicated in cellular responses to ethylene and ABA, α amylase synthesis in aleurone cells, stomatal closure, pathogen responses, leaf senescence and drought tolerance.⁵¹ Plants possess several PLD isoforms that differ in their affinity for Ca²⁺ and their modulation by phosphoinositides, free fatty acids and lysolipids.⁵¹ These biochemical modulators of PLD activity are the substrates or products of PLC, which generates IP₃, DAG, phospholipase A₂ and DAG-kinase, both of which are regulated by CaM. It is suggested that [Ca²⁺]_{cyt} signaling cascades might coordinate the activities of these diverse enzymes to effect specific responses to the environmental stimuli.⁵²

Annexins. These are a family of proteins in plants and animals that bind phospholipids in a Ca²⁺ dependent manner and contain four to eight repeats of approximately 70 amino acids.⁵³ Although exact function of annexins is not known, plant annexins are implicated in secretory processes and some have ATPase, peroxidase activities.

Calreticulin. It is a Ca²⁺ sequestering protein in the ER and functions as a molecular chaperone.⁵⁴ The function of calreticulin is also implicated in Ca²⁺ homeostasis.⁵⁵

Pistil expressed Ca²⁺-Binding Protein (PCP). A 19 kDa novel Ca²⁺ binding protein (PCP) expressed in anthers and pistil was isolated.⁵⁶ PCP is high capacity (binds 20 mol of Ca²⁺ per mol of PCP), low affinity Ca²⁺ binding protein. Role of PCP has been implicated in pollen-pistil interactions and/or pollen development.

References

- Mahajan S, Tuteja N. Cold, salinity and drought stresses: an overview. *Arch Biochem Biophys* 2005; 444:139-58.
- Mahajan S, Sopoy SK, Tuteja N. CBL-CIPK paradigm: role in calcium and stress signaling in plants. *Proc Indian Nat Sci Acad* 2006; 72:63-78.
- Xiong TC, Bourque S, Lecourieux D, Amelot N, Grat S, Briere C, Mazars C, Pugin A, Ranjeva R. Calcium signaling in plant cell organelles delimited by a double membrane. *Biochim Biophys Acta*. 2006; 1763:1209-15.
- Kaplan B, Davydov O, Knight H, Galon Y, Knight MR, Fluhr R, Fromm H. Rapid transcriptome changes induced by cytosolic Ca²⁺ transients reveal ABRE-related sequences as Ca²⁺-responsive cis elements in *Arabidopsis*. *Plant Cell*. 2006; 18:2733-48.
- Harada A, Shimazaki KI. Phototropins and blue light-dependent calcium signaling in higher plants. *Photochem Photobiol* 2006; Mar 1; [Epub ahead of print].
- Martinez-Noel G, Tognetti J, Nagaraj V, Wiemken A, Pontis H. Calcium is essential for fructan synthesis induction mediated by sucrose in wheat. *Planta*. 2006; 225:183-91.
- Muller JF, Hulster AA, Papke OC, Ball MC, Marschner H. Transfer of PCDD/PCDF from contaminated soils into carrots, lettuce and peas. *Chemosphere*. 1994; 29:2175-81.
- Sanders D, Pelloux J, Brownlee C, Harper JF. Calcium at the crossroads of signaling. *Plant Cell* 2002; 14:S401-17.
- Burström HG. Calcium and plant growth. *Biol Rev* 1968; 43:287-316.
- Bush DS. Calcium regulation in plant cells and its role in signaling. *Ann Rev Plant Physiol Plant Mol Biol* 1995; 46:95-122.
- Marschner H. *Mineral Nutrition of Higher Plants*. Academic Press, London, 1986; pp. 252-254.
- Loneragan JF, Snowball K. Calcium requirements of plants. *Aust J Agric Res* 1969; 20:465-78.
- Hirschi K. Vacuolar H⁺/Ca²⁺ transport: who's directing the traffic? *Trends Plant Sci* 2001; 6:100-4.
- Harper JF. Dissecting calcium oscillators in plant cells. *Trends Plant Sci* 2001; 6:395-7.
- Denis V, Cyert MS. Internal Ca²⁺ release in yeast is triggered by hypertonic shock and mediated by a TRP channel homologue. *J Cell Biol* 2002; 156:29-34.
- Axelsen KB, Palmgren MG. Inventory of the superfamily of P-type ion pumps in *Arabidopsis*. *Plant Physiol* 2001; 126:696-706.
- Hwang I, Sze H, Harper JF. A calcium-dependent protein kinase can inhibit a calmodulin-stimulated Ca²⁺ pump (ACA2) located in the endoplasmic reticulum of *Arabidopsis*. *Proc Natl Acad Sci USA*. 2000; 97:6224-9.
- Rasi-Caldogno F, Carnelli A, DeMicheli MI. Identification of the plasma membrane Ca²⁺ ATPase and of its autoinhibitory domain. *Plant Physiol* 1995; 108:105-13.
- Garcia-Abblas B, Benito B, Rodríguez-Navarro A. Plant cells express several stress calcium ATPases but apparently no sodium ATPase. *Plant Soil* 2001; 235:81-192.
- Hirschi KD, Zhen RG, Cunningham KW, Rea PA, Fink GR. CAX1, an H⁺/Ca²⁺ antiporter from *Arabidopsis*. *Proc Natl Acad Sci USA*, 1996; 93:8782-6.
- Chanson A. Characterization of the tonoplast Ca²⁺/H⁺ antiport system from maize roots. *Plant Physiol Biochem* 1994; 32:341-6.
- Macklon AES. Calcium fluxes at plasmalemma and tonoplast. *Plant Cell Environ* 1984; 7:407-13.
- Rudd JJ, Franklin-Tong VE. Unravelling response-specificity in Ca²⁺ signalling pathways in plant cells. *New Phytol* 2001; 151:7-33.
- Trewavas A. Le calcium, c'est la vie: calcium makes waves. *Plant Physiol* 1999; 120:1-6.
- Drøbak BK, Watkins PAC. Inositol (1,4,5) trisphosphate production in plant cells: an early response to salinity and hyperosmotic stress. *FEBS Lett* 2000; 481:240-4.
- Franklin-Tong VE, Holdaway-Clarke TL, Straatman KR, Kunkel JG, Hepler PK. Involvement of extracellular calcium influx in the self-incompatibility response of *Papaver rhoeas*. *Plant J* 2002; 29:333-45.
- Allen GJ, Kwak JM, Chu SP, Llopis J, Tsien RY, Harper JF, Schroeder JI. Cameleon calcium indicator reports cytoplasmic calcium dynamics in *Arabidopsis* guard cells. *Plant J* 1999; 19:735-47.
- Schroeder JI, Allen GJ, Hugouvieux V, Kwak JM, Waner D. Guard cell signal transduction. *Ann Rev Plant Physiol Plant Mol Biol* 2001; 52:627-58.
- Fasano JM, Massa GD, Gilroy S. Ionic signaling in plant responses to gravity and touch. *J Plant Growth Regul* 2002; 21:71-88.
- Moore CA, Bowen HC, Scrase-Field S, Knight MR, White PJ. The deposition of suberin lamellae determines the magnitude of cytosolic Ca²⁺ elevations in root endodermal cells subjected to cooling. *Plant J* 2002; 30:457-66.
- Knight MR, Read ND, Campbell AK, Trewavas AJ. Imaging calcium dynamics in living plants using semisynthetic recombinant aequorins. *J Cell Biol* 1993; 12:83-90.
- Campbell AK, Trewavas AJ, Knight MR. Calcium imaging shows differential sensitivity to cooling and communication in luminous transgenic plants. *Cell Calcium* 1996; 19:211-8.
- Levine A, Pennell RI, Alvarez ME, Palmer R, Lamb C. Calcium mediated apoptosis in a plant hypersensitive disease resistance response. *Curr Biol* 1996; 6:427-37.
- Knight H, Trewavas AJ, Knight MR. Cold calcium signaling in *Arabidopsis* involves two cellular pools and a change in calcium signature after acclimation. *Plant Cell* 1996; 8:489-503.
- Blume B, Nürmberger T, Nass N, Scheel D. Receptor-mediated increase in cytoplasmic free calcium required for activation of pathogen defense in parsley. *Plant Cell* 2000; 12:1425-40.
- Price AH, Taylor A, Ripley SJ, Griffiths A, Trewavas AJ, Knight MR. Oxidative signals in tobacco increase cytosolic calcium. *Plant Cell* 1994; 6:1301-10.
- Kretsinger RH, Nockolds CE. Carp muscle calcium-binding protein. II. Structure determination and general description. *J Biol Chem* 1973; 248:3313-26.
- Strynadka NCJ, James MNG. Crystal structures of the helix-loop-helix calcium-binding proteins. *Ann Rev Biochem* 1989; 58:951-98.
- Luan S, Kudla J, Rodríguez-Concepción M, Yalovsky S, Grissem W. Calmodulins and calcineurin B-like proteins: calcium sensors for specific signal response coupling in plants. *Plant Cell* 2002; 14:S389-400.
- Zielinski RE. Calmodulin and calmodulin-binding proteins in plants. *Ann Rev Plant Physiol Plant Mol Biol* 1998; 49:697-725.
- Bouché N, Scharlat A, Snedden W, Bouchez D, Fromm H. A novel family of calmodulin-binding transcription activators in multicellular organisms. *J Biol Chem* 2002; 277: 21851-61.
- Cheng SH, Willmann MR, Chen HC, Sheen J. Calcium signaling through protein kinases. The *Arabidopsis* calcium-dependent protein kinase gene family. *Plant Physiol* 2002; 129:469-85.
- Harper JF, Huang JF, Lloyd SJ. Genetic identification of an auto inhibitor in CDPK, a protein kinase with a calmodulin like domain. *Biochemistry* 1994; 33:7278-87.
- Xiong L, Schumaker KS, Zhu JK. Cell signaling during cold, drought, and salt stress. *Plant Cell* 2002; 14:S165-83.
- Harmon AC, Gribskov M, Gubrium E, Harper JF. The CDPK superfamily of protein kinases. *New Phytol* 2001; 151:175-83.
- Zhang L, Lu YT. Calmodulin-binding protein kinases in plants. *Trends Plant Sci* 2003; 8:123-7.
- Lu YT, Hidaka H, Feldman LJ. Characterization of a calcium/calmodulin protein kinase homologue from maize roots showing light-regulated gravitropism. *Planta* 1996; 199:18-24.
- Mahajan S, Sopoy SK, Tuteja N. Cloning and characterization of CBL-CIPK signaling components from a legume (*Pisum sativum*) FEBS J. 2006; 273:907-25.
- Liu J, Zhu JK. A calcium sensor homolog required for plant salt tolerance. *Science* 1998; 280:1943-5.
- Kudla J, Xu Q, Harter K, Grissem W, Luan S. Genes for calcineurin B-like proteins in *Arabidopsis* are differentially regulated by stress signals. *Proc Natl Acad Sci USA* 1999; 96:4718-23.
- Wang X. Plant phospholipases. *Ann Rev Plant Physiol Plant Mol Biol* 2001; 52:211-31.
- Ritchie SM, Swanson SJ, Gilroy S. From common signalling components to cell specific responses: insights from the cereal aleurone. *Physiol Plant* 2002; 115:342-51.
- Clark GB, Roux SJ. Annexins of plant cells. *Plant Physiol* 1995; 1133-9.
- Baluška F, Šamaj J, Napier R, Volkmann D. Maize calreticulin localizes preferentially to plasmodesmata in root apex. *Plant J* 1999; 19: 481-8.
- Michalak M, Mariani P, Opas M. Calreticulin, a multifunctional Ca²⁺ binding chaperone of the endoplasmic reticulum. *Biochem Cell Biol* 1998; 76:779-85.
- Furuyama T, Dzelzkalns VA. A novel calcium-binding protein is expressed in Brassica pistils and anthers late in flower development. *Plant Mol Biol* 1999; 39:729-37.