



Published in final edited form as:

AIDS. 2008 October 18; 22(16): 2069–2074. doi:10.1097/QAD.0b013e32830edd44.

A Call for Randomized Controlled Trials of Antiretroviral Therapy for HIV-2 Infection in West Africa

Geoffrey S. Gottlieb¹, Serge-Paul Eholié², John N. Nkengasong³, Sabelle Jallow^{4,6}, Sarah Rowland-Jones⁴, Hilton C. Whittle⁴, and Papa Salif Sow⁵

1 Department of Medicine-Allergy & Infectious Diseases, University of Washington, Seattle, Washington, USA;

2 Service des Maladies Infectieuses et Tropicales, Centre Hospitalier Universitaire de Treichville, Abidjan, Côte d'Ivoire

3 Centers for Disease Control and Prevention, Atlanta, Georgia, USA

4 Medical Research Council Laboratories, Fajara, The Gambia

5 Clinique des Maladies Infectieuses Ibrahima Diop Mar - CHU de Fann, Université Cheikh Anta Diop de Dakar, Senegal

6 Department of Microbiology, Institute of Tropical Medicine, Antwerp, Belgium

Keywords

HIV-2; Africa; Resource-Limited; Controlled Trial; ARV

HIV-2 is endemic in West Africa, but unlike HIV-1 has had limited spread to other locales [1]. HIV-1 has emerged in West Africa more recently and because HIV-1 and HIV-2 co-circulate in this region, some individuals become dually infected with both viruses [2,3]. Compared to HIV-1 infection, HIV-2 is characterized by a much longer asymptomatic stage, lower plasma viral loads, slower decline in CD4 count, decreased mortality rate due to AIDS, lower rates of mother to child transmission, genital tract shedding, and sexual transmission [1,4–12]. Nonetheless, a significant proportion of HIV-2 infected individuals eventually progress to AIDS [6,13–15]. The clinical consequences of dual infection with both HIV-1 and HIV-2 need to be more fully understood, but current data suggest the majority of such patients also eventually progress to AIDS [2,16–20].

Antiretroviral therapy (ART) is becoming increasingly available in West Africa where HIV-2 infects up to 1–2 million people [21]. As ART “scale-up” programs proliferate in West Africa, significant numbers of HIV-2 and dually infected individuals will have access to and will be treated with antiretrovirals (ARV) developed against HIV-1 [22–25]. However, HIV-2 is intrinsically resistant to the non-nucleoside reverse transcriptase inhibitors (NNRTI) and T-20 (enfuvirtide) and reports suggest that HIV-2 may be partially resistant to some protease inhibitors (PI) (e.g. Amprenavir, atazanavir and nelfinavir) and have a low genetic barrier to nucleoside reverse transcriptase inhibitors (NRTI) resistance [26–39]. In addition, at least one recent report from Burkina Faso suggested the NRTI mutations M184V and Q151M may rarely

*Corresponding author: Geoffrey S. Gottlieb, MD PhD, Division of Allergy & Infectious Diseases, Department of Medicine, University of Washington, Seattle, WA 98195, Tel: 206 732-6150, Fax: 206 732-6167, Mailstop: 358070, Email: gottlieb@u.washington.edu.

Author Contributions: Geoffrey S. Gottlieb and Papa Salif Sow conceived of the opinion piece. All authors contributed to writing the paper.

be found in ARV-naïve individuals [40]. The presence of HIV-2 in dually infected patients complicates treatment, requiring drugs that are active against both HIV-1 and HIV-2.

Several observational cohort studies in developed countries have shown variable but generally poor outcomes of ART for HIV-2 infection [15,41–49] with similar results reported from the few small cohort studies from resource limited settings in Senegal, The Gambia and Cote d'Ivoire, West Africa [24,25,50].

Recently, there has been an effort in Europe to develop an international consortium (ACHIEV₂E (A Collaboration on HIV-2 Infection)) of HIV-2 cohorts, to better study HIV-2 infection and standardize its management, including HIV-2 viral load testing, and treatment [51]. These efforts should ultimately lead to better care of HIV-2 infected individuals in the developed world and West Africa.

To date there has not been a single randomized clinical trial of ART for HIV-2 infection. This is despite greater than 20 years since the discovery of HIV-2 [52] and advent of AZT, greater than 10 years since the landmark studies showed the benefit of HAART for HIV-1 [53–55], and currently 6 different classes of ARV available for HIV-1 (NRTI, NNRTI, PI, Fusion Inhibitors, CCR5-coreceptor blockers, and integrase inhibitors) of which 4 show *in vitro* and/or *in vivo* activity against HIV-2 [37,38,56–58]. The WHO currently recommends for first line therapy of HIV-2 infected patients, 2 NRTI and a protease inhibitor boosted by ritonavir [59]; they make no explicit recommendations for dually infected patients or for second line HIV-2 (or dual) infection regimens and little such data are available. In resource limited West Africa biological monitoring (CD4 counts) is limited and HIV-2 viral load and genotypic or phenotypic resistance testing are not commercially available, standardized nor routinely obtainable; thus assessing regimen failure is often made on clinical criteria and data suggest that CD4 count recovery with ART in HIV-2 infection is often poor [48,49]. Consequently, an assessment of the relative efficacy of potential first line and salvage regimens including the newer generation of protease inhibitors, integrase inhibitors and CCR5 co-receptor blockers among HIV-2 (and dually) infected patients is needed. A key issue in West Africa where HIV and TB are common is to find the ART regimens for HIV-2 that can be given with concurrent treatment for TB, given interactions between protease inhibitors and rifampicin [59]. Finally, albeit rare, proven regimens to prevent mother to child transmission of HIV-2 (or dual infection) are needed [60].

We believe the 1–2 million HIV-2 infected people and communities in West Africa most affected by HIV-2 (and dual infection) deserve the same evidence-based medicine and efforts that those with HIV-1 have justly demanded [61]. Rigorous, well designed randomized controlled trials, that demonstrate which ARV regimens are effective in treating HIV-2 infected people, are urgently needed to guide patients and the clinicians who care for them. The outcome of such trials will also be beneficial in treating patients dually infected with HIV-1 and HIV-2, although ultimately trials of ART in dually patients are also needed. However, the apparent decline of HIV-2 in some West African locales [3,61,62] highlights the importance of involving multiple West African countries to establish large multi-center collaborative studies and trials. We believe the best place to carry out such trials is in West Africa, the locale where HIV-2 is endemic and effective treatment is most likely to serve the local population. We, therefore call on the wider AIDS community to support efforts for evidenced based treatment of HIV-2 infection.

Acknowledgements

We thank the editors and anonymous reviewers for their helpful comments.

References

1. De Cock KM, Adjorlolo G, Ekpini E, Sibailly T, Kouadio J, Maran M, et al. Epidemiology and transmission of HIV-2. Why there is no HIV-2 pandemic [published erratum appears in JAMA 1994 Jan 19;271(3):196] [see comments]. *JAMA* 1993;3(270):2083–2086. [PubMed: 8147962]
2. Evans LA, Moreau J, Odehouri K, Seto D, Thomson-Honnebier G, Legg H, et al. Simultaneous isolation of HIV-1 and HIV-2 from an AIDS patient. *Lancet* 1988;2:1389–1391. [PubMed: 2904524]
3. Hamel DJ, Sankale JL, Eisen G, Meloni ST, Mullins C, Gueye-Ndiaye A, et al. Twenty years of prospective molecular epidemiology in Senegal: changes in HIV diversity. *AIDS Res Hum Retroviruses* 2007;23:1189–1196. [PubMed: 17961103]
4. Andreasson PA, Dias F, Naucier A, Andersson S, Biberfeld G. A prospective study of vertical transmission of HIV-2 in Bissau, Guinea-Bissau. *Aids* 1993;7:989–993. [PubMed: 8357558]
5. Simon F, Matheron S, Tamalet C, Loussert-Ajaka I, Bartczak S, Pepin JM, et al. Cellular and plasma viral load in patients infected with HIV-2. *Aids* 1993;7:1411–1417. [PubMed: 7904166]
6. Marlink R, Kanki P, Thior I, Travers K, Eisen G, Siby T, et al. Reduced rate of disease development after HIV-2 infection as compared to HIV-1. *Science* 1994;265:1587–1590. [PubMed: 7915856]
7. Kanki PJ, Travers KU, Mboup S, Hsieh CC, Marlink RG, Gueye-Ndiaye A, et al. Slower heterosexual spread of HIV-2 than HIV-1. *Lancet* 1994;343:943–946. [PubMed: 7909009]
8. Popper SJ, Sarr AD, Travers KU, Gueye-Ndiaye A, Mboup S, Essex ME, Kanki PJ. Lower human immunodeficiency virus (HIV) type 2 viral load reflects the difference in pathogenicity of HIV-1 and HIV-2. *J Infect Dis* 1999;180:1116–1121. [PubMed: 10479138]
9. Ariyoshi K, Jaffar S, Alabi AS, Berry N, Schim van der Loeff M, Sabally S, et al. Plasma RNA viral load predicts the rate of CD4 T cell decline and death in HIV-2-infected patients in West Africa. *Aids* 2000;14:339–344. [PubMed: 10770535]
10. Gottlieb GS, Sow PS, Hawes SE, Ndoeye I, Redman M, Coll-Seck AM, et al. Equal plasma viral loads predict a similar rate of CD4+ T cell decline in human immunodeficiency virus (HIV) type 1- and HIV-2-infected individuals from Senegal, West Africa. *J Infect Dis* 2002;185:905–914. [PubMed: 11920314]
11. Gilbert PB, McKeague IW, Eisen G, Mullins C, Gueye NA, Mboup S, Kanki PJ. Comparison of HIV-1 and HIV-2 infectivity from a prospective cohort study in Senegal. *Stat Med* 2003;22:573–593. [PubMed: 12590415]
12. Gottlieb GS, Hawes SE, Agne HD, Stern JE, Critchlow CW, Kiviat NB, Sow PS. Lower levels of HIV RNA in semen in HIV-2 compared with HIV-1 infection: implications for differences in transmission. *Aids* 2006;20:895–900. [PubMed: 16549974]
13. Odehouri K, De Cock KM, Krebs JW, Moreau J, Rayfield M, McCormick JB, et al. HIV-1 and HIV-2 infection associated with AIDS in Abidjan, Cote d'Ivoire. *AIDS* 1989 1989;3:509–512.
14. De Cock KM, Odehouri K, Colebunders RL, Adjorlolo G, Lafontaine M-F, Porter A, et al. A comparison of HIV-1 and HIV-2 infections in hospitalized patients in Abidjan, Cote d'Ivoire. *AIDS* 1990 1990;4:443–448.
15. Matheron S, Pueyo S, Damond F, Simon F, Lepretre A, Campa P, et al. Factors associated with clinical progression in HIV-2 infected-patients: The French ANRS cohort. *Aids* 2003;17:2593–2601. [PubMed: 14685053]
16. Nkengasong JN, Kestens L, Ghys PD, Koblavi-Deme S, Otten RA, Bile C, et al. Dual infection with human immunodeficiency virus type 1 and type 2: impact on HIV type 1 viral load and immune activation markers in HIV-seropositive female sex workers in Abidjan, Ivory Coast. *AIDS Res Hum Retroviruses* 2000;16:1371–1378. [PubMed: 11018856]
17. Andersson S, Norrgren H, da Silva Z, Biague A, Bamba S, Kwok S, et al. Plasma viral load in HIV-1 and HIV-2 singly and dually infected individuals in Guinea-Bissau, West Africa: significantly lower plasma virus set point in HIV-2 infection than in HIV-1 infection. *Arch Intern Med* 2000;160:3286–3293. [PubMed: 11088091]
18. Schutten M, van der Ende ME, Osterhaus AD. Antiretroviral therapy in patients with dual infection with human immunodeficiency virus types 1 and 2. *N Engl J Med* 2000;342:1758–1760. [PubMed: 10858105]

19. Alabi AS, Jaffar S, Ariyoshi K, Blanchard T, Schim Van Der Loeff M, Awasana AA, et al. Plasma viral load, CD4 cell percentage, HLA and survival of HIV-1, HIV-2, and dually infected Gambian patients. *Aids* 2003;17:1513–1520. [PubMed: 12824789]
20. van der Sande MA, Luong TN, Schim van der Loeff MF, Sabally S, Aveika AA, Corrah T, et al. Dual HIV-1 and HIV-2 infection in a West African infant. *Ann Trop Paediatr* 2004;24:277–278. [PubMed: 15479581]
21. UNAIDS AIDS epidemic update 2006. Geneva, Switzerland: Joint United Nations Programme on HIV/AIDS (UNAIDS) and World Health Organization (WHO) 2004; 2006. p. 1-94.
22. WHO. Scaling Up Antiretroviral Therapy in Resource-Limited Settings. 2002.
23. Desclaux A, Ciss M, Taverne B, Sow PS, Egrot M, Faye MA, et al. Access to antiretroviral drugs and AIDS management in Senegal. *Aids* 2003;17 Suppl 3:S95–101. [PubMed: 14565615]
24. Adje-Toure CA, Cheingsong R, Garcia-Lerma JG, Eholie S, Borget MY, Bouchez JM, et al. Antiretroviral therapy in HIV-2-infected patients: changes in plasma viral load, CD4+ cell counts, and drug resistance profiles of patients treated in Abidjan, Cote d'Ivoire. *Aids* 2003;17 Suppl 3:S49–54. [PubMed: 14565609]
25. Ndour CT, Batista G, Manga NM, Gueye NF, Badiane NM, Fortez L, Sow PS. [Efficacy and tolerance of antiretroviral therapy in HIV-2 infected patients in Dakar: preliminary study]. *Med Mal Infect* 2006;36:111–114. [PubMed: 16480843]
26. Ren J, Bird LE, Chamberlain PP, Stewart-Jones GB, Stuart DI, Stammers DK. Structure of HIV-2 reverse transcriptase at 2.35-Å resolution and the mechanism of resistance to non-nucleoside inhibitors. *Proc Natl Acad Sci U S A* 2002;99:14410–14415. [PubMed: 12386343]
27. Descamps, D.; Matheron, S.; Damond, F.; Peytavin, G.; Campa, P.; Pueyo, S., et al. Polymorphism of HIV-2 Protease Gene and Selection of Drug Resistance Mutations in HIV-2 Infected Patients Included in The French ANRS Cohort And Treated with Protease Inhibitors. The 2nd IAS Conference on HIV Pathogenesis and Treatment, 2003; Paris. 2003.
28. Pieniazek D, Rayfield M, Hu DJ, Nkengasong JN, Soriano V, Heneine W, et al. HIV-2 protease sequences of subtypes A and B harbor multiple mutations associated with protease inhibitor resistance in HIV-1. *Aids* 2004;18:495–502. [PubMed: 15090802]
29. Witvrouw M, Pannecouque C, Switzer WM, Folks TM, De Clercq E, Heneine W. Susceptibility of HIV-2, SIV and SHIV to various anti-HIV-1 compounds: implications for treatment and postexposure prophylaxis. *Antivir Ther* 2004;9:57–65. [PubMed: 15040537]
30. Poveda E, Rodes B, Toro C, Soriano V. Are fusion inhibitors active against all HIV variants? *AIDS Res Hum Retroviruses* 2004;20:347–348. [PubMed: 15117459]
31. Colson P, Henry M, Tourres C, Lozachmeur D, Gallais H, Gastaut JA, et al. Polymorphism and drug-selected mutations in the protease gene of human immunodeficiency virus type 2 from patients living in Southern France. *J Clin Microbiol* 2004;42:570–577. [PubMed: 14766818]
32. Damond F, Brun-Vezinet F, Matheron S, Peytavin G, Campa P, Pueyo S, et al. Polymorphism of the human immunodeficiency virus type 2 (HIV-2) protease gene and selection of drug resistance mutations in HIV-2-infected patients treated with protease inhibitors. *J Clin Microbiol* 2005;43:484–487. [PubMed: 15635022]
33. Maniar JK, Damond F, Kamath RR, Mandalia S, Surjushe A. Antiretroviral drug-resistant HIV-2 infection--a new therapeutic dilemma. *Int J STD AIDS* 2006;17:781–782. [PubMed: 17062187]
34. Boyer PL, Sarafianos SG, Clark PK, Arnold E, Hughes SH. Why do HIV-1 and HIV-2 use different pathways to develop AZT resistance? *PLoS Pathog* 2006;2:e10. [PubMed: 16485036]
35. Rodes B, Sheldon J, Toro C, Jimenez V, Alvarez MA, Soriano V. Susceptibility to protease inhibitors in HIV-2 primary isolates from patients failing antiretroviral therapy. *J Antimicrob Chemother* 2006;57:709–713. [PubMed: 16464891]
36. Rodes B, Toro C, Sheldon JA, Jimenez V, Mansinho K, Soriano V. High rate of proV47A selection in HIV-2 patients failing lopinavir-based HAART. *Aids* 2006;20:127–129. [PubMed: 16327332]
37. Smith RA, Gottlieb GS, Anderson DJ, Pyrak CL, Preston BD. Human immunodeficiency virus types 1 and 2 exhibit comparable sensitivities to Zidovudine and other nucleoside analog inhibitors in vitro. *Antimicrob Agents Chemother* 2008;52:329–332. [PubMed: 17967913]

38. Desbois D, Roquebert B, Peytavin G, Damond F, Collin G, Benard A, et al. In Vitro Phenotypic Susceptibility of Human Immunodeficiency Virus Type 2 Clinical Isolates to Protease Inhibitors. *Antimicrob Agents Chemother* 2008;52:1545–1548. [PubMed: 18227188]
39. Brower ET, Bacha UM, Kawasaki Y, Freire E. Inhibition of HIV-2 protease by HIV-1 protease inhibitors in clinical use. *Chem Biol Drug Des* 2008;71:298–305. [PubMed: 18312292]
40. Ruelle J, Sanou M, Liu HF, Vandenbroucke AT, Duquenne A, Goubau P. Genetic polymorphisms and resistance mutations of HIV type 2 in antiretroviral-naïve patients in Burkina Faso. *AIDS Res Hum Retroviruses* 2007;23:955–964. [PubMed: 17725411]
41. Clark NM, Dieng Sarr A, Sankale JL, Kanki PJ, Kazanjian P, Winfield R, Markovitz DM. Immunologic and virologic response of HIV-2 infection to antiretroviral therapy. *Aids* 1998;12:2506–2507. [PubMed: 9875594]
42. Rodes B, Holguin A, Soriano V, Dourana M, Mansinho K, Antunes F, Gonzalez-Lahoz J. Emergence of drug resistance mutations in human immunodeficiency virus type 2-infected subjects undergoing antiretroviral therapy. *J Clin Microbiol* 2000;38:1370–1374. [PubMed: 10747109]
43. Soriano V, Gomes P, Heneine W, Holguin A, Doruana M, Antunes R, et al. Human immunodeficiency virus type 2 (HIV-2) in Portugal: clinical spectrum, circulating subtypes, virus isolation, and plasma viral load. *J Med Virol* 2000;61:111–116. [PubMed: 10745242]
44. Brandin E, Lindborg L, Gyllensten K, Brostrom C, Hagberg L, Gisslen M, et al. pol Gene Sequence Variation in Swedish HIV-2 Patients Failing Antiretroviral Therapy. *AIDS Res Hum Retroviruses* 2003;19:543–550. [PubMed: 12908931]
45. Smith NA, Shaw T, Berry N, Vella C, Okorafor L, Taylor D, et al. Antiretroviral therapy for HIV-2 infected patients. *J Infect* 2001;42:126–133. [PubMed: 11531319]
46. van der Ende ME, Prins JM, Brinkman K, Keuter M, Veenstra J, Danner SA, et al. Clinical, immunological and virological response to different antiretroviral regimens in a cohort of HIV-2-infected patients. *Aids* 2003;17 Suppl 3:S55–61. [PubMed: 14565610]
47. Mullins C, Eisen G, Popper S, Dieng Sarr A, Sankale JL, Berger JJ, et al. Highly active antiretroviral therapy and viral response in HIV type 2 infection. *Clin Infect Dis* 2004;38:1771–1779. [PubMed: 15227626]
48. Matheron S, Damond F, Benard A, Taieb A, Campa P, Peytavin G, et al. CD4 cell recovery in treated HIV-2-infected adults is lower than expected: results from the French ANRS CO5 HIV-2 cohort. *Aids* 2006;20:459–462. [PubMed: 16439883]
49. Drylewicz J, Matheron S, Lazaro E, Damond F, Bonnet F, Simon F, et al. Comparison of viro-immunological marker changes between HIV-1 and HIV-2-infected patients in France. *Aids* 2008;22:457–468. [PubMed: 18301058]
50. Jallow S, Kaye S, Alabi A, Aveika A, Sarge-Njie R, Sabally S, et al. Virological and immunological response to Combivir and emergence of drug resistance mutations in a cohort of HIV-2 patients in The Gambia. *Aids* 2006;20:1455–1458. [PubMed: 16791023]
51. Damond, F.; Bénard, A. ACHeIV2E Study Group. Quality Control Assessment of HIV-2 Viral Load Quantification Assays. Results from an International Collaboration on HIV-2 Infection. 14th Conference on Retroviruses and Opportunistic Infections; Los Angeles, CA, USA. 2007.
52. Clavel F, Guetard D, Brun-Vezinet F, Chamaret S, Rey MA, Santos-Ferreira MO, et al. Isolation of a new human retrovirus from West African patients with AIDS. *Science* 1986;233:343–346. [PubMed: 2425430]
53. Markowitz M, Saag M, Powderly W, Hurley A, Hsu A, Valdes J, et al. A preliminary study of ritonavir, an inhibitor of HIV-1 protease, to treat HIV-1 infection. *N Engl J Med* 1995;333:1534. [PubMed: 7477168]
54. Collier AC, Coombs RW, Schoenfeld DA, Bassett RL, Timpone J, Baruch A, et al. Treatment of human immunodeficiency virus infection with saquinavir, zidovudine, and zalcitabine. AIDS Clinical Trials Group. *N Engl J Med* 1996;334:1011–1017. [PubMed: 8598838]
55. Carpenter C, Fischl M, Hammer S, et al. Antiviral therapy for HIV infection in 1997. Updated recommendations of the international AIDS society - USA panel. *JAMA* 1997;277:1962–1969. [PubMed: 9200638]
56. Willey S, Peters PJ, Sullivan WM, Dorr P, Perros M, Clapham PR. Inhibition of CCR5-mediated infection by diverse R5 and R5X4 HIV and SIV isolates using novel small molecule inhibitors of

- CCR5: effects of viral diversity, target cell and receptor density. *Antiviral Res* 2005;68:96–108. [PubMed: 16157392]
57. Roquebert B, Damond F, Collin G, Matheron S, Taieb A, Peytavin G, et al. Polymorphism of HIV-2 integrase gene and in vitro phenotypic susceptibility of HIV-2 clinical isolates to integrase inhibitors: raltegravir and evitegravir. *Antivir Ther* 2007;12:S92.
58. Damond F, Lariven S, Roquebert B, Males S, Peytavin G, Morau G, et al. Virological and immunological response to HAART regimen containing integrase inhibitors in HIV-2-infected patients. *Aids* 2008;22:665–666. [PubMed: 18317013]
59. WHO. Recommendations for a public health approach. 2006. Antiretroviral therapy for HIV infection in adults and adolescents.
60. WHO. Recommendations for a public health approach. 2006. Antiretroviral drugs for treating pregnant women and preventing HIV infection in infants: towards universal access.
61. Eholie S, Anglaret X. Commentary: decline of HIV-2 prevalence in West Africa: good news or bad news? *Int J Epidemiol* 2006;35:1329–1330. [PubMed: 16847021]
62. van der Loeff MF, Awasana AA, Sarge-Njie R, van der Sande M, Jaye A, Sabally S, et al. Sixteen years of HIV surveillance in a West African research clinic reveals divergent epidemic trends of HIV-1 and HIV-2. *Int J Epidemiol* 2006;35:1322–1328. [PubMed: 16543363]