

## Research

# Can non-physician health-care workers assess and manage cardiovascular risk in primary care?

Dele O Abegunde,<sup>a</sup> Bakuti Shengelia,<sup>a</sup> Anne Luyten,<sup>a</sup> Alexandra Cameron,<sup>a</sup> Francesca Celletti,<sup>a</sup> Sania Nishtar,<sup>b</sup> Vasu Pandurangi<sup>c</sup> & Shanthi Mendis<sup>a</sup>

**Objective** To ascertain the reliability of applying the WHO Cardiovascular Risk Management Package by non-physician health-care workers (NPHWs) in typical primary health-care settings.

**Methods** Based on an a priori 80% agreement level between the NPHWs and the "expert" physicians (gold standard), 649 paired (matched) applications of the protocol were obtained for analysis using Kappa statistic and multivariate logit regression.

**Findings** Results indicate over 80% agreement between raters, from moderate to perfect levels of agreement in almost all of the sections in the package. The odds of obtaining a difference between raters and a benchmark are not statistically significant.

**Conclusion** Applying the WHO Cardiovascular Risk Management Package, NPHWs can be retrained to reliably and effectively assess and manage cardiovascular risks in primary health-care settings where there are no attending physicians. The package could be a useful tool for scaling up the management of cardiovascular diseases in primary health care.

Bulletin of the World Health Organization 2007;85:432–440.

Une traduction en français de ce résumé figure à la fin de l'article. Al final del artículo se facilita una traducción al español. الترجمة العربية لهذه الخلاصة في نهاية النص الكامل لهذه المقالة.

## Introduction

Chronic noncommunicable diseases, especially cardiovascular diseases, are a major and increasing cause of death and disability worldwide, and may have retarding effects on the economies of affected individuals, households and countries.<sup>1,2</sup> The epidemiological and economic effects of cardiovascular diseases (specifically stroke and heart diseases) and diabetes, are especially pervasive in low- and middle-income countries<sup>1–3</sup> where health systems are less likely to adequately respond to the challenges of the increasing burden. Socioeconomic barriers and inequalities in access to treatment, suboptimal staffing of health-care facilities and limited capacity for ancillary investigations that complement cardiovascular risk profiling are some of the common problems limiting these countries' control of chronic diseases, especially at the primary health-care level.<sup>4</sup> This situation is worsened by the brain-drain syndrome<sup>5–9</sup> resulting in shortages of skilled workers.

The absolute-risk approach for the clinical management of cardiovascular diseases has been advocated<sup>10–14</sup> as a cost-effective approach to cardiovascular disease (CVD) management with improved patient outcomes as compared to the treatment of individual risk factors. The Framingham and other similar studies (e.g. PROCAM [Munster],<sup>15</sup> Seven Countries Study, SCORE<sup>16</sup> and Progetto CUORE<sup>17</sup> studies) provide the basis for the equations upon which many of the existing cardiovascular risk-profiling packages<sup>11,18–28</sup> have been developed.<sup>29</sup> However, such risk profiling protocols lack universal applicability<sup>11,13,14,29–42</sup> and may be of limited applicability in developing countries, whose populations were not sampled for the Framingham<sup>31,32</sup> and other studies. Uncritical adoption of such protocols may result in negative clinical and economic consequences.<sup>43</sup>

To address the absence of a CVD risk profiling tool for developing countries, WHO in 2000 developed a package for the assessment and management

of cardiovascular risk in low-resource settings.<sup>44</sup> The package, developed through consultations with experts from all WHO regions, was designed as an adaptable, cost-effective tool for systematic case management at all health-care levels, and consequently for scaling up countries' health systems. The expert panel based the design of the package on the graded evidence available.

The package includes three scenarios that reflect commonly encountered resource availability strata in low- and medium-resource settings. While the basic elements remain the same across the three scenarios, the specific thresholds for clinical intervention differ according to the level of personnel and facilities available. Each scenario begins with cardiovascular risk screening using hypertension as an entry point, though each can be adapted for use with diabetes or smoking as entry points. The protocols in each scenario consist of algorithms for patient history (of heart attack, angina, stroke, transient ischaemic attack (TIA), diabetes and patients' lifestyle);

<sup>a</sup> World Health Organization, 20 avenue Appia, 1211 Geneva 27, Switzerland. Correspondence to Shanthi Mendis (e-mail: mendiss@who.int).

<sup>b</sup> Heartfile, Islamabad, Pakistan.

<sup>c</sup> CAMHADD Project, Bangalore Healthy City Initiative for Preventive Healthcare, Bangalore, India.

doi: 10.2471/BLT.06.032177

(Submitted: 5 April 2006 – Final revised version received: 17 November 2006 – Accepted: 23 November 2006)

examination (particularly two systolic blood pressure measurements at 5–10 minute intervals); and resultant systematic treatment and follow-up (Fig. 1). Patients are stratified into one of five possible treatment tracks according to their levels of cardiovascular risk. The decisions in each of the treatment tracks include one or more of the following: referral to the next care level; counselling on diet, physical activity and ceasing tobacco use; prescription of low-dose thiazides; and follow-up. High-risk patients are immediately referred to the next level of care, leaving only patients who can be managed appropriately at the primary health-care level. The decision algorithms extend to the second and third follow-up visits, spaced at 1–3, 2–3, or 4–6 month intervals depending on the patients' risk state.

Scenario I (Fig. 1) is applicable to the least-resourced section of health systems, which is usually staffed by non-physician health-care workers (NPHWs) in low- and middle-income countries. The algorithms of Scenario I are therefore designed to assist scientifically sound case-by-case decisions by NPHWs. They should guide the NPHW to make evidence-based and cost-effective patient management decisions comparable to those that would be made by skilled physicians. The overall patient management goal is to improve the patient's absolute cardiovascular risk profile in addition to aiding timely and appropriate referral decisions in high-risk cases.

A necessary precondition for adoption is proof of reliability when applied by non-physicians. We assumed face, construct and content validity for the package, since it was developed through a rigorous WHO-supervised expert consultation process. However, it would be necessary to establish criterion validity (application of the package by the NPHW correlating with a criterion of "true" value). The objective of this study was to ascertain the reliability of applying Scenario I of the package by NPHW when compared to "expert" physicians in typical primary health-care settings (primary health-care centres in Bangalore, India, and Islamabad, Pakistan). As is sometimes the case where the "true values" against which to test such protocols are not clearly established such that a strict standard of expert care is available, correlating raters' estimates with surrogate endpoints could reasonably

approximate a test of accuracy. A second stage to test the effectiveness of the package is under way in 11 countries.

## Methods

The study's sample size was based on an assumption that the NPHW's application of each component of the Scenario I protocol must agree or correlate up to 80% with those of the physicians when compared. NPHWs applied the algorithm on patients attending sampled clinics; this was followed by an independent application of the same algorithm on the same patient by a physician, thereby matching the observations. There were 111 paired observations from sampled NPHW-staffed primary health-care centres from the Bangalore region of India, and 538 pairs of observations from similar centres in Islamabad, Pakistan. All observations from the NPHWs and the physicians were recorded on standardized visit record forms. Observers were blinded to each other. An investigating physician conducted exit interviews on the patients after they had received treatment. Data were converted to electronic format using the Enter suite of EPI INFO version 6.04b, and analysed with STATA software intercooled version 8.<sup>45</sup>

The NPHWs and physicians participated jointly in an initial three-day training exercise to acquaint participants with the CVD-Risk Management Package and its mode of application.

Participants learned how the package should assist in improving and standardizing patient management in a primary health-care environment. The training materials consisted of a protocol application guide and information materials designed to increase knowledge of CVD risk assessment and management. This training cost US\$ 88.30 per participant (30 in all) in Pakistan including transportation, training materials, per diem and trainers' fee.

## Analysis

Our data consisted primarily of dichotomous choices with a few continuous biometric measurements; therefore analysis was two-pronged. The first was a pairwise comparison of each variable of interest in the protocol; this enabled the detection of variables that were problematic for the NPHW to elucidate. To observe the inter-rater agreement in their choices on the six possible decisions points, we employed Kappa statistic (see Box 1), scaled to zero when the amount of agreement is less than what would be expected by chance, and scaled to one when there is perfect agreement. Intermediate values which are possible, are usually (as we have done) interpreted as follows:  $K < 0$ , poor;  $0.0 < K < 0.20$ , slight;  $0.21 < K < 0.40$ , fair;  $0.41 < K < 0.60$ , moderate;  $0.61 < K < 0.80$ , substantial; and  $0.81 < K < 1.00$ , almost perfect.<sup>46</sup> Correlation analysis was employed to compare agreement in

### Box 1. Description of Kappa statistic

If  $r$  = number of raters (in this case 2), the observed proportion of agreement ( $P_o$ ) is given as

$$P_o = \sum_{i=1}^k \sum_{j=1}^k w_{ij} p_{ij}$$

where  $p_{ij}$  is the fraction of ratings  $i$  (by the non-physician health-care worker) and  $j$  (by the expert physician), and  $w_{ij}$  is the weight assigned to the raters. The expected proportion of the agreement ( $P_e$ ) if the raters agree at random is also given by

$$P_e = \sum_{i=1}^k \sum_{j=1}^k w_{ij} p_i p_j$$


where

$$p_i = \sum_j p_{ij} \text{ and } p_j = \sum_i p_{ij}$$

Kappa ( $K$ ) is given by

$$k = (P_o - P_e) / (1 - P_e)$$

Fig. 1. Patient management algorithm in Scenario I: WHO CVD-Risk Management Package



CVD, cardiovascular disease; BMI, body mass index; BP, blood pressure; SBP, systolic blood pressure, TIA, transient ischaemic attack.

<sup>a</sup> In areas where coronary artery diseases rates exceed stroke rates.

<sup>b</sup> Thiazide diuretic: Hydrochlorothiazide starting dose 12.5 mg (low-dose) to be increased up to 25 mg (maximum dose).

<sup>c</sup> Second drug option: use the cheapest out of beta-blockers or calcium-channel blockers or ACE-inhibitors.


If drugs given in footnotes (b) and (c) are not available: use methyldopa or reserpine or fixed dose combination.

Source: WHO CVD-Risk Management Package for low- and medium-resource settings.

the continuous numeric variables in the data: age, blood pressure, weight, body mass index (BMI) and waist circumference, setting the least acceptable level of agreement to 80%.

The second analysis was multivariate, in which the agreement between raters in applying the treatment decision section of the protocol was tested. This analysis was conducted because it was possible that the NPHWs reliably and accurately elucidated the decision section, but performed less well in the treatment section. There could also be several other possibilities, such as the expert physicians' (EPs') and the NPHWs' estimates erroneously agreeing in some cases, and/or systematic errors in the ratings of both raters. Further, the EPs may have been less than fully accurate, or as easily prone to errors in the protocol application as the NPHWs. It is also possible that NPHWs applied the protocol more precisely and accurately than did the EPs, for instance, if the NPHWs followed the decision and treatment logic of the protocol more thoroughly. To account for possible imprecision and inaccuracy in the gold standard, we compared the decisions taken by the NPHW and the EP with what the decision should have been if both raters had accurately applied the protocol. To this end, a third unbiased rater (the benchmark) was constructed by programming what the treatment decision should be (given the design of the Scenario I treatment protocol) for all observations in the history and examination sections where the EP and the

Fig. 2. Correlations between estimates of NPHWs and EPs for age



EPs, expert physicians; NPHWs, non-physician health-care workers.  
Source: WHO Cardiovascular Disease Unit.

NPHW agreed perfectly. This eliminated the outliers. Similarly to the analysis of matched multirater case control studies,<sup>47,48</sup> we estimated the odds of differences in following the correct treatment track of the protocol for each rater when compared with the benchmark. We used multinomial (conditional) logit regression for matched case control groups,<sup>47-50</sup> with the raters as the nominal dependent variable, and the choices they made in the treatment track as the independent variables. For each  $i^{\text{th}}$  rater and  $j$  choices, the probability ( $\text{Prob}(Y_j = j)$ ) of choice  $j$  is modelled as:

$$\text{Prob}(Y_j = j) = \frac{e^{Z_{ij}\beta}}{\sum_{j=1}^j e^{Z_{ij}\beta}}$$

$j = (0, 1 \dots n), i = (1, 2 \& 3)$


where  $Z$  represents the attributes of the treatment choices and raters' characteristics.<sup>48,49</sup> Though our data did not contain background information on the raters, we assumed that the administration of the instrument within the same visit period and location mitigates the potential impact of possible maturation (actual and time-related changes in patients' conditions). In addition, we assumed that the initial training reduced the possible impact of inter-rater differences. The model estimated the propensity for change ( $\beta$ : difference or slope coefficient) in raters' choices. The purpose was to assess which aspects of the treatment algorithm have valuations between the three raters (NPHW, EP or the benchmark) tended to differ with respect to the benchmark. That is, when all the choices were jointly correlated, if the odds of observing a difference in a particular value are not statistically significant, then all three rater groups largely agreed. A stepwise regression analysis was conducted to eliminate the variables where raters' agreement was perfect, setting  $P$ -values to 0.05. All statistics were implemented using the routines in STATA<sup>45</sup> statistical software.

## Results

### Correlation between rates: age, blood pressure, weight, BMI and waist circumference

We report the results of combined data from the two study sites because there were no significant differences in the results for India and Pakistan when analysed independently. The similarity

Fig. 3. Correlations between estimates of NPHWs and EPs for blood pressure



EPs, expert physicians; NPHWs, non-physician health-care workers.  
Source: WHO Cardiovascular Disease Unit.

may be largely due to the simplicity of the algorithms and the initial training. Pooling the data also appeals to the aim of universal application of the package in situations where there are no physicians.

Patients' ages as elicited by the NPHWs correlated strongly with those of the EPs (correlation coefficient: 0.9146). Similarly, raters' measurements of blood pressure (correlation coefficient: 0.8836) and body weight (correlation coefficient: 0.8912) correlated beyond the a priori 80% level (Figs. 2–4). Similar correlations are observable for waist circumference and BMI, except for a few outlying and omitted measurements on the part of both the EPs and the NPHWs (Figs. 5 and 6). In the Pakistan data, there were two BMI outliers of over 150 kg/m<sup>2</sup> from the EP, and one BMI value of over 80 kg/m<sup>2</sup> from the NPHW (Fig. 5). In addition, the EP in Pakistan and the NPHW in India each had an outlying value for waist circumference of > 600 cm and > 150 cm, respectively (Fig. 6). These outliers are likely due to computational, recording or data entry errors. There were also missing entries in a few cases. All of these, in addition to the inherent problems with measuring waist circumference in a culturally sensitive environment, will affect the true levels of correlation, particularly for the waist circumference measurements. Overall, correlation coefficients of over 0.80 were obtained by excluding these outliers.

**Inter-rater agreement (Kappa statistic)**


**I. The history and risk mapping section**

Over 80% agreement levels were obtained in all the variables in the history section of the protocol (Table 1). Kappa values varied from fair (history of stroke), moderate (history of TIA), substantial (history of heart attack and angina) to perfect agreement (sex of patient and history of diabetes).

**II. The health behaviour section (Table 1)**


Although elucidating lifestyle history could be characterized as imprecise, inter-rater agreement of over 80% between the EP and the NPHW was obtained for all the questions except that of physical activity (78.90% inter-rater agreement) though a moderate degree of agreement

Fig. 4. Correlations between estimates of NPHWs and EPs for body weight



EPs, expert physicians; NPHWs, non-physician health-care workers. Source: WHO Cardiovascular Disease Unit.

Fig. 5. Correlations between estimates of NPHWs and EPs for BMI



BMI, body mass index; EPs, expert physicians; NPHWs, non-physician health-care workers; Pak, Pakistan. Source: WHO Cardiovascular Disease Unit.

above chance was achieved. Overall, the degree to which the agreement levels were not due to chance varied from fair to substantial.

**III. The "treatment tracks" section of the algorithm (Table 2)**

Not surprisingly, given the high agreement in the history and examination

sections, the inter-rater agreement achieved for treatment decisions taken ranged from 88% (prescription in track F) to 99.5% (referral in track F). Compared to other sections of the protocol, inter-rater agreements were highest for the treatments tracks. Raters' agreement ranged from 93% to 99.5% for the six treatment tracks (A–F). Of the three

treatment choices, agreements on prescription were the highest in each of the treatment tracks except track F, where according to the protocol there should have been no prescription but only referral to a higher level of care. Kappa levels range from moderate to perfect agreement, indicating that the levels of agreement were far beyond those that are possible by chance.


### Multivariate analysis result

The results indicate an overall agreement in almost all of the treatment decision tracks, with the exception of counselling decisions in tracks D and C, and referral decisions in tracks A and E (see Table 3). These differences suggest that both the EPs and the NPHWs, when compared to the independent benchmark, may have rendered more counselling and less referral than was necessary. This observation may be related to some omitted values as noted previously, or to the influence of local practice.

### Discussion

With respect to the primary objective of this study, results indicate that NPHWs employed Scenario I of the WHO CVD-Risk Management Package comparably to physicians, who are arguably better skilled. Over 80% agreement was achieved for almost all of the items in the protocol and Kappa statistics indicate that the agreements were largely not due to chance.

Fig. 6. Correlations between estimates of NPHWs and EPs for waist circumference



EPs, expert physicians; Ind, India; NPHWs, non-physician health-care workers; Pak, Pakistan. Source: WHO Cardiovascular Disease Unit.

A view often held is that stepping up specialist training and investing in diagnostic technologies are a panacea for the increasing burden of cardiovascular diseases, even in low- to middle-income countries. However, in these settings

such strategies may be neither affordable nor cost-effective. An alternative viewpoint is that currently available resources and skills could be readapted to respond sufficiently and efficiently to the changing health needs in many

Table 1. Level of agreement (Kappa statistic) between EPs and NPHWs on past medical history and health behaviour sections of the WHO CVD-Risk Management Package

Variable	Inter-rater agreement, $P_o$ (%)	Kappa statistic		Comments
		$(P_o - P_e)/(1 - P_e)$ Kappa	Prob > z	
<b>Patient's cardiovascular history</b>				
Sex	97.75	0.9023	0.000	Perfect
Heart attack	99.53	0.7977	0.000	Substantial
Angina	90.22	0.6182	0.000	Substantial
Diabetes	97.67	0.8328	0.000	Perfect
Transient ischaemic attack	98.44	0.5376	0.000	Moderate
Stroke	99.38	0.3306	0.000	Fair
<b>Health/risk behaviour</b>				
Tobacco	89.88	0.7480	0.000	Substantial
Physical activity	78.90	0.5744	0.000	Moderate
Eating less salt	83.13	0.6624	0.000	Substantial
Fruits	92.50	0.3287	0.000	Fair
Fish	97.97	0.2294	0.000	Fair
Fatty foods	74.14	0.4560	0.000	Moderate
Alcohol	93.69	0.7733	0.000	Substantial

CVD, cardiovascular disease; EPs, expert physicians; NPHWs, non-physician health-care workers.

of these countries. The results of this exercise indicate that NPHWs could be retrained and assisted to be more effective in assuming primary roles in the care of patients with chronic noncommunicable diseases, especially where there are no physicians. This could be an initial step in incorporating the management of chronic diseases into the health-care setting in developing countries, which has traditionally focused mainly on the management of acute communicable diseases.

Though engaging low-skilled workers in the management of chronic disease risk may raise some ethical concerns, the reality in many countries is that NPHWs may have the only health-care skills available to a large proportion of the population. It could be that NPHWs are already compelled to care for cardiovascular diseases, though they may not be trained to recognize them or make robust, timely and life-saving treatment decisions. This is due, in part, to the “brain drain” of skilled health-care workers, which has compounded skill shortages and impacted negatively on health care.<sup>5-9</sup> Although brain drain has recently attracted international attention and concern from policy-makers, there are yet to be clear solutions to mitigate its impact on health care. It is therefore reasonable that available skills be re-adapted to the challenges on the ground. This package could aid such adaptation aimed at scaling up the health system response to chronic diseases. The package incorporates the principle of triage in its design, does not allow the NPHW to treat high-risk cases, and allows only the prescription of low-dose thiazide, mitigating ethical concerns.

The results of this study have implications for the country-level policy response to the increasing burden of

Table 2. Level of agreement (Kappa statistic) between EPs and NPHWs on counselling, drug treatment and referral sections of the WHO CVD-Risk Management Package

Variable	Inter-rater agreement, $P_o$ (%)	Kappa statistic		Comments
		$(P_o - P_e)/(1 - P_e)$ Kappa	Prob > z	
<b>Track A</b>	96.14	0.7537	0.000	Substantial
Counselling	96.30	0.6560	0.000	Substantial
Prescription	95.99	0.4796	0.000	Moderate
Referral	96.14	0.6984	0.000	Substantial
<b>Track B</b>	99.54	0.9067	0.000	Perfect
Counselling	96.14	0.7215	0.000	Substantial
Prescription	97.22	0.7484	0.000	Substantial
Referral	96.30	0.5940	0.000	Moderate
<b>Track C</b>	94.91	0.7470	0.000	Substantial
Counselling	94.19	0.7470	0.000	Substantial
Prescription	95.83	0.7152	0.000	Substantial
Referral	92.44	0.4877	0.000	Moderate
<b>Track D</b>	93.21	0.7861	0.000	Substantial
Counselling	92.90	0.7736	0.000	Substantial
Prescription	95.37	0.6976	0.000	Substantial
Referral	94.44	0.4300	0.000	Moderate
<b>Track E</b>	96.60	0.6680	0.000	Substantial
Counselling	96.60	0.6680	0.000	Substantial
Prescription	97.07	0.5215	0.000	Moderate
Referral	96.91	0.5317	0.000	Moderate
<b>Track F</b>	94.60	0.8914	0.000	Perfect
Counselling	94.14	0.8820	0.000	Perfect
Prescription	87.65	0.7145	0.000	Substantial
Referral	98.61	0.8452	0.000	Perfect

CVD, cardiovascular disease; EPs, expert physicians; NPHWs, non-physician health-care workers.

cardiovascular disease in low-resource settings. For instance, one of the main problems encountered in these settings is that, for reasons that include cost and busy schedules, opportunities for health workers to attend lengthy retraining programmes are limited. In this protocol test, the NPHW only underwent three days of training to achieve the observed

high level of agreement with the EP in the application of the risk management protocol. Health-care managers may find such three-day training acceptable to enable systematic, phased programming of regional or even countrywide retraining programmes. In general, readapting available health human resources to address the emerging chronic disease problem could result in health-care cost savings (apart from the motivational value for the workforce of retraining). Increased management of patients with cardiovascular risk in primary health-care settings could avoid more costly trips to higher care levels. Further, the use of NPHWs in managing CVD patients will free physicians to focus on high-risk cases, thereby resulting in increased efficiency of primary health-care resources. Low-resource countries could avoid unaffordable health-care costs and capital flight that would be incurred by investing in capital-intensive care and unaffordable medical equipment.

Table 3. Result of matched (stepwise) logistic regression<sup>a</sup>

Variable	Odds ratio	Standard error
Counselling in track C	2.469	1.032
Counselling in track D	66.770	48.339
Referral in track A	0.042	0.015
Referral in track E	0.075	0.031
Log likelihood	-238.13	
Log likelihood ratio chi <sup>2</sup>	422.07	
Prob > chi <sup>2</sup>	0	
Pseudo R <sup>2</sup>	0.4698	

<sup>a</sup> Expert physicians and non-physician health-care workers compared with independent benchmark.

## Conclusion

From this study, we conclude that NPHWs can easily be retrained to make reasonably safe and appropriate treat-

ment decisions with the aid of the WHO CVD-Risk Management Package within the scope of their available skills. Work is under way to test the impact of

the package on population cardiovascular well-being in several countries. ■

**Competing interests:** None declared.

## Résumé

### Les agents de santé non médecins sont-ils en mesure d'évaluer et de prendre en charge le risque cardiovasculaire dans le cadre des soins de santé primaire ?

**Objectif** S'assurer de la fiabilité de l'utilisation du module de prise en charge du risque cardiovasculaire de l'OMS par des agents de santé non médecins, dans le cadre d'établissements de soins de santé primaire ordinaires.

**Méthodes** Sur la base d'un accord de 80 % entre les résultats de l'utilisation du module par des agents de santé non médecins et par des médecins compétents (référence), on a obtenu 649 paires d'applications (applications appariées) du protocole par analyse statistique Kappa et régression logistique multivariée (logit).

**Résultats** L'étude indique un accord global de plus de 80 % entre les évaluateurs, la concordance allant d'un niveau moyen

à celui de la perfection pour presque tous les volets du module. La différence entre les évaluateurs et une évaluation de référence n'était pas statistiquement significative.

**Conclusion** Il est possible de renforcer la formation des agents de santé de manière à ce qu'ils soient en mesure, en utilisant le module OMS, d'évaluer et de prendre en charge de manière fiable et efficace le risque cardiovasculaire dans les établissements de soins de santé primaire en l'absence de médecin. Ce module pourrait être utile au développement de la prise en charge des maladies cardiovasculaires dans le contexte des soins de santé primaire.

## Resumen

### ¿Puede el personal sanitario no médico evaluar y controlar el riesgo cardiovascular en la atención primaria?

**Objetivo** Determinar la fiabilidad de la aplicación del Módulo de Gestión del Riesgo Cardiovascular de la OMS por personal sanitario no médico (PSNM) en los entornos de atención primaria habituales.

**Métodos** Partiendo de un nivel de concordancia a priori del 80% entre el PSNM y los médicos «expertos» (criterio de referencia), se reunieron 649 aplicaciones del protocolo emparejadas para analizarlas mediante el estadístico Kappa y un modelo de regresión logit multifactorial.

**Resultados** Los resultados muestran una concordancia de más del 80% entre los evaluadores, con niveles de coincidencia entre

moderados y perfectos en casi todas las secciones del módulo. Las diferencias entre los evaluadores y la referencia utilizada no son estadísticamente significativas.

**Conclusión** Aplicando el Módulo de Gestión del Riesgo Cardiovascular de la OMS, es posible formar al PSNM para que evalúe y controle de manera fiable y eficaz el riesgo cardiovascular en entornos de atención primaria en los que no hay ningún médico. El módulo podría ser un valioso instrumento para expandir el manejo de las enfermedades cardiovasculares en los entornos de atención primaria.

## ملخص

### هل بمقدور العاملين في الرعاية الصحية من غير الأطباء تقييم وتدبير الأخطار القلبية والوعائية في الرعاية الأولية؟

جميع الفقرات المندرجة ضمن حزمة المعلومات وبين التوافق المتوسط. ولم تكن قيمة الأرجحية التي حصلنا عليها من الفروق بين واضعي المعدلات من جهة وبين العلامات المميزة ذات اعتداد إحصائي.

**الاستنتاج:** يمكن إعادة تدريب العاملين في الرعاية الصحية من غير الأطباء على تطبيق حزمة معلومات منظمة الصحة العالمية حول تدبير الأخطار القلبية والوعائية وعلى تقييم وتدبير الأخطار القلبية والوعائية بشكل فعال وموثوق به في مواقع تقديم الرعاية الصحية الأولية التي تفتقد الأطباء. ويمكن لحزمة معلومات منظمة الصحة العالمية حول تدبير الأخطار القلبية والوعائية أن تكون أداة مفيدة للنهوض بتدبير الأمراض القلبية والوعائية في الرعاية الصحية الأولية.

**الهدف:** التأكد من مدى موثوقية تطبيق العاملين في الرعاية الصحية من غير الأطباء للحزمة التي أعدتها منظمة الصحة العالمية لتدبير الأخطار القلبية والوعائية في المواقع النموذجية للرعاية الصحية الأولية.

**الطريقة:** استناداً إلى مستوى الاتفاق المبدئي الذي تبلغ نسبته 80% بين العاملين في الرعاية الصحية من غير الأطباء وبين الأطباء الخبراء (الذين يعدون بمثابة المعيار الذهبي)، جمعت الدراسة 649 من الاستمارات المزدوجة (المتقارنة) التي أعدها الفريقان من العاملين الصحيين من غير الأطباء ومن الأطباء) حول البروتوكول المطبق لتقييم وتدبير الأخطار القلبية والوعائية، ثم حللتها باستخدام التحوف اللوغاريتمي المتعدد المتغيرات والتحوف الإحصائي كإجراء.

**الموجودات:** أشارت النتائج إلى وجود ما يزيد على 80 بالمائة من التوافق بين واضعي المعدلات، ويتراوح هذا التوافق بين مستوى التوافق الكامل في



## References

1. *Preventing chronic diseases: a vital investment*. Geneva, WHO, 2005.
2. Liu T. The Lancet's chronic diseases series. *Lancet* 2006;367:565-6.
3. Strong K, et al. Preventing chronic diseases: how many lives can we save? *Lancet* 2005;366:1578-82.
4. Mendis S, et al. Barriers to management of cardiovascular risk in a low-resource setting using hypertension as an entry point. *J Hypertens* 2004; 22:59-64.
5. Oberoi SS, Lin V. Brain drain of doctors from southern Africa: brain gain for Australia. *Aust Health Rev* 2006;30:25-33.
6. Muula AS. Is there any solution to the "brain drain" of health professionals and knowledge from Africa? *Croat Med J* 2005;46:21-9.
7. Martineau T, et al. "Brain drain" of health professionals: from rhetoric to responsible action. *Health Policy* 2004;70:1-10.
8. Scott ML, et al. "Brain drain" or ethical recruitment? *Med J Aust* 2004; 180:174-6.
9. Marchal B, Kegels G. Health workforce imbalances in times of globalization: brain drain or professional mobility? *Int J Health Plann Manage* 2003;18: 589-101.
10. Whitworth JA. 2003 World Health Organization (WHO)/International Society of Hypertension (ISH) statement on management of hypertension. *J Hypertens* 2003;21:1983-92.
11. Menotti A, et al. The estimate of cardiovascular risk. Theory, tools and problems. *Ann Ital Med Int* 2002;17:81-94.
12. 2003 European Society of Hypertension-European Society of Cardiology guidelines for the management of arterial hypertension. *J Hypertens* 2003; 21:1011-53.
13. Lenz M, Muhlhauser I. [Cardiovascular risk assessment for informed decision making. Validity of prediction tools]. *Med Klin (Munich)* 2004;99:651-61.
14. Neuhauser HK, et al. A comparison of Framingham and SCORE-based cardiovascular risk estimates in participants of the German National Health Interview and Examination Survey 1998. *Eur J Cardiovasc Prev Rehabil* 2005;12:442-50.
15. Assmann G, et al. Simple scoring scheme for calculating the risk of acute coronary events based on the 10-year follow-up of the prospective cardiovascular Munster (PROCAM) study. *Circulation* 2002;105:310-5.
16. Conroy RM, et al. Estimation of ten-year risk of fatal cardiovascular disease in Europe: the SCORE project. *Eur Heart J* 2003;24:987-1003.
17. Ferrario M, et al. Prediction of coronary events in a low incidence population. Assessing accuracy of the CUORE Cohort Study prediction equation. *Int J Epidemiol* 2005;34:413-21.
18. May M, et al. Cardiovascular disease risk assessment in older women - can we improve on Framingham?: British Women's Heart and Health prospective cohort study. *Heart* 2006.
19. Yip YB, et al. Cardiovascular disease: application of a composite risk index from the Telehealth System in a district community. *Public Health Nurs* 2004;21:524-32.
20. Milne R, et al. Discriminative ability of a risk-prediction tool derived from the Framingham Heart Study compared with single risk factors. *N Z Med J* 2003;116:U663.
21. Marchioli R, et al. Assessment of absolute risk of death after myocardial infarction by use of multiple-risk-factor assessment equations: GISSI-Prevenzione mortality risk chart. *Eur Heart J* 2001;22:2085-103.
22. Menotti A, et al. Riskard 2005. New tools for prediction of cardiovascular disease risk derived from Italian population studies. *Nutr Metab Cardiovasc Dis* 2005;15:426-40.
23. Stephens JW, et al. Cardiovascular risk and diabetes. Are the methods of risk prediction satisfactory? *Eur J Cardiovasc Prev Rehabil* 2004;11:521-8.
24. Menotti A, et al. An Italian chart for cardiovascular risk prediction. Its scientific basis. *Ann Ital Med Int* 2001;16:240-51.
25. Mehta RH, et al. Elderly patients at highest risk with acute myocardial infarction are more frequently transferred from community hospitals to tertiary centers: reality or myth? *Am Heart J* 1999;138:688-95.
26. Krumholz HM, et al. Validation of a clinical prediction rule for left ventricular ejection fraction after myocardial infarction in patients > or = 65 years old. *Am J Cardiol* 1997;80:11-5.
27. Haq IU, et al. Prediction of coronary risk for primary prevention of coronary heart disease: a comparison of methods. *QJM* 1999;92:379-85.
28. Rabindranath KS, et al. Comparative evaluation of the new Sheffield table and the modified joint British societies coronary risk prediction chart against a laboratory based risk score calculation. *Postgrad Med J* 2002;78:269-72.
29. Jurgensen JS. The value of risk scoring. *Heart* 2006.
30. Liu J, et al. Predictive value for the Chinese population of the Framingham CHD risk assessment tool compared with the Chinese Multi-Provincial Cohort Study. *JAMA* 2004;291:2591-9.
31. Brindle PM, et al. The accuracy and impact of risk assessment in the primary prevention of cardiovascular disease: A systematic review. *Heart* (2006).
32. Stork S, et al. Prediction of mortality risk in the elderly. *Am J Med* 2006; 119:519-25.
33. Bhopal R, et al. Predicted and observed cardiovascular disease in South Asians: application of FINRISK, Framingham and SCORE models to Newcastle Heart Project data. *J Public Health (Oxf)* 2005;27:93-100.
34. Hense HW, et al. Framingham risk function overestimates risk of coronary heart disease in men and women from Germany — results from the MONICA Augsburg and the PROCAM cohorts. *Eur Heart J* 2003;24:937-45.
35. Liao Y, et al. How generalizable are coronary risk prediction models? Comparison of Framingham and two national cohorts. *Am Heart J* 1999; 137:837-45.
36. Bastuji-Garin S, et al. The Framingham prediction rule is not valid in a European population of treated hypertensive patients. *J Hypertens* 2002; 20:1973-80.
37. Marrugat J, et al. An adaptation of the Framingham coronary heart disease risk function to European Mediterranean areas. *J Epidemiol Community Health* 2003;57:634-8.
38. D'Agostino RB Sr, et al. Validation of the Framingham coronary heart disease prediction scores: results of a multiple ethnic groups investigation. *JAMA* 2001;286:180-7.
39. Brindle PM, et al. The accuracy of the Framingham risk-score in different socioeconomic groups: a prospective study. *Br J Gen Pract* 2005;55:838-45.
40. Brindle P, et al. Predictive accuracy of the Framingham coronary risk score in British men: prospective cohort study. *BMJ* 2003;327:1267.
41. Empana JP, et al. Are the Framingham and PROCAM coronary heart disease risk functions applicable to different European populations? The PRIME Study. *Eur Heart J* 2003;24:1903-11.
42. Menotti A, et al. Comparison of the Framingham risk function-based coronary chart with risk function from an Italian population study. *Eur Heart J* 2000;21:365-70.
43. Fornasini M, et al. Consequences of using different methods to assess cardiovascular risk in primary care. *Fam Pract* 2006;23:28-33.
44. *WHO CVD-Risk Management Package for low- and medium-resource settings*. Geneva: WHO; 2002.
45. *Stata Statistical Software: Release 8.0*. College Station: Stata Corp.; 2003.
46. Lindis JR, Kock GG. The measurement of observer agreement for categorical data. *Biometrics* 1977;33:159-74.
47. Cook RJ, Farewell VT. Conditional inference for subject-specific and marginal agreement: two families of agreement measures. *Can J Stat* 1995;23:333-44.
48. Hosmer DW, Lemeshow S. *Applied Logistic Regression*. John Wiley & Sons, 1989.
49. Greene W. *Econometric Analysis*. New Jersey, Prentice Hall, 2003.
50. Bishop YMM, et al. *Discrete multivariate analysis: theory and practice*. Cambridge, MIT Press, 1975.