

# Maize AMEIOTIC1 is essential for multiple early meiotic processes and likely required for the initiation of meiosis

Wojciech P. Pawlowski<sup>a,1,2</sup>, Chung-Ju Rachel Wang<sup>b,1</sup>, Inna N. Golubovskaya<sup>b,c</sup>, Jessica M. Szymaniak<sup>a</sup>, Liang Shi<sup>d</sup>, Olivier Hamant<sup>b,3</sup>, Tong Zhu<sup>d</sup>, Lisa Harper<sup>b</sup>, William F. Sheridan<sup>e</sup>, and W. Zacheus Cande<sup>b,2</sup>

<sup>a</sup>Department of Plant Breeding and Genetics, Cornell University, Ithaca, NY 14853; <sup>b</sup>Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720; <sup>c</sup>N. I. Vavilov Research Institute of Plant Industry, St. Petersburg, Russia; <sup>d</sup>Syngenta Biotechnology, Inc., Research Triangle Park, NC 27709; and <sup>e</sup>Department of Biology, University of North Dakota, Grand Forks, ND 58202

Edited by Hugo K. Dooner, Rutgers, The State University of New Jersey, Piscataway, NJ, and approved December 22, 2008 (received for review October 13, 2008)

Molecular mechanisms that initiate meiosis have been studied in fungi and mammals, but little is known about the mechanisms directing the meiosis transition in other organisms. To elucidate meiosis initiation in plants, we characterized and cloned the *ameiotic1* (*am1*) gene, which affects the transition to meiosis and progression through the early stages of meiotic prophase in maize. We demonstrate that all meiotic processes require *am1*, including expression of meiosis-specific genes, establishment of the meiotic chromosome structure, meiosis-specific telomere behavior, meiotic recombination, pairing, synapsis, and installation of the meiosis-specific cytoskeleton. As a result, in most *am1* mutants premeiotic cells enter mitosis instead of meiosis. Unlike the genes involved in initiating meiosis in yeast and mouse, *am1* also has a second downstream function, whereby it regulates the transition through a novel leptotene–zygotene checkpoint, a key step in early meiotic prophase. The *am1* gene encodes a plant-specific protein with an unknown biochemical function. The AM1 protein is diffuse in the nucleus during the initiation of meiosis and then binds to chromatin in early meiotic prophase I when it regulates the leptotene–zygotene progression.

chromosomes | plant development | genetics | recombination

A number of processes taking place in early stages of meiotic prophase I, such as chromosome pairing and recombination, are known as critical for the chromosome number reduction in meiosis (1). These processes are specific to meiosis and must be regulated by mechanisms that control the transition from the somatic (mitotic) cell cycle to meiosis. However, in contrast to the overall evolutionary conservation of the basic meiotic processes, the mechanisms controlling the transition to meiosis show substantial diversity among species (2). Initiation of meiosis has so far been studied in budding and fission yeasts and more recently also in mouse. In budding yeast, meiosis is initiated by expression of the *IME1* gene encoding a meiosis-specific transcription factor, which upregulates expression of the *IME2* gene (3). *IME2* encodes a protein kinase, directing the activation of a cyclin-dependent kinase that promotes entry into meiosis (4). In fission yeast, the master switch for meiosis initiation is the *Mei2* gene encoding an RNA-binding protein (5). The *Mei2* protein acts as a regulator of a complex system that sequesters meiotic transcript during vegetative growth (6). To initiate meiosis, *Mei2* turns off this regulatory system so that meiosis-specific RNAs are no longer intercepted. In the mouse, *Stras8*, a vertebrate-only gene, is required for the transition to meiosis (7, 8). *Stras8* expression is specifically induced in the germ cells by retinoic acid, a derivative of vitamin A, which is produced in the cells of the excretory organ of the embryo, and from there enters the gonads (8, 9).

Because the meiosis-initiating mechanisms in budding yeast, fission yeast, and mouse show little in common, it is likely that the transition to meiosis in other taxa is also regulated by species-dependent mechanisms. Although mammals and plants are both

multicellular eukaryotes with complex developmental patterns, the transition from the mitotic cell cycle to meiosis in plants takes place in a very different developmental context than in mammals. To elucidate the mechanisms regulating the transition to meiosis in maize we examined the *ameiotic1* (*am1*) mutants that disrupt the entry into meiosis (10, 11). We cloned the *am1* gene and determined that it encodes a plant-specific protein, indicating that meiosis in plants is initiated by a novel, plant-specific mechanism.

## Results

**Chromosome Behavior in *am1* Mutants.** The 5 known mutants at the maize *am1* locus can be grouped into 2 classes (Table 1). In the first class, comprising *am1-1*, *am1-2*, *am1-485*, and *am1-489*, male meiocytes, and in most cases also female meiocytes, do not enter a normal meiosis but instead undergo 2 or 3 rounds of equational division (11). The *am1-pra1* mutant allele confers a dramatically different phenotype, in which male and female meiocytes enter meiotic prophase and arrest during its early stages (10). To understand the nature of the nuclear divisions in *am1* mutants and discern whether the equational division is a mitosis or a defective meiosis, we used 3-dimensional deconvolution microscopy to examine chromosome behavior in wild-type meiocytes, as well as meiocytes from the *am1-1* and *am1-pra1* mutants. In wild-type meiocytes, prophase chromosomes are first visible in early leptotene as thin, elongated chromatin threads that surround a centrally located nucleolus. In late leptotene, the nucleolus moves to the side of the nucleus, and the bulk of the chromatin is located off-center during zygotene and most of pachytene (see chromosomes in Fig. 1A). In the *am1-1* mutant, chromosomes also surround a centrally located nucleolus, but they do not move to the side of the nucleus at any time during prophase (Fig. 1A). Overall, chromosomes in *am1-1* display behavior resembling the prophase of a mitotic division rather than meiosis. Male meiocytes in the *am1-2*, *am1-485*, and *am1-489* mutants also behave in this way (supporting information Fig. S1). In contrast to the other *am1* mutants, chromosomes in *am1-pra1*

Author contributions: W.P.P., C.-J.R.W., I.N.G., and W.Z.C. designed research; W.P.P., C.-J.R.W., I.N.G., J.M.S., L.S., O.H., and T.Z. performed research; W.F.S. contributed new reagents/analytic tools; W.P.P., C.-J.R.W., I.N.G., J.M.S., L.S., O.H., T.Z., and L.H. analyzed data; and W.P.P., C.-J.R.W., and W.Z.C. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

<sup>1</sup>W.P.P. and C.-J.R.W. contributed equally to this work.


<sup>2</sup>To whom correspondence may be addressed. E-mail: wp45@cornell.edu or zcande@berkeley.edu.

<sup>3</sup>Present address: Plant Reproduction and Development Laboratory, École Normale Supérieure de Lyon, France.

This article contains supporting information online at [www.pnas.org/cgi/content/full/0810115106/DCSupplemental](http://www.pnas.org/cgi/content/full/0810115106/DCSupplemental).

© 2009 by The National Academy of Sciences of the USA

**Table 1. The series of *am1* mutant alleles**

	Allele	Phenotype	
		male meiocyte	female meiocyte
	<b>Dominant</b>	wild type	meiosis
		<i>am1-pral</i>	arrest in meiotic prophase
		<i>am1-1</i>	equational division
		<i>am1-2</i>	equational division
		<i>am1-485</i>	equational division
	<b>Recessive</b>	<i>am1-489</i>	equational division

Arrow indicates the dominance relationship of the phenotype in female meiocytes (11). *am1-489* is the most recessive; *am1-pral* is the most dominant of all *am1* mutant alleles. *am1-2* and *am1-485* are codominant.

meiocytes exhibit behavior similar to that of chromosomes in early stages of a normal meiosis (Fig. 1A).

***am1* Mutants Show Defective Telomere Bouquet Formation.** Clustering of telomeres on a single site on the nuclear envelope, known as telomere bouquet formation, is a central event of a general repositioning of chromosomes during zygotene (12). Bouquet formation is thought to facilitate homologous chromosome pairing by bringing the chromosome ends together and confining chromosomes to a limited region within the nucleus. To examine bouquet formation, we used FISH with a telomere-specific probe (Fig. 1A). In wild-type maize meiocytes, telomeres attached to the nuclear envelope during leptotene, clustered in early zygotene, and remained clustered until late zygotene. In contrast, telomeres in *am1-1* mutant meiocytes did not attach to the nuclear envelope or cluster but were scattered throughout the nuclear volume (Fig. 1A). In *am1-pral* some bouquet formation was observed, although it did not progress to completion. A tight bouquet was present in approximately 8% of *am1-pral* meiocytes ( $n = 24$ ), whereas another 37.5% of meiocytes displayed a “loose” bouquet or multiple bouquet clusters (Fig. 1A). The remaining *am1-pral* meiocytes showed no bouquet.

**Homologous Chromosomes Do Not Pair in *am1* Mutants.** To determine whether homologous chromosomes pair in *am1* mutant meiocytes, we monitored chromosome behavior using FISH with the 5S rRNA locus probe. The maize genome contains a single 5S rRNA locus (13). In wild-type maize meiocytes, the 2 unpaired 5S foci are visible at the beginning of zygotene. At the beginning of pachytene, a single focus representing the homologously paired 5S loci is observed in each cell (Fig. 1B). In contrast, 2 separate foci were always visible in *am1-1* meiocytes (Fig. 1B), indicating that homologous chromosomes did not pair. In *am1-pral* meiocytes, the 5S rRNA locus probe also showed only unpaired 5S loci (Fig. 1B).

**Sister-Chromatid Cohesion and Meiotic Chromosome Axis Formation Are Defective in *am1* Mutants.** Correct chromosome segregation in meiosis requires meiosis-specific sister-chromatid cohesion. A meiosis-specific  $\alpha$ -kleisin Rec8 is a key component of the meiotic cohesin complex (14). Rec8 is recruited to chromosomes during the premeiotic S phase and is essential for both arm and centromere cohesion (15). To discern whether the meiotic cohesion complex is installed on chromosomes in *am1* mutants, we used an antibody against the AFD1 protein, the maize homologue of Rec8 (16). In wild-type meiocytes, the antibody exhibits weak diffuse staining in premeiotic interphase (Fig. S2A) (16). In leptotene, it shows strong staining mostly localized to chromatin (Fig. S2B). In zygotene and pachytene, AFD1 is seen as mostly continuous stretches along chromosome cores (Figs. 1C and S2C and D). We observed no AFD1 staining in mitotically dividing anther tapetal cells (Fig. S2F and G). Although AFD1 staining was present in nuclei of dividing root meristem cells (Fig. S2H and J), it was diffuse and

did not localize to chromatin, indicating that just the presence of AFD1 in the nucleoplasm is not sufficient to ensure meiosis-specific cohesion.


In *am1-1* mutant meiocytes, the AFD1 antibody showed weak diffuse staining that did not localize to chromatin (Fig. 1C) and was comparable, although less intense, to the staining in wild-type root meristem cells. In *am1-pral* meiocytes, the antibody showed strong staining that was partly localized to chromatin and similar to the staining in wild-type leptotene cells (Fig. 1C). These results indicate that AM1 is required for proper recruitment of AFD1 to meiotic chromosomes.

As meiocytes enter prophase I, the establishment of the leptotene chromosome structure relies at least partially on the installation of axial elements onto the chromosomes (17). In budding yeast, the HOP1 protein is an essential component of the axial element (18). ASY1, a meiosis-specific protein that associates with chromosome axes, has been proposed as the HOP1 homologue in *Arabidopsis* (19). We used an antibody against the *Arabidopsis* ASY1 protein to investigate the axial element assembly in *am1-1*, *am1-pral*, and wild-type meiocytes. In wild-type meiosis, the ASY1 antibody showed punctate staining along chromosomes when the axial elements form in leptotene (16, 19). In early zygotene, the signal became associated with the entire length of the axis (Fig. 1D). In *am1-1* meiocytes, we detected no ASY1 staining (Fig. 1D) suggesting that a proper chromosome axis does not form. In contrast, *am1-pral* meiocytes showed punctuate ASY1 signals and some continuous stretches as in wild-type leptotene meiocytes (Fig. 1D), suggesting partial formation of the chromosome axis.

***am1* Is Required for the Initiation and Progression of Meiotic Recombination.** Meiotic recombination, which includes the formation and repair of meiotic DSBs, is required for segregation of chromosomes in the first meiotic division and is one of the hallmarks of meiotic prophase I. To examine whether the meiotic recombination machinery is installed on chromosomes in *am1* mutant meiocytes, we analyzed nuclear distribution of the recombination protein RAD51, which is involved in repair of meiotic DSBs and forms distinct foci on chromosomes on the sites of DSBs (20). In somatic plant cells, the foci are absent (20). In wild-type maize meiocytes, RAD51 foci are first visible in early zygotene and reach their peak number of  $500 \pm 47$  per nucleus ( $n = 9$ ) at mid-zygotene (Fig. 1E). In contrast, we did not find any RAD51 foci in male meiocytes in plants homozygous for the *am1-1* and *am1-pral* mutations (Fig. 1E) or any of the other *am1* mutant alleles (data not shown). These data indicate that the *am1* gene is required for the installation of the recombination machinery on meiotic chromosomes.

To establish whether the absence of RAD51 foci was caused by the absence of meiotic DSBs, we used a modification of the TUNEL assay (21), which allows cytologic detection of the presence of meiotic DNA breaks. In wild-type maize meiocytes, the TUNEL





**Fig. 1.** Chromosome behavior in male meiocytes in wild-type maize and in the *am1-1* and *am1-pral* mutants. (A) Telomere bouquet analysis using FISH with the telomere probe (green). (B) Chromosome pairing analyzed using FISH with the 5S rRNA locus probe (cyan). (C) Installation of meiotic cohesin AFD1 (green), the maize Rec8 homologue, on chromosomes. The pictured wild-type meiocyte is in zygotene. (D) Installation of ASY1 (green), a meiosis-specific protein associated with chromosome axis. (E) Installation of the RAD51 recombination protein (green) on meiotic chromosomes. (F) Cytologic detection of meiotic DSB formation using a modified TUNEL assay. Green: TUNEL assay staining meiotic DSBs. (A–F) Red: DAPI-stained chromatin. (Scale bars, 10  $\mu$ m.)

assay produces a strong signal on chromosomes from leptotene through early pachytene (Fig. 1F). In contrast, we did not observe TUNEL staining in *am1-1* meiocytes, implying the absence of DSBs (Fig. 1F). In *am1-pral* meiocytes, the staining pattern resembled that of the wild-type meiocytes, indicating the presence of meiotic DSBs (Fig. 1F).

***am1* Regulates Meiocyte Cytoskeleton Organization.** During the G2 phase of the mitotic cell cycle, microtubules in plant cells are

**Table 2. Expression analysis of known meiotic genes in *am1* mutants in maize**

Gene	Wild-type meiosis	<i>am1-1</i>	<i>am1-pral</i>
<i>dmc1</i>	1,230 $\pm$ 88	485 $\pm$ 86	1,346 $\pm$ 150
<i>spo11-1</i>	217 $\pm$ 10	227 $\pm$ 32	185 $\pm$ 47
<i>hop2</i>	297 $\pm$ 81	297 $\pm$ 67	204 $\pm$ 90
<i>rad51A1</i>	517 $\pm$ 40	530 $\pm$ 129	341 $\pm$ 226
<i>rad51A2</i>	262 $\pm$ 55	175 $\pm$ 10	200 $\pm$ 14

Expression levels were measured using a custom GeneChip array and are in arbitrary units (mean  $\pm$  SE). Number of biologic replicates: wild-type meiosis  $n = 3$ , *am1-1*  $n = 4$ , and *am1-pral*  $n = 3$ .

organized into a dense preprophase band that marks the future cell division plane. The preprophase band is, however, absent from meiotic cells (22). It has been reported previously that the preprophase band is present in *am1-1* mutant meiocytes before they enter division (22). We have now determined that the *am1-2*, *am1-485*, and *am1-489* meiocytes also show the mitotic pattern of microtubule organization (Fig. S3). In contrast, in *am1-pral* mutant meiocytes, the microtubules form an extensive network filling the cytoplasm, similar to what is observed in wild-type meiocytes (Fig. S3). Thus, different alleles of *am1* regulate meiocyte microtubule behavior in a cell cycle stage-specific manner.

***am1* Affects Meiotic Expression of *dmc1*.** In addition to examining the cellular processes that take place during meiotic prophase I, we investigated whether the *am1* mutation affects expression of meiotic genes. To do this, we analyzed expression of *dmc1*, *hop2*, *rad51*, and *spo11-1* in wild-type male flowers undergoing meiosis and in male flowers of the *am1-1* and *am1-pral* mutants. Products of these 4 genes encode homologues of key meiotic recombination proteins (1). SPO11-1 is responsible for creating DSBs in chromosomal DNA that initiate meiotic recombination (23), and RAD51 are involved in repair of meiotic DSBs (24–27). DMC1, HOP2, and SPO11-1 act exclusively during meiosis (23–25). RAD51 is, in addition, thought to act in somatic DNA repair, although *rad51* mutants in maize and *Arabidopsis* show only defects in meiosis (20, 26, 27). In contrast to their strictly meiotic roles, the HOP2 and SPO11-1 genes are expressed in a range of somatic tissues, at least in *Arabidopsis* (23, 25), suggesting that some meiotic genes in plants are not regulated at the transcriptional level but perhaps at the protein level. RAD51 is also expressed in somatic tissues in maize and *Arabidopsis*, but this expression pattern is consistent with its role in somatic DNA repair. In contrast to HOP2, SPO11-1, and RAD51, the DMC1 gene in *Arabidopsis* shows exclusively meiotic expression (28).

To compare expression of *dmc1*, *hop2*, *rad51*, and *spo11-1* in wild-type and mutant flowers, we used a custom maize oligonucleotide GeneChip array containing 82,000 probe sets to maize genes and EST clusters. Among the tested genes, only *dmc1* showed a significantly reduced (by approximately 60%) level of expression in *am1-1* mutant flowers, although its expression in the *am1-pral* mutant was similar to that in wild-type flowers (Table 2). In contrast, *hop2*, *rad51*, and *spo11-1* did not exhibit significant changes in their expression levels in either *am1-1* or *am1-pral* as compared with the wild type. The most plausible explanation for these data is that genes showing meiosis-specific expression are under the control of *am1*, whereas *am1* has no effect on genes whose expression is not under meiosis-specific control.

***am1* Is a Plant-Specific Gene.** To further understand the regulation of meiotic prophase initiation and progression in maize, we cloned the *am1* gene. The *am1-485* and *am1-489* alleles were recovered in a screen of maize populations with a high activity of the *Mutator* (*Mu*) transposon. We discovered that a DNA fragment containing a terminal inverted repeat of the transposon cosegregated with the

mutant phenotype in a population of plants segregating for the *am1-489* allele. Analysis of the adjacent genomic region revealed a *Mu* copy inserted into the ninth exon of a 10-exon-long ORF. The putative *am1* gene encodes a 780-aa protein with a predicted molecular mass of 85.6 kDa. BLAST searches showed that the AM1 protein does not exhibit significant similarity to any protein with a known biochemical function. However, the AM1 sequence contains 2 putative coiled-coil domains (AA 260–287 and AA 441–547; Fig. 2A), which may be involved in protein–protein interactions.


We found sequence homologues of AM1 in several plant species but not outside of plants. The genomes of a number of monocot and dicot species, including maize, rice, and *Arabidopsis*, harbor 2 copies of *am1*-like genes. This is a result of a gene duplication event that, interestingly, took place independently in both the monocot and dicot lineages after the 2 groups diverged (Fig. 2B). One of the 2 AM1 homologues in *Arabidopsis*, SWITCH1 (SWI1)/DYAD, has been previously characterized and proposed to regulate meiotic chromosome structure and cohesion (29–32). AM1 and SWI1/DYAD share approximately 30% identity, mostly in the central portion of the protein. This region also shows strong similarity to several other plant proteins (Fig. S4), including MS1 and MMD1 in *Arabidopsis*, both involved in regulating meiotic gene expression (33, 34), and 2 predicted proteins in *Arabidopsis* and rice.

We identified the lesions in the *am1-pral*, *am1-485*, and *am1-489* alleles of *am1*. The *am1-pral* allele carries a point mutation that results in an arginine-to-tryptophan substitution at AA position 358 (Fig. 2A). This position is conserved in both monocots and dicots (Fig. S4). The *am1-485* allele carries a deletion in the 5' region of the gene introducing a premature stop codon, which would result in a severely truncated protein (Fig. 2A). The *am1-489* allele contains a *Mu* transposon insertion in the 3' end of the *am1* coding region (Fig. 2A), which would produce an abnormal protein consisting of truncated AM1 fused to a short protein resulting from translation of a portion of the *Mu* sequence. The *am1-485* and *am1-489* transcripts are stable, at least in anthers of the mutant plants (Fig. 2C). The *am1-1* and *am1-2* mutations have no lesions in the *am1* ORF, but anthers from the 2 mutants showed significantly reduced levels of the *am1* transcript (Fig. 2C), suggesting that the mutant lesions may be in the *am1* regulatory regions.

#### The AM1 Protein Is Localized to Chromosomes in Meiotic Prophase I.

Although phenotypes of the *am1* mutants are confined to meiosis, we found that the *am1* gene is expressed throughout the plant and at most stages of development (Fig. 2C). To better understand the spatial and temporal pattern of *am1* function, we generated an antibody against the central portion of the AM1 protein (AA 233–501). Western blot analysis showed that the protein was present in anthers but not in root, shoot, and leaf tissues (Fig. 2D), suggesting that translation and/or stability of the protein are developmentally regulated. To further understand the function of the protein, we conducted immunolocalization experiments to identify where it localizes during meiosis (Fig. 3). The protein was absent from the tapetal (somatic) cells in the anther and only present in the meiocytes, where it accumulated starting with the premeiotic interphase. During the premeiotic interphase, the anti-AM1 produced diffuse antibody staining inside the nucleus. During leptotene and early zygotene, AM1 localized to the chromatin, forming punctuated foci, with more elongated patches in some chromosome regions. The protein largely disappeared by late-zygotene, except for pericentromeric regions of 5 to 6 out of the 10 maize chromosomes, where it persisted until late pachytene.


**AM1 Does Not Localize to Chromosomes in *am1-pral* Meiocytes.** We examined accumulation of the AM1 protein in anthers of *am1* mutants using Western blot analysis (Fig. 2D) and immunolocalization (Fig. S5). In the *am1-pral* mutant, we found that the size and amount of the AM1 protein in the anthers was similar to that seen in wild type (Fig. 2D). The protein showed diffuse staining in nuclei



**Fig. 2.** (A) Diagram of the AM1 protein with marked positions of lesions in the *am1-pral*, *am1-485*, and *am1-489* mutant alleles. Also noted are positions of 2 predicted coiled-coil domains and the protein region used to raise the anti-AM1 antibody. (B) Bayesian reconstruction of phylogeny of AM1 homologues in plants. Numbers next to branches are clade credibility values. (C) Expression of the *am1* gene detected by RT-PCR in various wild-type maize tissues and in the 5 *am1* mutants. (D) Western blot analysis of AM1 protein accumulation in wild-type maize tissues and in the 5 *am1* mutants.

in premeiotic interphase, similar to that seen in wild type. However, during early leptotene, when chromosome threads become clearly visible, the defective AM1 protein remained diffuse in most *am1-pral* nuclei, and staining of AM1 in nucleoli was observed (Fig. S5).





**Fig. 3.** Immunolocalization of the AM1 protein in male meiocytes in wild-type maize. Red: DAPI-stained chromatin; green: AM1; purple: CENPC, which is a constitutive component of the inner kinetochore and marks centromeres. (Inset) Magnification of a centromeric region of one of the pachytene chromosomes, with CENPC identifying the kinetochore and AM1 staining in the pericentromeric region. (Scale bars, 10  $\mu\text{m}$ ; small bar Inset, 1  $\mu\text{m}$ .)

Only a fraction of cells (8 of 54) showed faint AM1 staining on chromosomes.

In the *am1-1*, *am1-2*, *am1-485*, and *am1-489* mutants, we did not detect the AM1 protein in Western blot analyses of meiotic anthers (Fig. 2D) or immunolocalization experiments on male meiocytes (Fig. S5). These results correlate well with previous observations that *am1-1*, *am1-2*, *am1-485*, and *am1-489* show similar mutant phenotypes in male meiosis (Table 1) (11). In female meiosis, *am1* alleles form an allelic series (Table 1). We hypothesize that the differences between the male and female meocyte defects are caused by a sex-specific regulation of *am1* translation and/or by a difference in how the AM1 protein acts in male and female meiocytes. However, we could not examine *am1* gene expression or AM1 protein accumulation in female meiocytes, because these cells in maize are not easily accessible to analysis.

## Discussion

Our analyses of *am1* mutants indicate that the *am1* gene plays 2 distinct roles in regulating meiosis. First, it is required for the transition from mitotic cell cycle to meiosis, and in its absence meiocytes continue undergoing mitotic divisions. Later, *am1* also regulates the transition from leptotene to zygotene.

***am1* Controls the Transition from the Mitotic Cell Cycle to Meiosis in Maize.** Previous observations suggested that chromosomes in *am1-1*, *am1-2*, *am1-485*, and *am1-489* meiocytes undergo an equational, “mitosis-like” division (10, 11, 35). However, they did not establish whether this equational division was mitosis or an aberrant meiosis. If the former is true, then the *am1* gene can be regarded as a major regulator of meiotic transition in maize. However, if the latter were the case, it would mean that the *am1* gene is involved in early meiotic processes but acts downstream from meiosis initiation. To discern between the 2 possibilities, we conducted an

in-depth analysis of several key events of meiotic prophase I in *am1-1* mutant meiocytes. These analyses established that chromosomes in *am1-1* mutant meiocytes (*i*) exhibit arrangement in the nucleus that resembles the prophase of mitosis rather than meiosis, (*ii*) do not show the meiosis-specific telomere bouquet formation, (*iii*) do not exhibit homologous chromosome pairing, (*iv*) lack installation of AFD1, (*v*) lack installation of ASY1, (*vi*) do not show formation of meiotic DSBs in chromosomal DNA and lack chromosomal foci of RAD51, (*vii*) exhibit cytoskeleton features typical for mitosis rather than meiosis, and (*viii*) show reduced levels of expression of a meiosis marker *dmc1*. Overall, this evidence suggests that the cell division in *am1-1* male meiocytes is a normal mitosis rather than an abnormal meiosis. Analyses of male meiocytes in the *am1-2*, *am1-485*, and *am1-489* mutants show that they exhibit chromosome behavior similar to that of *am1-1*, suggesting that they too undergo normal mitotic divisions. All these data indicate that *am1* is required in multiple early meiosis processes and is likely involved in the initiation of meiosis.

It has been suggested that the decision to initiate meiosis in budding yeast, fission yeast, and mouse is made during, or before, the premeiotic S phase (5, 7, 36, 37). Our observations suggest that the timing of meiosis initiation in maize may be similar. This hypothesis is supported by (*i*) the nuclear localization of the AM1 protein during the premeiotic interphase, (*ii*) the role of AM1 in recruiting AFD1/Rec8, (*iii*) interphase arrest in female meiocytes in the *am1-2*, *am1-485*, and *am1-489* mutants (11), and (*iv*) the presence of the mitosis-specific preprophase cortical microtubule band in *am1-1*, *am1-2*, *am1-485*, and *am1-489* meiocytes.

***am1* Controls Transition Through a Novel Leptotene–Zygotene Checkpoint.** In contrast to the other *am1* mutants, *am1-pra1* meiocytes enter meiosis and arrest during early meiotic prophase (10). Our analyses show now that the prophase arrest in *am1-pra1* occurs specifically at the leptotene–zygotene transition [the prezygotene stage (38)]. This timing is indicated by (*i*) the incomplete formation of the telomere bouquet, which in wild-type meiocytes starts at the end of leptotene and is completed in early zygotene, (*ii*) the absence of chromosome pairing, which normally starts in early zygotene, (*iii*) the patterns of AFD1 and ASY1 localization, which resemble those of the wild-type leptotene cells, (*iv*) the presence of meiotic DSBs, which are formed in leptotene in wild-type meiocytes, and (*v*) the absence of RAD51 foci, which are normally installed on maize chromosomes in early zygotene. On the basis of these analyses, we propose that AM1 has a second function in meiosis in regulating the meocyte progression through prezygotene. The leptotene–zygotene transition is a known major transition point in meiotic prophase, associated with a dramatic structural reorganization of chromosomes (38). The leptotene–zygotene transition arrest in *am1-pra1* suggests the presence of a novel checkpoint. The existence of this checkpoint is intriguing because plants are thought to lack the typical meiotic checkpoints (39). It is likely that the conserved region in the AM1 protein sequence, which stretches from AA 324 to 436 and includes the arginine residue mutated in *am1-pra1*, functions as a domain controlling meiotic prophase I progression. The arginine 358 residue is also required for the chromosomal localization of the AM1 protein in leptotene and zygotene.

**Evolution of Mechanisms Regulating Meiosis Initiation.** The mechanisms that initiate meiosis have been studied extensively in fission and budding yeasts and in the mouse (5–9). The mechanisms of meiosis initiation in these species show little in common, in terms of the cues triggering meiosis, the signaling cascades that initiate the transition to meiosis, or the sequence conservation of the proteins involved in meiosis initiation. The fact that AM1 homologues are only found in plants suggests that the mechanisms of meiosis initiation in plants share little with analogous mechanisms in yeast or mammals. Also unique to the mechanism of meiosis initiation in maize is the fact that the AM1 protein required for the transition

from the mitotic cell cycle to meiosis also plays a role in regulating the progression of meiotic prophase I once meiosis is initiated.

One of the 2 AM1 homologues in *Arabidopsis*, SWI1/DYAD, also affects the events of meiotic prophase I. However, its role in meiosis seems to be different from the role of AM1 (29–32). Chromosomes in *swi1/dyad* mutants in some cases undergo an equational-like segregation, but this division is an abnormal meiosis rather than a normal mitosis, based on chromosome behavior, installation of the meiosis-specific cohesin Rec8, and presence of the meiosis-specific expression of *DMC1* (30, 31). This suggests that the SWI1/DYAD protein affects early stages of meiosis that are downstream from meiosis initiation but does not act directly to initiate meiosis in *Arabidopsis*. In addition, the SWI1/DYAD protein is present in the nucleus only during the premeiotic interphase, and it is not known to have a role in regulating progression of meiotic prophase I (30–32). Characterizing *At5g23610*, the second *am1*-like gene in *Arabidopsis* and the second *am1* homologue in maize, should clarify the similarities and differences between the meiosis initiation mechanisms in the 2 species. Overall, our data imply that meiosis-initiating mechanisms show substantial variation not only between distant taxa but also among relatively more closely related groups like monocots and dicots.

## Materials and Methods

**am1 Mutants.** All *am1* mutant alleles were reported previously (10, 11). The *am1-485* and *am1-489* mutants were isolated and initially characterized at the University of North Dakota.

**Microscopy.** Immunolocalization using antibodies against RAD51 (40), AFD1 (16), ASY1 (19),  $\alpha$ -tubulin, and AM1 was performed as previously described (16, 41). FISH and meiotic DSB detection were also performed as published (12, 21), except

that the meiocyte-containing polyacrylamide pads were permeabilized in 0.1% sodium citrate + 0.1% Triton X-100 before the TUNEL reaction.

**Analysis of Meiotic Gene Expression.** Expression analysis was performed using a Syngenta-designed maize GeneChip genome array (SYNG007a520046). The array contains 82,661 probe sets, designed on the basis of 87,572 maize transcripts, of which 50,058 had protein similarity and 37,514 were unknown. Staged flowers of wild-type, *am1-1*, and *am1-pral* plants were collected and immediately frozen in liquid nitrogen to prevent expression of stress-response genes. Preparation of labeled cDNA, array hybridization, and data acquisition were carried out as previously described (42).

**Cloning the am1 Gene.** Ligation-based expression analysis display (LEAD) (45) was used to PCR-amplify genomic DNA fragments adjacent to *Mutator* transposon insertions in the *am1-489* mutant. Screening of a population of 82 plants from families segregating for the *am1-489* mutation identified a fragment that co-segregated with the mutant gene. The sequence of the rest of the *am1* ORF was obtained using 5' and 3' RACE. The GenBank accession number for the *am1* gene is DQ663482.

**RT-PCR.** RT-PCR was performed with the SuperScript One-Step RT-PCR kit (Invitrogen). To control for the amount of the RNA template, we ran 1.0  $\mu$ g of each RNA sample in a 1% agarose gel and compared the intensities of the rRNA bands between samples.

**AM1 Antibody Production.** A partial *am1* cDNA was cloned into the pGEX-4T-3 plasmid. The fusion protein was expressed in *Escherichia coli* BL21 cells and purified with the GST purification kit (Amersham Biosciences). The tag was cleaved by thrombin digestion, and the purified antigen was used to raise an antibody in rabbits (Covance).

**ACKNOWLEDGMENTS.** We thank Tomoko Ogawa, Chris Franklin, and Kelly Dawe for antibodies, Charles Chilcott for technical assistance, and members of the Pawlowski and Cande laboratories for comments on the manuscript. This study was supported by National Institutes of Health, National Science Foundation, Syngenta, and Cornell University.

- Ronceret A, Sheehan MJ, Pawlowski WP (2007) Chromosome dynamics in meiosis. *Cell Division Control in Plants*, eds Verma DPS, Hong Z (Springer-Verlag, Heidelberg), pp 103–124.
- Pawlowski WP, Sheehan MJ, Ronceret A (2007) In the beginning: The initiation of meiosis. *Bioessays* 29:511–514.
- Chu S, et al. (1998) The transcriptional program of sporulation in budding yeast. *Science* 282:699–705.
- Benjamin KR, Zhang C, Shokat KM, Herskowitz I (2003) Control of landmark events in meiosis by the CDK Cdc28 and the meiosis-specific kinase Ime2. *Genes Dev* 17:1524–1539.
- Marston AL, Amon A (2004) Meiosis: Cell-cycle controls shuffle and deal. *Nat Rev Mol Cell Biol* 5:983–997.
- Harigaya Y, et al. (2006) Selective elimination of messenger RNA prevents an incidence of untimely meiosis. *Nature* 442:45–50.
- Baltus AE, et al. (2006) In germ cells of mouse embryonic ovaries, the decision to enter meiosis precedes premeiotic DNA replication. *Nat Genet* 38:1430–1434.
- Koubova J, et al. (2006) Retinoic acid regulates sex-specific timing of meiotic initiation in mice. *Proc Natl Acad Sci USA* 103:2474–2479.
- Bowles J, et al. (2006) Retinoid signaling determines germ cell fate in mice. *Science* 312:596–600.
- Golubovskaya I, Grebennikova ZK, Avalkina NA, Sheridan WF (1993) The role of the *ameiotic1* gene in the initiation of meiosis and in subsequent meiotic events in maize. *Genetics* 135:1151–1166.
- Golubovskaya I, Avalkina N, Sheridan WF (1997) New insights into the role of the maize *ameiotic1* locus. *Genetics* 147:1339–1350.
- Golubovskaya IN, Harper LC, Pawlowski WP, Schichnes D, Cande WZ (2002) The *pam1* gene is required for meiotic bouquet formation and efficient homologous synapsis in maize (*Zea mays*, L.). *Genetics* 162:1979–1993.
- Mascia PN, Rubenstein I, Phillips RL, Wang AS, Xiang LZ (1981) Localization of the 5S rRNA genes and evidence for diversity in the 5S rDNA region of maize. *Gene* 15:7–20.
- Watanabe Y (2005) Sister chromatid cohesion along arms and at centromeres. *Trends Genet* 21:405–412.
- Watanabe Y, Nurse P (1999) Cohesin Rec8 is required for reductional chromosome segregation at meiosis. *Nature* 400:461–464.
- Golubovskaya IN, et al. (2006) Alleles of *afd1* dissect REC8 functions during meiotic prophase I. *J Cell Sci* 119:3306–3315.
- Zickler D, Kleckner N (1999) Meiotic chromosomes: Integrating structure and function. *Annu Rev Genet* 33:603–754.
- Hollingsworth NM, Byers B (1989) HOP1: A yeast meiotic pairing gene. *Genetics* 121:445–462.
- Armstrong SJ, Caryl AP, Jones GH, Franklin FC (2002) Asy1, a protein required for meiotic chromosome synapsis, localizes to axis-associated chromatin in *Arabidopsis* and *Brassica*. *J Cell Sci* 115:3645–3655.
- Franklin AE, et al. (1999) Three-dimensional microscopy of the Rad51 recombination protein during meiotic prophase. *Plant Cell* 11:809–824.
- Pawlowski WP, et al. (2004) Coordination of meiotic recombination, pairing, and synapsis by PHS1. *Science* 303:89–92.
- Staiger CJ, Cande WZ (1992) *Ameiotic*, a gene that controls meiotic chromosome and cytoskeletal behavior in maize. *Dev Biol* 154:226–230.
- Grelon M, Vezon D, Gendrot G, Pelletier G (2001) *AtSPO11-1* is necessary for efficient meiotic recombination in plants. *EMBO J* 20:589–600.
- Couteau F, et al. (1999) Random chromosome segregation without meiotic arrest in both male and female meiocytes of a *dmc1* mutant of *Arabidopsis*. *Plant Cell* 11:1623–1634.
- Schommer C, Beven A, Lawrenson T, Shaw P, Sablowski R (2003) AHP2 is required for bivalent formation and for segregation of homologous chromosomes in *Arabidopsis* meiosis. *Plant J* 36:1–11.
- Li W, et al. (2004) The *Arabidopsis* ATRAD51 gene is dispensable for vegetative development but required for meiosis. *Proc Natl Acad Sci USA* 101:10596–10601.
- Li J, et al. (2007) Functional analysis of maize RAD51 in meiosis and double-strand break repair. *Genetics* 176:1469–1482.
- Klimyuk VI, Jones JDG (1997) ATRAD51, the *Arabidopsis* homologue of the yeast *DMC1* gene: Characterization, transposon-induced allelic variation and meiosis-associated expression. *Plant J* 11:1–14.
- Mercier R, et al. (2001) SWITCH1 (SWI1): A novel protein required for the establishment of sister chromatid cohesion and for bivalent formation at meiosis. *Genes Dev* 15:1859–1871.
- Mercier R, et al. (2003) The meiotic protein SWI1 is required for axial element formation and recombination initiation in *Arabidopsis*. *Development* 130:3309–3318.
- Agashe B, Prasad CK, Siddiqi I (2002) Identification and analysis of *DYAD*: A gene required for meiotic chromosome organization and female meiotic progression in *Arabidopsis*. *Development* 129:3935–3943.
- Siddiqi I, Ganesh G, Grossniklaus U, Subbiah V (2000) The *dyad* gene is required for progression through female meiosis in *Arabidopsis*. *Development* 127:197–207.
- Wilson ZA, Morroll SM, Dawson J, Swarup R, Tighe PJ (2001) The *Arabidopsis* MALE STERILITY1 (*MS1*) gene is a transcriptional regulator of male gametogenesis, with homology to the PHD-finger family of transcription factors. *Plant J* 28:27–39.
- Yang X, Makaroff CA, Ma H (2003) The *Arabidopsis* MALE MEIOTIC DEATH1 gene encodes a PHD-finger protein that is required for male meiosis. *Plant Cell* 15:1281–1295.
- Rhoades MM (1956) Genic control of chromosomal behavior. *Maize Genetics Cooperation Newsletter* 30:38–42.
- Watanabe Y, Yokobayashi S, Yamamoto M, Nurse P (2001) Pre-meiotic S phase is linked to reductional chromosome segregation and recombination. *Nature* 409:359–363.
- Borde V, Goldman AS, Lichten M (2000) Direct coupling between meiotic DNA replication and recombination initiation. *Science* 290:806–809.
- Dawe RK, Sedat JW, Agard DA, Cande WZ (1994) Meiotic chromosome pairing in maize is associated with a novel chromatin organization. *Cell* 76:901–912.
- Caryl AP, Jones GH, Franklin FC (2003) Dissecting plant meiosis using *Arabidopsis thaliana* mutants. *J Exp Bot* 54:25–38.
- Terasawa M, Shinohhara A, Hotta Y, Ogawa H, Ogawa T (1995) Localization of RecA-like protein in chromosomes of the lily at various meiotic stages. *Genes Dev* 9:925–934.
- Pawlowski WP, Golubovskaya IN, Cande WZ (2003) Altered nuclear distribution of recombination protein RAD51 in maize mutants suggests involvement of RAD51 in the meiotic homology recognition. *Plant Cell* 15:1807–1816.
- Guimil S, et al. (2005) Comparative transcriptomics of rice reveals an ancient pattern of response to microbial colonization. *Proc Natl Acad Sci USA* 102:8066–8070.
- Li BY, et al. (2002) Analysis of differential gene expression by ligation specificity-based transcript profiling. *Omic* 6:175–185.