

Published in final edited form as:

Biochemistry. 2009 February 10; 48(5): 850–862. doi:10.1021/bi801897p.

Effects of Ca²⁺, Mg²⁺ and Myristoylation on GCAP1 Structure and Stability†

Sunghyuk Lim¹, Igor Peshenko², Alexander Dizhoor², and James B. Ames^{1,*}

¹Department of Chemistry, University of California, Davis, CA 95616

²Basic Sciences, Pennsylvania College of Optometry, Salus University, Elkins Park, PA 19027.

Abstract

Guanylyl cyclase activating protein 1 (GCAP1), a member of the neuronal calcium sensor (NCS) subclass of the calmodulin superfamily, confers Ca²⁺-dependent activation of retinal guanylyl cyclase (RetGC) during phototransduction in vision. Here we analyze the energetics of Ca²⁺ and Mg²⁺ binding to the individual EF-hands, characterize metal-induced conformational changes, and evaluate structural effects of myristoylation as studied by isothermal titration calorimetry (ITC), differential scanning calorimetry (DSC) and NMR. GCAP1 binds cooperatively to Ca²⁺ at EF3 and EF4 ($\Delta H_{EF3} = -3.5$ kcal/mol and $\Delta H_{EF4} = -0.9$ kcal/mol) with nanomolar affinity ($K_{EF3} = 80$ nM and $K_{EF4} = 200$ nM), and a third Ca²⁺ binds entropically at EF2 ($\Delta H_{EF2} = +3.1$ kcal/mol and $K_{EF2} = 0.9$ μ M). GCAP1 binds functionally to Mg²⁺ at EF2 ($\Delta H_{EF2} = +4.3$ kcal/mol and $K_{EF2} = 0.7$ mM) required for RetGC activation. Ca²⁺ and/or Mg²⁺ binding to GCAP1 dramatically alter DSC and NMR spectra, indicating metal-induced protein conformational changes in EF2, EF3 and EF4. Myristoylation of GCAP1 does not significantly alter its metal binding energetics or NMR spectra, suggesting that myristoylation does not influence the structure of the metal-binding EF-hands. Myristoylation also has almost no effect on protein folding stability measured by DSC. NMR resonances of myristate attached to GCAP1 are exchange broadened, upfield shifted and insensitive to Ca²⁺, consistent with the myristoyl group being sequestered inside the protein as seen in the crystal structure. We conclude that the protein environment near the myristate is not influenced by Mg²⁺ or Ca²⁺ binding, but instead is constitutively dynamic and may play a role in promoting GCAP1 interactions with the cyclase.

Keywords

GCAP1; calcium; NMR; EF-hand; RetGC; phototransduction; retina; neuronal calcium sensor

Guanylyl cyclase activating proteins (GCAPs) belong to the neuronal calcium sensor (NCS) branch of the calmodulin superfamily (1-3) and regulate Ca²⁺-sensitive activity of retinal guanylyl cyclase (RetGC) in rod and cone cells (4,5). Phototransduction in retinal rods and cones is modulated by intracellular Ca²⁺ sensed by GCAPs (6,7) and defects in Ca²⁺ signaling by GCAPs are linked to retinal diseases (8). Light excitation of photoreceptor cells triggers a

†This work was supported by NIH Grants NS045909, EY012347 (J.B.A.) and RR11973 (UC Davis NMR).

*To whom correspondence should be addressed. Tel: (530) 752–6358. Fax: (530) 752–8995. E-mail: ames@chem.ucdavis.edu.

SUPPORTING INFORMATION AVAILABLE

Summary of DSC data of myristoylated and unmyristoylated recoverin; fluorescence spectra of myristoylated and unmyristoylated GCAP1; ¹³C(F3)-filtered/¹³C(F1)-edited NOESY-HMQC spectrum of GCAP1 containing a ¹³C-labeled myristoyl group; and a plot of the relative NMR intensities of amide ¹H resonances from Gly 105 (EF3) and Gly 149 (EF4) vs. Ca²⁺ are available free of charge via the Internet at <http://pubs.acs.org>.

phototransduction cascade, leading to the closure of cGMP-gated channels, which hyperpolarizes the plasma membrane and generates a neural signal (9). The recovery of the dark state requires activation of RetGC to restore the cytosolic cGMP level. The activity of RetGC is Ca^{2+} -sensitive (10) and is mediated by GCAPs (4,5,7). Channel closure during phototransduction blocks the entry of Ca^{2+} and lowers the cytosolic Ca^{2+} concentration from ~250 nM in the dark down to ~25 nM in the light (11). This drop in Ca^{2+} causes the formation of Ca^{2+} -free/ Mg^{2+} -bound GCAPs that activate RetGC (12), whereas Ca^{2+} -bound GCAPs inhibit RetGC at high Ca^{2+} levels generated in the dark (7).

The GCAPs (GCAP1 (5), GCAP2 (13), GCAP3 (14) and GCAP4–8 (15)) are all ~200-residue proteins containing a covalently attached N-terminal myristoyl group and four EF-hand motifs (Fig. 1). Mg^{2+} binds to three EF-hands (EF2, EF3 and EF4) when cytosolic Ca^{2+} levels are low and Mg^{2+} -bound GCAP1 activates RetGC (12,16). Ca^{2+} also binds functionally to these three EF-hands (12,17) when Ca^{2+} levels are high in the dark. The first EF-hand (EF1) is unable to bind Ca^{2+} or Mg^{2+} due to conserved substitutions in the binding loop (Cys29 and Pro30 in GCAP1). The x-ray crystal structure of Ca^{2+} -bound GCAP1 (18) and NMR structure of GCAP2 (19) clearly showed that Ca^{2+} is bound at EF2, EF3 and EF4. In addition, the N-terminal myristoyl group in GCAP1 is buried inside the Ca^{2+} -bound protein flanked by hydrophobic residues at the N- and C-termini (see italicized residues in Fig. 1). The structure of the physiologically active form of GCAPs (Mg^{2+} -bound/ Ca^{2+} -free state) is currently unknown.

What are the Ca^{2+} - and Mg^{2+} -dependent conformational changes in GCAPs that promote activation of RetGC? Recoverin is the only NCS protein whose structure is known in both the Ca^{2+} -free and Ca^{2+} -bound states (Fig. 1) (20,21). Ca^{2+} -free recoverin contains a highly sequestered myristoyl group buried inside the protein that interacts intimately with residues from EF1, EF2 and EF3 (22,23). Ca^{2+} binding at EF2 and EF3 leads to large overall conformational changes in recoverin that promotes the extrusion of the fatty acyl group outward, enabling it to interact with membrane targets (20,24). Since the target for GCAPs (RetGC) is a membrane protein, we wondered if GCAPs might undergo a similar Ca^{2+} -induced conformational change and if Ca^{2+} and/or Mg^{2+} binding to GCAPs alter the protein environment around the myristoyl group. Here, we perform ITC, DSC and NMR studies on GCAP1 to analyze the thermodynamics of Ca^{2+} and Mg^{2+} binding to the individual EF-hands, characterize metal-induced protein conformational changes, and evaluate the Ca^{2+} - and Mg^{2+} -dependent structural environment around the myristoyl group.

EXPERIMENTAL PROCEDURES

GCAP1 Protein Expression and Purification

Plasmids (pET11d) encoding wild-type GCAP1 and its mutants for disabling both Ca^{2+} and Mg^{2+} binding (D68G/E75Q (EF2-), D100N/D102G (EF3-), D144N/D148G (EF4-), and D100N/D102G/D144N/D148G (EF34-)) were prepared as described previously (12). All plasmids contain the D6S mutation for the recognition of yeast N-myristoyl-transferase (NMT) needed for expressing recombinant myristoylated GCAP1 in *E. coli*. Plasmid vector (pET11d) harboring GCAP1 or its mutants were co-transformed into BL21(DE3) cells with or without pBB131 vector encoding NMT. The bacterial cells were typically pre-cultured until optical density at 600 nm (A_{600}) reached 1.0 in 20 ml Luria-Bertani (LB) medium at 37 °C with antibiotics (100 µg/ml ampicillin or along with 50 µg/ml kanamycin). They were then inoculated into 1 L LB medium with the antibiotics and grown until $A_{600} = 1.3$. GCAP1 and NMT proteins were co-expressed upon adding isopropyl- β -D-1-thiogalactopyranoside (IPTG) into the cell culture to a final concentration of 0.5 mM at 25 °C or at 37 °C for 4 ~ 16 hours. GCAP1 was expressed in either the soluble fraction or as inclusion bodies by adjusting the induction temperature: Soluble GCAP1 expression occurs at 25 °C, whereas GCAP1

expression in inclusion bodies occurred at 37 °C. For the expression of myristoylated proteins, 5 mg/L (final concentration) myristic acid (from 5mg/ml ethanol stock) was added to the culture 20 minutes prior to the induction. The purification of myristoylated GCAP1 and mutants (EF2⁻, EF3⁻, EF4⁻ and EF34⁻) expressed in inclusion bodies has been described previously (12).

The purification of recombinant unmyristoylated GCAP1 (expressed in soluble lysate) was as follows. Cells were harvested by centrifugation and the cell pellet was resuspended and sonicated in the lysis buffer containing 20 mM Tris-HCl (pH 7.5), 0.1 M KCl, 2 mM EGTA, 1 mM DTT, 10% glycerol and 0.1 mM PMSF. Cell lysate was centrifuged down at 100,000 g by ultra centrifugation. The supernatant was mixed with CaCl₂ to a final concentration of 3 mM and then applied onto butyl-Sepharose column (HiPrep 16/10 Butyl FF, Amersham) which was pre-equilibrated with buffer A containing 20 mM Tris (pH 7.5), 0.3 M KCl, 2 mM CaCl₂, 1 mM MgCl₂ and 1 mM DTT. The elution of the protein was done by removing Ca²⁺ ions by adding the buffer B containing 20 mM Tris (pH 7.5), 0.1 M KCl, 2 mM EGTA and 1 mM DTT. The fractions containing target proteins were pooled and diluted 3-fold with ddH₂O lowering ionic strength for anion exchange column application. For myristoylated GCAP1, a similar purification procedure was followed except that the lysis buffer contained 0.5 M ammonium sulfate instead of 0.1 M KCl. This modification was necessary to cause more efficient binding and elution of myristoylated GCAP1 using Butyl-Sepharose chromatography. GCAP1 protein samples were then further purified by DEAE anion exchange chromatography (HiTrap 5ml DEAE FF, Amersham) pre-equilibrated with 20 mM Tris (pH 7.5) buffer containing 1 mM EGTA and 1 mM DTT. GCAP1 was eluted using a linear NaCl gradient (0 to 0.5 M over 20 column volumes). Lastly, proteins were purified by a size exclusion chromatography (HiLoad 26/60 Superdex 200, Amersham) pre-equilibrated with 20 mM Tris (pH 7.5) buffer containing 1 mM EGTA and 1 mM DTT. The final purity (>95%) of the GCAP1 sample was verified by SDS-PAGE and mass spectrometry.

Metal-free GCAP1 was prepared as described previously (16). Briefly, the final purified GCAP1 samples in the presence of 5 mM EGTA were concentrated 10-fold (Amicon-10, 10 kDa cut-off), diluted 10-fold with de-calcified buffer (prepared as described previously (25, 26)) and then concentrated again. This procedure was repeated four times to completely exchange GCAP1 into de-calcified and EGTA-free buffer used in the metal-binding studies. The free Ca²⁺ concentration in the apo-GCAP1 sample was verified to be less than 10 nM using fluo-3 fluorescent indicator dye analysis (27) and atomic absorption measurements. The lack of Ca²⁺ in apo GCAP1 was also verified by monitoring the absence of downfield-shifted amide proton NMR resonances of Gly 69, Gly 105 and Gly 149 that report on Ca²⁺ binding to EF2, EF3 and EF4, respectively. The apo-GCAP1 samples were also verified by NMR to not contain any lingering EGTA.

Isothermal titration calorimetry (ITC)

All ITC experiments were performed using a VP-ITC calorimeter (MicroCal) and the data were processed with the Origin 7 software package from MicroCal as described previously (28). GCAP1 samples (myristoylated, unmyristoylated, wildtype and mutants) for ITC studies were prepared in 20 mM Tris buffer (pH 7.5), 100 mM NaCl and 1 mM β-mercaptoethanol. The protein concentration was either 25 μM or 50 μM determined by measuring optical density at 280 nm. A series of 5 μL aliquots of 2 mM CaCl₂ were injected into the protein sample (1.6 mL) in the presence and absence of 2 mM Mg²⁺ ions and corresponding heat signals were monitored calorimetrically. For the Mg²⁺ titration, 10 μL aliquots of 40 mM MgCl₂ were titrated into the protein sample. All titrations were typically performed at 30 °C. Additional titrations were also performed at 23, 25 and 27 °C for measuring ΔC_p (29,30). The baseline from a buffer blank titration was subtracted from the raw data. A sequential binding sites model

was used to fit ITC data using a non-linear least squares minimization method (31) and calculate the dissociation constant (K_d) and enthalpy change (ΔH) for each site.

Fluorescence

Tryptophan fluorescence emission spectra of myrGCAP1 were observed with a Varian Cary Eclipse spectrometer. Protein samples were prepared in 20 mM Tris-HCl (pH 7.5) buffer containing 5 mM DTT in a 1 cm path-length cell with increasing amounts of guanidine HCl (0 – 6 M) in the presence and absence of 5 mM Ca^{2+} or Mg^{2+} ions. Protein concentration was 10 μM . The emission spectra were recorded in range of 300 – 500 nm with the excitation wavelength at 295 nm at room temperature. The protein samples were incubated at room temperature for 30 min before each measurement.

NMR spectroscopy

For ^1H - ^{15}N -HSQC experiments, samples of myristoylated and unmyristoylated GCAP1 were prepared in 5 mM Tris- d_{11} -HCl (pH 7.4) buffer containing 10 % D_2O , 1 mM Dithiothreitol- d_{10} (DTT) and either 5 mM CaCl_2 or 5 mM MgCl_2 . For ^1H - ^{13}C HMQC experiments, NMR sample were prepared in the similar condition as above except the sample was prepared in 100 % D_2O . The reading of the pH meter was not corrected for the deuterium isotope effect. The sample concentration was typically around 0.3 mM.

All NMR spectra were acquired using a Bruker Advance 600 MHz spectrometer equipped with a triple resonance cryo-probe and z gradient. Typically, spectra were obtained at 300 K, 310 K and 320 K with 8 ~ 32 scans per increment. For ^{15}N -labeled myristoylated GCAP1 (apo-, Mg^{2+} -bound or Ca^{2+} -bound form), two dimensional ^1H - ^{15}N -HSQC spectra with total number of 256 increments were acquired. For samples of GCAP1 with ^{13}C -labeled myristic group covalently attached to N-terminus, 2D ^1H - ^{13}C -HMQC spectra were recorded with 1024 \times 256 points. Three dimensional ($^{13}\text{C}/\text{F1}$)-edited $^{13}\text{C}/\text{F3}$ -filtered HMQC-NOESY spectra (120 ms mixing time) were obtained with 1024 \times 64 \times 180 points (23,32).

Differential scanning calorimetry

A VP-DSC calorimeter from MicroCal was used for all DSC measurements and the data were processed with the Origin 7 software package from MicroCal as described previously (28). Scanning was done in the temperature range of 10 – 120 $^\circ\text{C}$ at a scan rate of 60 $^\circ\text{C}/\text{h}$. A buffer baseline was subtracted from each scan. Myristoylated and unmyristoylated GCAP1 with concentrations of 50 ΔM were prepared in 20 mM Tris buffer (pH 7.5) containing 100 mM NaCl and 1 mM β -mercaptoethanol with/without 2 mM CaCl_2 or MgCl_2 . Samples were degassed before each scan.

RESULTS

Thermodynamics of Ca^{2+} -binding to GCAP1

Isothermal titration calorimetry (ITC) was used in this study to analyze the energetics of Ca^{2+} and Mg^{2+} binding to the individual EF-hands of GCAP1 (Fig. 1). Previous equilibrium binding studies revealed that GCAP1 binds Ca^{2+} at EF2, EF3 and EF4 in the nanomolar range (16), but precise values of dissociation constants for the individual EF-hands were difficult to resolve because their binding appeared somewhat overlapped. ITC can resolve dissociation constants (K_d) of multiple sites on the basis of differences in their binding enthalpy (ΔH). Hence, two or more sites with similar dissociation constants can be resolved if their binding enthalpies are sufficiently different or vice-versa.

The Ca^{2+} binding properties of both myristoylated and unmyristoylated GCAP1 were monitored by ITC (Fig. 2). N-terminal myristoylation of GCAP1 had a very small influence

on Ca^{2+} binding (see circles in Figs. 2A-B) unlike that of recoverin (26,33), and all subsequent analyses below were performed on myristoylated GCAP1. Titration of CaCl_2 into myristoylated GCAP1 resulted in a binding isotherm that is multiphasic and best fit by a 3-site sequential binding model (solid lines in Figs. 2A-B and Table 1), consistent with Ca^{2+} binding at EF2, EF3 and EF4 as suggested by previous fluorescence equilibrium binding studies (12, 16). The ITC Ca^{2+} -binding isotherm of apo-GCAP1 (absence of Mg^{2+} , see Fig. 2A) exhibited exothermic binding of two Ca^{2+} ($\Delta H_1 = -3.5$ kcal/mol and $\Delta H_2 = -0.9$ kcal/mol) in the nanomolar range followed by endothermic binding of one Ca^{2+} ($\Delta H = +3.1$ kcal/mol) with much lower affinity ($K_d = 0.9$ μM). The high resolution ITC data clearly indicated three binding sites having distinct values of ΔH and K_d . To assign each ITC binding phase as binding by a particular EF-hand, we have constructed various mutants that disable functional Ca^{2+} binding to the individual EF-hands (Fig. 1): D68G/E75Q (EF2-), D100N/D102G (EF3-), D144N/D148G (EF4-), and D100N/D102G/D144N/D148G (EF34-). In each mutant, a negatively charged Glu or Asp at the beginning and end of the EF-hand loop has been substituted with a neutral residue (Gln or Asn). Also the second negatively charged residue in the loop (Asp) was replaced by Gly. These substitutions dramatically lower the Ca^{2+} and/or Mg^{2+} binding affinity of the respective EF-hand outside of the physiological range as previously described in detail (16,34).

The ITC isotherms for wildtype GCAP1 and each of the EF-hand mutants were fit to a sequential binding model (solid lines in Fig. 2) and optimal Ca^{2+} -binding parameters (ΔH and K_d) are shown in Table 1. All ITC Ca^{2+} binding measurements were performed both in the presence (Fig. 2B) or absence (Fig. 2A) of physiological Mg^{2+} levels (2 mM Mg^{2+}) at 30 °C. We show that Mg^{2+} has relatively small effects on Ca^{2+} binding, indicating that GCAP1 binds selectively to Ca^{2+} with at least 500-fold preference over Mg^{2+} (Table 1). The ITC Ca^{2+} -binding isotherm of the EF2- mutant (Fig. 2C) exhibited exothermic binding of 2 Ca^{2+} ions but lacked the endothermic phase seen in wildtype, indicating that EF2 must be the low affinity endothermic site. Correspondingly, the ITC isotherm of EF34- (Fig. 2E) exhibited only endothermic binding ($\Delta H = +2.1$ kcal/mol), further confirming that EF2 is the endothermic site. The ITC Ca^{2+} -binding isotherm of EF4- (Fig. 2D) exhibited an exothermic phase ($\Delta H = -3.4$ kcal/mol) assigned as Ca^{2+} binding at EF3 followed by a lower affinity, endothermic phase ($K_d = 69$ μM and $\Delta H = +0.8$ kcal/mol) assigned above to EF2. By the process of elimination, the lower affinity exothermic site in the wildtype isotherm ($\Delta H = -0.9$ kcal/mol and $K_d = 200$ nM), therefore, must be assigned to EF4.

Our ITC Ca^{2+} -binding analysis summarized in Table 1 reveals that EF2 is an endothermic site ($K_{\text{EF2}} = 0.9$ μM and $\Delta H_{\text{EF2}} = +3.1$ kcal/mol); EF3 and EF4 exhibit exothermic binding in the nanomolar range ($K_{\text{EF3}} = 80$ nM and $\Delta H_{\text{EF3}} = -3.5$ kcal/mol; $K_{\text{EF4}} = 200$ nM and $\Delta H_{\text{EF4}} = -0.9$ kcal/mol). The Ca^{2+} -binding dissociation constants of EF3 and EF4 are in the physiological range of free Ca^{2+} concentration measured in rod outer segments (11) and are similar to binding constants measured previously by direct equilibrium Ca^{2+} binding assays (12,16). The relatively low apparent Ca^{2+} -binding affinity of EF2 measured by ITC ($K_{\text{EF2}} = 0.9$ μM) is ~ 10 -fold weaker than the intrinsic binding affinity determined by fluorescent Ca^{2+} binding assays (16). The discrepancy is explained in part by a protein conformational change in GCAP1 that is thermodynamically coupled to Ca^{2+} binding at EF2. The intrinsic Ca^{2+} binding at EF2 measured by the fluorescence binding assay ($K_a = 10^7$ M^{-1}) coupled to an unfavorable conformational change ($K_{\text{eq}} \sim 10^{-1}$) probed by ITC yields an overall equilibrium constant of $K_{\text{tot}} = K_a * K_{\text{eq}} \sim 10^6$ M^{-1} , consistent with the overall K_d measured by ITC. Hence, the overall low apparent binding affinity of EF2 suggests that Ca^{2+} binding at this site drives an unfavorable protein conformational change. Consistent with this conformational change are large Ca^{2+} -induced NMR chemical shift changes for residues in EF2 (see below) and the relatively steep temperature dependence of ΔH_{EF2} ($\Delta C_p = -139$ cal mol^{-1} K^{-1}),

suggesting a change in the solvent accessible hydrophobic surface area caused by Ca^{2+} binding at EF2 (29,30).

Ca^{2+} -induced conformational changes in EF2 are further supported by its Mg^{2+} -dependent Ca^{2+} binding properties. Physiological Mg^{2+} levels (~ 1 mM in rod outer segments (35)) are required for GCAP1 to activate RetGC at low Ca^{2+} , implying that Ca^{2+} -free GCAP1 must bind Mg^{2+} (see below) (16). The presence of physiological Mg^{2+} levels caused a ~ 1.8 -fold increase in K_{EF2} and two-fold decrease in ΔH_{EF2} with smaller effects on binding to EF3 and EF4 (Table 1). This suggests that Mg^{2+} competes with Ca^{2+} for binding at EF2 even at high Ca^{2+} concentrations (~ 1 μM in dark-adapted rods), but Mg^{2+} does not effectively compete for binding at EF3 and EF4 under these conditions. We conclude that both Mg^{2+} and Ca^{2+} -induced conformational changes in EF2 are consistent with recent studies showing that binding of GCAP1 to RetGC requires Mg^{2+} binding to EF2 under conditions that mimic light adaptation of rods (12,17).

We do not see any evidence of Ca^{2+} binding at EF1. The primary structure of EF1 contains unfavorable substitutions (e.g. Cys29 and Pro30 in Fig. 1) known to abolish physiological Ca^{2+} binding at this site in other NCS proteins (3,36,37). Accordingly, our ITC Ca^{2+} binding measurements on GCAP1 do not detect any heat signal from Ca^{2+} binding at EF1, suggesting that either EF1 does not bind Ca^{2+} under physiological conditions or the enthalpy of binding is zero.

Mg^{2+} Binding to GCAP1

The presence of physiological levels of Mg^{2+} (2 mM Mg^{2+}) is required for activation of RetGC by GCAP1 at low Ca^{2+} levels (16), suggesting that Mg^{2+} binds to Ca^{2+} -free GCAP1. Indeed, we showed the Ca^{2+} -binding affinity of EF2 is Mg^{2+} -dependent (Table 1). These Mg^{2+} -dependent effects prompted us to quantitate the direct Mg^{2+} binding to GCAP1 using ITC. Titration of MgCl_2 into apo GCAP1 resulted in a biphasic binding isotherm best fit by a sequential 2-site model (solid lines in Fig. 3A). The binding isotherm shows endothermic Mg^{2+} binding with high micromolar affinity ($\Delta H = +4.3$ kcal/mol and $K_d = 700$ μM) followed by weaker binding in the millimolar range ($\Delta H_{\text{low}} = +0.9$ kcal/mol and $K_{\text{low}} = 2$ mM). Mg^{2+} binding experiments were also performed on the EF-hand mutants to assign each Mg^{2+} binding phase.

The ITC isotherms for wildtype and the EF-hand mutants were fit to a sequential binding model (solid lines in Fig. 3) and optimal Mg^{2+} -binding parameters (ΔH and K_d) are shown in Table 2. The ITC Mg^{2+} -binding isotherm of EF34- (Fig. 3D) exhibits a single endothermic phase representing binding of Mg^{2+} with high micromolar affinity, suggesting that EF2 must be the highest affinity endothermic site ($\Delta H_{\text{EF2}} = +4.3$ kcal/mol and $K_{\text{EF2}} = 0.7$ mM). Correspondingly, the isotherm of EF2- (Fig. 3B) exhibits millimolar binding of 1 or more Mg^{2+} , but lacks the micromolar phase seen in the wildtype isotherm, confirming that EF2 is the highest affinity site. The lower affinity endothermic phase in the wildtype isotherm likely represents Mg^{2+} binding to EF3 and/or EF4. However, individual values of ΔH and K_d for Mg^{2+} binding at EF3 and EF4 could not be accurately resolved due to their low affinity and non-stoichiometric binding ($K_d \gg$ protein concentration).

Our ITC Mg^{2+} -binding analysis reveals that EF2 is an endothermic site with sub-millimolar affinity ($K_{\text{EF2}} = 0.7$ mM and $\Delta H_{\text{EF2}} = +4.3$ kcal/mol); EF3 and/or EF4 bind Mg^{2+} in the millimolar range. Mg^{2+} competes somewhat with Ca^{2+} for binding at EF2, explaining its Mg^{2+} -dependent Ca^{2+} binding affinity and enthalpy (Table 1). By contrast, EF3 and EF4 each bind Ca^{2+} with at least 10000-fold higher affinity than Mg^{2+} , explaining why these sites are more selective for Ca^{2+} and less sensitive to Mg^{2+} . Mg^{2+} binding at EF2 may be functionally important and explain why the activation of RetGC by Ca^{2+} -free GCAP1 requires physiological

levels of Mg^{2+} . Indeed recent experiments have shown that apo GCAP1 does not regulate RetGC, and Mg^{2+} binding at EF2 is required for light-dependent activation of the cyclase (17).

Folding Stability vs. Ca^{2+} , Mg^{2+} and Myristoylation

Differential scanning calorimetry (DSC) experiments were performed on GCAP1 to quantitatively assess the effect of Ca^{2+} , Mg^{2+} and myristoylation on protein folding stability. Representative DSC scans of GCAP1 are shown in Fig. 4. The peak maximum of unmyristoylated metal-free (apo) GCAP1 (transition temperature, $T_m = 47$ °C) is lower than the peak maximum of Mg^{2+} -bound ($T_m = 55$ °C) and Ca^{2+} -bound ($T_m = 104$ °C) unmyristoylated GCAP1, indicating that both Mg^{2+} and particularly Ca^{2+} increase the folding stability quite substantially. N-terminal myristoylation of GCAP1 has a much smaller effect on the folding stability. The DSC peak maxima of myristoylated GCAP1 in the apo (transition temperature, $T_m = 52$ °C), Mg^{2+} -bound ($T_m = 58$ °C) and Ca^{2+} -bound ($T_m = 103$ °C) forms are slightly higher than the corresponding melting temperatures of unmyristoylated GCAP1, indicating that myristoylated and unmyristoylated forms of GCAP1 have similar folding stabilities (Fig. 4). The transition peaks in all the thermograms did not fully reappear upon re-scanning each of the samples, suggesting irreversible unfolding due to aggregation and/or denaturation. In addition, the DSC thermograms of apo, Mg^{2+} -bound, and Ca^{2+} -bound GCAP1 were all quite broad with steeply sloped post-transitional baselines, making it impossible to accurately fit these thermograms by any quantitative models. The cause of the sloping and highly curved baselines is not fully understood, but may be explained in part by visible protein aggregation that occurred during unfolding at the very high melting temperatures.

In summary, the DSC analysis on GCAP1 reveals that Mg^{2+} and Ca^{2+} binding both increase the protein folding stability. N-terminal myristoylation of GCAP1 also stabilizes the protein fold slightly as expected based on the sequestered myristoyl group observed in the x-ray crystal structure (18). However, the increased folding stability conferred by myristoylation of GCAP1 is much smaller than that observed by myristoylated recoverin (Supplemental Fig. 1). We suggest that the vast majority of the folding stability in GCAP1 is derived by structural interactions among the EF-hands, which would explain the strong dependence of T_m on Mg^{2+} and Ca^{2+} binding. Therefore, the myristoyl group interaction with the N- and C-terminal helices observed in the crystal structure appears to contribute very little to the overall protein stability. By contrast, the myristoyl group in Ca^{2+} -free recoverin interacts intimately with hydrophobic residues in EF1, EF2 and EF3 (22), which explains how Ca^{2+} -binding is coupled to extrusion of the myristoyl group and why myristoylation of recoverin has a much more dramatic effect on the overall protein folding stability.

Mg^{2+} and Ca^{2+} Alter the Tertiary Structure of GCAP1

NMR spectroscopy was used to probe protein conformational changes in GCAP1 induced by Mg^{2+} and/or Ca^{2+} binding (Fig. 5). The peaks in the 1H - ^{15}N HSQC NMR spectra of GCAP1 represent main chain and side-chain amide groups and provide a residue-specific fingerprint of the overall protein conformation. HSQC spectra of myristoylated (red) and unmyristoylated (black) forms of GCAP1 look very similar, indicating once again that myristoylation has very little effect on the overall main chain structure of GCAP1. The structural similarity between myristoylated and unmyristoylated GCAP1 is also supported by tryptophan fluorescence emission spectra that have nearly the same emission wavelength maxima (Supplemental Fig. 2). This is in stark contrast to the spectral properties of Ca^{2+} -free myristoylated recoverin, in which myristoylation has a profound influence on both NMR and fluorescence spectra (32, 33). The myristoyl group of recoverin interacts intimately with the various EF-hands and induces large overall structural changes in the protein (20). By contrast, the myristoyl group of GCAP1 appears to have very little effect on the overall protein structure.

The two-dimensional ^1H - ^{15}N HSQC spectrum of apo GCAP1 (both myristoylated and unmyristoylated) exhibited poorly resolved and overlapping peaks with narrow chemical shift dispersion in the amide proton dimension (Fig. 5A). The number of observed peaks was far less than the expected number of amide groups and the intensities of many peaks were quite weak due to broadening caused by dimerization or conformational heterogeneity. The poor chemical shift dispersion suggests that apo GCAP1 may form an unstructured molten-globule state similar to that described for the apo states of many other Ca^{2+} -binding proteins such as GCAP-2 (19), Frq1 (38), CIB (39), calexcitin (40), protein S (41), and DREAM (25). However, circular dichroism analysis (data not shown) suggests that apo GCAP1 adopts a high degree of helical content, consistent with the formation of the four EF-hands, and that the helical content does not change much upon binding Mg^{2+} and/or Ca^{2+} . Together, our structural studies suggest that apo GCAP1 forms native secondary structure but lacks stable tertiary structure.

The HSQC spectrum of GCAP1 changed upon the addition of saturating Mg^{2+} (Fig. 5B). Mg^{2+} caused a greater number of peaks to appear and the NMR intensities were in general more uniform than those of apo GCAP1. Mg^{2+} -binding to GCAP1 increased the NMR chemical shift dispersion and increased the number of observable long-range NOEs, demonstrating that Mg^{2+} -bound GCAP1 adopts stable tertiary structure. Unfortunately, the number of observable HSQC peaks for Mg^{2+} -bound GCAP1 was still only ~50% of the expected number of amide groups, making it impossible to obtain enough sequence specific assignments necessary to solve the structure by NMR. Also, the average peak width in the spectrum appears much broader than expected for a monomeric protein, consistent with protein dimerization in the NMR samples as determined by size-exclusion chromatography studies (SEC). Pulsed-field gradient diffusion NMR studies (42) determined a hydrodynamic radius of 4.0 nm (corresponding to a spherical molecular mass of ~60 kDa) for Mg^{2+} -bound GCAP1, consistent with mostly dimer or higher order species present under conditions for NMR (500- μM protein concentration). However, this GCAP1 dimerization appears only at protein concentrations above 100 μM (as judged by SEC analysis) and therefore GCAP1 is monomeric at physiological concentrations in the rod cell (10–50 μM).

The HSQC spectrum of GCAP1 changed even further upon the addition of saturating Ca^{2+} (Fig. 5C). Spectral changes induced by the addition of saturating Ca^{2+} to the Mg^{2+} -bound protein sample indicated that Ca^{2+} induced conformational changes are distinct and separate from the Mg^{2+} -induced changes (Figs. 5B-C). Three downfield-shifted peaks near 10.5 ppm are characteristic of conserved glycine residues at the 6-position of Ca^{2+} occupied EF-hands (Gly69, Gly105 and Gly149), consistent with Ca^{2+} bound at EF2, EF3 and EF4. Additional unique peaks of Ca^{2+} -bound GCAP1 (observed between 9 and 10 ppm) represent amino acid residues in EF3 and EF4 altered structurally by Ca^{2+} -binding. The HSQC spectrum of Ca^{2+} -saturated GCAP1 exhibited somewhat broadened peaks (like that of Mg^{2+} -bound protein) with variable NMR intensities, consistent with protein dimerization as measured by SEC. Pulsed-field gradient diffusion NMR studies determined a hydrodynamic radius of 3.8 nm, corresponding to a Ca^{2+} -bound protein dimer under NMR conditions.

The NMR peaks of Ca^{2+} -bound GCAP1 sharpened quite substantially upon adding octylglucoside detergent and raising the sample temperature to 47 °C (Fig. 5D). Under these conditions the protein became monomeric, which greatly improved the overall NMR sensitivity. Also, the chemical shifts remained unperturbed by the detergent, consistent with a monomeric and properly folded protein. Initially, this improvement in spectral quality suggested that it might be feasible to determine the full three-dimensional structure by NMR. However upon closer inspection, we were only able to identify ~80% of the main chain amide resonances in HSQC spectra and less than 50% of the resonances could be reliably assigned in triple resonance experiments. The resonances with strongest NMR intensities could be assigned mostly to residues in EF3 and EF4. However, many other resonances (associated with

EF1 and EF2) had much weaker NMR intensities in triple resonance experiments and could not be accurately assigned. Furthermore, the first 16 residues from the N-terminus and a stretch of ~20 residues following EF4 at the C-terminus have very weak NMR intensities due to exchange broadening as evidenced by their sharp temperature and magnetic field dependence. These N- and C-terminal residues (italicized in Fig. 1) are exchange broadened more so for myristoylated GCAP1 compared to that of the unmyristoylated protein. Interestingly, many of these same residues are found in helices that interact with the myristoyl group in the x-ray structure. The exchange broadening for the N- and C-terminal residues suggests that these residues undergo dynamical motions on the chemical shift time-scale to facilitate their interactions with the myristoyl group, and such dynamical behavior might be functionally important for recognition of RetGC.

Myristoyl Group Structural Environment vs. Ca^{2+} and Mg^{2+}

The recent x-ray crystal structure of GCAP1 indicates that the myristoyl group is buried inside the Ca^{2+} -bound protein (18). This is in contrast to a solvent exposed myristoyl group observed for GCAP2 (43), yeast frequenin (38), FCaBP (44), and recoverin (20). We performed NMR experiments on samples of apo, Mg^{2+} -bound and Ca^{2+} -bound forms of GCAP1 with a covalently attached ^{13}C -labeled myristoyl group to probe its structural environment inside the protein and to look for any effects of Mg^{2+} or Ca^{2+} -binding. Previously, two-dimensional (^1H - ^{13}C HMQC) and three-dimensional (^{13}C -filtered NOESY-HMQC) NMR experiments on samples of recoverin that contained a ^{13}C -labeled myristoyl group were used to selectively probe Ca^{2+} -induced changes to the chemical environment around the amino-terminal myristoyl group (23,32). These studies revealed that the covalently-attached fatty acyl chain in recoverin is sequestered in a hydrophobic pocket in the Ca^{2+} -free protein (22) and that binding of Ca^{2+} leads to conformational changes that extrude the myristoyl group into solvent.

Similar NMR experiments were performed on samples of GCAP1 that contained a covalently attached ^{13}C -labeled myristoyl group (Fig. 6). Because the HMQC experiment selectively probes protons that are covalently attached to ^{13}C , only the methylene and methyl resonances of the fatty acyl chain are expected to appear in these spectra. Extraneous peaks near 1.0, 1.4, 1.8 and 2.2 ppm (^1H dimension) are due to background signals from the protein. The NMR spectrum of the myristoyl group of GCAP1 looks nearly identical for apo, Mg^{2+} -bound and Ca^{2+} -bound GCAP1 (Figs. 6A-C), indicating that the structural environment around the myristoyl group does NOT depend on Ca^{2+} , in stark contrast to the large Ca^{2+} -induced spectral changes for the myristoyl group in recoverin (20).

The NMR chemical shifts of the myristoyl resonances of GCAP1 are consistent with the attached myristoyl group being sequestered inside a protein environment like was seen in the crystal structure (Table 3). Assignments of NMR resonances of the myristoyl group attached to GCAP1 were derived from assignments of free myristic acid in solution determined previously (32). The proton chemical shifts of the C_{14} methyl (0.62–0.73 ppm) and C_4 - C_{12} methylene resonances (0.9–1.1 ppm) in GCAP1 are all somewhat upfield shifted compared to the corresponding chemical shifts of free myristic acid in solution (32). These upfield chemical shifts in GCAP1 may be explained in part by aromatic groups (F39, F43, F63 and Y76) close to the myristoyl group in the crystal structure that might impose an aromatic ring-current effect on the nearby myristoyl group. Indeed, the myristoyl group methyl and methylene proton chemical shifts in GCAP1 became downfield shifted like those of myristate in solution upon unfolding GCAP1 by adding 8 M urea (Fig. 6D and Table 3).

The spectral widths of the myristoyl NMR resonances in GCAP1 are strikingly broad (50–80 Hz) compared to the widths of the natural abundance protein peaks as well as the widths of myristoyl resonances in recoverin (32) and related proteins (38,44). The very broad NMR resonances of the myristoyl group in GCAP1 suggest that the attached myristoyl group may

be located in a dynamic and/or heterogeneous protein environment. For free myristic acid in solution, a single proton resonance of the C₁₄ methyl group is quite sharp because free rotation of the methyl group about the C₁₃-C₁₄ single bond causes the three methyl protons to experience a time-averaged, uniform environment. By contrast, the C₁₄ methyl resonance (¹H dimension) in GCAP1 is quite broad and is partially split into two resolved components at 0.73 and 0.62 ppm with approximately equal intensity (Fig. 6). This splitting (66 Hz) suggests that the C₁₄ methyl group may occupy at least two different structural environments. These two chemical shift environments represent distinct conformational states of the protein (e.g. activator vs. inhibitor forms), but the relative intensities of the two components did not change much upon Ca²⁺ binding. The two methyl resonances might also represent the myristoyl group in folded and misfolded protein forms or in two subunits of an asymmetric dimer. However, the spectral heterogeneity persisted in the presence of 25 mM octylglucoside detergent (which causes GCAP1 to be monomeric), arguing against any artifacts caused by dimerization.

The apparent spectral heterogeneity and broadening implies that the myristoyl group must be in at least two different protein environments that are exchanging on the chemical shift time scale. The doublet methyl resonance (66 Hz splitting) collapsed into a single broad peak (80 Hz linewidth) when the temperature was raised from 32 to 45 °C, consistent with an exchange rate on the millisecond time scale. The exchange behaviour of the myristoyl resonances could also relate and help explain why the ¹H-¹⁵N HSQC peaks of the N- and C-terminal residues in GCAP1 (italicized in Fig. 1) are exchanged broadened as described above (Fig. 5). We propose that the N- and C-terminal residues, known to interact with the myristoyl group in the crystal structure (italicized in Fig. 1), can undergo conformational fluctuations on the chemical shift timescale and therefore give rise to the spectral heterogeneity observed around the myristoyl C₁₄ methyl group.

To further probe the protein structural environment around the myristoyl group of GCAP1, three-dimensional (¹³C/F₁)-edited and (¹³C/F₃)-filtered NOESY experiments (32) were performed on unlabeled GCAP1 protein containing a ¹³C-labeled myristate. These spectra selectively probed atoms of residues in the protein that lie within 5 Å of the labeled CH₃-group of the myristoyl chain. Nuclear Overhauser effect (NOE) dipolar interactions between the myristate methyl group and the protein could not be detected (data not shown), because the exchange broadening of the methyl resonances described above severely attenuates any NOE signals. However, the myristate methylene resonances at C₃ and C₄ did exhibit detectable NOEs to aromatic protons from the protein (Supplemental Fig. 3), providing further evidence that the myristoyl group is indeed surrounded by a hydrophobic and aromatic protein environment like what is observed in the crystal structure. The addition of Ca²⁺ had no effect on the filtered NOESY spectrum, further confirming that the myristate remains sequestered in a protein environment in both Ca²⁺-bound and Ca²⁺-free GCAP1. This Ca²⁺-independent sequestration of the myristate in GCAP1 is quite different from what is seen in recoverin that exhibits a sequestered myristoyl environment only in the Ca²⁺-free protein (20,24).

DISCUSSION

In this study, we determined the energetics of Mg²⁺ and Ca²⁺ binding to GCAP1 as well as structural effects of myristoylation. Ca²⁺ binds enthalpically to both EF3 and EF4 in the nanomolar range with more than 1000-fold selectivity over Mg²⁺. EF2 binds entropically to both Ca²⁺ and Mg²⁺ and undergoes metal-induced conformational changes related to cyclase regulation (17). Myristoylation has almost no influence on the Ca²⁺/Mg²⁺ binding properties of GCAP1. Furthermore, the Ca²⁺-induced structural changes in the EF-hands do not influence the structural environment that surrounds the sequestered myristoyl group. Unlike recoverin, the myristoyl group attached to GCAP1 remains buried inside the protein regardless of Ca²⁺ level and GCAP1 does NOT possess a functional Ca²⁺-myristoyl switch.

The Ca^{2+} dissociation constants ($K_{\text{EF}2}$, $K_{\text{EF}3}$ and $K_{\text{EF}4}$) of wildtype GCAP1 are somewhat different from those of the mutants (Table 1), implying that the three Ca^{2+} -binding sites must interact and possess some cooperativity. Indeed, the EF3- mutation in GCAP1 abolishes Ca^{2+} binding to EF4 (16), suggesting both ordered and cooperative Ca^{2+} binding at EF3 and EF4. In essence, Ca^{2+} must bind at EF3 first to facilitate subsequent binding at EF4. Such ordered Ca^{2+} binding is consistent with our measurement that EF3 has the highest binding affinity ($K_{\text{EF}3} = 80 \text{ nM}$), whereas EF4 has intermediate affinity ($K_{\text{EF}4} = 200 \text{ nM}$) and EF2 has lowest affinity ($K_{\text{EF}2} = 0.9 \text{ }\mu\text{M}$). Cooperative Ca^{2+} binding between EF3 and EF4 is also consistent with their intimate structural contacts observed in the x-ray crystal structure (18), and concerted Ca^{2+} -induced NMR spectral changes observed for Gly105 (EF3) and Gly149 (EF4) in Supplemental Fig. 4. This cooperativity seems at odds with the overall Ca^{2+} -binding isotherm of GCAP1 having a Hill slope of 1.0 (16). The Hill coefficient in this case, however, is not always a reliable measure of cooperativity, particularly if the individual sites have different intrinsic affinities (45). The free energy of interaction between the two sites ($\Delta G_{\text{interact}}$) is a more stringent definition of cooperativity and this interaction energy can be inferred by comparing how the dissociation constant of EF3 ($K_{\text{EF}3} = \exp(-\Delta G^\circ/RT)$) changes when EF4 is disabled (EF4-). For GCAP1, $K_{\text{EF}3}$ increases substantially by disabling EF4 (Table 1), which means that $\Delta G_{\text{interact}} > 0$ and the two sites have positive cooperativity. We propose that this positive cooperativity of Ca^{2+} binding between EF3 and EF4 could help explain how GCAP1 confers such a steep Ca^{2+} -dependent activation of RetGC with a Hill coefficient = 2 (46). The cooperative cyclase activation may also arise in part by Ca^{2+} -induced dimerization of RetGC (47).

The results of this study provide structural insights into the regulatory mechanism of RetGC. The activation of RetGC by Ca^{2+} -free GCAP1 requires physiological Mg^{2+} levels (16). Our NMR structural analysis shows that the completely metal-free (apo) GCAP1 lacks organized tertiary structure, explaining why the apo state is physiologically inactive (Fig. 5A). Mg^{2+} binding induces structural changes in the EF-hands and stabilizes protein tertiary structure as evidenced by DSC (Fig. 4) and NMR (Fig. 5). Mg^{2+} binds most tightly to GCAP1 at EF2 (Table 2), which we suggest might constitute an important interaction site for cyclase activation. Indeed, recent biochemical and cell biology studies have shown that EF2 is required for the cyclase interaction (12,17) and promotes docking of Mg^{2+} -bound GCAP1 on RetGC (17). In addition, Mg^{2+} binding at EF3 appears to facilitate the docking of GCAP1 with RetGC, because cyclase binding to GCAP1 is weakened by a mutation that blocks Mg^{2+} binding at EF3 (12). Finally, Ca^{2+} binding at EF4 is necessary and sufficient for GCAP1 to inhibit RetGC when Ca^{2+} rises in the dark (12,16). It seems likely that Ca^{2+} binding at EF4 may promote the docking of Ca^{2+} bound GCAP1 onto the cyclase (48,49).

The replacement of bound Ca^{2+} by Mg^{2+} in GCAP1 is functionally important for switching GCAP1 from a Ca^{2+} -bound inhibitor to a Mg^{2+} -bound activator conformational state during light activation. The relatively low apparent Ca^{2+} binding affinity of EF2 measured by ITC is explained by its thermodynamic coupling to an unfavorable protein conformational change probed structurally by NMR (Fig. 5). The intrinsic Ca^{2+} binding to EF2 measured by a direct Ca^{2+} binding assay ($K_a = 10^7 \text{ M}^{-1}$) coupled to an unfavorable conformational change ($K_{\text{eq}} \sim 10^{-1}$) yields an overall binding constant of $K_{\text{tot}} = K_a * K_{\text{eq}} \sim 10^6 \text{ M}^{-1}$, consistent with the overall $K_{\text{EF}2}$ measured by ITC. Thus, the intrinsic Ca^{2+} -binding constants inferred from the ITC analysis are consistent with those measured by direct equilibrium binding assays (16) and consistent with the functional behavior of GCAP1 in physiological analyses (16,17). The relative binding affinities of Mg^{2+} ($K_{\text{EF}2} = 0.7 \text{ mM}$) and Ca^{2+} ($K_{\text{EF}2} = 0.9 \text{ }\mu\text{M}$) at EF2 suggest that a fraction of GCAP1 will be occupied with Mg^{2+} even in dark adapted rods (250 nM Ca^{2+} and 1 mM Mg^{2+} (35)) and virtually all of the EF2 sites are occupied by Mg^{2+} upon illumination, when the Ca^{2+} concentration decreases below 100 nM (50). For EF3 and EF4, the relatively weak Mg^{2+} binding ($K_d = 2 \text{ mM}$) and strong Ca^{2+} binding ($K_{\text{EF}3} = 80 \text{ nM}$ and

$K_{EF4} = 200$ nM) means that both of these sites are fully occupied by Ca^{2+} in the dark. Mg^{2+} can bind to these sites only under bright light illumination, when the free Ca^{2+} concentration in rod outer segment falls drastically to ~ 25 nM (50). Mg^{2+} binding at EF3 is required for RetGC activation (12,16,17), whereas Mg^{2+} binding at EF4 is not essential (12,16), suggesting that EF4 does not bind functionally to Mg^{2+} .

Ca^{2+} binding to GCAP1 leads to protein conformational changes that cause inhibition of RetGC activity at high Ca^{2+} levels in the rod cell, which is necessary to maintain a steady cGMP level in the dark that otherwise may lead to retinal disease (8,51). Our ITC and NMR studies suggest that GCAP1 binds cooperatively to Ca^{2+} at EF3 and EF4 that promotes a Ca^{2+} -induced conformational change (Fig. 5). Indeed, a number of residues in EF3 and EF4 (e.g. Gly105 and Gly149) exhibit Ca^{2+} -induced NMR spectral changes that occur in a concerted fashion (Suppl Fig. 4). Also, Ca^{2+} binding at EF3 and EF4 both display large negative ΔC_p values (-379 cal mol $^{-1}$ K $^{-1}$ and -101 cal mol $^{-1}$ K $^{-1}$) suggesting that Ca^{2+} binding at both sites promotes an overall burial of hydrophobic residues (29). This seems somewhat at odds with Ca^{2+} -dependent changes of intrinsic Trp fluorescence from GCAP1, especially for Trp21 in EF1 and Trp94 in EF3 (46,52). Trp21 and Trp94 shift into polar environments in both the Mg^{2+} -bound and Ca^{2+} -bound GCAP1 relative to its apo-conformation, whereas Trp94 shifts back into a more hydrophobic environment when Ca^{2+} replaces Mg^{2+} at EF3 (16). Our ITC analysis detects an overall Ca^{2+} -induced decrease in exposed hydrophobic surface area, because ITC detects signals from all residues of the protein (and not just Trp). Therefore, Ca^{2+} binding at EF3 and EF4 causes conformational changes leading to a net burial of hydrophobic residues in GCAP1 that presumably compensates for the exposure of Trp21 and Trp94.

What is the structural and functional role of the N-terminal myristoyl group of GCAP1? Myristoylation of GCAP1 is thought to be important for the regulation of RetGC (53,54), and the recent crystal structure of GCAP1 shows the myristate to be buried inside the protein hydrophobic core (18). In this study, our NMR analyses confirmed that the myristoyl group in GCAP1 is indeed sequestered inside a protein environment in the native conformation of the protein in solution. Upfield chemical shifts and NOE patterns of the myristoyl resonances (Fig. 6, Table 3) are consistent with ring-currents imposed by aromatic residues (F39, F43, F63 and Y76) shown to be near the myristate in the crystal structure. Ca^{2+} - and/or Mg^{2+} -binding to GCAP1 has very little influence on the chemical shifts and NOE patterns of the myristoyl group (Fig. 6), suggesting that the myristoyl group remains in a similar environment in both the Mg^{2+} -bound activator and Ca^{2+} -bound inhibitor forms of GCAP1. Surprisingly, however, myristoylation of GCAP1 has only a small effect on the overall protein folding stability (Fig. 4) and structure of the EF-hands (Fig. 5). Thus, it seems that myristoylation interacts most strongly with residues outside the EF-hand regions; namely, the N- and C-terminal helices that contact the myristate in the crystal structure (italicized in Fig. 1). Interestingly, the NMR resonances of these N- and C-terminal residues and those of myristate are all spectrally quite broad (linewidth, $\Delta\nu = 50$ –80 Hz) due to exchange processes on the millisecond time scale ($\tau_{ex} \propto 1/\Delta\nu$), suggesting a highly dynamic environment around the myristate. These millisecond exchange processes (activation energy ~ 30 kJ/mol) occur at physiological temperatures but are essentially frozen out under conditions for x-ray crystallography (77 K), explaining why this dynamical behavior apparently was not observed in the crystal structure. We propose that conformational fluctuations at the myristoyl binding site might regulate the cyclase by enabling specific recognition of RetGC or by facilitating secondary interactions required for RetGC activation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

We are grateful to Dr. Jeff de Ropp for help with NMR experiments, Dr. Frits Abildgaard for providing NMR pulse-sequence programs, and Frank Delaglio for writing computer software for NMR data processing and analysis. A.M.D. is The Martin and Florence Hafter Chair Professor of Pharmacology.

1 Abbreviations

Ca²⁺, calcium ion
 Mg²⁺, magnesium ion
 DSC, differential scanning calorimetry
 GCAP1, guanylyl cyclase activating protein 1
 EDTA, ethylenediaminetetraacetic acid
 HMQC, heteronuclear multiple quantum coherence
 HSQC, heteronuclear single quantum coherence
 IPTG, isopropylthiogalactoside
 ITC, isothermal titration calorimetry
 NOE, nuclear Overhauser effect
 NOESY, nuclear Overhauser effect spectroscopy
 RetGC, retinal guanylyl cyclase
 SDS PAGE, sodium dodecylsulfate polyacrylamide gel electrophoresis

REFERENCES

- Burgoyne RD. Neuronal calcium sensor proteins: generating diversity in neuronal Ca²⁺ signalling. *Nat. Rev. Neurosci* 2007;8:182–193. [PubMed: 17311005]
- Burgoyne RD, Weiss JL. The neuronal calcium sensor family of Ca²⁺-binding proteins. *Biochem. J* 2001;353:1–12. [PubMed: 11115393]
- Ames JB, Tanaka T, Stryer L, Ikura M. Portrait of a myristoyl switch protein. *Curr. Opin. Struct. Biol* 1996;6:432–438. [PubMed: 8794166]
- Dizhoor AM, Lowe DG, Olsevskaya EV, Laura RP, Hurley JB. The human photoreceptor membrane guanylyl cyclase, RetGC, is present in outer segments and is regulated by calcium and a soluble activator. *Neuron* 1994;12:1345–1352. [PubMed: 7912093]
- Palczewski K, Subbaraya I, Gorczyca WA, Helekar BS, Ruiz CC, Ohguro H, Huang J, Zhao X, Crabb JW, Johnson RS. Molecular cloning and characterization of retinal photoreceptor guanylyl cyclase-activating protein. *Neuron* 1994;13:395–404. [PubMed: 7520254]
- Palczewski K, Polans AS, Baehr W, Ames JB. Ca(2+)-binding proteins in the retina: structure, function, and the etiology of human visual diseases. *Bioessays* 2000;22:337–350. [PubMed: 10723031]
- Stephen R, Filipek S, Palczewski K, Sousa MC. Ca²⁺-dependent regulation of phototransduction. *Photochem. Photobiol* 2008;84:903–910. [PubMed: 18346093]
- Payne AM, Downes SM, Bessant DA, Taylor R, Holder GE, Warren MJ, Bird AC, Bhattacharya SS. A mutation in guanylate cyclase activator 1A (GUCA1A) in an autosomal dominant cone dystrophy pedigree mapping to a new locus on chromosome 6p21.1. *Hum. Mol. Genetics* 1998;7:273–277.
- Baylor D. How photons start vision. *Proc. Natl. Acad. Sci. USA* 1996;93:560–565. [PubMed: 8570595]
- Koch KW, Stryer L. Highly cooperative feedback control of retinal rod guanylate cyclase by calcium ions. *Nature* 1988;334:64–66. [PubMed: 2455233]
- Olsevskaya EV, Calvert PD, Woodruff ML, Peshenko IV, Savchenko AB, Makino CL, Ho YS, Fain GL, Dizhoor AM. The Y99C mutation in guanylyl cyclase-activating protein 1 increases intracellular Ca²⁺ and causes photoreceptor degeneration in transgenic mice. *J. Neurosci* 2004;24:6078–6085. [PubMed: 15240799]
- Peshenko IV, Dizhoor AM. Activation and inhibition of photoreceptor guanylyl cyclase by guanylyl cyclase activating protein 1 (GCAP-1): the functional role of Mg²⁺/Ca²⁺ exchange in EF-hand domains. *J. Biol. Chem* 2007;282:21645–21652. [PubMed: 17545152]

13. Dizhoor AM, Olshevskaya EV, Henzel WJ, Wong SC, Stults JT, Ankoudinova I, Hurley JB. Cloning, sequencing and expression of a 24-kDa Ca²⁺-binding protein activating photoreceptor guanylyl cyclase. *J. Biol. Chem* 1995;270:25200–25206. [PubMed: 7559656]
14. Imanishi Y, Li N, Sokal I, Sowa ME, Lichtarge O, Wensel TG, Saperstein DA, Baehr W, Palczewski K. Characterization of retinal guanylate cyclase-activating protein 3 (GCAP3) from zebrafish to man. *Eur. J. Neurosci* 2002;15:63–78. [PubMed: 11860507]
15. Imanishi Y, Yang L, Sokal I, Filipek S, Palczewski K, Baehr W. Diversity of guanylate cyclase-activating proteins (GCAPs) in teleost fish: characterization of three novel GCAPs (GCAP4, GCAP5, GCAP7) from zebrafish (*Danio rerio*) and prediction of eight GCAPs (GCAP1–8) in pufferfish (*Fugu rubripes*). *J. Mol. Evol* 2004;59:204–217. [PubMed: 15486694]
16. Peshenko IV, Dizhoor AM. Ca²⁺ and Mg²⁺ binding properties of GCAP-1. Evidence that Mg²⁺-bound form is the physiological activator of photoreceptor guanylyl cyclase. *J. Biol. Chem* 2006;281:23830–23841. [PubMed: 16793776]
17. Peshenko IV, Olshevskaya EV, Dizhoor AM. Binding of guanylyl cyclase activating protein 1 (GCAP1) to retinal guanylyl cyclase (RetGC1). The role of individual EF-hands. *J. Biol. Chem* 2008;283:21747–21757. [PubMed: 18541533]
18. Stephen R, Bereta G, Golczak M, Palczewski K, Sousa MC. Stabilizing function for myristoyl group revealed by the crystal structure of a neuronal calcium sensor, guanylate cyclase-activating protein 1. *Structure* 2007;15:1392–1402. [PubMed: 17997965]
19. Ames JB, Dizhoor AM, Ikura M, Palczewski K, Stryer L. Three-dimensional structure of guanylyl cyclase activating protein-2, a calcium-sensitive modulator of photoreceptor guanylyl cyclases. *J Biol Chem* 1999;274:19329–19337. [PubMed: 10383444]
20. Ames JB, Ishima R, Tanaka T, Gordon JI, Stryer L, Ikura M. Molecular mechanics of calcium-myristoyl switches. *Nature* 1997;389:198–202. [PubMed: 9296500]
21. Ames JB, Ikura M, Stryer L. Molecular structure of membrane-targeting calcium sensors in vision: recoverin and guanylate cyclase-activating protein 2. *Methods Enzymol* 2000;316:121–132. [PubMed: 10800672]
22. Tanaka T, Ames JB, Harvey TS, Stryer L, Ikura M. Sequestration of the membrane-targeting myristoyl group of recoverin in the calcium-free state. *Nature* 1995;376:444–447. [PubMed: 7630423]
23. Tanaka T, Ames JB, Kainosho M, Stryer L, Ikura M. Differential isotope labeling strategy for determining the structure of myristoylated recoverin by NMR spectroscopy. *J Biomol NMR* 1998;11:135–152. [PubMed: 9679292]
24. Valentine K, Mesleh M, Ikura M, Ames JB, Opella S. Structure, topology and dynamics of myristoylated recoverin bound to phospholipid bilayers. *Biochemistry* 2003;42:6333–6340. [PubMed: 12767213]
25. Osawa M, Dace A, Tong KI, Valiveti A, Ikura M, Ames JB. Mg²⁺ and Ca²⁺ differentially regulate DNA binding and dimerization of DREAM. *J Biol Chem* 2005;280:18008–18014. [PubMed: 15746104]
26. Ames JB, Hamasaki N, Molchanova T. Structure and calcium-binding studies of a recoverin mutant (E85Q) in an allosteric intermediate state. *Biochemistry* 2002;41:5776–5787. [PubMed: 11980481]
27. Tsien RY. A new generation of Ca²⁺ indicators with greatly improved fluorescence properties. *Methods Cell Biol* 1989;30:127–156. [PubMed: 2538708]
28. Wingard JN, Chan J, Bosanac I, Haeseleer F, Palczewski K, Ikura M, Ames JB. Structural analysis of Mg²⁺ and Ca²⁺ binding to CaBP1, a neuron-specific regulator of calcium channels. *J Biol Chem* 2005;280:37461–37470. [PubMed: 16147998]
29. Gomez J, Hilser VJ, Xie D, Freier E. The heat capacity of proteins. *Proteins* 1995;22:404–412. [PubMed: 7479713]
30. Gomez J, Freier E. Thermodynamic mapping of the inhibitor site of the aspartic protease endothiapepsin. *J. Mol. Biol* 1995;252:337–350. [PubMed: 7563055]
31. Wiseman T, Williston S, Brandts JF, Lin LN. Rapid measurement of binding constants and heats of binding using a new titration calorimeter. *Anal Biochem* 1989;179:131–137. [PubMed: 2757186]
32. Ames JB, Tanaka T, Ikura M, Stryer L. Nuclear magnetic resonance evidence for Ca(2+)-induced extrusion of the myristoyl group of recoverin. *J. Biol. Chem* 1995;270:30909–30913. [PubMed: 8537345]



33. Ames JB, Porumb T, Tanaka T, Ikura M, Stryer L. Amino-terminal myristoylation induces cooperative calcium binding to recoverin. *J Biol Chem* 1995;270:4526–4533. [PubMed: 7876221]
34. Maune JF, Klee CB, Beckingham K. Ca²⁺ binding and conformational change in two series of point mutations to the individual Ca(2+)-binding sites of calmodulin. *J Biol Chem* 1992;267:5286–5295. [PubMed: 1544911]
35. Chen C, Nakatani K, Koutalos Y. Free magnesium concentration in salamander photoreceptor outer segments. *J. Physiol* 2003;553:125–135. [PubMed: 14500766]
36. Flaherty KM, Zozulya S, Stryer L, McKay DB. Three-dimensional Structure of Recoverin, a Calcium Sensor in Vision. *Cell* 1993;75:709–716. [PubMed: 8242744]
37. Zhou W, Qian Y, Kunjilwar K, Pfaffinger PJ, Choe S. Structural insights into the functional interaction of KChIP1 with Shal-type K(+) channels. *Neuron* 2004;41:573–586. [PubMed: 14980206]
38. Ames JB, Hendricks KB, Strahl T, Huttner IG, Hamasaki N, Thorner J. Structure and calcium-binding properties of Frq1, a novel calcium sensor in the yeast *Saccharomyces cerevisiae*. *Biochemistry* 2000;39:12149–12161. [PubMed: 11015193]
39. Yamniuk AP, Nguyen LT, Hoang TT, Vogel HJ. Metal ion binding properties and conformational states of calcium- and integrin-binding protein. *Biochemistry* 2004;43:2558–2568. [PubMed: 14992593]
40. Gombos Z, Durussel I, Ikura M, Rose DR, Cox JA, Chakrabarti A. Conformational coupling of Mg²⁺ and Ca²⁺ on the three-state folding of calmodulin. *Biochemistry* 2003;42:5531–5539. [PubMed: 12731896]
41. Qi XF, Bagby S, Gombos Z, Ikura M, Chakrabarti A. Alternate routes to conformational specificity in a Greek key beta barrel protein. *Eur. J. Biochem* 2001;268:4653–4663. [PubMed: 11532002]
42. Altieri AS, Hinton DP, Byrd RA. Association of biomolecular systems via pulsed field gradient NMR self-diffusion measurements. *J. Am. Chem. Soc* 1995;117:7566–7567.
43. Vogel A, Schroder T, Lange C, Huster D. Characterization of the myristoyl lipid modification of membrane-bound GCAP-2 by 2H solid-state NMR spectroscopy. *Biochim. Biophys. Acta* 2007;1768:3171–3181. [PubMed: 17936244]
44. Wingard JN, Ladner J, Vanarotti M, Fisher AJ, Buchanan KT, Engman DM, Ames JB. Structural insights into membrane targeting by the flagellar calcium-binding protein (FCaBP), a myristoylated and palmitoylated calcium sensor in *Trypanosoma cruzi*. *J. Biol. Chem* 2008;283:23388–23396. [PubMed: 18559337]
45. Forsen S, Linse S. Cooperativity: over the Hill. *Trends. Biochem. Sci* 1995;20:495–497. [PubMed: 8571449]
46. Peshenko IV, Dizhoor AM. Guanylyl cyclase-activating proteins (GCAPs) are Ca²⁺/Mg²⁺ sensors: implications for photoreceptor guanylyl cyclase (RetGC) regulation in mammalian photoreceptors. *J. Biol. Chem* 2004;279:16903–16906. [PubMed: 14993224]
47. Ramamurthy V, Tucker C, Wilkie SE, Daggett V, Hunt DM, Hurley JB. Interactions within the coiled-coil domain of RetGC-1 guanylyl cyclase are optimized for regulation rather than for high affinity. *J. Biol. Chem* 2001;276:26218–26229. [PubMed: 11306565]
48. Dizhoor AM, Boikov SG, Olshevskaya EV. Constitutive activation of photoreceptor guanylate cyclase by Y99C mutant of GCAP-1. Possible role in causing human autosomal dominant cone degeneration. *J. Biol. Chem* 1998;273:17311–17314. [PubMed: 9651312]
49. Krylov DM, Niemi GA, Dizhoor AM, Hurley JB. Mapping sites in guanylyl cyclase activating protein-1 required for regulation of photoreceptor membrane guanylyl cyclases. *J. Biol. Chem* 1999;274:10833–10839. [PubMed: 10196159]
50. Woodruff ML, Sampath AP, Mathews HR, Krasnoperova NV, Lem J, Fain GL. Measurement of cytoplasmic calcium concentration in the rods of wild-type and transducin knock-out mice. *J. Physiol* 2002;542:843–854. [PubMed: 12154183]
51. Woodruff ML, Janisch KM, Peshenko IV, Dizhoor AM, Tsang SH, Fain GL. Modulation of phosphodiesterase6 turnoff during background illumination in mouse rod photoreceptors. *J. Neurosci* 2008;28:2064–2074. [PubMed: 18305241]
52. Sokal I, Otto-Bruc AE, Surgucheva I, Verlinde CL, Wang CK, Baehr W, Palczewski K. Conformational changes in guanylyl cyclase-activating protein 1 (GCAP1) and its tryptophan

- mutants as a function of calcium concentration. *J. Biol. Chem* 1999;274:19829–19837. [PubMed: 10391927]
53. Hwang JY, Koch KW. Calcium- and myristoyl-dependent properties of guanylate cyclase-activating protein-1 and protein-2. *Biochemistry* 2002;41:13021–13028. [PubMed: 12390029]
54. Hwang JY, Lange C, Helten A, Hoppner D, Duda T, Sharma RK, Koch KW. Regulatory modes of rod outer segment membrane guanylate cyclase differ in catalytic efficiency and Ca(2+)-sensitivity. *Eur. J. Biochem* 2003;270:3814–3821. [PubMed: 12950265]

	1	10	20	30	40	50	
GCAP1	MGNIMDGKSV	<i>EEL</i>SSTECHQ	WYKKFMTECP	<u>SGQLTLYEFR</u>	<u>QFFGLKNLSP</u>	
GCAP2	MGQQFSWEEA	EE...NGAV	GAADAAQLQE	WYKKFLEECF	<u>SGTLFMHEFK</u>	<u>RFFKVPD.NE</u>	
GCAP3	MGNKSIAGDQ	KAVPTQETHV	WYRTFMMEYP	<u>SGLQTLHEFK</u>	<u>TLLGLQGLNQ</u>	
recoverin	MGNSKSGALS	KEILEELQLN	TKFTEEELSS	WYQSFLKECP	<u>SGRITRQEFQ</u>	<u>TIYSKFFPEA</u>	
					EF1		
	60	70	80	90	100	110	
GCAP1	WASQYVEQMF	<u>ETFDENK</u> DGY	<u>IDFMEY</u> VAAAL	SLVLKKGKVEQ	KLRWYFKLYD	<u>VDGNGCIDRD</u>	
GCAP2	EATQYVEAMF	<u>RAFDTNG</u> DNT	<u>IDFLEY</u> VAAAL	NLVLRGTLLEH	KLKWTFKIYD	<u>KDRNGCIDRQ</u>	
GCAP3	KANKHIDQVY	<u>NTFDTNK</u> DGF	<u>IDFLEF</u> IAAV	NLIMQEKMEQ	KLKWFYFKLYD	<u>ADGNLSIDKN</u>	
recoverin	DPKAYAQHVF	<u>RSFDANS</u> DGT	<u>LDFKEY</u> VIAL	HMTSAGKTNQ	KLEWAFSLYD	<u>VDGNGTISKN</u>	
		EF2				EF3	
	120	130	140	150	160		
GCAP1	ELLTIIRAIR	AINP.C....SDSTMT	AEEFTDVFVS	KIDVNGD GEL	<u>SLEEFMEGVQ</u>	
GCAP2	ELLDIVESIY	KLKACSV	EAEQQGKLLT	PEEVVDRIFL	<u>LVDENG</u> DGQL	<u>SLNEFVEGAR</u>	
GCAP3	ELLDMFMAVQ	ALNG.....QQTLS	PEEFINLVFH	KIDINNDGEL	<u>TLEEFINGMA</u>	
recoverin	EVLEIVTAIF	KMISPEDTKH	L...PEDENT	PEKRAEKIWG	<u>FFGKKDDDKL</u>	<u>TEKEFIEGTL</u>	
					EF4		
	170	180	190	200			
GCAP1	KDQMLLDTLT	<i>RSLDLTRIVR</i>	<i>RLQNGEQ</i> .DE	EGASGR....	ETEAAEADG	204	
GCAP2	RDKWVMKMLQ	MDLNPSSWIS	QQR.....RKSAMF	205	
GCAP3	KDQDLLEIVY	KSFDNSNVL	VICNGKQPDM	ETDSSKSPDK	AGLGKVKMK	210	
recoverin	ANKEILRLIQ	FE..PQKVKE	K.....LKEKKL	202	

Figure 1.

Amino acid sequence alignment of GCAP1 with various NCS proteins. Secondary structural elements are indicated schematically. The four EF-hands (EF1, EF2, EF3 and EF4) are underlined. Mutated residues in the EF-hand loops are indicated in bold. N- and C-terminal residues that interact with myristate are shown in italics.






Figure 2.

ITC Ca^{2+} titration of apo GCAP1^{WT} (A), Mg^{2+} -bound GCAP1^{WT} (B), apo EF2⁻ (C), apo EF4⁻ (D) and apo EF34⁻ (E). The upper panels show representative scans of ITC data from the Ca^{2+} titrations. The lower panels show plots of the corresponding integrated binding isotherms fit to a sequential binding model with 3-sites (wildtype), 2-sites (EF2⁻ and EF4⁻), or 1-site (EF34⁻) as described in Methods (solid lines). Optimal fitting parameters are displayed in Table 1. Ca^{2+} binding isotherms of unmyristoylated GCAP1 are overlaid as circles in A and B.




Figure 3. ITC Mg^{2+} titration of myristoylated GCAP1^{WT} (A), EF2⁻ (B) and EF34⁻ (C). The upper panels show representative scans of ITC data from the Mg^{2+} titrations. The lower panels show integrated binding isotherms fit to 2-site (WT) or 1-site (EF2⁻ and EF34⁻) model as described in Methods (solid lines). Optimal fitting parameters are displayed in Table 2.




Figure 4.

DSC scans showing thermal denaturation of GCAP1. Data were obtained for unmyristoylated GCAP1 (A) and myristoylated GCAP1 (B). Thermograms of apo, Mg²⁺-bound and Ca²⁺-bound GCAP1 are indicated by solid, dashed and dotted lines, respectively.




Figure 5.

NMR spectra of myristoylated and unmyristoylated GCAP1. Two-dimensional (^1H - ^{15}N HSQC) NMR spectra of ^{15}N -labeled unmyristoylated (black) and myristoylated GCAP1 (red) in the apo (A), Mg^{2+} -bound (B), Ca^{2+} -bound (C), and detergent solubilized, Ca^{2+} -bound (D) states. Spectra in A-C were obtained at 37 °C and the spectrum in D was obtained in the presence of 25 mM octylglucoside at 47 °C.




Figure 6.

Two-dimensional ^1H - ^{13}C -HMQC NMR spectra of ^{13}C -labeled myristoyl group attached to unlabeled GCAP1 in the (A) Ca^{2+} -bound state, (B) Mg^{2+} -bound state, (C) apo state (1 mM EGTA), and (D) apo state in the presence of 8 M urea. The $^{13}\text{C}_2$ resonance of myristate is folded and aliased shown in the upper left corner of the spectra. Extraneous peaks represent ^{13}C natural abundance signals from the protein as determined by recording ^1H - ^{13}C HMQC spectra on GCAP1 samples with an unlabeled myristoyl group.

Table 1
Thermodynamic Parameters of Ca²⁺ Binding to GCAP1 and Mutants at 30 °C.

	K _{EF2}	K _{EF3}	K _{EF4}	ΔH _{EF2}	ΔH _{EF3}	ΔH _{EF4}
0 Mg²⁺						
Wildtype	0.9 ±0.2	0.08 ±0.05	0.2 ±0.05	+3.1 ±0.1	-3.5 ±0.1	-0.9 ±0.2
EF2-	-	0.2 ±0.1	0.7 ±0.5	-	-3.0 ±0.1	-0.05±0.1
EF4-	69 ±20	4.9 ±0.9	-	+0.8 ±0.1	-3.4 ±0.1	-
EF34-	11 ±1	-	-	+2.1 ±0.2	-	-
2 mM Mg²⁺						
Wildtype	1.6 ±0.3	0.1 ±0.05	0.2 ±0.05	+1.4 ±0.1	-3.9 ±0.1	-0.6 ±0.2
EF2-	-	0.2 ±0.1	2.0 ±0.4	-	-4.2 ±0.1	-0.5±0.2
EF4-	1.1 ±0.2	0.4 ±0.1	-	+0.4 ±0.1	-4.0 ±0.1	-
EF34-	50 ±20	-	-	+1.6 ±0.1	-	-

Dissociation constants (K_{EF2}, K_{EF3}, K_{EF4}) are expressed in micromolar. The enthalpy differences (ΔH_{EF2}, ΔH_{EF3}, ΔH_{EF4}) are in units of kcal/mol.

Table 2Thermodynamic Parameters of Mg²⁺ Binding to GCAP1 and Mutants at 30 °C.

	K_{EF2}	K_{low}	ΔH_{EF2}	ΔH_{low}
Wildtype	0.7 ±0.2	2 ±0.8	+4.3 ±0.1	+0.9 ±0.5
EF2-	-	2 ±0.8	-	+2.9 ±0.2
EF4-	0.7 ±0.2	2 ±0.8	+0.5 ±0.3	+3.9 ±0.3
EF34-	0.7 ±0.2	-	+4.2 ±0.1	-

Dissociation constants (K_{EF2} and K_{low} - low-affinity phase) are expressed in millimolar. The enthalpy differences (ΔH_{EF2} and ΔH_{low}) are in units of kcal/mol.

Table 3

^1H and ^{13}C (in parentheses) chemical shift assignments of myristoyl resonances.

Position	Mg^{2+} -GCAP1 ^a	Ca^{2+} -GCAP1 ^b	denatured GCAP1 ^c
C2	2.34 (38.5)	2.34(38.4)	2.35(38.2)
C3	1.5 (27.8)	1.5 (27.5)	1.55 (27.0)
C4 - C11	0.9–1.1 (31–33)	0.9–1.1 (31–33)	1.2–1.3 (32–33)
C12	0.88 (34.0)	0.92 (33.8)	1.27 (33.5)
C13	0.95 (24.4)	0.98 (24.2)	1.0 (25)
C14	0.62, 0.73 (16.5)	0.62, 0.73 (16.8)	0.92 (13)

The NMR sample conditions were 0.3 mM GCAP1 in 5 mM Tris at pH 7.4 containing either 5 mM Mg^{2+} ^a, 5 mM Ca^{2+} ^b, or 8 M urea^c.