

Protection From Clinical Peripheral Sensory Neuropathy in Alström Syndrome in Contrast to Early-Onset Type 2 Diabetes

RICHARD B. PAISEY, MD¹
 ROSAMUND M. PAISEY, RGN¹
 MARY P. THOMSON, PHD¹
 LYNNE BOWER¹

PIETRO MAFFEI, MD³
 JULIAN P.H. SHIELD, MD²
 SUE BARNETT, PHD²
 JAN D. MARSHALL⁴

OBJECTIVE — Alström syndrome, with type 2 diabetes, and blindness could confer a high risk of foot ulceration. Clinical testing for neuropathy in Alström syndrome and matched young-onset type 2 diabetic subjects was therefore undertaken.

RESEARCH DESIGN AND METHODS — Fifty-eight subjects with Alström syndrome (18 insulin-resistant nondiabetic and 40 diabetic; aged 8–43 years) and 30 young-onset diabetic subjects (aged 13–35 years) were studied. Neuropathy symptom questionnaires were administered. Graded monofilament and 128-MHz tuning fork vibration perception were assessed in both feet.

RESULTS — Neuropathic symptoms, loss of monofilament, and/or vibration perception were reported by 12 of the 30 young-onset type 2 diabetic subjects (6 had neuropathic ulceration) but none of the subjects with Alström syndrome.

CONCLUSIONS — The striking preservation of protective foot sensation in Alström syndrome may provide a clue to the causes of differential susceptibility to neuropathy in the wider diabetic population.

Diabetes Care 32:462–464, 2009

Alström syndrome is a rare autosomal recessive condition characterized by cone-rod dystrophy and childhood blindness, obesity and insulin resistance leading to type 2 diabetes in adolescence, and hyperlipidemia (1,2). Mutations in the *ALMS1* gene have been described in the majority of cases (3,4). The accompanying blindness might increase the risk of foot ulceration if peripheral neuropathy were to develop. Therefore, a systematic foot examination of 58 subjects with Alström syndrome and 30 young-onset type 2 diabetic subjects was undertaken.

RESEARCH DESIGN AND METHODS

Ethics committees of Torbay Hospital, Bristol Royal Hospital

for Children, and The Jackson Laboratory gave study approval. Subjects with Alström syndrome were studied at Alström Syndrome U.K. and Alström Syndrome International clinics. Fifty-eight subjects with Alström syndrome and 30 young-onset (<25 years of age at diagnosis) type 2 diabetic subjects; 10 of 12 patients from the adolescent type 2 diabetes register at Bristol Royal Hospital for Children; and 20 of 22 patients from the diabetes retinal screening register of South Devon Healthcare, Torbay Hospital were included.

Clinical protocol

A validated questionnaire (available in an online appendix at <http://dx.doi.org/10.2337/dc08-1584>) seeking symptoms of bilateral neuropathic pain was adminis-

tered to all subjects by three trained investigators. Research-grade 2-, 4-, 6-, 8-, 10-, and 15-g monofilaments (Bailey Instruments, Manchester, U.K.) and 128-MHz tuning forks were used to test for protective sensation in subjects with eyes closed (5) (see Clinical Protocol available in the online appendix). Briefly, monofilaments were bounced to warm them up and applied firmly to six sites on each foot with a variable pause between tests to exclude false-positive responses. Calloused sites were avoided. Preservation of vibration perception was recorded if the subject sensed vibration at the tip of the hallux for 3 s and correctly identified when damped. Serum lipids and A1C were taken from patient records in the U.S. and from clinical laboratories at Torbay Hospital and Bristol Royal Hospital for Children in the U.K.

Statistics

For statistical testing, parametric ANOVA was used for BMI and A1C and the Kruskal-Wallis test was used for monofilaments. For pulses, 128 V, and reflexes, Fisher's exact test was used. Statistical computations were performed with Foundation for Statistical Computing software (R2.6.1; <http://www.r-project.org>). The significance threshold was set at $P \leq 0.05$.

RESULTS — The Alström diabetic subjects ($n = 40$) and young-onset type 2 diabetic patients ($n = 30$) are well matched (Table A1 of the online appendix) except for duration of diabetes, which is longer in the Alström diabetic group: for type 2 diabetes vs. Alström syndrome diabetes, respectively, duration of diabetes 4.6 ± 3.2 vs. 13.8 ± 2.8 years; A1C 8.6 ± 2.5 vs. $9.1 \pm 1.5\%$, $P = 0.14$; serum cholesterol 5.3 ± 0.9 vs. 6.1 ± 1.3 mmol/l, $P = 0.25$; and serum triglyceride 3.7 ± 1.9 vs. 0.8 ± 6.0 mmol/l, log-transformed $P = 0.15$. Table 1 shows the prevalence of neuropathic symptoms and loss of protective sensation in each group of subjects. No Alström syndrome subjects manifested typical neuropathic pain or absence of vibration perception. All perceived 6-g monofilament stimuli at all

From ¹Torbay Hospital, Torquay, U.K.; the ²Bristol Royal Hospital for Children, Bristol, U.K.; the ³Department of Medical and Surgical Science, University School of Medicine, Padua, Italy; and ⁴The Jackson Laboratory, Bar Harbor, Maine.

Corresponding author: Richard B. Paisey, richard.paisey@nhs.net.

Received 29 August 2008 and accepted 4 December 2008.

Published ahead of print at <http://care.diabetesjournals.org> on 17 December 2008. DOI: 10.2337/dc08-1584.

© 2009 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Table 1—Significance testing between young-onset type 2 diabetic subjects and Alström syndrome subjects with diabetes

Parameter	Alström syndrome with no diabetes	Alström syndrome with diabetes	Type 2 diabetes	Type 2 diabetes vs. Alström syndrome with diabetes (P)
n	18	40	30	
Neuropathic symptoms (%)	0	0	30.1	<0.0001
Perception				
all 2-g monofilament (%)	80	95	36	0.001
all 10-g monofilament (%)	100	100	38	0.001
128-MHz vibration (%)	100	97	72	0.004
Neuropathy (%)	0	0	34.6	0.0006
Foot ulceration	0	0	6	<0.0001

The Kruskal-Wallis test for monofilament perception and Fisher's exact test for 128 V (128-MHz tuning fork perception) were used. See text for definition of neuropathy and characteristics of symptoms.

sites, and >80% perceived all 2-g tests. None had present or past foot ulcers.

In contrast, neuropathic symptoms, absent vibration perception, or impairment of 6-g or 10-g monofilament perception was found in 14, 11, 7, and 12 of the control young-onset type 2 diabetic subjects, respectively. Six had neuropathic ulcers, and one had a bilateral ulcer.

Statistical analysis showed highly significant differences between Alström syndrome diabetic and young-onset type 2 diabetic patients with respect to presence of neuropathic symptoms ($P < 0.0001$), absence of vibration perception ($P = 0.004$), and mean lightest perceived monofilament ($P = 0.0001$).

CONCLUSIONS— Impairment of vibration sense and/or monofilament perception at 10 g is strongly predictive of future ulceration in diabetes (6,7). Our findings with graded monofilaments and vibration perception have confirmed the high prevalence of peripheral sensory loss, neuropathy, and ulceration in a small group of young/adolescent-onset type 2 diabetic individuals.

Alström syndrome diabetic subjects maintained good protective sensation despite comparable hyperglycemia and dyslipidemia (online appendix). This finding is encouraging, as it confirms that Alström syndrome subjects can undertake exercise and domestic activities with low risk of foot ulceration. The freedom from clinical signs of neuropathy suggests the possibility of a protective factor, associated with the syndrome, that may, when identified, increase understanding of the causes of diabetic neuropathy and suggest novel therapeutic interventions. Studies of nerve conduction in these patients are needed to confirm these findings.

Alström syndrome patients could be protected from clinical diabetic neuropathy because of their short stature (8,9), though the mean height for Alström syndrome patients was not significantly different from that for control subjects. It was recently reported that patients with Alström syndrome have subtle impairments in the growth hormone-IGF axis with a reduction in acid labile subfraction and IGF binding protein-1, whereas IGF binding protein-2 was shown to be markedly increased (10). These alterations could protect against microvascular complications as in the sex-linked form of ateliotic dwarfism (11,12).

The finding that the ALMS1 protein localizes intracellularly to the centrosome and may therefore influence microtubular function has led to speculation that transport of GLUT1-5 receptors to the cell surface may be impaired in Alström syndrome (13). Underexpression of GLUT1 receptors might protect neurons from hyperglycemic metabolic insult in those with diabetes and Alström syndrome. Further studies evaluating the roles of ALMS1 protein and microtubular function in normal neuronal function and neuropathies are strongly indicated.

Acknowledgments— Funding was received from Alström Syndrome International, Alström Syndrome U.K., the special medical grant fund of Torbay Hospital, and the National Clinical Specialist Services Group of the Department of Health, U.K.

No potential conflicts of interest relevant to this article were reported.

We thank all of the patients and families who helped with the study and Alström Syndrome International and Alström Syndrome U.K.

References

- Alstrom CH, Hallgren B, Nilsson LB, Asander H: Retinal degeneration combined with obesity, diabetes mellitus and neurogenous deafness: a specific syndrome (not hitherto described) distinct from the Laurence-Moon-Bardet-Biedl syndrome: a clinical, endocrinological and genetic examination based on a large pedigree. *Acta Psychiatr Neurol Scand Suppl* 129:1-35, 1959
- Marshall JD, Bronson RT, Collin GB, Nordstrom AD, Maffei P, Paisey RB, Carey C, Macdermott S, Russell-Eggitt I, Shea SE, Davis J, Beck S, Shatirishvili G, Mihai CM, Hoeltzenbein M, Pozzan GB, Hopkinson I, Siculo N, Naggert JK, Nishina PM: New Alström syndrome phenotypes based on the evaluation of 182 cases. *Arch Intern Med* 165:675-683, 2005
- Hearn T, Renforth GL, Spalluto C, Hanley NA, Piper K, Brickwood S, White C, Connolly V, Taylor JF, Russell-Eggitt I, Bonneau D, Walker M, Wilson DI: Mutation of ALMS1, a large gene with a tandem repeat encoding 47 amino acids, causes Alström syndrome. *Nat Genet* 31:79-83, 2002
- Collin GB, Marshall JD, Ikeda A, So WV, Russell-Eggitt I, Maffei P, Beck S, Boerkoel CF, Siculo N, Martin M, Nishina PM, Naggert JK: Mutations in ALMS1 cause obesity, type 2 diabetes and neurosensory degeneration in Alström syndrome. *Nat Genet* 31:74-78, 2002
- Thomson MP, Potter J, Finch PM, Paisey RB: Threshold for detection of diabetic peripheral sensory neuropathy using a range of research grade monofilaments in persons with Type 2 diabetes mellitus. *J Foot Ankle Res* 1:9, 2008
- Pham H, Armstrong DG, Harvey C, Harkless LB, Giurini JM, Veves A: Screening techniques to identify people at high risk for diabetic foot ulceration: a prospective multicenter trial. *Diabetes Care* 23:606-611, 2000
- Abbott CA, Carrington AL, Ashe H, Bath S, Every LC, Griffiths J, Hann AW, Hussein A, Jackson N, Johnson KE, Ryder CH, Tork-

- ington R, Van Ross ER, Whalley AM, Widdows P, Williamson S, Boulton AJ, North-West Diabetes Foot Care Study: The North-West Diabetes Foot Care Study: incidence of, and risk factors for, new diabetic foot ulceration in a community-based patient cohort. *Diabet Med* 19:377–384, 2002
8. Sosenko JM, Gadia MT, Fournier AM, O'Connell MT, Aguiar MC, Skyler JS: Body stature as a risk factor for diabetic sensory neuropathy. *Am J Med* 80:1031–1034, 1986
9. Sosenko JM, Boulton AJ, Gadia MT, Ward JD, Skyler JS: The association between symptomatic sensory neuropathy and body stature in diabetic patients. *Diabetes Res Clin Pract* 4:95–98, 1988
10. Maffei P, Boschetti M, Marshall JD, Paisey RB, Beck S, Resmini E, Collin GB, Naggert JK, Milan G, Vettor R, Minuto F, Siculo N, Barreca A: Characterization of the IGF system in 15 patients with Alström syndrome. *Clin Endocrinol (Oxf)* 66:269–275, 2007
11. Merimee TJ, Hall JD, Rimoin DL, McKusick VA: A metabolic and hormonal basis for classifying ateliotic dwarfs. *Lancet* 1:963–965, 1969
12. Merimee TJ, Siperstein MD, Hall JD, Fineberg SE: Capillary basement membrane structure: a comparative study of diabetics and sexual ateliotic dwarfs. *J Clin Invest* 49:2161–2164, 1970
13. Hearn T, Spalluto C, Phillips VJ, Renforth GL, Copin N, Hanley NA, Wilson DI: Subcellular localization of ALMS1 supports involvement of centrosome and basal body dysfunction in the pathogenesis of obesity, insulin resistance, and type 2 diabetes. *Diabetes* 54:1581–1587, 2005