

Silencing of the Transforming Growth Factor- β (TGF β) Receptor II by Krüppel-like Factor 14 Underscores the Importance of a Negative Feedback Mechanism in TGF β Signaling^{*§}

Received for publication, October 8, 2008, and in revised form, December 8, 2008. Published, JBC Papers in Press, December 15, 2008, DOI 10.1074/jbc.M807791200

Mark J. Truty^{‡§1}, Gwen Lomber[‡], Martin E. Fernandez-Zapico^{‡2}, and Raul Urrutia^{‡3}

From the [‡]Department of Biochemistry and Molecular Biology and Department of Physiology and Biophysics, Gastroenterology Research Unit, and the [§]Department of Surgery, Clinician Investigator Training Program, Mayo Clinic, Rochester, Minnesota 55905

The role of non-Smad proteins in the regulation of transforming growth factor- β (TGF β) signaling is an emerging line of active investigation. Here, we characterize the role of KLF14, as a TGF β -inducible, non-Smad protein that silences the TGF β receptor II (*TGF β RII*) promoter. Together with endocytosis, transcriptional silencing is a critical mechanism for down-regulating TGF β receptors at the cell surface. However, the mechanisms underlying transcriptional repression of these receptors remain poorly understood. KLF14 has been chosen from a comprehensive screen of 24 members of the Sp/KLF family due to its TGF β inducibility, its ability to regulate the *TGF β RII* promoter, and the fact that this protein had yet to be functionally characterized. We find that KLF14 represses the *TGF β RII*, a function that is augmented by TGF β treatment. Mapping of the *TGF β RII* promoter, in combination with site-directed mutagenesis, electromobility shift, and chromatin immunoprecipitation assays, have identified distinct GC-rich sequences used by KLF14 to regulate this promoter. Mechanistically, KLF14 represses the *TGF β RII* promoter via a co-repressor complex containing mSin3A and HDAC2. Furthermore, the TGF β pathway activation leads to recruitment of a KLF14-mSin3A-HDAC2 repressor complex to the *TGF β RII* promoter, as well as the remodeling of chromatin to increase histone marks that associate with transcriptional silencing. Thus, these results describe a novel negative-feedback mechanism by which TGF β RII activation at the cell surface induces the expression of KLF14 to ultimately silence the *TGF β RII* and further

expand the network of non-Smad transcription factors that participate in the TGF β pathway.

The family of cytokines composed of TGF β ,⁴ bone morphogenetic proteins, activins, inhibins, connective tissue growth factors (CCN family), along with their corresponding signaling molecules, are master regulators of normal homeostasis and development (1–10). Consequently, alterations in these pathways lead to severe malformations and diseases, including cancer. TGF β is the best characterized pathway within this family of cytokines. Recent studies reveal, for instance, the existence of two types of membrane-to-nucleus TGF β signaling mechanisms, namely the Smad-dependent and non-Smad protein-mediated cascades, although evidence of cross-talk between these two cascades is also emerging (4, 9). Therefore, even though our understanding of the complexity underlying TGF β signaling continues to grow, classification into these two types of mechanisms has helped to organize the nascent theoretical framework for advancing this field of research by the integration of new findings into easily understandable paradigms.

The canonical Smad-mediated TGF β pathway is activated by binding of TGF β 1, -2, and/or -3 cytokines to the *TGF β RII*, which then dimerizes with and activates the TGF β receptor I through serine phosphorylation of the regulatory GS-domain. The Type I receptor, in turn, phosphorylates receptor-bound Smad (Smad2/3) at the C-terminal SXS motif, releasing them from retention in the cytoplasm and allowing their translocation into the nucleus. Smad4 acts as a common partner of activated Smads to help execute their function. In this manner, TGF β signaling is transduced through the cytoplasm into the nucleus to form complexes with distinct transcriptional regulators for specific gene promoters.

The role of non-Smad protein-mediated pathways in the regulation of TGF β signaling is also an active line of investigation. For instance, the Sp/KLF family of proteins is emerging as important non-Smad protein-mediated pathway cascades and, under certain circumstances, a cross-talk regulator with Smads

* This work was supported, in whole or in part, by National Institutes of Health Grant DK 52913. This work was also supported by Mayo Clinic Pancreatic SPORE (Grant P50 CA102701 to R. U.). The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

We dedicate this manuscript to the memory of Kazimierz Truty and to all the other patients who have succumbed to pancreatic cancer.

§ Author's Choice—Final version full access.

§ The on-line version of this article (available at <http://www.jbc.org>) contains supplemental Table S1.

¹ Supported by the Mayo Clinic Clinician Investigator Training Program and the Dept. of Surgery.

² Supported by National Institute of Health Grant CA125127, Mayo Clinic Pancreatic SPORE Career Development Award (Grant P50 CA102701), in honor of Carole and Bob Daly, Mayo Clinic Cancer Center, and the Division of Gastroenterology (Mayo Clinic College of Medicine).

³ To whom correspondence should be addressed: 200 First St. SW, GI Research Unit, Guggenheim 10, Mayo Clinic, Rochester, MN 55905. Tel.: 507-255-6029; Fax: 507-255-6318; E-mail: urrutia.raul@mayo.edu.

⁴ The abbreviations used are: TGF β , transforming growth factor- β ; *TGF β RII*, TGF β receptor type II; CMV, cytomegalovirus; RT, reverse transcription; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; B2M, β_2 -microglobulin; TUBB2, β_2 -Tubulin; HPRT1, hypoxanthine phosphoribosyltransferase 1; GST, glutathione S-transferase; HDAC, histone deacetylase; WT, wild type; KLF14, Krüppel-like factor 14; ChIP, chromatin immunoprecipitation.

KLF14, a Non-Smad Repressor of the TGF β Receptor II

to achieve distinct cellular functions. Sp1 is the founding member this expanding group of Sp/KLF proteins. The structure of these proteins is defined by the presence of three highly conserved and homologous C-terminal Cys₂His₂ zinc finger domains, which are responsible for DNA binding, and a variable N-terminal domain, which is responsible for transcriptional regulation (11–13). However, the identification and characterization of this family of proteins has revealed that many bind to GC-rich target sequences similar, if not identical to, the “Sp1 sites” through which they can either activate or repress gene expression (11–13). Therefore, the discovery of repressors within this Sp/KLF family of transcriptional regulators has challenged the early paradigm that Sp1 activates all GC-rich sites. As a result, these Sp/KLF transcriptional repressors provide a novel mechanism for silencing a large number of genes that are already known to be activated by Sp1, particularly in response to TGF β .

TGF β RII has been previously shown to be activated by Sp1 (14). However, since these elegant studies were done, 24 Sp/KLF transcription factors have been discovered with some members acting as activators while others as repressors via the same type of GC-rich *cis* regulatory sequences used by Sp1. Thus, some Sp/KLF transcription factors are excellent candidates that may play a role in silencing of *TGF β RII*. Indeed, fortunately, in the current study, we describe, for the first time, the functional characterization of KLF14 as a novel non-Smad regulatory protein of the TGF β pathway. Our results outline a novel, biochemically significant role for KLF14 in the silencing of the *TGF β RII* via Sp/KLF sites. This pathway provides a well characterized example of how Sp/KLF proteins are emerging as important non-Smad proteins that can directly regulate TGF β signaling by regulating the expression of key molecules from this pathway.

EXPERIMENTAL PROCEDURES

Cell Culture—Tissue culture reagents were purchased from commercial sources (Invitrogen). The human pancreatic epithelial cancer cell lines, PANC-1, ASPC-1, Capan-1, Capan-2, BxPC-3, L3.6, MiaPaCa-2, and CFPAC-1, were obtained from American Type Culture Collection and maintained according to the supplier's suggestions. All cells were grown at 37 °C in a humidified incubator under 5% CO₂.

Plasmid Construction—Standard molecular biology techniques were used to clone *KLF4*, *KLF5*, *KLF7*, *KLF9*, *KLF11*, *KLF14*, and *KLF15* into the pcDNA3.1/His (Invitrogen) and pCMVtag2 (Stratagene) vectors for expression as His-tagged or FLAG-tagged proteins, respectively, as well as the truncated (263 bp) *TGF β RII* promoter into the pGL3-Lux vector (Promega). The p3TP-Lux reporter plasmid containing TGF β -responsive elements was kindly provided by Dr. Anita Roberts (National Institutes of Health) as a positive control for TGF β 1 stimulation experiments (data not shown). Full-length *TGF β RII* reporter was kindly provided by Dr. David Danielpour (Case Western).

Semi-quantitative RT-PCR—Total RNA was extracted from cells according to the manufacturer's instructions using an RNeasy Kit (Qiagen), and 5 μ g was used for cDNA synthesis using oligo(dT) primer using the SuperScriptTM III First-

Strand Synthesis System for RT-PCR (Invitrogen) per the manufacturer's protocol. RT-PCR was performed using LA TaqDNA Polymerase with a GC Buffers kit (Takara) per the manufacturer protocols. Semiquantitative RT-PCR analysis was performed with the primer sets provided in supplemental Table S1. Each experiment was done in triplicate. Amplification of four human housekeeping genes, *GAPDH* (UniGene Hs.544577), β_2 -microglobulin (*B2M*, Hs.709313), β_2 -Tubulin (*TUBB2*, Hs.300701), and hypoxanthine phosphoribosyltransferase 1 (*HPRT1*, Hs.412707) was used for all samples as an internal control. Densitometric values were obtained and normalized to the average of the housekeeping cDNAs for each individual sample using Scion Image Beta 4.02 software (Scion Corp.).

To represent the TGF β inducibility of individual *KLF* transcripts, each TGF β -treated sample was compared with an untreated control that was originally plated and ultimately collected at the same time to minimize compounding factors, which can often influence gene expression (*i.e.* cell cycle stage, cell density, potential unknown paracrine, and/or autocrine stimuli). At each time point, values were determined by first normalizing the TGF β -treated sample (T, treated) to the average of the four aforementioned housekeeping genes in the same sample (TC, treated housekeeping gene control). This resultant value was divided by the corresponding untreated sample (UT, untreated) normalized in the same manner (UTC, untreated housekeeping gene control), to express the fold of TGF β induction ($[T/TC]/[UT/UTC]$ = fold TGF β -induction). Another housekeeping gene, β -actin (*ACTB*, Hs.520640), was not used in our analyses, because it showed significant differences with TGF β treatment over control values (data not shown).

Western Blot—Total protein extracts were prepared by lysing cells in radioimmune precipitation assay buffer supplemented with Complete protease inhibitor mixture (Roche Applied Science). Cellular lysates were subjected to 10% SDS-PAGE and then separated proteins are transferred to polyvinylidene difluoride membranes (Millipore). Membranes were incubated overnight at 4 °C in blocking solution (Tris-buffered saline solution containing 5% nonfat dried milk and 0.1% Tween 20). Subsequently, membranes were incubated with specified primary antibodies overnight at 4 °C. Immune complexes were visualized by enhanced chemiluminescence (Pierce) and exposed to x-ray film. An antibody against β 2-actin (Sigma) was used as loading control.

Transcriptional Reporter Assays—Cells were transfected with specified reporter constructs along with expression constructs and/or empty vector using electroporation (2×10^6 cells/0.4-cm microcuvette, 360 V, and 10 ms) and subsequently serum-starved overnight. Transfection was performed with equimolar concentrations of DNA, and expression was quantified with Western blotting directed against epitope-tagged proteins as described. Cells were stimulated with TGF β 1 (R&D Systems) as specified and assayed at various specified time points. At 24 or 48 h after transfection and treatment as noted, cells were lysed, and luciferase measurements were performed using a 20/20 luminometer (Turner Designs) according to manufacturer's suggestions (Promega). Data were normalized as relative light units and normalized to the protein concentra-

tion as the mean \pm S.D. All experiments were performed in triplicate at least three independent times.

Immunoprecipitation—Cells were transfected with FLAG-tagged constructs. At 24 h post-transfection, cells were washed and lysed in lysis buffer (150 mM NaCl, 0.5% Nonidet P-40, 50 mM Tris-HCl, pH 7.5, 20 mM MgCl₂) supplemented with Complete protease inhibitor tablets (Roche Applied Science) for 30 min at 4 °C. Immunoprecipitations were performed using anti-FLAG M2 agarose-conjugated antibodies (Sigma) for 2 h at 4 °C. To detect interaction with endogenous co-repressors, immunocomplexes were collected by centrifugation, washed with lysis buffer, and analyzed by Western blot as described above using anti-mSin3a and HDAC2 antibodies (Santa Cruz Biotechnology).

Chromatin Immunoprecipitation—ChIP assays were performed using the EZ-ChIP kit. The following primer set for the 263-bp TGF β RII promoter was used for PCR: 5'-GCA GAT GTT CTG ATC TAC TA-3' (forward); 5'-AGC TGG GCA GGA CCT CTC TC-3' (reverse) using TaKaRa LA *Taq* according to the manufacturer's protocol (Mirus).

Site-directed Mutagenesis—Site-directed mutagenesis was performed with QuikChange® II site-directed mutagenesis kits per the manufacturer's protocol (Stratagene). All constructs were sequenced by the Mayo Clinic Molecular Biology Core Facility.

GST Fusion—The *KLF14* cDNA fragment encoding amino acids 191–323 corresponding to the DNA-binding zinc finger region was cloned into the GST fusion vector pGEX 5X-1 (Amersham Biosciences) using standard techniques. GST fusion protein expression was induced in BL21 cells (Stratagene) by the addition of 1 mM isopropyl-D-thiogalactopyranoside and incubation for 2 h. Cells were lysed and subsequently purified by using glutathione-Sepharose 4B affinity chromatography as previously described (15).

Electromobility Shift Assay—Gel shift assays were performed as previously described (16). Briefly, 1.75 pmol of double-stranded oligonucleotides were end-labeled with [γ -³²P]ATP using 10 units of T4 polynucleotide kinase and appropriate buffer (700 mM Tris-HCl, pH 7.6, 100 mM MgCl₂, 50 mM dithiothreitol) according to the manufacturer's instructions (Promega). The reaction was incubated at 37 °C for 10 min and halted with addition of TE plus EDTA (0.5 M EDTA, 1 M Tris, pH 8.0, H₂O). Protein lysates included 0.5 μ g of purified GST-KLF14/ZF fusion protein and in some experiments rhSP1 (Promega) at the indicated dilutions. A 5 \times ZnCl₂ buffer was used in this reaction (100 mM Hepes, pH 7.5, 250 mM KCl, 25 mM MgCl₂, 50 μ M ZnCl₂, 30% glycerol, 1 mg/ml bovine serum albumin, 250 μ g/ml poly(dI-dC), H₂O) for 10 min at room temperature. The γ -³²P-labeled oligonucleotides were added for 20 min. In some cases, an excess of cold probe, at the indicated dilutions, was added concomitant with the addition of radiolabeled probe in addition to anti-Sp1 polyclonal rabbit antibody purchased from commercial sources (Millipore). The mixtures were electrophoresed in a 4% nondenaturing polyacrylamide gel in a Hoeffer midi-gel using 0.5 \times Tris borate-EDTA for ~4 h at 160 V. Gels were then transferred to blotting paper (Whatman 3MM), covered in plastic wrap, and vacuum dried for 1.5 h at 65 °C. Dried gels were then analyzed using a Storm Scanner




FIGURE 1. *A*, repression of the TGF β RII promoter occurs after 24 h of TGF β stimulation in Panc1 cells. Panc1 cells were transfected with a full-length and a truncated 263-bp TGF β RII promoter luciferase reporter and then stimulated with TGF β 1 after overnight serum starvation. Luciferase levels were obtained at specified time points after treatment and compared with untreated controls. Values were normalized to lysate protein concentrations and relative to untreated controls. Data are the mean \pm S.D. from three independent experiments, with triplicates for each experiment. *B*, TGF β RII promoter has five putative "Sp1" binding sites. The diagram represents a schematic of the human TGF β RII proximal promoter containing five GC-rich Sp1-like elements.

860 PhosphorImager (Amersham Biosciences). Sp1 consensus and Sp1 mutant double-stranded oligonucleotides were obtained from commercial sources (Santa Cruz Biotechnology) with the following sequences: Sp1 consensus, 5'-ATT CGA TCG GGG CGG GGC GAG C-3' (forward); 5'-GCT CGC CCC GCC CCG ATC GAA T-3' (reverse) and Sp1 mutant, 5'-ATT CGA TCG GTT CGG GGC GAG C-3' (forward); 5'-GCT CGC CCC GAA CCG ATC GAA T-3' (reverse).

RESULTS

A Yet Undefined Sp/KLF Repressor Protein Plays a Role in Silencing of the Type II TGF β -receptor Promoter—Previous studies have described the regulation of the TGF β RII by TGF β ligands, primarily showing a bimodal response consisting of an Sp1-dependent up-regulation (14, 17) and a subsequent down-regulation of receptor transcript levels upon this stimulation (18–20). Interestingly, although the activation of TGF β RII promoter, in particular by Sp1, has received precise attention, how this receptor is repressed remains poorly understood. Consequently, the major goal of the current study has been to characterize the role of a specific family of non-Smad proteins (Sp/KLF transcription factors), which may functionally explain the down-regulation TGF β RII through GC-rich Sp1-like sequences. Our studies began with examination of the transcriptional activity of the TGF β RII gene promoter in PANC1 epithelial cells, a widely used model for studying TGF β signaling. We have performed an initial series of reporter assays using full-length TGF β RII and the previously described, TGF β -sensitive, 263-bp core promoter, which is located 5' of the transcriptional start site (14). Treatment of PANC1 cells with exogenous TGF β 1 leads to a marked reduction in activity of the full-length TGF β RII reporter when compared with untreated control cells (Fig. 1A). This silencing effect is recapitulated in the 263-bp TGF β RII core promoter, suggesting that both the

KLF14, a Non-Smad Repressor of the TGF β Receptor II

previously described activation pathway (14) and the negative transcriptional regulatory mechanism characterized further here (Fig. 1A), are operational at this core promoter level. Using bioinformatics analyses (TRANSFAC Public), we have identified five putative Sp/KLF binding sites within this TGF β -sensitive, 263-bp core promoter region (Fig. 1B). Four of these sites (#1–4) have been previously identified, although an additional site (#5) has not been previously reported (14, 21). Some of these previously identified sites have been shown to be activated by Sp1. However, whether novel Sp/KLF silencer proteins can also bind to this sequence to reverse the activation by Sp1 remained unknown. Therefore, this analysis has led us to the hypothesis that a yet undefined Sp/KLF repressor protein plays a role in the silencing of this promoter.

Novel TGF β -inducible Non-Smad, Sp/KLF Proteins Are Identified as Candidate Regulators of the Type II TGF β -receptor—To test our hypothesis, we have developed a four-tier screening approach. This approach includes first, testing which of the 24 known Sp/KLF proteins are expressed in TGF β -sensitive epithelial cells and, thus, can be considered initial candidates to target the TGF β RII promoter. Our experimental cell model, PANC1, a human epithelial cell line, is an optimal model for our studies, because they have adequate expression of TGF β RII mRNA and display growth inhibition to exogenous TGF β 1 stimulation (22). In addition, as shown in Fig. 2A, each of the 24 known Sp/KLF transcription factors are consistently expressed in these cells, which made it ideal for performing a comprehensive screen. Second, we determine whether any Sp/KLF genes are TGF β -inducible with a kinetic that is consistent with playing a role in the down-regulation of the TGF β RII. Third, by utilizing transfection studies combined with reporter assays, we test the potential of distinct members of this family to repress TGF β RII promoter activity and then, by electromobility shift assays, determine which sites on the promoter are utilized by our candidate KLF protein. Finally, we examine whether the repressor that is isolated according to these criteria binds to the endogenous TGF β RII gene and can remodel chromatin on this target, suggesting the bona fide target status of the candidate KLF protein. Thus, by applying this comprehensive screening, our study has been robust in evaluating the KLF protein family in the TGF β response and regulation of the TGF β RII.

The genomic axiom that functionally related genes follow a similar pattern of expression suggested that, hypothetically, the protein that represses the TGF β RII may be expressed in a similar manner after TGF β treatment, in particular, during the repression response to this cytokine. Consequently, we have evaluated which, if any, of these Sp/KLF members are inducible by TGF β 1 treatment. Thus, using RNA from PANC1 cells either untreated or treated with TGF β 1, we have performed RT-PCR at various time points (Fig. 2B). As a positive control for TGF β -mediated transcriptional induction, we monitor the expression of *p21*, a known TGF β -inducible gene within the pathway. Of the 24 known Sp/KLF transcription factors, we have identified 7 that were markedly induced with exogenous TGF β 1 treatment, suggesting that these could be potential candidate transcriptional repressors of the TGF β RII promoter (Fig. 2B). These results are not only consistent with our previous work, which identified *KLF11* as a TGF β -inducible gene




FIGURE 2. A, eight pancreatic cancer cell lines were screened by RT-PCR for expression of 24 members of the Sp/KLF family of transcription factors. The figure shows that all 24 Sp/KLF members are expressed in the TGF β -sensitive PANC1 cell line. B, TGF β 1 induces the expression of several KLF family members. PANC1 cells were serum-starved overnight, treated with 10 ng/ml TGF β 1, and screened for the expression of each KLF gene by semi-quantitative RT-PCR. The expression levels for each KLF gene from TGF β 1-treated samples were compared with untreated controls and normalized to housekeeping genes as described under "Material and Methods." The expression of the *p21* gene was used as a positive control for TGF β 1 treatment. Four different housekeeping genes, namely *GAPDH*, *B2M*, *TUBB2*, and *HPRT1*, were used as internal standards for normalization. C, quantification of TGF β induction of transcript levels over time. At each time point, values were determined by first normalizing the densitometric measurement of the TGF β -treated sample to the average of four housekeeping genes (*GAPDH*, *B2M*, *TUBB2*, and *HPRT1*) in the same sample. This resulting value was divided by the corresponding value of the untreated sample normalized in the same manner in order to express the -fold of TGF β induction.

(16, 23), but, more importantly, it characterizes previously unidentified KLF targets for this cascade, namely *KLF4*, *-5*, *-7*, *-9*, *-14*, and *-15*. Thus, based upon their expression patterns, these




FIGURE 3. *A*, Panc1 cells were transfected with KLF(FLAG/His) epitope-tagged expression constructs or control empty vector to test *TGF β RII* promoter activity. *B*, Panc1 cells were transfected with indicated KLF(FLAG/His) epitope-tagged expression constructs or control vector and then stimulated with 10 ng/ml TGF β 1 after overnight serum starvation to test *TGF β RII* promoter activity. *C*, Panc1 cells were transfected with various concentrations of epitope-tagged KLF14(FLAG) expression construct or control vector and then stimulated with 10 ng/ml TGF β 1 after overnight serum starvation to observe the effect on *TGF β RII* promoter activity. Western controls (FLAG/His) were shown for epitope-tagged KLF expression in all experiments. Expression of FLAG-tagged KLF4, KLF7, KLF14, and KLF15 was confirmed by an anti-FLAG antibody (Sigma), whereas expression of His-tagged KLF5, KLF9, and KLF11 was verified by the OMNI D8 antibody (Santa Cruz Biotechnology). Data are the mean \pm S.D. from three independent experiments, with triplicates for each experiment.

seven genes are good candidates to further investigate their regulation of the *TGF β RII* promoter.

Subsequently, we have tested the transcriptional activity of these candidates on the *TGF β RII* promoter using reporter assays by co-transfecting the 263-bp core *TGF β RII* promoter-luciferase construct with cDNAs encoding each of the seven TGF β -inducible KLF candidates. Out of these, five candidates induced a marked decrease in *TGF β RII* promoter activity, namely KLF4, -7, -11, -14, and -15 (Fig. 3*A*). KLF5 did not affect *TGF β RII* promoter activity above control levels, whereas KLF9 appeared to slightly activate this promoter, therefore these two proteins were not continued in subsequent experiments due to our objective of identifying *TGF β RII* repressors. To confirm whether this observed repression of *TGF β RII* promoter regulation is consistent with TGF β pathway activation, these five proteins were further tested in PANC1 cells treated with exogenous TGF β 1 treatment. Interestingly, upon treatment, these KLF proteins were capable of further repressing *TGF β RII* promoter activity, with the largest repression achieved by KLF14 (Fig. 3*B*). These data also indicate that a second TGF β -dependent mechanism, directly (signal-induced KLF expression or post-translational modifications) or indirectly (induction of another transcription factor), cooperates with KLF proteins to additionally repress this gene. Upon increasing concentrations of *KLF14* cDNA in transfection studies, we found a concentration-dependent repression of the core *TGF β RII* promoter (Fig. 3*C*). Together, these results strongly identify KLF14 as a good

candidate to be a regulator of *TGF β RII* promoter activity and expression.

KLF14, a Novel Non-Smad Protein, Regulates the Type II TGF β -receptor—The gene encoding KLF14 has been previously identified by our group and named *BTEB5* due to its sequence similarities to members of this subfamily of KLF silencing proteins.⁵ Although, recently, genetic studies have reported the genomic structure, intronless nature, and potential imprinted status of this gene (24), a functional characterization of this protein at the cellular and biochemical level has never been performed. Because our data indicate that KLF14 appears to be a potent regulator of the *TGF β RII* in promoter assays combined with this existing gap in knowledge on this KLF family member and its targets, the choice to further investigate the role of this particular KLF in *TGF β RII* regulation would significantly expand the current knowledge on the functional properties of members of this family.

Noteworthy, we have validated these *in vitro* reporter results *in vivo* by assessing whether this protein

has a regulatory effect on endogenous *TGF β RII* levels in PANC1 cells. Initial correlative experiments demonstrate that the levels of *TGF β RII* mRNA levels decrease at a time in which the amount of KLF14 increases (repression phase of the bimodal expression pattern of *TGF β RII* in response to TGF β), raising the possibility that *KLF14* is induced to subsequently down-regulate the *TGF β RII* (Fig. 4*A*). To mechanistically support this correlation, we have performed RT-PCR on cells overexpressing KLF14 to determine *TGF β RII* levels in comparison to mock transfected cells in the presence or absence of exogenous TGF β 1 stimulation (Fig. 4*B*). KLF14 overexpression alone is sufficient to decrease *TGF β RII* mRNA, interestingly to the same extent as TGF β 1 stimulation alone. Furthermore, subsequent TGF β 1 treatment in cells transfected with *KLF14* leads to further down-regulation of *TGF β RII* transcripts. Because *KLF14* is a TGF β -inducible gene, the further decrease in *TGF β RII* mRNA levels observed upon TGF β 1 treatment likely results from the induction of endogenous *Sp/KLF* transcription factors by this cytokine. These results suggest that TGF β down-regulates *TGF β RII* transcripts through *KLF14* expression with subsequent negative regulation of promoter activity. Next, we have tested whether the effect of KLF14 on *TGF β RII* expression interfered with TGF β -induced signals that target downstream genes, such as *p21* (25). Indeed, as

⁵ Reported by J. Kaczynski and R. Urrutia to NCBI, 2002.

KLF14, a Non-Smad Repressor of the TGF β Receptor II




FIGURE 4. *A*, TGF β RII and KLF14 transcript levels were analyzed by determination of density, and relative transcript levels were represented by the ratio TGF β RII cDNA/GAPDH cDNA and KLF14 cDNA/GAPDH cDNA. Data were the mean \pm S.D. from three independent experiments. *B*, KLF14 inhibits endogenous TGF β RII transcript levels. Panc1 cells were transfected with KLF14 or control empty vector and then stimulated with 10 ng/ml TGF β 1 for 24 h after overnight serum starvation. Densitometry was performed after normalizing to the average of GAPDH, B2M, TUBB2, and HPRT1 levels and untreated control cDNA. *C*, Panc1 cells were transfected with KLF14 FLAG epitope-tagged constructs or control empty vector along with a p21 promoter reporter construct and then stimulated with 10 ng/ml TGF β 1 after overnight serum starvation. Western control for KLF14(FLAG) epitope-tagged protein expression is shown. *D*, KLF14 inhibits endogenous p21 transcript levels. Panc1 cells were transfected with FLAG-KLF14 or control empty vector and then stimulated with 10 ng/ml TGF β 1 for 24 h after overnight serum starvation. Densitometry was obtained after normalizing to the average of GAPDH, B2M, TUBB2, and HPRT1 levels and untreated control cDNA.

expected, TGF β 1 treatment increases p21 promoter activity, as observed via reporter assays, as well as its mRNA levels (Fig. 4, *C* and *D*), whereas KLF14 reduces both p21 promoter activity and mRNA levels upon TGF β 1 treatment (Fig. 4, *C* and *D*). These results suggest that KLF14 interferes with the activation of downstream TGF β signaling effects, at least in part, by its ability to repress the TGF β RII.

KLF14 Represses the TGF β RII Promoter via Distinct Sp1-like GC-rich Sequences and Competition with Sp1—Repressor KLF proteins, such as KLF14, have been previously shown to compete with the canonical Sp1 protein for overall transcriptional activity of a promoter, such as the *CYP1A1* and *LDLR* promoters (26, 27). Therefore, herein we test the ability of KLF14 to repress transcription of the TGF β RII promoter in the presence




FIGURE 5. A, KLF14 and Sp1 compete to regulate *TGFBR11* promoter activity. Panc1 cells were transfected with KLF14(FLAG) and Sp1(His) epitope-tagged expression constructs or control empty vector as indicated to evaluate *in vivo* competition on *TGFBR11* promoter activity. Western control (FLAG/His) is shown for epitope-tagged expression. Data are the mean \pm S.D. from three independent experiments, each done in triplicate. B, KLF14 binds to four of the putative Sp1 sites and competes with Sp1 on the *TGFBR11* promoter. Electromobility shift assay was performed using KLF14-ZF GST fusion proteins and radiolabeled oligonucleotides for each of the five putative Sp1 sites. Lane 1: control-Sp1 consensus oligonucleotide; lane 2: control-Sp1 mutant oligonucleotide; lane 3: Sp1 consensus oligonucleotide and rhSP1 protein; lane 4: WT oligonucleotide; lane 5: mut#1 oligonucleotide; lane 6: mut#2 oligonucleotide; lane 7: WT oligonucleotide and 25 \times cold probe; lane 8: WT oligonucleotide and 100 \times cold probe; lane 9: WT oligonucleotide and anti-GST; lane 10: WT oligonucleotide and 1 \times rhSp1 protein; lane 11: WT oligonucleotide and 5 \times rhSp1 protein. KLF14 band (open arrowhead). Sp1 band (closed arrowhead). The depicted blots are representative of triplicate experiments. C, KLF14 utilizes four Sp/KLF sites to repress the *TGFBR11* promoter *in vivo*. Panc1 cells were transfected with *TGFBR11*-luciferase with mutated Sp1 sites as indicated and stimulated with 10 ng/ml TGF β 1 after overnight serum starvation, and transcriptional activity was measured and normalized to control empty vector. Data are the mean \pm S.D. from three independent experiments, with triplicates for each experiment.

of exogenous *Sp1* expression. Indeed, as shown in Fig. 5A, the expression of exogenous *Sp1* relieves the repression mediated by KLF14 in a dose-dependent manner. Due to the direct, Sp1-dependent up-regulation of the *TGFBR11* promoter as part of its bimodal response during activation, we dissect whether the repression of the *TGFBR11* promoter via KLF14 is also acting through a direct mechanism on the promoter rather than indirect due to KLF14 mediating a secondary effector.

Thus, to first delineate the DNA binding properties of KLF14 to the promoter of *TGFBR11* *in vitro*, we characterize the five putative Sp/KLF sites (shown in Fig. 1B) by electromobility shift assay (Fig. 5B). In this experiment, radiolabeled oligonucleotides for each of the 5 Sp1 sites have been created as well as mutants for each site containing either 2- or 4-nucleotide substitutions (mut#1 or mut#2, respectively) within the GC-rich binding sequence. Sp1-consensus oligonucleotides and respective mutants have been used as internal binding controls (Fig. 5B, lanes 1–3). The interaction between a recombinant KLF14 zinc finger DNA-binding domain and four of the Sp/KLF site

probes (#1–3 and 5) is specific as indicated by the fact that an excess of unlabeled probe competes with the radiolabeled probes (Fig. 5B, lanes 7 and 8). Furthermore, we were able to abolish this binding by using antibodies against the recombinant KLF protein (anti-GST antibody), thus confirming specificity of the complex (Fig. 5B, lane 9). Further specificity is evidenced by the impaired ability of recombinant KLF14 protein to bind to mutated Sp/KLF probes (Fig. 5B, lanes 5 and 6). Visible but reduced binding is noted with the 2-bp mutant probes (mut#1, lane 5) for sites #1–3, however a significant loss of binding was observed with the 4-bp mutant probes (mut#2, lane 6) at all sites. Moreover, through addition of recombinant Sp1 protein to the reaction, an increase in the concentration of Sp1 diminished the binding of KLF14, demonstrating that KLF14 and Sp1 compete for these GC-rich binding sites (Fig. 5B, lanes 10 and 11), complementing the competition results we observed *in vivo* (Fig. 5A). Only a weak complex was visualized with site #4, indicating that this site was less specific for KLF14 binding than the other four sites.

To functionally complement these studies, we have performed *TGFBR11* reporter assays with both wild-type and mutant promoter-luciferase constructs. Site-directed mutagenesis has been utilized to create a 2-nucleotide substitution in each of the 5 GC-rich Sp/KLF sites (Fig. 5C). These assays resulted in a marked loss of the repression observed with the wild-type *TGFBR11* promoter upon KLF14 overexpression with mutation in Sp/KLF sites #1, 2, 3, and 5, particularly with a significant gain of transcriptional activity upon mutating site #1. These findings demonstrate that these four are operational to silence the transcriptional activity of the *TGFBR11* promoter by KLF14. Furthermore, we did not observe a significant loss of repression with mutation of site #4 with KLF14 overexpression, which supports our finding of negligible binding of KLF14 to this specific Sp/KLF site (Fig. 5B). All together, KLF14 appears to bind and act on four of the GC-rich sites of the *TGFBR11* promoter to repress its transcriptional activity, and this occurs through a mechanism that includes, at least in part, competition with Sp1.

Repression of the TGFBR11 Gene by KLF14 Occurs via Mechanisms Involving Chromatin-modifying Enzymes—Careful examination of KLF14 sequence reveals a 26-amino acid

KLF14, a Non-Smad Repressor of the TGF β Receptor II




FIGURE 6. *A*, alignment displaying the 26-amino acid Sin3-interacting domain of KLF14 similar to other KLF repressors, KLF9, KLF13, and KLF16. *B*, HDAC2 and mSin3a form a complex with KLF14 *in vivo*. Panc1 cells were transfected with FLAG-tagged KLF14 or control empty vector constructs and FLAG-immunoprecipitated. Immunoprecipitated complexes were then probed with antibodies against various known corepressors. *WCE*, whole cell extract. *C*, KLF14 occupies the *TGFBR2* promoter *in vivo*. Panc1 cells were transfected with FLAG-tagged KLF14 or control empty vector constructs and subject to chromatin immunoprecipitation. A 263-bp fragment of the *TGFBR2* promoter was amplified by PCR from anti-FLAG or mock immunoprecipitated DNA samples. Note that the *TGFBR2* promoter was amplified from anti-FLAG (α -FLAG) but not mock immunoprecipitated samples (*mock*) transfected with KLF14. The *input* shows the presence of the *TGFBR2* promoter prior to immunoprecipitation. *D*, TGF β 1 leads to repressive chromatin modifications upon KLF14 occupation of the *TGFBR2* promoter. Panc1 cells were transfected with FLAG-tagged KLF14 and treated with TGF β 1 and chromatin landscape ChIP assay was performed using specified antibodies.

domain that is highly similar to repression domains of KLF9 (BTEB1), KLF13 (BTEB3), and KLF16 (BTEB4) proteins (Fig. 6A), which we have previously shown to repress transcription by recruiting Sin3a via its paired amphipathic helix 2 domain, as well as HDAC (15, 28). Interestingly, other studies have found that HDAC inhibition leads to transcriptional activation of the *TGFBR2* promoter in various cancer cell lines, raising the possibility that a repressor of this type may be operational to achieve this effect (29, 30). Thus, we have tested whether KLF14 does indeed interact with Sin3a and HDAC by immunoprecipitating full-length FLAG-tagged KLF14 from PANC1 cells. The presence of Sin3a and HDAC complexed to KLF14 is detected by Western-blot analysis (Fig. 6B). Thus, these results indicate that KLF14 interacts with Sin3a and HDAC in mammalian cells, implying that these proteins are part of a repressor complex. Subsequently, we have investigated whether this repressor complex occupies the endogenous *TGFBR2* promoter *in vivo* via ChIP assays. First, we find that KLF14 indeed binds to the *TGFBR2* promoter (Fig. 6C). Moreover, congruent with our data using promoter assays (Fig. 6C), treatment with TGF β 1 increases the amount of KLF14 bound to the *TGFBR2* promoter. In addition, the treatment with TGF β 1 coincides with the appearance of Sin3a on the same region of the promoter as KLF14, further implicating this repressor complex in the mechanism of KLF14 repression of the *TGFBR2* promoter (Fig. 6D). To gain insight into the type of chromatin remodeling that accompanies KLF14 and Sin3a occupation, we also have performed ChIP using antibodies to various histone marks. Inter-

estingly, we find that, under native conditions, acetylated histones H3 and H4 are present on the *TGFBR2* promoter; however, upon TGF β 1 treatment, these fall to negligible levels, indicating a change from a transcriptionally “active,” acetylated state to a relatively “inactive,” non-acetylated state (Fig. 6D). Moreover, we observe that, although methylated K20 H4, a mark of repression (31), is not present under native conditions, it occupies the promoter upon TGF β 1 treatment along with Sin3a (Fig. 6D). These types of experiments are in agreement with the “histone code hypothesis” (32), revealing that the state of chromatin in the *TGFBR2* promoter appears to switch from “active” to “repressed” upon KLF14 occupation after treatment with TGF β 1, all consistent with the idea that KLF14 binds to the *TGFBR2* promoter to cause repression (Fig. 6, C and D). Therefore, both competition with Sp1 and direct repression via the N-terminal domain, are likely to behave as a dual mechanism of repression that would make it more difficult to

reverse than if one of them was operational. This redundancy would ensure that the promoter remains silent even under circumstances that inactivate either of the single mechanisms (Fig. 7).

DISCUSSION

The current study provides us with several novel mechanistic insights on the regulation of the TGF β pathway. For instance, the data reported here represent the first functional characterization of the KLF14 protein, outline a novel pathway for the silencing of the *TGFBR2* promoter, provide insight into the molecular mechanisms by which these phenomena are achieved, and report additional TGF β -inducible, KLF proteins that are good candidates to regulate this promoter. These findings are of significant relevance to the areas of KLF proteins, TGF β signaling, the maintenance of cell homeostasis, and their potential contribution to disease states.

TGF β 1 ligand itself may play a role in regulating signaling by down-regulating the cell surface receptor and thus abrogating downstream messages. Others have also shown that the TGF β receptors are under autoregulation control by ligand stimulation (18). It has been previously shown that treatment with TGF β 1 leads to down-regulation of TGFBR2 levels (19). Derynck and colleagues (20) have previously described a process in osteoblastic differentiation in which at first there is a marked up-regulation of TGF β 1 and sensitivity followed by marked down-regulation of receptors and insensitivity to TGF β 1. Furthermore, recent *in vitro* studies on the biosynthe-




FIGURE 7. Model of KLF14-mediated TGF β RII silencing. Activation of the TGF β RII would occur via TGF β 1 ligand initiating an up-regulation of the TGF β RII promoter by Sp1. In our model, activation of the TGF β pathway also causes an induction of KLF14 expression (A), which would, in turn, bind to the TGF β RII promoter through its GC-rich sites (B), competing away Sp1 (C), recruiting the Sin3a/HDACs complex (D), and remodeling chromatin from a transcriptionally "active," acetylated state to an "inactive" state with silencing methyl marks (E). Thus, KLF14 mediates a negative feedback loop to down-regulate the type II TGF β receptor upon TGF β stimulation.

sis of the receptors indicate that the half-life of TGF β RII is \sim 60 min and is further reduced to 45 min in the presence of exogenous TGF β 1 (33). Therefore, understanding how this down-regulation of the TGF β RII promoter occurs is of primary biological importance for better understanding TGF β signaling.

In this regard, the current study reports several novel observations of significant biochemical relevance for extending our knowledge of TGF β signaling regulation by non-Smad proteins. For instance, this study represents the first functional characterization of KLF14 as a novel TGF β -inducible repressor protein that mediates silencing of the TGF β RII, and the mechanisms by which this phenomenon occurs, in particular the role of corepressors and chromatin modifications on the TGF β RII promoter. Previous promoter studies have primarily focused on the role of Sp1 in the activation of this promoter. However, recent studies have uncovered a paradoxical behavior for this promoter, namely that histone deacetylase inhibitors are needed to activate this promoter (34, 35). These data point to the existence of a repressive state of this promoter, which is less likely to be regulated by Sp1 but rather by Sp/KLF repressors of the type previously described by our laboratory (11). However, in the current study, rather than looking for a candidate gene within the family of KLF repressors, we have performed an unbiased screening for the TGF β inducibility of each of the 24 KLF transcription factors and find that KLF14 is up-regulated upon TGF β treatment and also can efficiently repress the TGF β RII. Interestingly, we demonstrate that there is a strong inverse correlation between expression patterns in a temporal manner of TGF β RII mRNA and KLF14 mRNA in human pancreatic cancer cells. Expression of exogenous KLF14 decreases the level of transcription from the TGF β RII promoter, implying a repressive role for KLF14 in TGF β RII expression.

Further analysis reveals the mechanism by which this protein works, indicating that KLF14 is a member of the Sin3-depend-

ent KLF repressors. This is important in light of there being two types of repressors in this family as recently reviewed by us (11), the Sin3a-dependent proteins, which include KLF9, -10, -11, -16, and now -14, and the CtBP-dependent proteins, such as KLF3 and -8. This knowledge can now be very useful for performing rapid screening of the proteins that may be acting as repressors of particular genes, besides the traditional HDAC inhibitor experiments. This screening, we propose, can utilize siRNA to either target CtBP or Sin3a. An abolition of the silencing activity would then rapidly focus the investigations on a reduced number of candidates belonging to each of these groups. In addition to the recruitment of a KLF14-mSin3A-HDAC2 repressor complex to the TGF β RII promoter, we observed a concurrent remodeling of chromatin, which involves loss of transcriptionally "active" acetylated histone marks and an increase in histone marks that associate with transcriptional silencing. Finally, we defined binding sites involved in KLF14-mediated repression of the TGF β RII promoter, namely sites -143/138 (#1), -101/-95 (#2), -59/52 (#3) and -11/-5 (#5), which are consistent with previously reported Sp1 sites, and these same sites, when mutated, lose the repressive effect of KLF14.

Overall, however, the major relevance of this study is the fact that it contributes to organize our thoughts on how TGF β signaling is regulated. It has recently been appropriately proposed to classify these mechanisms by whether they are directly mediated by Smads (the best studied to date) or those less understood events regulated by non-Smad proteins (4). In fact, our laboratory has previously described a role for KLF repressors, namely KLF10 (TIEG1) and KLF11 (TIEG2), as non-Smad effectors of the TGF β response in cell growth (9, 36). Now, due to the findings reported in the current study, we can propose a model wherein TGF β cytokines activate and repress the TGF β RII promoter using the very same GC-rich sites utilized by this family of proteins. Activation would occur in an initial phase via Sp1, whereas inhibition would follow activation and require the action of KLF14, competition of Sp1, recruitment of Sin3a, HDAC, and distinct chromatin modifications on the promoter (see model in Fig. 7). Therefore, KLF14 becomes another important non-Smad protein, in which at least one of its functions is to silence the TGF β RII. This novel transcriptional pathway thus becomes an important step for modulating the activity of TGF β signaling.

REFERENCES

- Zavadil, J., and Bottinger, E. (2005) *Oncogene* **24**, 5764–5774
- Truty, M., and Urrutia, R. (2007) *Pancreatology* **7**, 423–435
- Raftery, L., and Sutherland, D. (1999) *Dev. Biol.* **210**, 251–268
- Moustakas, A., and Heldin, C. (2005) *J. Cell Sci.* **118**, 3573–3584
- Miyazono, K., Maeda, S., and Imamura, T. (2005) *Cytokine & Growth Factor Rev. Bone Morph. Prot.* **16**, 251–263
- Massague, J., and Gomis, R. (2006) *FEBS Lett.* **580**, 2811–2820, Istanbul Special Issue
- Knight, P., and Glistler, C. (2006) *Reproduction* **132**, 191–206
- Itman, C., Mendis, S., Barakat, B., and Loveland, K. (2006) *Reproduction* **132**, 233–246
- Ellenrieder, V., Fernandez Zapico, M., and Urrutia, R. (2001) *Curr. Opin. Gastroenterol.* **17**, 434–440
- Bachman, K., and Park, B. (2005) *Curr. Opin. Oncol.* **17**, 49–54
- Lomber, G., and Urrutia, R. (2005) *Biochem. J.* **392**, 1–11

KLF14, a Non-Smad Repressor of the TGF β Receptor II

12. Black, A. R., Black, J. D., and Azizkhan-Clifford, J. (2001) *J. Cell Physiol.* **188**, 143–160
13. Turner, J., and Crossley, M. (1999) *Trends Biochem. Sci.* **24**, 236–240
14. Bae, H. W., Geiser, A. G., Kim, D. H., Chung, M. T., Burmester, J. K., Sporn, M. B., Roberts, A. B., and Kim, S. J. (1995) *J. Biol. Chem.* **270**, 29460–29468
15. Zhang, J. S., Moncrieffe, M. C., Kaczynski, J., Ellenrieder, V., Prendergast, F. G., and Urrutia, R. (2001) *Mol. Cell Biol.* **21**, 5041–5049
16. Cook, T., Gebelein, B., Mesa, K., Mladek, A., and Urrutia, R. (1998) *J. Biol. Chem.* **273**, 25929–25936
17. Venkatasubbarao, K., Ammanamanchi, S., Brattain, M. G., Mimari, D., and Freeman, J. W. (2001) *Cancer Res.* **61**, 6239–6247
18. Woodward, T. L., Dumont, N., O'Connor-McCourt, M., Turner, J. D., and Philip, A. (1995) *J. Cell Physiol.* **165**, 339–348
19. Nishikawa, Y., Wang, M., and Carr, B. I. (1998) *J. Cell Physiol.* **176**, 612–623
20. Gazit, D., Ebner, R., Kahn, A. J., and Derynck, R. (1993) *Mol. Endocrinol.* **7**, 189–198
21. Jennings, R., Alsarraj, M., Wright, K. L., and Munoz-Antonia, T. (2001) *Oncogene* **20**, 6899–6909
22. Schneider, D., Kleeff, J., Berberat, P. O., Zhu, Z., Korc, M., Friess, H., and Buchler, M. W. (2002) *Biochim. Biophys. Acta* **1588**, 1–6
23. Subramaniam, M., Harris, S. A., Oursler, M. J., Rasmussen, K., Riggs, B. L., and Spelsberg, T. C. (1995) *Nucleic Acids Res.* **23**, 4907–4912
24. Parker-Katirae, L., Carson, A. R., Yamada, T., Arnaud, P., Feil, R., Abu-Amero, S. N., Moore, G. E., Kaneda, M., Perry, G. H., Stone, A. C., Lee, C., Meguro-Horike, M., Sasaki, H., Kobayashi, K., Nakabayashi, K., and Scherer, S. W. (2007) *PLoS Genet.* **3**, 665–678
25. Koutsodontis, G., Moustakas, A., and Kardassis, D. (2002) *Biochemistry* **41**, 12771–12784
26. Kaczynski, J., Conley, A. A., Fernandez-Zapico, M. E., Delgado, S. M., Zhang, J. S., and Urrutia, R. (2002) *Biochem. J.* **366**, 873–882
27. Natesampillai, S., Fernandez-Zapico, M. E., Urrutia, R., and Veldhuis, J. D. (2006) *J. Biol. Chem.* **281**, 3040–3047
28. Kaczynski, J., Zhang, J. S., Ellenrieder, V., Conley, A., Duenes, T., Kester, H., van Der Burg, B., and Urrutia, R. (2001) *J. Biol. Chem.* **276**, 36749–36756
29. Huang, W., Zhao, S., Ammanamanchi, S., Brattain, M., Venkatasubbarao, K., and Freeman, J. W. (2005) *J. Biol. Chem.* **280**, 10047–10054
30. Lee, B. I., Park, S. H., Kim, J. W., Sausville, E. A., Kim, H. T., Nakanishi, O., Trepel, J. B., and Kim, S. J. (2001) *Cancer Res.* **61**, 931–934
31. Nishioka, K., Rice, J. C., Sarma, K., Erdjument-Bromage, H., Werner, J., Wang, Y., Chuikov, S., Valenzuela, P., Tempst, P., Steward, R., Lis, J. T., Allis, C. D., and Reinberg, D. (2002) *Mol. Cell* **9**, 1201–1213
32. Jenuwein, T., and Allis, C. D. (2001) *Science* **293**, 1074–1080
33. Koli, K. M., and Arteaga, C. L. (1997) *J. Biol. Chem.* **272**, 6423–6427
34. Park, S. H., Lee, S. R., Kim, B. C., Cho, E. A., Patel, S. P., Kang, H. B., Sausville, E. A., Nakanishi, O., Trepel, J. B., Lee, B. I., and Kim, S. J. (2002) *J. Biol. Chem.* **277**, 5168–5174
35. Zhao, S., Venkatasubbarao, K., Li, S., and Freeman, J. W. (2003) *Cancer Res.* **63**, 2624–2630
36. Cook, T., and Urrutia, R. (2000) *Am. J. Physiol.* **278**, G513–G521