



Published in final edited form as:

Anesth Analg. 2008 September ; 107(3): 832–848. doi:10.1213/ane.0b013e318182aedb.

Is a New Paradigm Needed to Explain How Inhaled Anesthetics Produce Immobility?

Edmond I Eger II, MD,

Department of Anesthesia and Perioperative Care, University of California, San Francisco, CA 94143-0464

Douglas E. Raines, MD,

Department of Anesthesia and Critical Care, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts

Steven L. Shafer, MD,

Department of Anesthesia, Columbia University, New York, NY 10032

Hugh C. Hemmings Jr., MD, PhD, and

Departments of Anesthesiology and Pharmacology, Weill Cornell Medical College, New York, NY 10065

James M. Sonner, MD

Department of Anesthesia and Perioperative Care, University of California, San Francisco, CA 94143-0464

Abstract

A paradox arises from present information concerning the mechanism(s) by which inhaled anesthetics produce immobility in the face of noxious stimulation. Several findings, such as additivity, suggest a common site at which inhaled anesthetics act to produce immobility. However, two decades of focused investigation have not identified a ligand- or voltage-gated channel that alone is sufficient to mediate immobility. Indeed, most putative targets provide minimal or no mediation. For example, opioid, 5-HT₃, gamma-aminobutyric acid type A and glutamate receptors, and potassium and calcium channels appear to be irrelevant or play only minor roles. Furthermore, no combination of actions on ligand- or voltage-gated channels seems sufficient. A few plausible targets (e.g., sodium channels) merit further study, but there remains the possibility that immobilization results from a nonspecific mechanism.

Introduction

Immobility of the surgical patient is a cardinal feature of general anesthesia. All inhaled anesthetics (including all clinically useful anesthetics) share the capacity to produce immobility in the face of noxious stimulation. Immobility forms part of the standard unit of anesthetic potency, MAC, the minimum alveolar concentration of inhaled anesthetic that abolishes movement in response to noxious stimulation in 50% of subjects.^{1–3} The mechanism of inhaled anesthetic action provides one of the oldest problems in pharmacology, and the most challenging. The present essay reviews the evidence for possible mechanisms by which inhaled anesthetics produce immobility.

Address correspondence to Dr. Edmond I Eger II: Department of Anesthesia, S-455, University of California, San Francisco, California 94143-0464. Telephone: (415) 476-6927; Fax (415) 502-5375. egere@anesthesia.ucsf.edu.
Dr. Eger is a paid consultant to Baxter Healthcare Corp.

A few observations might serve the reader before embarking on that review. As will be emphasized below, the spinal cord is the primary site affected by inhaled anesthetics to produce immobility.^{4,5} However, immobility does not result from a lack of processed input into the cord, nor is it necessarily consequent to the ability of the cord to respond to impulses from the brain that would provoke movement. Although the amnesia produced by inhaled anesthetics precludes extracting from surgical patients whether they felt pain during surgery, a surrogate measure, autonomic responses to noxious stimulation, indicates that immobility is not a consequence of a limitation on sensory input. At 1.0 MAC, incision produces an increase in arterial blood pressure and heart rate,⁶ and it increases ventilation, even at twice MAC.⁷ MAC-BAR for sevoflurane exceeds MAC by a factor of 2.2.⁸ Sensory-evoked potentials in humans can be recorded at concentrations well in excess of MAC.⁹ Transmission of impulses through the dorsal horn of rats continues during halothane, isoflurane and propofol anesthesia,^{10,11} and propofol anesthesia does not prevent spinal cord c-fos expression in mice subjected to an intraplantar injection of formalin.¹² Similarly, immobility is not a consequence of paralysis, nor does it necessarily result from the inability of motor nerves in the anterior horn to respond to impulses from the brain. At least some studies find that motor-evoked potentials continue at 1.0 MAC.¹³

If inhaled anesthetics at MAC allow both transmission of sensory input and cerebral control over motor movement, how do they suppress movement in response to noxious stimulation? How do they accomplish that at the level of the spinal cord? The answer is not known for certain, but one possibility is that they depress central pattern generators in the cord, local command-posts that coordinate movement.¹⁴ Consistent with this notion, studies with nitrous oxide suggest that its action lies in ventral portions of the cord.¹⁵

The essay first considers evidence suggesting that inhaled anesthetics act at a common site to produce immobility. The evidence includes the absence of synergy in inhaled anesthetic interactions, similarities in steric and electrostatic properties of inhaled anesthetics, the correlation of MAC with affinity to the membrane bilayer interface or its surrogate, and the evolutionary conservation of the site at which anesthetics act.

The essay continues with an examination of the potential contributions of specific ligand-gated channels, concluding that one or two such channels (e.g., glycine) might play a role, but that present evidence suggests that no one channel can explain more than a portion of anesthetic-induced immobility. Voltage-gated potassium channels seem unable to explain the production of immobility, but the voltage-gated sodium channels remain a plausible candidate. How inhaled anesthetics act to block this and other sites remains a mystery, but some new concepts are proposed.

Evidence for a Common Site of Inhaled Anesthetic Action

Additivity

Observations made below indicate that inhaled anesthetic actions on potassium channels or a single ligand-gated channel can explain only a minor part of the capacity of inhaled anesthetics to produce immobility. Even summing the effects of inhaled anesthetics on several channels appears to be insufficient to explain immobility. Could synergistic inhaled anesthetic effects on ligand- and voltage-gated channels magnify their actions sufficiently to produce immobility?

We have found¹⁶ that inhaled anesthetic pairs that act on different channels (i.e., at MAC, anesthetic A acts potently on receptor X but weakly on receptor Y, whereas anesthetic B acts weakly on X but potently on Y) combine in an additive, never synergistic, manner to produce immobility (Fig. 1). As the recent review by Hendrickx et al. shows,¹⁷ this is unusual. Most

drugs that act on separate channels (i.e., as anesthetics A and B) show synergy when combined. A recent theoretical analysis of additivity and synergy demonstrated that there are only two mechanisms by which additivity can be observed: two drugs competing at the same site of action, or two drugs acting at different sites of action at concentrations causing very low levels of receptor occupancy.¹⁸ The implication is that, if there is more than one biological target for inhaled anesthetic action, then binding of an inhaled anesthetic to those sites of action must be very weak. This effectively excludes high affinity targets as potential sites of anesthetic action.

Since clinical concentrations are typically in the range of 0.3 mM (especially see Table 1 of the following cited article),¹⁹ it follows that if anesthetics act at more than one site of action, then Kd must exceed 0.3 mM, possibly many-fold. This is weak binding compared to IV anesthetics, which are typically effective at μM or nM concentrations,²⁰ and have Kd values also in the nM range.^{21,22} Such observations effectively exclude the relatively high affinity targets of IV anesthetic action as targets of inhaled anesthetic action.

Although additivity argues for a single site of action, it is not definitive. Although synergy more commonly results from concurrent actions of drugs on different sites, such pairs can produce additivity.¹⁷ A report concerning the anesthetic actions of propofol plus sevoflurane provides an example.²³ Thus, additivity constitutes but part of a broader argument for a single site.

Common Steric and Electrostatic Properties Define Anesthetic Action

Bertaccini et al. found common chemical motifs within various anesthetic binding sites.²⁴ Similarly, Sewell and Sear asked if volatile halogenated anesthetics have electrostatic (charge) and steric (shape and size) properties that define their potencies as anesthetics, specifically their capacities to produce immobility.²⁵ They applied comparative molecular field analysis (CoMFA) to 69 structurally diverse halogenated anesthetics, randomly divided into a training-set (N=52) used to derive their model and a test-set (N=17) used to independently assess the model's predictive power. The method maximized similarity in molecular shape and electrostatic potential. The predicted and observed activities of the training set had a correlation coefficient squared of 94% (i.e., the model explained 94% of the variance in the observed activities) and 70%–84% of the test-set. Similar correlations were found for non-halogenated volatile anesthetics,²⁶ with considerable overlap, particularly for certain steric characteristics. The demonstration that CoMFA can predict anesthetic potency from anesthetic structures suggests that potency and binding affinity (which depends on anesthetic structure) are tightly linked as it is unlikely that one could develop a single 3-D pharmacophoric map for MAC with high predictive ability if there was little relationship between anesthetic binding affinity and potency. The CoMFA results might also be taken to imply a single rather than multiple sites at which inhaled anesthetics might act. However, Sewell and Sear caution that "It should be noted that a common molecular basis for immobilizing activity does not necessarily imply a common site of action."

Implications of Stereoselectivity

The (+) isomer of isoflurane is 53% more potent than the (–) isomer in rats, "consistent with a receptor-mediated anesthetic mechanism by volatile anesthetics."²⁷ Small stereoselective effects are found with 2-butanol and 2-pentanol (but not 2-hexanol or 2-heptanol.)²⁸ Although stereoselective effects can be found *in vitro* (in frog oocytes) for the actions of these alcohols on TRESK potassium channels, gamma-aminobutyric acid type A (GABA_A) receptors, and/or N-methyl-D-aspartate (NMDA) receptors, the effect does not correlate with the *in vivo* differences in MAC.²⁹ Stereospecificity is an important determinant of potency for anesthetics such as ketamine.³⁰ It implies a binding site, and that inhaled anesthetics conform to a specific pharmacophore. If stereoselectivity is important to inhaled anesthetics, does that indicate a

specific (read single) site of action? How would xenon, a nicely rounded simple atom, dock in a protein in the same highly specific manner that more complex ligands would dock? Does xenon bind, and cause profound effects, at the same place isoflurane binds? Perhaps so,³¹ and yet a single portion of a protein acting stereoselectively would seem an unlikely site for a common place at which xenon and isoflurane would act; perhaps a lipid bilayer (which has chiral centers) would form a more attractive site.

An Updated Meyer-Overton Relationship

More than one-hundred years ago, Meyer³² and Overton³³ demonstrated a correlation between anesthetic affinity for lipid and anesthetic potency. This correlation guided studies of anesthetic mechanisms for 80 years, focusing work on the lipid bilayer. Several investigators, notably Franks and Lieb,³⁴ shifted that focus to proteins. Part of the shift resulted from the failure of a bilayer focus to produce a verifiable theory. Part resulted from evidence against the correlation. For example, non-immobilizers are inhaled compounds that do not produce immobility despite possessing a lipophilicity that would indicate anesthetic capability.³⁵ However, if another factor, polarity, is added to the calculus of factors determining potency, a modified Meyer-Overton relationship remains defensible.^{36,37} That is, the anesthetic interaction with proteins implies amphipathicity.²⁴ The correlation can be much improved by selecting a solvent that has an element of polarity. Abraham et al. suggested methanol.³⁸ The correlation also can be much improved by selecting a lipid-like phase that more closely resembles the membrane bilayer [e.g., one that includes phospholipoproteins].^{39,40} If such a modified Meyer-Overton relationship is correct, it implies similar sites of action for multiple receptor/channels. It would be remarkable if specific pockets in various channel proteins (both inhibitory and excitatory) share relatively common characteristics.

Go a step further and demand a correlation with a slope of 1.0. This results in MAC times some index of solubility (e.g., solubility in methanol or some other solvent) equals a constant. A key point is not that anesthetic potency has something to do with methanol, but that exactly the same number of anesthetic molecules at the site of action are required to produce MAC and that site of action resembles methanol. How could that be across 5–7 orders of MAC values, unless some fundamental, highly conserved, process was at work?

Evolution and Conservation of the Anesthetic Site of Action

MAC or its equivalent varies little (perhaps two- or three-fold) among different vertebrate classes, again suggesting conservation of the site at which anesthetics act. Such a site has no apparent survival benefit since anesthetics do not form part of the natural environment. Thus, the constancy implies that the susceptibility to anesthetics develops from a parallel process, some essential aspect of ion channel function that fortuitously leads to this sensitivity and confers a survival benefit. It is difficult to see how this might result from an invariant protein structure of inhibitory and excitatory receptors. Is it reasonable to believe that there is a single such structure in all plausible channels? These issues are discussed in a recent symposium in this issue of *Anesthesia & Analgesia*.^{41–46}

In that symposium, Sonner hypothesized that one-celled organisms selected a beneficial trait that also fortuitously produced the capacity of inhaled anesthetics to increase in currents through inhibitory channels, and decrease currents through excitatory channels.⁴¹ The beneficial trait arose as a response to compounds present in the environment that influenced the conformational equilibrium of ion channels and otherwise would have promoted the entry of positive charges that might damage the cell. That is, the trait increased the fitness (survival) of the organism by limiting the effect of compounds present in the environment that might otherwise reduce electrochemical potentials across the cell membrane through their effects on channel function. Consistent with this view, exposure to inhaled anesthetics changes the

membrane composition of one-celled organisms.^{47–50} The finding that surfactants modulate anesthetic-sensitive channels in a manner similar to inhaled anesthetics⁵¹ is consistent with the notion that the response to anesthetics arose as an adaptation to environmental conditions which influenced channel function by perturbing bilayer properties.

This evolutionary narrative correctly predicted that certain nonvolatile compounds have anesthetic-like modulatory effects on ion channels and in animals. These may include endogenous compounds increased in disease [e.g., ammonia⁵² and ketoacids.]⁵³ Such compounds modulate ion channel function in a manner similar to inhaled anesthetics.⁵¹ Cantor proposed that the slow adsorption and desorption of high (higher than those arising at synapses?) concentrations of neurotransmitter onto and off the membrane may produce a parallel, membrane-mediated effect manifested as receptor desensitization.⁵⁴ This process may provide a selective pressure for receptors to respond to membrane-mediated effects of inhaled anesthetics in multicellular organisms. The anesthetic-like modulatory effect of non-native neurotransmitters on receptors (e.g., acetylcholine on an NMDA receptor) supports this hypothesis.⁵⁵

Tests of Relevance

Inhaled anesthetics affect many ligand- and voltage-gated ion channels in ways that plausibly explain immobility (e.g., enhancement of inhibitory channels; blockade of excitatory channels). How can we test whether plausible translates to relevant? Exclusion or inclusion of a target as a likely mediator of anesthesia may require the concurrence of results from several tests of relevance.

Location of effect

An explanation for MAC must consider the importance of the spinal cord in this process. Two seminal studies, differing in experimental design, demonstrated that the spinal cord, not the brain, is the primary site at which inhaled anesthetics produce immobility.^{4,56,57} One study finds that direct application of sevoflurane to the spinal cord can produce reversible immobility in response to noxious stimulation of the hindlimbs,⁵⁸ and a second study finds that epidural administration of an emulsion of 8% isoflurane can produce a reversible epidural anesthetic in rabbits but does not appear to affect the level of consciousness.⁵⁹ A limitation of these investigations is that one cannot tell if the cord anesthetic concentrations far exceed those needed during conventional anesthesia.

Non-Immobilizers

Some inhaled compounds do not produce immobility in the face of noxious stimulation, nor do they decrease the requirement for anesthesia by conventional inhaled anesthetics.³⁵ These "non-immobilizers" do not produce or contribute to anesthesia despite having a lipophilicity that predicts that they should.^{32,33} Non-immobilizers should not influence a relevant site of anesthesia.

Correlation of Physiological Changes on the Channel and on MAC

Physiological variables that affect a channel should produce an effect that is consistent with the effect of those variables on MAC. For example, an increase in temperature can open an inhibitory potassium channel.⁶⁰ An increase in inhibitory current should decrease MAC but, in fact, MAC increases with temperature,⁶¹ diminishing the likelihood that this channel is a relevant mediator.

Agonists and Antagonists as Pharmacological Probes

Drugs that block or enhance putative targets of inhaled anesthetic action can be used to test the relevance of those targets. An advantage of this approach over some genetic approaches is that the effects leave less time for compensation. However, the interpretation of pharmacological interactions can be complex.

Suppose administration of a given receptor antagonist does not affect MAC. Blockade of that receptor by the inhaled anesthetic cannot be the sole cause of anesthesia or the antagonist would have produced anesthesia. The initial interpretation of an absence of blocker effect might be that the receptor is not relevant. However, what if MAC concentrations potently block the receptor in question, but concurrent blockade of a second receptor is required for immobility? Blockade of both receptors would be needed to produce anesthesia. Thus, administration of a blocker of one might not decrease the need for the inhaled anesthetic to block the second, leaving the concentration required for anesthesia unchanged.

To find that the receptor was irrelevant might require demonstration that blockade had no effect on an anesthetic whose action was known to be caused solely by alteration of a different receptor. Inhaled anesthetics block specific neuronal acetylcholine receptors at concentrations much less than MAC.⁶² Neuronal nicotinic acetylcholine receptors are probably irrelevant to immobilization because administration of mecamylamine (an antagonist of neuronal nicotinic acetylcholine receptors) does not affect MAC for several inhaled anesthetics, and because mecamylamine does not influence the anesthetic requirement for etomidate (which produces immobility solely by enhancing the effect of GABA on GABA_A receptors).⁶³

Parallel arguments may apply to inhibitory receptors. Consider GABA, the major inhibitory transmitter in the nervous system, as an example. Suppose blockade of GABA_A receptors (e.g., with picrotoxin) increases the MAC of test anesthetics. One might argue that MAC increased because anesthetic-induced enhancement of the GABA_A receptor response to GABA had been abolished. However, relevance demands that the increases in MAC consequent to blockade be proportional to the separately measured capacity of the anesthetic to enhance the receptor response *in vitro*. In other words, suppose that one MAC of test anesthetic X minimally enhances GABA_A receptors *in vitro*, but one MAC of anesthetic Y has a major effect. This would mean that the anesthetic effects of X do not result from enhancement of GABA_A receptors, but the anesthetic effects of Y might, in part, be explained by their action on GABA_A receptors. If that were true, then blockade of GABA_A receptors might minimally increase the MAC of X, but substantially increase the MAC of Y. That is what is found for blockade of glycine receptors with strychnine but not for blockade of GABA_A receptors with picrotoxin (see below). Thus GABA_A receptors are not relevant as mediators of immobility, but glycine receptors could be.

One might also suppose blockade of an excitatory receptor decreases MAC. In itself, that proves nothing since many factors that do not mediate anesthesia (e.g., administration of opioids) can modulate anesthesia. As with blockade of inhibitory receptors, relevance demands that the effect on MAC be inversely proportional to the capacity of the test anesthetics to block the excitatory receptor *in vitro*. For example, if the anesthetic completely blocks the receptor at one MAC, then administration of a blocker cannot increase blockade and cannot affect MAC. In contrast, administration of a blocker can decrease MAC of an anesthetic that minimally affects the receptor because the blockade can add to the decrease in excitatory neurotransmission. By this test, we can exclude NMDA receptors as relevant mediators of the effects of conventional inhaled anesthetics (see below).

Genetic Tests

Global knockouts—If global knockout (genetic inactivation) of a receptor markedly changes MAC, we might suspect that the receptor mediates anesthesia. While global knockout studies have advanced our understanding of anesthetic mechanisms,⁶⁴ two problems can confound results obtained using this approach. First, gene inactivation can alter expression of other genes and thereby compensate for the loss of the targeted gene. Second, global knockout affects all neurons and thus can influence MAC by an effect on cerebral rather than spinal cord neurons, an effect we believe is irrelevant since the brain, at most, provides only a minor contribution to the immobility produced by inhaled anesthetics.

Knockins—An alternative approach relies on the production of receptor knockins, wherein an "engineered" receptor that responds normally to its ligand (e.g., GABA) but is not influenced by anesthetics replaces the natural ("wild-type") receptor. If a receptor knockin animal no longer responds to anesthetics, that receptor likely mediates anesthesia. A major advantage of this strategy is that it should not provoke compensatory effects because it permits a normal response to the natural ligand. The knockin mouse has powerfully contributed to our appreciation of the GABA_A receptor to the action of anesthetics such as etomidate.⁶⁴ Of note, this knockin does not materially alter isoflurane MAC,⁶⁵ but the knockin approach can have problems. The replacement can have a different distribution from the wild-type receptor and this may provoke compensation, or the engineered receptor might not have completely normal function. Finally, both the knockin and knockout approaches usually alter only one subtype of a receptor (although multiple mutations are possible). A given family (e.g., GABA_A receptors) can include tens or hundreds of receptors and perhaps the critical subtype was not chosen!

Variations on a Theme—Some genetic approaches minimize the potential for compensation. For example, the receptor can be targeted for expression of a receptor deletion (induction of deletion) at a specific time in life or in a specific region of the body/brain.

Cautions

Absence of movement in response to noxious stimuli underlies MAC, which has obvious clinical importance. Other end-points, such as loss of righting reflex, do not substitute for immobility. Loss of righting reflex results from cerebral as well as spinal cord actions. There is no necessary correlation between the concentrations that suppress immobility versus righting.^{66,67} Similar restrictions limit the usefulness of animals that have the attraction of allowing considerable genetic manipulation (e.g., fruit flies and worms).

Relevance of Plausible Specific Targets (Alphabetical Order) to MAC

A present consensus holds that multiple ligand-gated and/or voltage-gated ion channels and possibly other targets mediate the immobility produced by inhaled anesthetics. These include receptors for GABA, glycine, acetylcholine (neuronal nicotinic), various glutamate (e.g., NMDA and AMPA-kainate), opioid, adrenergic, and serotonin receptors, nitric oxide synthase (NOS) (and nitric oxide), sodium and potassium channels, and gap junctions. The present essay will not assemble the considerable evidence for the plausibility of these targets as mediators of immobility; we⁶⁸ and others^{20,69–71} have done that previously, with suggestions of specific receptor locations needed to produce anesthesia.^{72,73} Instead we will discuss evidence that suggests that no single target can explain immobility produced by inhaled anesthetics.

Acetylcholine Receptors

In rats, blockade of nicotinic (mecamylamine) or muscarinic (atropine or scopolamine) receptors does not modify anesthetic potency, either *in vivo*^{74,75} or *in vitro*.⁷⁶ Co-administration of large doses of mecamylamine and atropine does not affect the cerebral

concentration of etomidate required to produce immobility.⁶³ Intrathecal administration of atropine does not alter MAC of isoflurane.⁷⁵ Similarly, in mice, administration of nicotine [at concentrations that produce behavioral effects]⁷⁷ does not alter isoflurane MAC.⁷⁴ 1,1,2-trichlorocyclobutane (an anesthetic) and 1,2-dichlorohexafluorocyclobutane (a non-immobilizer) both inhibit neuronal nicotinic receptors.^{78–80} These observations suggest that acetylcholine receptors do not mediate inhaled anesthetic-induced immobility.

Acid-Sensitive Ion Channels (ASICs)

ASICs are proton-gated members of the family of degenerin channels⁸¹ and are expressed in the spinal cord,⁸² particularly in the dorsal horn.⁸³ They are important to the modulation of nociception.⁸⁴ Acidosis opens such channels, increasing excitatory currents but also increasing the rate of inactivation.⁸³ The latter may decrease repetitive firing. If these channels mediate the capacity of inhaled anesthetics to produce immobility, one might expect that the acidosis produced by inhalation of carbon dioxide might increase MAC by opening such channels. However, a rectilinear decrease in MAC is found.⁸⁵

Adenosine Receptors

Adenosine receptors have been proposed as mediators of the capacity of inhaled anesthetics to produce immobility. Administration of adenosine decreases MAC of halothane in dogs,⁸⁶ and intrathecal administration of the adenosine agonist R-phenylisopropyl-adenosine (R-PIA) decreases halothane MAC in rats.⁸⁷ However, aminophylline, an adenosine receptor blocker, does not increase halothane MAC in dogs,⁸⁸ rats,⁸⁸ or humans.⁸⁹ Similarly, administration of the A₁ adenosine receptor antagonist 8-cyclopentyl-1,3-dipropylxanthine (DPCPX) does not alter halothane MAC in rats, but DPCPX administration does prevent the decrease in MAC that otherwise is produced by the adenosine receptor agonist R-PIA.⁸⁷ Thus, adenosine receptors do not appear to mediate the immobility produced by inhaled anesthetics.

Adenosine Triphosphate Receptors

Administration of 100 $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ adenosine triphosphate to patients does not alter MAC awake or MAC of sevoflurane in humans.⁹⁰

α -2 Adrenoreceptors

Results from studies of α -2 adrenoreceptors expressed in *Xenopus* oocytes suggest that halothane does not act via such receptors.⁹¹ N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline (a catecholamine depleter) does not change halothane MAC in rats,⁹² and the α -2 adrenoreceptor blocker tolazoline does not affect halothane MAC in dogs.⁹³ The blocking drugs yohimbine and atipamezole at 0.8 mg/kg–1.0 mg/kg increase the MAC of isoflurane in rats by approximately 10%, but larger doses decrease MAC, particularly as lethal doses are reached.⁹⁴ The 10% increase in MAC probably results from suppression of the effect of normal tonic stimulation of α -2 adrenoreceptors. Since blockade and depletion of α -2 adrenoreceptors minimally affects MAC, these receptors are unlikely mediators of the capacity of inhaled anesthetics to produce immobility.

Calcium Channels

Although some studies support a role for Ca⁺⁺ channels,⁹⁵ several do not. Enflurane, halothane, and isoflurane concentrations exceeding MAC by an order of magnitude minimally affect Ca⁺⁺ channels *in vitro*.^{96,97} Halothane and sevoflurane exert opposite effects on Ca⁺⁺ release from sarcoplasmic reticulum,⁹⁸ but are nearly exactly additive in their capacities to produce immobility.¹⁶ Knockout of the α_{1G} T-type calcium channel does not alter MAC in mice.⁹⁹ Ca⁺⁺ channels seem unlikely mediators of immobility.

Catecholamine Receptors

Dopamine receptors do not materially mediate the immobility produced by inhaled anesthetics in rats.¹⁰⁰ However, depletion of catecholamine neurotransmitters (including norepinephrine, and epinephrine) with reserpine or other drugs decreases MAC for halothane and/or cyclopropane in dogs^{101,102} and rats.^{103,104} The decrease varies, one study finding a 20% decrease,^{102,103} a second a maximum of 30%,¹⁰¹ and others of 40%,^{102–104} but one study found no change in cyclopropane MAC in rats given a single dose of 10 mg/kg reserpine,¹⁰⁵ a dose that decreased whole brain norepinephrine to 3% of control values. Such results suggest that neurotransmission via catecholamine receptors might explain a minor portion of the immobility produced by inhaled anesthetics, but the decrease could be an epiphenomenon related to the nearly complete depletion of an excitatory neurotransmitter(s).

GABA_A Receptors

Propofol, alfaxalone and etomidate produce immobility by enhancing GABA_A receptor function (i.e., by increasing the effect of a given concentration of GABA).¹⁰⁶ A point mutation that minimizes the effect of propofol or etomidate on the GABA_A receptor also minimizes the capacity of these compounds to produce anesthesia.⁶⁴ Since inhaled anesthetics also enhance the action of GABA on GABA_A receptors,¹⁰⁷ a parallel effect would seem plausible. However, present evidence argues against the importance of this receptor to the immobility produced by inhaled anesthetics.

Not all drugs that augment the *in vivo* action of GABA produce immobility. Gabaculine, a GABA-transaminase inhibitor, produces a dose-related loss of righting reflex in mice but does not alter the MAC of halothane, enflurane, isoflurane, or sevoflurane.⁶⁶ The absence of a correlation of loss of righting reflex and MAC is consistent with the finding that the concentration of an inhaled anesthetic required to impair righting does not bear a constant relationship with the MAC for that anesthetic.⁶⁷ Others have noted a differential effect of sevoflurane on GABA_A receptors in the cortex versus the spinal cord, minimally affecting the latter.¹⁰⁸

In rats, the noncompetitive GABA_A receptor antagonist picrotoxin differs in its effects on the immobilization produced by propofol versus inhaled anesthetics. The direct antagonism of GABA_A receptors by picrotoxin progressively increases the immobilizing ED₅₀ of propofol up to at least 400%, showing no ceiling for this effect.¹⁰⁹ In contrast, picrotoxin (or gabazine, a competitive antagonist at GABA_A receptors) administration equally increases isoflurane MAC and the immobilizing ED₅₀ of the nonGABAergic anesthetic ketamine to a ceiling of approximately 60% (reflecting an indirect antagonism of the effect of naturally occurring tonic GABA release).¹⁰⁹

Cyclopropane and xenon minimally enhance the response of GABA_A receptors to GABA *in vitro*, whereas isoflurane causes substantial enhancement.^{20,110,111} If GABA_A receptors mediate the capacity of inhaled anesthetics to suppress movement in response to noxious stimuli, then blockade of GABA_A receptors should increase the MAC of isoflurane more than the MAC of xenon or cyclopropane. However, intrathecal administration of picrotoxin increases MAC equally for all three anesthetics (Fig. 2),^{68,112} indicating that GABA_A receptors do not mediate immobility produced by inhaled anesthetics.

Evidence from genetically engineered mice adds to the notion that enhancement of the effect of GABA does not underlie MAC. Global knockout of the α -1 GABA_A receptor does not change isoflurane MAC.¹¹³ Knockout of the β 3 subunit of the GABA_A receptor increases enflurane MAC by 26% but only increases halothane MAC by 9%,¹¹⁴ and even these small increases could be attributable to compensation for lack of the subunit. Such minor increases

agree with those found by Jurd et al. for $\beta 3$ N265M knockin mice unresponsive to the anesthetic effects of propofol but “normally” sensitive to GABA.⁶⁴ enflurane MAC was 15% greater than in wild-type mice, and halothane MAC was 21% greater. More importantly, this mutation equally increased the MAC of isoflurane and cyclopropane despite the enormous differences in enhancement of GABA_A receptors by these anesthetics.⁶⁵

Similarly, MAC for fluorinated alkanols¹¹⁵ does not correlate with their capacity to enhance the response of GABA_A receptors.¹¹⁶ Also, enflurane and halothane enhance GABA-mediated chloride conductance in rat hippocampal neurons,¹¹⁷ but a given MAC-multiple of enflurane has twice the effect of halothane (i.e., the result is not quantitatively consistent across anesthetics.)

Thus results from studies of blocking drugs and of knockin and knockout animals do not support a role for GABA_A receptors as mediators of the immobility produced by inhaled anesthetics.

Gap Junctions

Gap junctions are protein channels that form electrical synapses by directly connecting the cytosol of neighboring cells. Although these may be plausible targets of drugs, such as thiopental and propofol, 10 MAC, but not 2 MAC, halothane can block gap junction coupling in hippocampal slices.¹¹⁸ Others similarly report a low sensitivity of gap junctions to the effects of inhaled anesthetics.¹¹⁹

Glycine Receptors

Many IV anesthetics enhance glycine receptor function.^{107,120,121} Their spinal localization and their enhancement by volatile anesthetics suggest a role as a mediator of MAC. Evidence in rats supports this notion. Intravenous and intrathecal administration of strychnine, a glycine receptor antagonist, increases MAC.^{122,123} In contrast to the similar effect of intrathecal picrotoxin on the MAC of cyclopropane, xenon and isoflurane (see above), the increase in the MAC of cyclopropane, isoflurane, and halothane produced by intrathecal strychnine administration correlates with the enhancing effect of these anesthetics on glycine receptors *in vitro* (Fig. 2).^{68,123}

Results from studies of mice with genetic alterations provide conflicting evidence for the importance of glycine receptors to MAC. Spastic mice have decreased glycine receptor expression¹²⁴ and show a 30% increase in enflurane MAC but no increase in halothane MAC.¹²⁵ A missense mutation in the glycine receptor $\alpha 1$ subunit decreases the sensitivity of spasmodic mice to glycine,¹²⁶ but these mice show no difference in enflurane or halothane MAC from control mice.¹²⁵

Overall, current data indicate that glycine receptors might mediate part of the immobility produced by some inhaled anesthetics (e.g., halothane and isoflurane) but not by other anesthetics (e.g., cyclopropane).

Glutamate Receptors

AMPA Receptors—Blockade of AMPA receptors can decrease MAC by approximately 60%,¹²⁷ a maximum decrease similar to that found with NMDA receptor blockade.^{128,129} AMPA receptor blockade can augment the capacity of blockade of NMDA receptors to decrease MAC.¹³⁰ Enflurane inhibits the postsynaptic action of glutamate on AMPA receptors in mouse spinal cord,¹³¹ and halothane similarly affects the hippocampus, but at higher MAC values.¹³² Also, clinically-relevant concentrations of volatile anesthetics partially inhibit native and recombinant AMPA receptors activated by exogenous agonists.^{131,133–136}

However, results from studies of genetically modified mice do not support the notion that AMPA receptors mediate the immobility produced by inhaled anesthetics. GluR1-GluR4 subunit combinations form AMPA receptors. Mice lacking the GluR2 subunit have MAC values for halothane, isoflurane and sevoflurane similar to wild type littermates, despite altered concentrations for loss of righting and antinociception.¹³⁵ Clinical concentrations of isoflurane and halothane minimally inhibit AMPA receptors *in vitro* (both GluR2-containing and deficient).¹³⁵ Motor neurons in the ventral spinal cord (i.e., that could mediate MAC) lack GluR2 subunits.^{137–139} Thus, knockout of the GluR2-containing receptors should not change MAC but might change righting reflex and antinociception.

Although the above data suggest that the GluR2 subunit does not affect MAC, this cannot exclude the possibility that other subunits of AMPA receptors contribute to MAC. This concern appears to be addressed by the failure of the intrathecal injection of a blocker of AMPA receptors to change the MAC of isoflurane.¹⁴⁰

Kainate Receptors—GluR5-7, KA1 and KA2 subunits combine to form the kainate subtype of ionotropic glutamate receptors.¹⁴¹ Although inhaled anesthetics enhance currents mediated by kainate receptors containing GluR6 *in vitro*,¹⁴² GluR6 knockout mice have normal desflurane, halothane and isoflurane MAC values.¹⁴³ GluR6 editing mutant mice also do not demonstrate consistent changes in MAC values for these anesthetics.¹⁴³ However, the findings for GluR6 mutations do not conclusively eliminate kainate receptors as mediators of immobility because kainate receptors can be assembled from other subunits, even in the absence of the GluR6 subunit. Finally, the non-immobilizer F6 blocks mGluR5 (a metabotropic receptor),^{78,79} a finding inconsistent with a role for this class of receptors as mediators of immobility. Intrathecal injection of a blocker of metabotropic glutamate receptors does not change the MAC of isoflurane.¹⁴⁰

NMDA Receptors—Blockade of NMDA receptors, can markedly decrease MAC.^{129,144,145} Ketamine, which largely produces anesthesia by inhibiting NMDA receptor function [but also acts on acetylcholine receptors],¹⁴⁶ can abolish movement in response to noxious stimulation. However, blockade of NMDA receptors alone does not appear to cause immobility.¹²⁹ Several lines of evidence suggest that NMDA receptors do not mediate the immobility produced by inhaled anesthetics. For example, knockout of the $\epsilon 1$ subunit in mice does not change the concentration of isoflurane or sevoflurane¹⁴⁷ or of ethanol¹⁴⁸ producing loss of righting. Approximately 10% of Caucasians are missing the NMDA receptor subunit NR3B,¹⁴⁹ but determinations of MAC do not reveal outliers and, clinically, 10% of Caucasian patients do not appear to have major decreases in inhaled anesthetic requirement.

Other indirect evidence further supports the notion that NMDA receptors do not mediate immobility. Like ketamine¹⁵⁰ and nitrous oxide,¹⁵¹ xenon potently inhibits NMDA receptors *in vitro*.¹⁵² but 1 MAC xenon decreases cerebral metabolism in humans¹⁵³, whereas ketamine¹⁵⁴ and nitrous oxide¹⁵⁵ increase metabolism. On the basis of such discrepancies, the authors of a human study of xenon suggested that "...inhibition of the glutamatergic system is likely to be of minor significance for the anesthetic action of xenon *in vivo*."¹⁵³

A shortening of intervals between repeated stimuli increases the collective effect of stimulation (temporal summation) and increases the likelihood of a motor response to stimulation.¹²⁸ Studies with isoflurane indicate that approximately 40% of the generation of movement evoked by noxious stimulation (MAC) depends on interstimulus interval, suggesting the persistence of temporal summation and transmission via NMDA pathways.¹⁵⁶ That is, if temporal summation persists, then administration of the NMDA blocker MK-801 should and does abolish summation.¹⁵⁶ However, another interpretation is possible: perhaps isoflurane causes suppression of NMDA receptor transmission, and blockade with MK-801 simply substitutes

for the blockade produced by isoflurane. Consistent with the interpretation that isoflurane does not block temporal summation, electrophysiologic studies of neuronal wind-up show that temporal summation can occur during anesthesia.¹⁵⁷

Finally, administration of drugs that block NMDA receptors can decrease MAC for conventional inhaled anesthetics by more than 60%. This decrease does not correlate with the extent of functional blockade that these anesthetics produce at MAC, although it does correlate with blockade that fluorinated aromatic anesthetics produce at MAC.^{158,159} This failure of correlation plus the evidence from temporal summation and knockout mice leads to the conclusion that NMDA receptors do not mediate the immobility produced by conventional inhaled anesthetics.

NOS and Nitric Oxide

Knockout of the type I NOS isoform significantly increases isoflurane MAC.¹⁶⁰ The neuronal NOS inhibitors L-NAME^{161–163} or 7-NI^{160,162,164,165} can decrease MAC by 35%–95%, but not all investigators find an effect of L-NAME on MAC.^{166,167} Peculiarly, 7-NI decreases MAC to the same extent in wild-type and NOS knockout mice,¹⁶⁰ and arginine reverses the effect of the 7-NI by the same amount in wild-type and NOS knockout mice.¹⁶⁰ Thus, diverse interactions may govern the anesthetic effects of changes in transmission mediated by NOS and nitric oxide, but evidence for a role as a mediator of the immobility produced by inhaled anesthetics is limited.

Opioid Receptors

Analgesia is thought to accompany anesthesia by inhaled anesthetics. Thus, inhaled anesthetics might enhance the release of endogenous opioids, and/or enhance sensitivity of opioid receptors and thereby contribute to the anesthetic state. However, inhaled anesthetics do not appear to increase endogenous opioid concentrations in cerebrospinal fluid,¹⁶⁸ and they do not prevent autonomic or ventilatory responses to surgical stimulation at concentrations that suppress movement.⁷ Small doses of opioids markedly decrease inhaled anesthetic concentrations that prevent movement.¹⁶⁹ That is, opioids supply something (analgesia) that inhaled anesthetics do not produce or minimally produce. Finally, several reports suggest that administration of naloxone, a μ opioid receptor antagonist, does not affect the MAC of various inhaled anesthetics.^{104,170–177} A single report suggested that mice lacking the μ opioid receptor gene ($\mu\text{OR}^{-/-}$) had a sevoflurane MAC 20% greater than wild-type mice bearing the normal ($\mu\text{OR}^{+/+}$) gene ($3.3 \pm 0.5\%$ vs. $2.7 \pm 0.2\%$ sevoflurane), and that 0.1 mg/kg naloxone did not increase MAC in the knockout mice but increased MAC 18% in wild-type mice.¹⁷⁸ However, in a replication of this study, 0.1 and 1.0 mg/kg naloxone did not change sevoflurane MAC in the same strain of wild-type mice.¹⁷⁹ This replication added a control group of mice, and MAC was determined in a blinded manner. We conclude that opioid receptors can modulate, but do not mediate, the immobility produced by inhaled anesthetics.

Potassium Channels

Knockout of the TREK-1 potassium channel increases the MAC of chloroform, desflurane, halothane and sevoflurane.¹⁸⁰ However, the increase is not consistent, varying from 7% (desflurane) to 48% (halothane). This effect on MAC does not parallel the effect of these and other anesthetics on the capacity of those anesthetics to increase opening (current through) the channels *in vitro* (Fig. 3),^{180,181} a disparity that is inconsistent with a causal connection. Finally, an increase in temperature opens TREK-1 channels,⁶⁰ and the consequent increase in inhibitory current should decrease MAC. However, an increase in body temperature increases MAC in diverse mammals,^{61,182–184} further diminishing the likelihood that this channel is a relevant mediator.

Knockout of the TASK-3 channel significantly increases halothane MAC (by 18%) but not isoflurane MAC (a nonsignificant 9% increase is found).¹⁸⁵ Increases in pH increase the opening of TASK potassium channels,¹⁸⁶ but decreases in PaCO₂ and thus increases in pH do not decrease MAC.⁸⁵ Reducing pH blocks TASK channels,¹⁸⁶ and thus should increase MAC, but decreasing pH by increasing PaCO₂ decreases MAC rectilinearly.¹⁸⁷

MAC does not increase in mice lacking KNCK5 or Kir3.2 potassium channels.¹⁸⁸ Intracerebroventricular administration of cromakalim or pinacidil [ATP-sensitive potassium channel (K_{ATP}) blocking drugs] does not affect isoflurane MAC,¹⁸⁹ but this does not indicate what action would result from application to the spinal cord. Systemic administration of the K⁺ channel blocker 4-aminopyridine does not decrease halothane MAC in rats.⁸⁷ Administration of the TASK-1 or TASK-3 potassium channel blocker doxapram to mice does not increase MAC.¹⁹⁰

Intravenous and intrathecal infusions of riluzole (a nonspecific activator of KCNK potassium channels) decreases isoflurane MAC in rats,¹⁹¹ but a given dose produces the same change regardless of the route of administration. This result indicates that activation of potassium channels might affect anesthetic requirement, but does so primarily by an effect on higher centers rather than the spinal cord. The non-specificity of riluzole further complicates the issue. Riluzole even more potently blocks non-inactivating sodium channels,¹⁹² channels important to repetitive firing (i.e., sustained activity)¹⁹³ and thus to MAC.

If potassium channels are important, an increase in extracellular potassium ion concentration might increase MAC by decreasing polarization. In dogs, an increase in serum K⁺ from 3.8 ± 0.2 mEq/L to 7.4 ± 0.5 mEq/L did not change halothane MAC (1.09 ± 0.04% and 1.09 ± 0.04%).¹⁹⁴ Although the concurrent increase in cerebrospinal fluid K⁺ was statistically significant, it was too small (2.5 ± 0.1 mEq/L to 2.7 ± 0.1 mEq/L) to test the importance of K⁺. Changes in intrathecal K⁺ in rats induced by infusing artificial cerebrospinal fluid with altered KCl concentrations ranging from zero to 24 times normal did not increase MAC.¹⁹⁵ The highest concentration infused caused post-infusion impairment of hindlimb function.

Overall, these findings are not consistent with a role for potassium channels as mediators of the capacity of inhaled anesthetics to produce immobility.

Serotonin (5HT) Receptors

5-HT_{2A} Receptors—Although inhaled anesthetic concentrations of approximately 1 MAC block the *in vitro* effect of 5-HT on 5HT_{2A} receptors⁷⁸ and the 5HT_{2A} receptor blocker ketanserin^{196–204} can decrease nociception by supraspinal and spinal^{200,203,205} effects, other evidence suggests that 5HT_{2A} receptors, and serotonin receptors in general, play a minimal role in producing immobility.

Systemic administration of ketanserin does not change²⁰⁶ or may decrease^{207,208} MAC by up to 60%, larger doses proving lethal,²⁰⁸ but the same study finds that intrathecal administration of ketanserin decreases MAC only 20%–25%. If 5HT₂ receptors mediate MAC, then intrathecal injection should produce the greater effect. The 20%–25% decrease in MAC could result from absorption and an effect on higher centers (i.e., this would indicate that spinal 5HT₂ receptors are not important mediators of the immobility produced by inhaled anesthetics).

Halothane and the nonimmobilizer 1,2-dichlorohexafluorocyclobutane equally affect the 5HT₂ receptor *in vitro*, at 1 MAC or concentrations predicted to equal 1 MAC (1,2-dichlorohexafluorocyclobutane).⁷⁸ Finally, administration of parachlorophenylalanine, which

depletes serotonin, does not decrease halothane MAC in dogs.²⁰⁹ Such a finding, alone, suggests the lack of relevance of 5HT3 receptors as mediators of MAC.

5-HT2C Receptors—Both 1,1,2-trifluorocyclobutane (an anesthetic) and 1,2-dichlorohexafluorocyclobutane (a non-immobilizer) inhibit 5-HT2C receptors,^{78,79} and thus these receptors are unlikely mediators of immobility.

5-HT3 Receptors—Blockade of the 5-HT3 receptor by systemic²¹⁰ or intrathecal (Pamela Flood, personal communication) administration of ondansetron does not affect the MAC of isoflurane.

Summary Regarding Ligand- and Voltage-Gated Channels as Mediators

No ligand- or voltage-gated channel or other targets discussed above appears capable of explaining more than a minor part of the immobility produced by inhaled anesthetics. One response to this observation is that it is the combined actions of numerous small effects on diverse targets that explains immobility. Perhaps 20 such targets, each contributing 5% to the production of anesthesia, add to produce a 100% effect, i.e., produce MAC. What does this hypothesis imply? Given that the spinal cord mediates the immobility produced by inhaled anesthetics,⁴ each of the 20 targets would have to be present in appreciable numbers in the spinal cord or, if in small numbers, such a target would have to contribute mightily. Anesthetics would have to affect each target at relevant (anesthetizing) concentrations in a way that plausibly explained anesthesia. Note the enormous variation in the differences in functional potencies, acetylcholine receptors nearly completely blocked at 0.1–0.2 MAC⁶² and NMDA receptors affected, the majority blocked, by some anesthetics only at concentrations 2–3 times MAC.¹⁵⁹ Targets affected at sub-MAC concentrations would contribute less than those requiring supra-MAC concentrations so that at MAC each would contribute just 5% to the MAC. That is, fortuitously, each would, indeed, make a 5% contribution without any one target providing a large, obvious, contribution (i.e., one that would exceed the 5% average contribution, and therefore be measurable). That is, the 20-target theory would have to argue that no one target would stand out, even for one anesthetic. The 20 targets would have to add to each other's effect and not be synergistic (lest we see wide differences in potency/MAC among anesthetics that did not reflect the Meyer-Overton correlation). Thus, we see the 20-target theory as possible, but unlikely.

Another response is that we have not found the channel that mediates a major part of the immobility produced by inhaled anesthetics. The sodium channel might fit that argument (see below), but we are at a loss to deal with the researcher who says there are tens or thousands of X (e.g., potassium) channels that have not been explored. We can only answer that we cannot disprove the existence of dragons.²¹¹

Sodium Channels as the Mediators of Immobility

Why Sodium Channels?

The sodium channel presents an attractive target in that it potentially affects all ligand-gated ion channels because depolarization of the nerve terminal (a process governed by the sodium channel) underlies neurotransmitter release at those terminals. The diversity of sodium channels adds to their attractiveness, and the difficulty of ascribing anesthesia to an action on a particular channel. Fast inactivating sodium channels mediate the fast depolarization-repolarization that underlies action potentials, while other slowly inactivating sodium channels contribute to repetitive firing (i.e., sustained activity)¹⁹³ and perhaps temporal summation, and thus to MAC. Still other sodium channels on dendrites influence the response to release of synaptic ligands.²¹² Thus, an effect on sodium channels allows for a multiplicity of

anesthetic actions that depends on the distribution of effects on various sodium channels. If immobility results from an action on sodium channels and, thereby, on multiple ligand-gated channels simultaneously, that might be consistent with the steep dose (anesthetic concentration)-response (movement) relationship seen for MAC,²¹³ although a more likely explanation for the steepness of the dose-response relationship that underlies MAC is low inter-subject variability in sensitivity.

Evidence Supporting Sodium Channels as Mediators of Immobility

Some findings indirectly support a role for sodium channels. An increase in central nervous system extracellular sodium rectilinearly increases MAC and conversely a decrease decreases MAC.¹⁹⁴ Systemic administration of the sodium channel blocker lidocaine progressively decreases MAC for several conventional anesthetics in rats.²¹⁴ Lidocaine shares one other property with some inhaled anesthetics [e.g., enflurane²¹⁵ and sevoflurane],²¹⁶ the capacity to produce convulsions.²¹⁷ Could this result from differential effects on the various sodium channels (e.g., decrease inhibitory output relative to excitatory output)? Intrathecal administration of veratridine, a compound that sustains the open state of sodium channels, increases isoflurane MAC in rats by a maximum of 21%.²¹⁸

Sodium channels have been given little attention because sodium channel-dependent axonal conduction continues at all levels of anesthesia, and many levels of synaptic transmission remain intact. However, this ignores the greater vulnerability of the bare nerve terminal in contrast to the considerable reserve in conduction in myelinated nerves. In further support of the importance of sodium channels, inhaled anesthetics inhibit multiple isoforms of Nav α -subunits (rat Nav_v1.2,²¹⁹ human Nav_v1.5,²²⁰ rat Nav_v1.2, rat 1.6, and human Nav_v1.4),²²¹ with small differences in anesthetic potencies and mechanism. Isoflurane at clinically relevant concentrations inhibits rat neuronal (Nav_v1.2), skeletal muscle (Nav_v1.4), and cardiac muscle (Nav_v1.5) voltage-gated Na⁺ channel α subunits heterologously expressed in Chinese hamster ovary cells with subtle isoform-dependent differences.²²² Although an earlier study found that the human Nav_v1.8 isoform heterologously expressed in *Xenopus* oocytes was insensitive,²²¹ recent evidence indicates that isoflurane inhibits Nav_v1.8 expressed in a neuroblastoma cell line (HCH, unpublished observations).

Potency differs among anesthetics with inhibition of rat Nav_v1.2 expressed in *Xenopus* oocytes by 2 MAC cyclopropane being two-thirds that of 2 MAC isoflurane and by 2 MAC halothane being one-third.²²¹ Rehberg et al. found that isoflurane and halothane were equally potent but that enflurane was much less potent.²¹⁹ The non-immobilizer 1,2-dichlorohexafluorobutane has no effect.²²¹

Sodium channel inhibition, and a consequently decreased nerve action potential, decreases release of neurotransmitters such as glutamate,²²³ with 1 MAC isoflurane having a greater effect than 1 MAC halothane, but at 2 MAC, the difference disappears. Both isoflurane and propofol produce a dose-related inhibition of sodium channels in neurohypophysial nerve terminals, approximately a 30% inhibition at 1 MAC or its equivalent,²²⁴ and progressively greater inhibition at larger concentrations. The non-immobilizer 1,2-dichlorohexafluorobutane does not affect sodium-channel-dependent glutamate release.²²⁵

A few factors suggest caution regarding the relevance of sodium channels. As noted above, the effect of inhaled anesthetics on most sodium channels is limited at MAC. *In vitro* studies suggest that the effect at MAC differs among conventional inhaled anesthetics and that propofol acts like isoflurane in decreasing current.²²⁴ If propofol acts solely by enhancing GABA_A receptors what does this imply? A partial answer might be that propofol is a much weaker blocker than isoflurane. A similar concern applies to barbiturates which, like propofol, act by enhancing the response of GABA_A receptors. Barbiturates can block sodium channels by a

pH-dependent mechanism acting on the inside of the cell.²²⁶ Finally, as noted above, IV and intrathecal infusions of riluzole equally decrease isoflurane MAC in rats.¹⁹¹ Riluzole potently blocks both fast inactivating and non-inactivating sodium channels,¹⁹² the latter channels being important to repetitive firing (i.e., sustained activity).¹⁹³ This result indicates that blockade of such sodium channels might affect anesthetic requirement by acting on higher centers rather than the spinal cord, the site of inhaled anesthetic action. This says nothing concerning sodium channels responsible for rapid nerve conduction. Finally, none of this indicates how anesthetics might block the sodium channel. Do they do it by an action directly on the channel or indirectly through an action on the membrane bilayer?

What Might a Nonspecific Mechanism Be?

Do inhaled anesthetics act by affecting surface properties of the membrane bilayer? Polyhydroxyalkanes are compounds confined to the membrane surface by virtue of their multiple hydroxyl groups, yet they have anesthetic effects in tadpoles.²²⁷ Their size also would preclude entrance of the whole molecule into the pockets hypothesized as binding sites for inhaled anesthetics. Still, other large alcohols can cause anesthesia.^{228,229} The molecular excess that does not fit into the pocket might simply extend from the pocket.

Cantor has suggested that anesthetics might act at the bilayer/membrane interface to alter the pressure profile within the membrane and thereby alter the function of proteins that reside there.²³⁰ Roth et al.²³¹ propose a different theory, one that has elements of Meyer and Overton's thesis. They suggest that water vapor naturally forms bubbles in small tubes (i.e., ion channels) and that organisms make use of such formation in the regulation of conduction through channels. Anesthetics influence the formation of the bubbles, thereby altering conduction and producing anesthesia. One appeal of this theory is that it would affect many channels, an action known for inhaled anesthetics.

Conclusions

The preceding discussion suggests that despite many plausible candidates, at present, no single target, including ligand- and voltage-gated channels, can explain the immobility produced by inhaled anesthetics, nor can any combination of effects currently explain immobility. (We consider "plausible candidates" to be channels conducting excitatory impulses that inhaled anesthetics block or channels conducting inhibitory impulses that anesthetics enhance.) As an aside, we note that none of the discussion necessarily applies to the capacity of inhaled anesthetics to suppress learning and memory. Future studies will continue to test the relevance of particular ligand- and voltage-gated channels to immobility and other anesthetic end-points. Future studies might need to test radically different theories, perhaps resurrecting Meyer and Overton's ideas in a drastically different form. Even if anesthetics act on lipid bilayers, they will exert their action through changes induced in protein and ion current conduction. No matter where they act, we still must explain how conformational changes in lipid or protein structure leads to anesthesia.

Acknowledgements

NIH grant 1P01GM47818 (UCSF) supported this work

References

1. Merkel G, Eger EI II. A comparative study of halothane and halopropane anesthesia. Including a method for determining equipotency. *Anesthesiology* 1963;24:346–357. [PubMed: 13935000]

2. Saidman LJ, Eger EI II. Effect of nitrous oxide and of narcotic premedication on the alveolar concentration of halothane required for anesthesia. *Anesthesiology* 1964;25:302–306. [PubMed: 14156570]
3. Antognini JF, Raines DE, Solt K, Barter LS, Atherley RJ, Bravo E, Laster MJ, Jankowska K, Eger EI 2nd. Hexafluorobenzene acts in the spinal cord, whereas o-difluorobenzene acts in both brain and spinal cord, to produce immobility. *Anesth Analg* 2007;104:822–828. [PubMed: 17377088]
4. Antognini JF, Schwartz K. Exaggerated anesthetic requirements in the preferentially anesthetized brain. *Anesthesiology* 1993;79:1244–1249. [PubMed: 8267200]
5. Rampil IJ. Anesthetic potency is not altered after hypothermic spinal cord transection in rats. *Anesthesiology* 1994;80:606–610. [PubMed: 8141455]
6. Roizen MF, Horrigan RW, Frazer BF. Anesthetic doses blocking adrenergic (stress) and cardiovascular responses to incision - MAC BAR. *Anesthesiology* 1981;54:390–398. [PubMed: 7224208]
7. France CJ, Plumer HM, Eger EI II, Wahrenbrock EA. Ventilatory effects of isoflurane (Forane) or halothane when combined with morphine, nitrous oxide and surgery. *Br J Anaesth* 1974;46:117–120. [PubMed: 4834474]
8. Katoh T, Kobayashi S, Suzuki A, Iwamoto T, Bito H, Ikeda K. The effect of fentanyl on sevoflurane requirements for somatic and sympathetic responses to surgical incision. *Anesthesiology* 1999;90:398–405. [PubMed: 9952144]
9. Rehberg B, Ruschner R, Fisher M, Ebeling BJ, Hoeft A. Concentration-dependent changes in the latency and amplitude of somatosensory-evoked potentials by desflurane, isoflurane and sevoflurane. *Anaesthesiol Intensivmed Notfallmed Schmerzther* 1998;33:425–429.
10. Antognini JF, Carstens E. Increasing isoflurane from 0.9 to 1.1 minimum alveolar concentration minimally affects dorsal horn cell responses to noxious stimulation. *Anesthesiology* 1999;90:208–214. [PubMed: 9915330]
11. Barter LS, Mark LO, Jinks SL, Carstens EE, Antognini JF. Immobilizing doses of halothane, isoflurane or propofol, do not preferentially depress noxious heat-evoked responses of rat lumbar dorsal horn neurons with ascending projections. *Anesth Analg* 2008;106:985–990. [PubMed: 18292450]
12. Merrill AW, Barter LS, Rudolph U, Eger EI 2nd, Antognini JF, Carstens MI, Carstens E. Propofol's effects on nociceptive behavior and spinal c-fos expression after intraplantar formalin injection in mice with a mutation in the gamma-aminobutyric acid-type(A) receptor beta3 subunit. *Anesth Analg* 2006;103:478–483. [PubMed: 16861438]
13. Reinacher PC, Priebe HJ, Blumrich W, Zentner J, Scheufler KM. The effects of stimulation pattern and sevoflurane concentration on intraoperative motor-evoked potentials. *Anesth Analg* 2006;102:888–895. [PubMed: 16492846]
14. Jinks SL, Atherley RJ, Dominguez CL, Sigvardt KA, Antognini JF. Isoflurane disrupts central pattern generator activity and coordination in the lamprey isolated spinal cord. *Anesthesiology* 2005;103:567–575. [PubMed: 16129982]
15. Antognini JF, Atherley RJ, Dutton RC, Laster MJ, Eger EI 2nd, Carstens E. The excitatory and inhibitory effects of nitrous oxide on spinal neuronal responses to noxious stimulation. *Anesth Analg* 2007;104:829–835. [PubMed: 17377089]
16. Eger EI II, Tang M, Liao M, Laster MJ, Solt K, Flood P, Jenkins A, Raines D, Hendrickx J, Shafer S, Yasumasa T, Sonner JM. Inhaled anesthetics do not combine to produce synergistic effects regarding MAC. *Anesth Analg* 2008;106In press AUGUST 2008 ISSUE
17. Hendrickx JFA, Eger EI II, Sonner JM, Shafer SL. Is synergy the rule? A review of anesthetic drug interactions producing hypnosis and immobility. *Anesth Analg* 2008;106In press AUGUST 2008 ISSUE
18. Shafer SL, Hendrickx JFA, Flood P, Sonner J, Eger EI II. Additivity, synergy, and anesthetic mechanisms. *Anesth Analg* 2008;106In press AUGUST 2008 ISSUE
19. Franks NP, Lieb WR. Temperature dependence of the potency of volatile general anesthetics. *Anesthesiology* 1996;84:716–720. [PubMed: 8659800]
20. Krasowski MD, Harrison NL. General anaesthetic actions on ligand-gated ion channels. *Cell Mol Life Sci* 1999;55:1278–1303. [PubMed: 10487207]

21. Squires RF, Saederup E. Additivities of compounds that increase the numbers of high affinity [3H] muscimol binding sites by different amounts define more than 9 GABA(A) receptor complexes in rat forebrain: implications for schizophrenia and clozapine research. *Neurochem Res* 2000;25:1587–1601. [PubMed: 11152388]
22. Davies M, Thuynsma RP, Dunn SM. Effects of propofol and pentobarbital on ligand binding to GABAA receptors suggest a similar mechanism of action. *Can J Physiol Pharmacol* 1998;76:46–52. [PubMed: 9564548]
23. Sebel LE, Richardson JE, Singh SP, Bell SV, Jenkins A. Additive effects of sevoflurane and propofol on gamma-aminobutyric acid receptor function. *Anesthesiology* 2006;104:1176–1183. [PubMed: 16732088]
24. Bertaccini EJ, Trudell JR, Franks NP. The common chemical motifs within anesthetic binding sites. *Anesth Analg* 2007;104:318–324. [PubMed: 17242087]
25. Sewell JC, Sear JW. Determinants of volatile general anesthetic potency: a preliminary three-dimensional pharmacophore for halogenated anesthetics. *Anesth Analg* 2006;102:764–771. [PubMed: 16492826]
26. Sewell JC, Sear JW. Derivation of preliminary three-dimensional pharmacophores for nonhalogenated volatile anesthetics. *Anesth Analg* 2004;99:744–751. [PubMed: 15333405]
27. Lysko GS, Robinson JL, Casto R, Ferrone RA. The stereospecific effects of isoflurane isomers in vivo. *Eur J Pharmacol* 1994;263:25–29. [PubMed: 7821359]
28. Won A, Oh I, Laster MJ, Popovich J, Eger EI 2nd, Sonner JM. Chirality in anesthesia I: minimum alveolar concentration of secondary alcohol enantiomers. *Anesth Analg* 2006;103:81–84. [PubMed: 16790631]
29. Brosnan R, Gong D, Cotten J, Keshavaprasad B, Yost CS, Eger EI 2nd, Sonner JM. Chirality in anesthesia II: stereoselective modulation of ion channel function by secondary alcohol enantiomers. *Anesth Analg* 2006;103:86–91. [PubMed: 16790632]
30. White PF, Schuttler J, Shafer A, Stanski DR, Horai Y, Trevor AJ. Comparative pharmacology of the ketamine isomers. *Studies in volunteers*. *Br J Anaesth* 1985;57:197–203. [PubMed: 3970799]
31. Dickinson R, Peterson BK, Banks P, Simillis C, Martin JC, Valenzuela CA, Maze M, Franks NP. Competitive inhibition at the glycine site of the N-methyl-D-aspartate receptor by the anesthetics xenon and isoflurane: evidence from molecular modeling and electrophysiology. *Anesthesiology* 2007;107:756–767. [PubMed: 18073551]
32. Meyer HH. Theorie der Alkoholnarkose. *Arch Exptl Pathol Pharmacol* 1899;42:109–118.
33. Overton E. Studien über die Narkose, Zugleich ein Beitrag zur allgemeinen Pharmakologie. Gustav Fischer, Jena 1901:1–195.
34. Franks NP, Lieb WR. Do general anaesthetics act by competitive binding to specific receptors? *Nature* 1984;310:599–601. [PubMed: 6462249]
35. Koblin DD, Chortkoff BS, Laster MJ, Eger EI II, Halsey MJ, Ionescu P. Polyhalogenated and perfluorinated compounds that disobey the Meyer-Overton hypothesis. *Anesth Analg* 1994;79:1043–1048. [PubMed: 7978424]
36. Chipot C, Wilson MA, Pohorille A. Interactions of anesthetics with the hexane-water interface. *J Phys Chem B* 1996;101:782–791. [PubMed: 11542402]
37. Pohorille A, Wilson MA, Cieplak P. Interaction of alcohols and anesthetics with the water-hexane interface: A molecular dynamics study. *Prog Colloid Polym Sci* 1997;103:29–40.
38. Abraham MH, Acree WE Jr, Mintz C, Payne S. Effect of anesthetic structure on inhalation anesthesia: Implications for the mechanism. *J Pharm Sci*. 2007Epub ahead of print
39. Janoff AS, Pringle MJ, Miller KW. Correlation of general anesthetic potency with solubility in membranes. *Biochem Biophys Acta* 1981;649:125–128. [PubMed: 7306543]
40. Pringle MJ, Brown KB, Miller KM. Can the lipid theories of anesthesia account for the cutoff in anesthetic potency in homologous series of alcohols. *Mol Pharmacol* 1981;19:49–55. [PubMed: 7207463]
41. Sonner JM. A hypothesis on the origin and evolution of the response to inhaled anesthetics. *Anesth Analg* 2008;107In press THIS ISSUE
42. Eckenhoff R. Why can all of biology be anesthetized? *Anesth Analg* 2008;107In press THIS ISSU

43. Lynch C. Meyer and overton revisited. *Anesth Analg* 2008;107In press THIS ISSUE
44. Sedensky MM, Morgan PG. Genetics and the evolution of the anesthetic response. *Anesth Analg* 2008;107In press THIS ISSUE
45. Crowder CM. Does natural selection explain the universal response of metazoans to volatile anesthetics? *Anesth Analg* 2008;107In press THIS ISSUE
46. Durieux ME. The evolution of anesthetic sensitivity and the evolution of a paper. *Anesth Analg* 2008;107In press THIS ISSUE
47. Nandini-Kishore SG, Mattox SM, Martin CE, Thompson GA Jr. Membrane changes during growth of *Tetrahymena* in the presence of ethanol. *Biochim Biophys Acta* 1979;551:315–327. [PubMed: 105757]
48. Nandini-Kishore SG, Kitajima Y, Thompson GA Jr. Membrane fluidizing effects of the general anesthetic methoxyflurane elicit an acclimation response in *Tetrahymena*. *Biochim Biophys Acta* 1977;471:157–161. [PubMed: 411511]
49. Koblin DD, Wang HH. Chronic exposure to inhaled anesthetics increases cholesterol content in *Acholeplasma laidlawii*. *Biochim Biophys Acta* 1981;649:717–725. [PubMed: 6274406]
50. Ingram LO. Adaptation of membrane lipids to alcohols. *J Bacteriol* 1976;125:670–678. [PubMed: 1107328]
51. Yang L, Sonner JM. Anesthetic-like modulation of receptor function by surfactants: A test of the interfacial theory of anesthesia. *Anesth Analg* 2008;106In press THIS ISSUE
52. Brosnan RJ, Yang L, Milutinovic PS, Zhao J, Laster MJ, Eger EI 2nd, Sonner JM. Ammonia has anesthetic properties. *Anesth Analg* 2007;104:1430–1433. [PubMed: 17513636]
53. Yang L, Zhao J, Milutinovic PS, Brosnan RJ, Eger EI 2nd, Sonner JM. Anesthetic properties of the ketone bodies beta-hydroxybutyric acid and acetone. *Anesth Analg* 2007;105:673–679. [PubMed: 17717222]
54. Cantor RS. Receptor desensitization by neurotransmitters in membranes: are neurotransmitters the endogenous anesthetics? *Biochemistry* 2003;42:11891–11897. [PubMed: 14556619]
55. Milutinovic PS, Yang L, Cantor RS, Eger EI 2nd, Sonner JM. Anesthetic-like modulation of a gamma-aminobutyric acid type A, strychnine-sensitive glycine, and N-methyl-d-aspartate receptors by coreleased neurotransmitters. *Anesth Analg* 2007;105:386–392. [PubMed: 17646495]
56. Antognini JF, Carstens E, Atherley R. Does the immobilizing effect of thiopental in brain exceed that of halothane? *Anesthesiology* 2002;96:980–986. [PubMed: 11964608]
57. Rampil IJ, Mason P, Singh H. Anesthetic potency (MAC) is independent of forebrain structures in the rat. *Anesthesiology* 1993;78:707–712. [PubMed: 8466071]
58. Garcia-Fernandez J, Parodi E, Garcia P, Matute E, I AG-d-S, Cediel R, Gilsanz F. Clinical actions of subarachnoid sevoflurane administration in vivo: a study in dogs. *Br J Anaesth* 2005;95:530–534. [PubMed: 16040635]
59. Chai YF, Yang J, Liu J, Song HB, Yang JW, Liu SL, Zhang WS, Wang QW. Epidural anaesthetic effect of the 8% emulsified isoflurane: a study in rabbits. *Br J Anaesth* 2008;100:109–115. [PubMed: 18042558]
60. Maingret F, Lauritzen I, Patel AJ, Heurteaux C, Reyes R, Lesage F, Lazdunski M, Honore E. TREK-1 is a heat-activated background K(+) channel. *EMBO J* 2000;19:2483–2491. [PubMed: 10835347]
61. Eger EI II, Johnson BH. MAC of I-653 in rats, including a test of the effect of body temperature and anesthetic duration. *Anesth Analg* 1987;66:974–976. [PubMed: 3631594]
62. Violet JM, Downie DL, Nakisa RC, Lieb WR, Franks NP. Differential sensitivities of mammalian neuronal and muscle nicotinic acetylcholine receptors to general anesthetics. *Anesthesiology* 1997;86:866–874. [PubMed: 9105231]
63. Zhang Y, Laster MJ, Eger EI II, Sharma M, Sonner JM. Blockade of acetylcholine receptors does not change the dose of etomidate required to produce immobility in rats. *Anesth Analg* 2007;104:850–852. [PubMed: 17377093]
64. Jurd R, Arras M, Lambert S, Drexler B, Siegart R, Crestani F, Zaugg M, Vogt KE, Ledermann B, Antkowiak B, Rudolph U. General anesthetic actions in vivo strongly attenuated by a point mutation in the GABA(A) receptor beta3 subunit. *FASEB J* 2003;17:250–252. [PubMed: 12475885]

65. Liao M, Sonner JM, Jurd R, Rudolph U, Borghese CM, Harris RA, Laster MJ, Eger EI 2nd. Beta3-containing gamma-aminobutyric acidA receptors are not major targets for the amnesic and immobilizing actions of isoflurane. *Anesth Analg* 2005;101:412–418. [PubMed: 16037154]
66. Katayama S, Irifune M, Kikuchi N, Takarada T, Shimizu Y, Endo C, Takata T, Dohi T, Sato T, Kawahara M. Increased gamma-aminobutyric acid levels in mouse brain induce loss of righting reflex, but not immobility, in response to noxious stimulation. *Anesth Analg* 2007;104:1422–1429. [PubMed: 17513635]
67. Deady J, Koblin DD, Eger EI II, Heavner JE, D'Aoust B. Anesthetic potencies and the unitary theory of narcosis. *Anesth Analg* 1981;60:380–384. [PubMed: 7195159]
68. Sonner JM, Antognini JF, Dutton RC, Flood P, Gray AT, Harris RA, Homanics GE, Kendig JJ, Orser BA, Raines DE, Trudell JR, Vissel B, Eger EI II. Inhaled anesthetics and immobility: Mechanisms, mysteries, and MAC. *Anesth Analg* 2003;97:718–740. [PubMed: 12933393]
69. Franks NP, Lieb WR. Molecular and cellular mechanisms of general anaesthesia. *Nature* 1994;367:607–614. [PubMed: 7509043]
70. Harris RA, Mihic SJ, Dildy-Mayfield JE, Machu TK. Actions of anesthetics on ligand-gated ion channels: role of receptor subunit composition. *FASEB J* 1995;9:1454–1462. [PubMed: 7589987]
71. Hemmings HC Jr, Akabas MH, Goldstein PA, Trudell JR, Orser BA, Harrison NL. Emerging molecular mechanisms of general anesthetic action. *Trends Pharmacol Sci* 2005;26:503–510. [PubMed: 16126282]
72. Krasowski MD, Harrison NL. The actions of ether, alcohol and alkane general anaesthetics on GABAA and glycine receptors and the effects of TM2 and TM3 mutations. *Br J Pharmacol* 2000;129:731–743. [PubMed: 10683198]
73. Mihic SJ, Ye Q, Wick MJ, Koltchine VV, Krasowski MD, Finn SE, Mascia MP, Valenzuela CF, Hanson KK, Greenblatt EP, Harris RA, Harrison NL. Sites of alcohol and volatile anaesthetic action on GABA_A and glycine receptors. *Nature* 1997;389:385–389. [PubMed: 9311780]
74. Flood P, Sonner JM, Gong D, Coates KM. Heteromeric nicotinic inhibition by isoflurane does not mediate MAC or loss of righting reflex. *Anesthesiology* 2002;97:902–905. [PubMed: 12357157]
75. Eger EI II, Zhang Y, Laster MJ, Flood P, Kendig JJ, Sonner JM. Acetylcholine receptors do not mediate the immobilization produced by inhaled anesthetics. *Anesth Analg* 2002;94:1500–1504. [PubMed: 12032015]
76. Wang MY, Rampil IJ, Kendig JJ. Ethanol directly depresses AMPA and NMDA glutamate currents in spinal cord motor neurons independent of actions on GABAA or glycine receptors. *J Pharmacol Exp Ther* 1999;290:362–367. [PubMed: 10381800]
77. Do SH, Kamatachi GL, Durieux ME. The effects of isoflurane on native and chimeric muscarinic acetylcholine receptors: the role of protein kinase C. *Anesth Analg* 2001;93:375–381. [PubMed: 11473864]
78. Minami K, Minami M, Harris RA. Inhibition of 5-hydroxytryptamine type 2A receptor-induced currents by n-alcohols and anesthetics. *J Pharmacol Exp Ther* 1997;281:1136–1143. [PubMed: 9190846]
79. Minami K, Gereau RW, Minami M, Heinemann SF, Harris RA. Effects of ethanol and anesthetics on type 1 and 5 metabotropic glutamate receptors expressed in *Xenopus laevis* oocytes. *Mol Pharmacol* 1998;53:148–156. [PubMed: 9443943]
80. Raines DE, Claycomb RJ, Forman SA. Nonhalogenated anesthetic alkanes and perhalogenated nonimmobilizing alkanes inhibit α 4 β 2 neuronal nicotinic acetylcholine receptors. *Anesth Analg* 2002;95:573–577. [PubMed: 12198040]
81. Pidoplichko VI, Dani JA. Acid-sensitive ionic channels in midbrain dopamine neurons are sensitive to ammonium, which may contribute to hyperammonemia damage. *Proc Natl Acad Sci U S A* 2006;103:11376–11380. [PubMed: 16847263]
82. Wu LJ, Duan B, Mei YD, Gao J, Chen JG, Zhuo M, Xu L, Wu M, Xu TL. Characterization of acid-sensing ion channels in dorsal horn neurons of rat spinal cord. *J Biol Chem* 2004;279:43716–43724. [PubMed: 15302881]
83. Baron A, Voilley N, Lazdunski M, Lingueglia E. Acid sensing ion channels in dorsal spinal cord neurons. *J Neurosci* 2008;28:1498–1508. [PubMed: 18256271]

84. Duan B, Wu LJ, Yu YQ, Ding Y, Jing L, Xu L, Chen J, Xu TL. Upregulation of acid-sensing ion channel ASIC1a in spinal dorsal horn neurons contributes to inflammatory pain hypersensitivity. *J Neurosci* 2007;27:11139–11148. [PubMed: 17928456]
85. Eisele JH, Eger EI II, Muallem M. Narcotic properties of carbon dioxide in the dog. *Anesthesiology* 1967;28:856–865. [PubMed: 6035017]
86. Seitz PA, ter Reit M, Rush W, Merrell W. Adenosine decreases the minimum alveolar concentration of halothane in dogs. *Anesthesiology* 1990;73:990–994. [PubMed: 2240689]
87. Ma HC, Wang YF, Feng CS, Zhao H, Dohi S. Effects of adenosine agonist R-phenylisopropyl-adenosine on halothane anesthesia and antinociception in rats. *Acta Pharmacol Sin* 2005;26:181–185. [PubMed: 15663896]
88. Nicholls EA, Louie GL, Prokocimer PG, Maze M. Halothane anesthetic requirements are not affected by aminophylline treatment in rats and dogs. *Anesthesiology* 1986;65:637–641. [PubMed: 3789436]
89. Govinda R, Kasuya Y, Orhan-Sungur M, Turan A, Sessler DI. Aminophylline does not increase the minimum alveolar concentration of desflurane. *Anesthesiology* 2007;107:A1919.(Abstract)
90. Suzuki A, Katoh T, Ikeda K. The effect of adenosine triphosphate on sevoflurane requirements for minimum alveolar anesthetic concentration and minimum alveolar anesthetic concentration-awake. *Anesth Analg* 1998;86:179–183. [PubMed: 9428875]
91. Hara K, Yamakura T, Sata T, Harris RA. The effects of anesthetics and ethanol on alpha2 adrenoceptor subtypes expressed with G protein-coupled inwardly rectifying potassium channels in *Xenopus* oocytes. *Anesth Analg* 2005;101:1381–1388. [PubMed: 16243998]
92. Rabin BC, Reid K, Guo TZ, Gustafsson E, Zhang C, Maze M. Sympatholytic and minimum anesthetic concentration-sparing responses are preserved in rats rendered tolerant to the hypnotic and analgesic action of dexmedetomidine, a selective α -2 adrenergic agonist. *Anesthesiology* 1996;85:565–573. [PubMed: 8853087]
93. Bloor BC, Flacke WE. Reduction in halothane anesthetic requirement by clonidine, and alpha-adrenergic agonist. *Anesth Analg* 1982;61:741–745. [PubMed: 6125112]
94. Eger EI II, Xing Y, Laster MJ, Sonner JM. α -2 adrenoceptors probably do not mediate the immobility produced by inhaled anesthetics. *Anesth Analg* 2003;96:1661–1664. [PubMed: 12760992]
95. Study RE. Isoflurane inhibits multiple voltage-gated calcium currents in hippocampal pyramidal neurons. *Anesthesiology* 1994;81:104–116. [PubMed: 8042778]
96. Lingamaneni R, Hemmings HC Jr. Differential interaction of anaesthetics and antiepileptic drugs with neuronal Na⁺ channels, Ca₂⁺ channels, and GABA(A) receptors. *Br J Anaesth* 2003;90:199–211. [PubMed: 12538378]
97. Hall AC, Lieb WR, Franks NP. Insensitivity of P-type calcium channels to inhalational and intravenous general anesthetics. *Anesthesiology* 1994;81:117–123. [PubMed: 8042779]
98. Graham MD, Hopkins PM, Harrison SM. Antagonistic actions of halothane and sevoflurane on spontaneous Ca₂⁺ release in rat ventricular myocytes. *Anesthesiology* 2006;105:58–64. [PubMed: 16809995]
99. Petrenko AB, Tsujita M, Kohno T, Sakimura K, Baba H. Mutation of alpha1G T-type Calcium Channels in Mice Does Not Change Anesthetic Requirements for Loss of the Righting Reflex and Minimum Alveolar Concentration but Delays the Onset of Anesthetic Induction. *Anesthesiology* 2007;106:1177–1185. [PubMed: 17525593]
100. Tanifuji Y, Zhang Y, Liao M, Eger EI 2nd, Laster MJ, Sonner JM. Do dopamine receptors mediate part of MAC? *Anesth Analg* 2006;103:1177–1181. [PubMed: 17056951]
101. Miller RD, Way WL, Eger EI II. The effects of alpha-methyl dopa, reserpine, guanethidine, and iproniazid on minimum alveolar anesthetic requirement (MAC). *Anesthesiology* 1968;29:1153–1158. [PubMed: 5726752]
102. Johnston R, White P, Eger EI II. Comparative effects of dextroamphetamine and reserpine on halothane and cyclopropane anesthetic requirements. *Anesth Analg* 1975;54:655–659. [PubMed: 1237254]
103. Raja SN, Moscicki JC, Woodside JR Jr, DiFazio CA. The effect of acute phencyclidine administration on cyclopropane requirement (MAC) in rats. *Anesthesiology* 1982;56:275–279. [PubMed: 6802034]

104. Segal IS, Vickery RG, Walton JK, Doze VA, Maze M. Dexmedetomidine diminishes halothane anesthetic requirements in rats through a postsynaptic α -2 adrenergic receptor. *Anesthesiology* 1988;69:818–823. [PubMed: 2848424]
105. Mueller RA, Smith RD, Spruill WA, Breese GR. Central monaminergic neuronal effects on minimum alveolar concentrations (MAC) of halothane and cyclopropane in rats. *Anesthesiology* 1975;42:143–152. [PubMed: 1115362]
106. Krasowski M, Jenkins A, Flood P, Kung A, Hopfinger A, Harrison N. General anesthetic potencies of a series of propofol analogs correlate with potency for potentiation of gamma-aminobutyric acid (GABA) current at the GABA(A) receptor but not with lipid solubility. *J Pharmacol Exp Ther* 2001;297:338–351. [PubMed: 11259561]
107. Harrison NL, Kugler JL, Jones MV, Greenblatt EP, Pritchett DB. Positive modulation of human γ -aminobutyric acid type A and glycine receptors by the inhalation anesthetic isoflurane. *Mol Pharmacol* 1993;44:628–632. [PubMed: 7690453]
108. Grasshoff C, Drexler B, Hentschke H, Thiermann H, Antkowiak B. Cholinergic Modulation of Sevoflurane Potency in Cortical and Spinal Networks In Vitro. *Anesthesiology* 2007;106:1147–1155. [PubMed: 17525590]
109. Sonner JM, Zhang Y, Stabernack C, Abaigar W, Xing Y, Sharma M, Eger EI II. GABA(A) receptor blockade antagonizes the immobilizing action of propofol but not ketamine or isoflurane in a dose-related manner. *Anesth Analg* 2003;96:706–712. [PubMed: 12598250]
110. Hara K, Eger EI II, Laster MJ, Harris RA. Nonhalogenated alkanes cyclopropane and butane affect neurotransmitter-gated ion channel and G-protein-coupled receptors: Differential actions on GABA (A) and glycine receptors. *Anesthesiology* 2002;97:1512–1520. [PubMed: 12459679]
111. Raines DE, Claycomb RJ, Scheller M, Forman A. Nonhalogenated alkane anesthetics fail to potentiate agonist actions on two ligand-gated channels. *Anesthesiology* 2001;95:470–477. [PubMed: 11506122]
112. Zhang Y, Sonner JM, Eger EI II, Stabernack CR, Laster MJ, Raines DE, Harris RA. GABAA receptors do not mediate the immobility produced by inhaled anesthetics. *Anesth Analg* 2004;99:85–90. [PubMed: 15281509]
113. Sonner JM, Cascio M, Xing Y, Fanselow MS, Kralic JE, Morrow AL, Korpi ER, Hardy S, Sloat B, Eger EI 2nd, Homanics GE. Alpha 1 subunit-containing GABA type A receptors in forebrain contribute to the effect of inhaled anesthetics on conditioned fear. *Mol Pharmacol* 2005;68:61–68. [PubMed: 15833735]
114. Quinlan JJ, Homanics GE, Firestone LL. Anesthesia sensitivity in mice that lack the β 3 subunit of the γ -aminobutyric acid type A receptor. *Anesthesiology* 1998;88:775–780. [PubMed: 9523823]
115. Eger EI II, Ionescu P, Laster MJ, Fang Z, Gong D, Hudlicky T, Kendig J, Harris A, Trudell J, Pohorille A. MAC of fluorinated alkanols in rats: Relevance to theories of narcosis. *Anesth Analg* 1999;88:867–876. [PubMed: 10195540]
116. Ueno S, Trudell JR, Eger EI II, Harris RA. Actions of fluorinated alkanols on GABAA receptors: Relevance to theories of narcosis. *Anesth Analg* 1999;88:877–883. [PubMed: 10195541]
117. Jones MV, Brooks PA, Harrison NL. Enhancement of gamma-aminobutyric acid-activated Cl⁻ currents in cultured rat hippocampal neurones by three volatile anaesthetics. *J Physiol (Lond)* 1992;449:279–293. [PubMed: 1326046]
118. Wentlandt K, Samoilova M, Carlen PL, El Beheiry H. General anesthetics inhibit gap junction communication in cultured organotypic hippocampal slices. *Anesth Analg* 2006;102:1692–1698. [PubMed: 16717311]
119. Johnston MF, Simon SA, Ramon F. Interaction of anaesthetics with electrical synapses. *Nature* 1980;286:498–500. [PubMed: 6250068]
120. Mascia MP, Machu TK, Harris RA. Enhancement of homomeric glycine receptor function by long-chain alcohols and anaesthetics. *Br J Pharmacol* 1996;119:1331–1336. [PubMed: 8968539]
121. Zimmerman SA, Jones MV, Harrison NL. Potentiation of γ -aminobutyric acid A receptor Cl⁻ current correlates with in vivo anesthetic potency. *J Pharmacol Exp Ther* 1994;270:987–991. [PubMed: 7932211]

122. Zhang Y, Wu S, Eger EI II, Sonner JM. Neither GABA(A) nor strychnine-sensitive glycine receptors are the sole mediators of MAC for isoflurane. *Anesth Analg* 2001;92:123–127. [PubMed: 11133613]
123. Zhang Y, Laster MJ, Hara K, Harris RA, Eger EI II, Stabernack CR, Sonner JM. Glycine receptors may mediate part of the immobility produced by inhaled anesthetics. *Anesth Analg* 2003;96:97–101. [PubMed: 12505932]
124. Kingsmore S, Giros B, Suh D, Bieniarz M, Caron MG, Seldin MF. Glycine receptor beta-subunit gene mutation in spastic mouse associated with LINE-1 element insertion. *Nat Genet* 1994;7:136–141. [PubMed: 7920630]
125. Quinlan JJ, Ferguson C, Jester K, Firestone LL, Homanics GE. Mice with glycine receptor subunit mutations are both sensitive and resistant to volatile anesthetics. *Anesth Analg* 2002;95:578–582. [PubMed: 12198041]
126. Ryan SG, Buckwalter MS, Lynch JW, Handford CA, Segura L, Shiang R, Wasmuth JJ, Camper SA, Schofield P, O'Connell P. A missense mutation in the gene encoding the alpha 1 subunit of the inhibitory glycine receptor in the spasmodic mouse. *Nat Genet* 1994;7:131–135. [PubMed: 7920629]
127. McFarlane C, Warner DS, Todd MM, Nordholm L. AMPA receptor competitive antagonism reduces halothane MAC in rats. *Anesthesiology* 1992;77:1165–1170. [PubMed: 1334638]
128. Dutton RC, Zhang Y, Stabernack CR, Laster MJ, Sonner JM, Eger EI II. Temporal summation governs part of the minimum alveolar concentration (MAC) of isoflurane anesthesia. *Anesthesiology* 2003;98:1372–1377. [PubMed: 12766645]
129. Stabernack C, Sonner JM, Laster M, Zhang Y, Xing Y, Sharma M, Eger EI II. Spinal NMDA receptors may contribute to the immobilizing action of isoflurane. *Anesth Analg* 2003;96:102–107. [PubMed: 12505933]
130. McFarlane C, Warner DS, Dexter F. Interactions between NMDA and AMPA glutamate receptor antagonists during halothane anesthesia in the rat. *Neuropharmacology* 1995;34:659–663. [PubMed: 7566502]
131. Cheng G, Kendig JJ. Enflurane directly depresses glutamate AMPA and NMDA currents in mouse spinal cord motor neurons independent of actions on GABAA or glycine receptors. *Anesthesiology* 2000;93:1075–1084. [PubMed: 11020764]
132. Perouansky M, Kirson ED, Yaari Y. Mechanism of action of volatile anesthetics: effects of halothane on glutamate receptors in vitro. *Toxicol Lett* 1998;100–101:65–69. [PubMed: 10049182]
133. Yamakura T, Harris RA. Effects of gaseous anesthetics nitrous oxide and xenon on ligand-gated ion channels: Comparison with isoflurane and ethanol. *Anesthesiology* 2000;93:1095–1101. [PubMed: 11020766]
134. Carla V, Moroni F. General anaesthetics inhibit the responses induced by glutamate receptor agonists in the mouse cortex. *Neurosci Lett* 1992;146:21–24. [PubMed: 1282227]
135. Joo DT, Gong D, Sonner JM, Jia Z, MacDonald JF, Eger EI II, Orser BA. Blockade of AMPA receptors and volatile anesthetics: reduced anesthetic requirements in GluR2 null mutant mice for loss of the righting reflex and antinociception but not minimum alveolar concentration. *Anesthesiology* 2001;94:478–488. [PubMed: 11374610]
136. Nishikawa K, MacIver MB. Excitatory synaptic transmission mediated by NMDA receptors is more sensitive to isoflurane than are non-NMDA receptor-mediated responses. *Anesthesiology* 2000;92:228–236. [PubMed: 10638920]
137. Bar-Peled O, O'Brien RJ, Morrison JH, Rothstein JD. Cultured motor neurons possess calcium-permeable AMPA/kainate receptors. *Neuroreport* 1999;10:855–859. [PubMed: 10208560]
138. Van Damme P, Van Den Bosch L, Van Houtte E, Callewaert G, Robberecht W. GluR2-dependent properties of AMPA receptors determine the selective vulnerability of motor neurons to excitotoxicity. *J Neurophysiology* 2002;88:1279–1287.
139. Petralia RS, Wang YX, Mayat E, Wenthold RJ. Glutamate receptor subunit 2-selective antibody shows a differential distribution of calcium-impermeable AMPA receptors among populations of neurons. *J Comp Neurol* 1997;385:456–476. [PubMed: 9300771]

140. Ishizaki K, Sasaki M, Karasawa S, Obata H, Nara T, Goto F. Intrathecal co-administration of NMDA antagonist and NK-1 antagonist reduces MAC of isoflurane in rats. *Acta Anaesthesiol Scand* 1999;43:753–759. [PubMed: 10456816]
141. Dingledine R, Borges K, Bowie D, Traynelis SF. The glutamate receptor ion channels. *Pharmacol Rev* 1999;51:7–61. [PubMed: 10049997]
142. Minami K, Wick MJ, Stern-Bach Y, Dildy-Mayfield JE, Brozowski SJ, Gonzales EL, Trudell JR, Harris RA. Sites of volatile anesthetic action on kainate (Glutamate receptor 6) receptors. *J Biol Chem* 1998;273:8248–8255. [PubMed: 9525931]
143. Sonner JM, Vissel B, Royle G, Maurer A, Gong D, Baron NV, Harrison N, Fanselow M, Eger EI 2nd. The effect of three inhaled anesthetics in mice harboring mutations in the GluR6 (kainate) receptor gene. *Anesth Analg* 2005;101:143–148. [PubMed: 15976221]
144. Ishizaki K, Yoon DM, Yoshida N, Tamazaki M, Arai K, Fujita T. Intrathecal administration of N-methyl-D-aspartate receptor antagonist reduces the minimum alveolar anaesthetic concentration of isoflurane in rats. *Br J Anaesth* 1995;75:636–638. [PubMed: 7577295]
145. Kuroda Y, Strebel S, Rafferty C, Bullock R. Neuroprotective doses of N-methyl-D-aspartate receptor antagonists profoundly reduce the minimum alveolar anesthetic concentration (MAC) for isoflurane in rats. *Anesth Analg* 1993;77:795–800. [PubMed: 8214668]
146. Flood P, Krasowski MD. Intravenous anesthetics differentially modulate ligand-gated ion channels. *Anesthesiology* 2000;92:1418–1425. [PubMed: 10781289]
147. Sato Y, Kobayashi E, Murayama T, Mishina M, Seo N. Effect of N-methyl-D-aspartate receptor epsilon1 subunit gene disruption of the action of general anesthetic drugs in mice. *Anesthesiology* 2005;102:557–561. [PubMed: 15731593]
148. Sato Y, Seo N, Kobayashi E. Ethanol-induced hypnotic tolerance is absent in N-methyl-D-aspartate receptor epsilon 1 subunit knockout mice. *Anesth Analg* 2006;103:117–120. [PubMed: 16790637]
149. Niemann S, Kanki H, Fukui Y, Takao K, Fukaya M, Hynynen MN, Churchill MJ, Shefner JM, Bronson RT, Brown RH Jr, Watanabe M, Miyakawa T, Itoharu S, Hayashi Y. Genetic ablation of NMDA receptor subunit NR3B in mouse reveals motoneuronal and nonmotoneuronal phenotypes. *Eur J Neurosci* 2007;26:1407–1420. [PubMed: 17880385]
150. Liu HT, Hollmann MW, Liu WH, Hoenemann CW, Durieux ME. Modulation of NMDA receptor function by ketamine and magnesium: Part I. *Anesth Analg* 2001;92:1173–1181. [PubMed: 11323343]
151. Yamakura T, Askalany AR, Petrenko AB, Kohno T, Baba H, Sakimura K. The NR3B subunit does not alter the anesthetic sensitivities of recombinant N-methyl-D-aspartate receptors. *Anesth Analg* 2005;100:1687–1692. [PubMed: 15920196]
152. de Sousa LM, Dickinson R, Lieb WR, Franks NP. Contrasting synaptic actions of the inhalational general anesthetics isoflurane and xenon. *Anesthesiology* 2000;92:1055–1066. [PubMed: 10754626]
153. Rex S, Schaefer W, Meyer PH, Rossaint R, Boy C, Setani K, Bull U, Baumert JH. Positron Emission Tomography Study of Regional Cerebral Metabolism during General Anesthesia with Xenon in Humans. *Anesthesiology* 2006;105:936–943. [PubMed: 17065887]
154. Langsjo JW, Salmi E, Kaisti KK, Aalto S, Hinkka S, Aantaa R, Oikonen V, Viljanen T, Kurki T, Silvanto M, Scheinin H. Effects of subanesthetic ketamine on regional cerebral glucose metabolism in humans. *Anesthesiology* 2004;100:1065–1071. [PubMed: 15114201]
155. Kaisti KK, Langsjo JW, Aalto S, Oikonen V, Sipila H, Teras M, Hinkka S, Metsahonkala L, Scheinin H. Effects of sevoflurane, propofol, and adjunct nitrous oxide on regional cerebral blood flow, oxygen consumption, and blood volume in humans. *Anesthesiology* 2003;99:603–613. [PubMed: 12960544]
156. Dutton RC, Laster MJ, Xing Y, Sonner JM, Raines DE, Solt K, Eger EI 2nd. Do N-methyl-D-aspartate receptors mediate the capacity of inhaled anesthetics to suppress the temporal summation that contributes to minimum alveolar concentration? *Anesth Analg* 2006;102:1412–1418. [PubMed: 16632819]
157. Dickenson AH, Sullivan AF. Electrophysiologic studies on the effects of intrathecal morphine on nociceptive neurons in the rat dorsal horn. *Pain* 1986;24:211–222. [PubMed: 3754322]

158. Eger EI II, Liao M, Laster MJ, Won A, Popovich J, Raines DE, Solt K, Dutton RC, Cobos FV, Sonner JM. Contrasting roles of the N-methyl-D-aspartate receptor in the production of immobilization by conventional and aromatic anesthetics. *Anesth Analg* 2006;102:1397–1406. [PubMed: 16632817]
159. Solt K, Eger EI II, Raines DE. Differential modulation of human n-methyl-D-aspartate receptors by structurally diverse general anesthetics. *Anesth Analg* 2006;102:1407–1411. [PubMed: 16632818]
160. Engelhardt T, Lowe PR, Galley HF, Webster NR. Inhibition of neuronal nitric oxide synthase reduces isoflurane MAC and motor activity even in nNOS knockout mice. *Br J Anaesth* 2006;96:361–366. [PubMed: 16431879]
161. Johns RA, Moscicki JC, DiFazio CA. Nitric oxide synthase inhibitor dose-dependently and reversibly reduces the threshold for halothane anesthesia. A role for nitric oxide in mediating consciousness? *Anesthesiology* 1992;77:779–784. [PubMed: 1384399]
162. Pajewski TN, DiFazio CA, Moscicki JC, Johns RA. Nitric oxide synthase inhibitors, 7-nitro indazole and nitroG-L-arginine methyl ester, dose dependently reduce the threshold for isoflurane anesthesia. *Anesthesiology* 1996;85:1111–1119. [PubMed: 8916829]
163. Ichinose F, Huang PL, Zapol WM. Effects of targeted neuronal nitric oxide synthase gene disruption and nitroG-L-arginine methylester on the threshold for isoflurane anesthesia. *Anesthesiology* 1995;83:101–108. [PubMed: 7541615]
164. Ichinose F, Mi WD, Miyazaki M, Onouchi T, Goto T, Morita S. Lack of correlation between the reduction of sevoflurane MAC and the cerebellar cyclic GMP concentrations in mice treated with 7-nitroindazole. *Anesthesiology* 1998;89:143–148. [PubMed: 9667304]
165. Kobayashi S, Katoh T, Iwamoto T, Bito H, Sato S. Effect of the neuronal nitric oxide synthase inhibitor 7-nitroindazole on the righting reflex ED50 and minimum alveolar concentration during sevoflurane anaesthesia in rats. *Eur J Anaesthesiol* 2003;20:212–219. [PubMed: 12650492]
166. Adachi T, Kurata J, Nakao S, Murakawa M, Shichino T, Shirakami G, Shinomura T, Mori K. Nitric oxide synthase inhibitor does not reduce minimum alveolar anesthetic concentration of halothane in rats. *Anesth Analg* 1994;78:1154–1157. [PubMed: 7515216]
167. Adachi T, Shinomura T, Nakao S, Kurata J, Murakawa M, Shichino T, Seo N, Mori K. Chronic treatment with nitric oxide synthase (NOS) inhibitor profoundly reduces cerebellar NOS activity and cyclic guanosine monophosphate but does not modify minimum alveolar anesthetic concentration. *Anesth Analg* 1995;81:862–865. [PubMed: 7574024]
168. Way WL, Hosobuchi Y, Johnson BH, Eger EI II. Anesthesia does not increase opioid peptides in cerebrospinal fluid of humans. *Anesthesiology* 1984;60:43–45. [PubMed: 6318607]
169. Katoh T, Ikeda K. The effects of fentanyl on sevoflurane requirements for loss of consciousness and skin incision. *Anesthesiology* 1998;88:18–24. [PubMed: 9447851]
170. Harper MH, Winter PM, Johnson BH, Eger EI II. Naloxone does not antagonize general anesthesia in the rat. *Anesthesiology* 1978;49:3–5. [PubMed: 666036]
171. Pace NL, Wong KC. Failure of naloxone and naltrexone to antagonize halothane anesthesia in the dog. *Anesth Analg* 1979;58:36–39. [PubMed: 219728]
172. Levine LL, Winter PM, Nemoto EM, Uram M, Lin MR. Naloxone does not antagonize the analgesic effects of inhalation anesthetics. *Anesth Analg* 1986;65:330–332. [PubMed: 3006550]
173. de Wolff MH, Leather HA, Wouters PF. Effects of tramadol on minimum alveolar concentration (MAC) of isoflurane in rats. *Br J Anaesth* 1999;83:780–783. [PubMed: 10690143]
174. Mantz J, Azerad J, Limoge A, Desmots JM. Transcranial electrical stimulation with Limoge's currents decreases halothane requirements in rats. Evidence for the involvement of endogenous opioids. *Anesthesiology* 1992;76:253–260. [PubMed: 1736702]
175. Tay AA, Tseng CK, Pace NL, Wong KC, Hong-Goka BC. Failure of narcotic antagonist to alter electroacupuncture modification of halothane anaesthesia in the dog. *Can Anaesth Soc J* 1982;29:231–235. [PubMed: 6978757]
176. Maze M, Vickery R, Merlone S, Gaba D. Anesthetic and hemodynamic effects of the alpha2-adrenergic agonist, azepexole, in isoflurane-anesthetized dogs. *Anesthesiology* 1988;68:689–694. [PubMed: 2897174]
177. Michelsen LG, Salmenpera M, Hug CC Jr, Szlam F, VanderMeer D. Anesthetic potency of remifentanyl in dogs. *Anesthesiology* 1996;84:865–872. [PubMed: 8638841]

178. Dahan A, Sarton E, Teppema L, Olievier C, Nieuwenhuijs D, Matthes HW, Kieffer BL. Anesthetic potency and influence of morphine and sevoflurane on respiration in muopioid receptor knockout mice. *Anesthesiology* 2001;94:824–832. [PubMed: 11388534]
179. Liao M, Laster MJ, Eger EI 2nd, Tang M, Sonner JM. Naloxone does not increase the minimum alveolar anesthetic concentration of sevoflurane in mice. *Anesth Analg* 2006;102:1452–1455. [PubMed: 16632825]
180. Heurteaux C, Guy N, Laigle C, Blondeau N, Duprat F, Mazzuca M, Lang-Lazdunski L, Widmann C, Zanzouri M, Romey G, Lazdunski M. TREK-1, a K(+) channel involved in neuroprotection and general anesthesia. *EMBO J* 2004;23:2684–2695. [PubMed: 15175651]
181. Patel AJ, Honore E, Lesage F, Fink M, Romey G, Lazdunski M. Inhalational anesthetics activate two-pore-domain background K⁺ channels. *Nat Neurosci* 1999;2:422–426. [PubMed: 10321245]
182. Eger EI II, Saidman LJ, Brandstater B. Temperature dependence of halothane and cyclopropane anesthesia in dogs: Correlation with some theories of anesthetic action. *Anesthesiology* 1965;26:764–770. [PubMed: 5844268]
183. Antognini JG. Hypothermia eliminates isoflurane requirements at 20 degrees C. *Anesthesiology* 1993;78:1152–1156. [PubMed: 8512109]
184. Liu M, Hu X, Liu J. The effect of hypothermia on isoflurane MAC in children. *Anesthesiology* 2001;94:429–432. [PubMed: 11374601]
185. Linden AM, Sandu C, Aller MI, Vekovischeva OY, Rosenberg PH, Wisden W, Korpi ER. TASK-3 knockout mice exhibit exaggerated nocturnal activity, impairments in cognitive functions, and reduced sensitivity to inhalation anesthetics. *J Pharmacol Exp Ther* 2007;323:924–934. [PubMed: 17875609]
186. Leonoudakis D, Gray AT, Winegar BD, Kindler CH, Harada M, Taylor DM, Chavez RA, Forsayeth JR, Yost CS. An open rectifier potassium channel with two pore domains in tandem cloned from rat cerebellum. *J Neurosci* 1998;18:868–877. [PubMed: 9437008]
187. Brosnan RJ, Eger EI II, Laster MJ, Sonner JM. Anesthetic properties of carbon dioxide in the rat. *Anesth Analg* 2007;105:103–106. [PubMed: 17578964]
188. Gerstin KM, Gong DH, Abdallah M, Winegar BD, Eger EI II, Gray AT. Mutation of KCNK5 or Kir3.2 potassium channels in mice does not change MAC. *Anesth Analg* 2003;96:1345–1349. [PubMed: 12707131]
189. Zucker JR. ATP-sensitive potassium channel agonists do not alter MAC for isoflurane in rats. *Anesthesiology* 1992;76:560–563. [PubMed: 1550281]
190. Cotten JF, Keshavaprasad B, Laster MJ, Eger EI II, Yost CS. The ventilatory stimulant doxapram inhibits TASK tandem pore (K_{2p}) potassium channel function but does not affect minimum alveolar anesthetic concentration. *Anesth Analg* 2006;102:779–785. [PubMed: 16492828]
191. Xing Y, Zhang Y, Stabernack CR, Eger EI II, Gray AT. The use of the potassium channel activator riluzole to test whether potassium channels mediate the capacity of isoflurane to produce immobility. *Anesth Analg* 2003;97:1020–1024. [PubMed: 14500151]
192. Urbani A, Belluzzi O. Riluzole inhibits the persistent sodium current in mammalian CNS neurons. *Eur J Neurosci* 2000;12:3567–3574. [PubMed: 11029626]
193. Kuo JJ, Lee RH, Zhang L, Heckman CJ. Essential role of the persistent sodium current in spike initiation during slowly rising inputs. *J Physiol*. 2006
194. Tanifuji Y, Eger EI II. Brain sodium, potassium, and osmolality: Effects on anesthetic requirement. *Anesth Analg* 1978;57:404–410. [PubMed: 568400]
195. Shnayderman D, Laster MJ, Eger EI II, Oh I, Zhang Y, Jinks SL, Antognini JF, Raines DE. Increases in spinal cerebrospinal fluid potassium concentration do not increase isoflurane MAC in rats. *Anesth Analg* 2008;107In Press THIS ISSUE
196. Vonvoigtlander P, Lewis RA, Neff GL. Kappa opioid analgesia is dependent on serotonergic mechanisms. *J Pharmacol Exp Ther* 1984;231:270–274. [PubMed: 6092612]
197. Kellstein DE, Malseed RT, Goldstein FJ. Opioid-monomine interactions in spinal antinociception: Evidence for serotonin but not norepinephrine reciprocity. *Pain* 1988;34:85–92. [PubMed: 2841636]
198. Paul D, Mana MJ, Pfaus JG, Pinel JP. Attenuation of morphine analgesia by the S2 antagonists, pirenperone and ketanserin. *Pharmacol Biochem Behav* 1988;31:641–647. [PubMed: 3251248]

199. Kiefel JM, Paul D, Bodnar RJ. Reduction in opioid and non-opioid forms of swim analgesia by 5-HT₂ receptor antagonists. *Brain Res* 1989;500:231–240. [PubMed: 2605492]
200. Giordano J. Analgesic profile of centrally administered 2-methylserotonin against acute pain in rats. *Eur J Pharmacol* 1991;199:233–236. [PubMed: 1954980]
201. Coimbra NC, Brandao ML. Effects of 5-HT₂ receptors blockade on fear-induced analgesia elicited by electrical stimulation of the deep layers of the superior colliculus and dorsal periaqueductal gray. *Behav Brain Res* 1997;87:97–103. [PubMed: 9331477]
202. Yamazake N, Umeno H, Kuraishi Y. Involvement of brain serotonergic terminals in the antinociceptive action of peripherally applied calcitonin. *Jpn J Pharmacol* 1999;81:367–374. [PubMed: 10669042]
203. Bardin L, Lavarenne J, Eschaliere A. Serotonin receptor subtypes involved in the spinal antinociceptive effect of 5-HT in rats. *Pain* 2000;86:11–18. [PubMed: 10779655]
204. Hentall ID, Andresen MJ, Taguchi K. Serotonergic, cholinergic and nociceptive inhibition or excitation of raphe magnus neurons in barbiturate-anesthetized rats. *Neuroscience* 1993;52:303–310. [PubMed: 8450948]
205. Zhang L. Effects of 5-hydroxytryptamine on cat spinal motoneurons. *Can J Physiol Pharmacol* 1991;69:154–163. [PubMed: 2054730]
206. Dringenberg HC. Serotonergic receptor antagonists alter responses to general anaesthetics in rats. *Br J Anaesth* 2000;85:904–906. [PubMed: 11732529]
207. Doherty TJ, McDonnell WN, Dyson DH, Black WD, Valliant AE. The effect of a 5-hydroxytryptamine antagonist (R51703) on halothane MAC in the dog. *J Vet Pharmacol Ther* 1995;18:153–155. [PubMed: 7629931]
208. Zhang Y, Laster MJ, Eger EI II, Stabernack CR, Sonner JM. Blockade of 5-HT_{2A} receptors may mediate or modulate part of the immobility produced by inhaled anesthetics. *Anesth Analg* 2003;97:475–479. [PubMed: 12873938]
209. Johnston RR, Way WL, Miller RD. The effect of CNS catecholamine-depleting drugs on dextroamphetamine-induced elevation of halothane MAC. *Anesthesiology* 1974;41:57–61. [PubMed: 4275974]
210. Rampil IJ, Laster MJ, Eger EII. Ondansetron does not alter isoflurane MAC in rats. *Anesthesiology* 2001;95:562–564. [PubMed: 11506137]
211. Eger EI II. Dragons and other scientific hazards. *Anesthesiology* 1979;50:1. [PubMed: 104640]
212. Fernandez FR, Mehaffey WH, Turner RW. Dendritic Na⁺ current inactivation can increase cell excitability by delaying a somatic depolarizing afterpotential. *J Neurophysiol* 2005;94:3836–3848. [PubMed: 16120659]
213. Eckenhoff RG, Johansson JS. On the relevance of "clinically relevant concentrations" of inhaled anesthetics in *in vitro* experiments. *Anesthesiology* 1999;91:856–860. [PubMed: 10485798]
214. Zhang Y, Laster MJ, Eger EI II, Sharma M, Sonner JM. Lidocaine, MK-801, and MAC. *Anesth Analg* 2007;104:1098–1102. [PubMed: 17456658]
215. Linde HW, Lamb VE, Quimby CW Jr, Homi J, Eckenhoff JE. The search for better anesthetic agents: clinical investigation of ethrane. *Anesthesiology* 1970;32:555–559. [PubMed: 4913148]
216. Jaaskelainen SK, Kaisti K, Suni L, Hinkka S, Scheinin H. Sevoflurane is epileptogenic in healthy subjects at surgical levels of anesthesia. *Neurology* 2003;61:1073–1078. [PubMed: 14581667]
217. Liu PL, Feldman HS, Giasi R, Patterson MK, Covino BG. Comparative CNS toxicity of lidocaine, etidocaine, bupivacaine, and tetracaine in awake dogs following rapid intravenous administration. *Anesth Analg* 1983;62:375–379. [PubMed: 6829942]
218. Zhang Y, Sharma M, Eger EI II, Laster MJ, Hemmings HC Jr, Harris RA. Intrathecal veratridine administration increases MAC in rats. *Anesth Analg* 2008;107:In press THIS ISSUE
219. Rehberg B, Xiao YH, Duch DS. Central nervous system sodium channels are significantly suppressed at clinical concentrations of volatile anesthetics. *Anesthesiology* 1996;84:1223–1233. [PubMed: 8624017]
220. Stadnicka A, Kwok WM, Hartmann HA, Bosnjak ZJ. Effects of halothane and isoflurane on fast and slow inactivation of human heart hH1a sodium channels. *Anesthesiology* 1999;90:1671–1683. [PubMed: 10360866]

221. Shiraishi M, Harris RA. Effects of alcohols and anesthetics on recombinant voltage-gated Na⁺ channels. *J Pharmacol Exp Ther* 2004;309:987–994. [PubMed: 14978193]
222. OuYang W, Hemmings HC Jr. Isoform-selective effects of isoflurane on voltage-gated Na⁺ channels. *Anesthesiology* 2007;107:91–98. [PubMed: 17585220]
223. Schlame M, Hemmings HC Jr. Inhibition by volatile anesthetics of endogenous glutamate release from synaptosomes by a presynaptic mechanism. *Anesthesiology* 1995;82:1406–1416. [PubMed: 7793654]
224. Ouyang W, Wang G, Hemmings HC Jr. Isoflurane and propofol inhibit voltage-gated sodium channels in isolated rat neurohypophysial nerve terminals. *Mol Pharmacol* 2003;64:373–381. [PubMed: 12869642]
225. Ratnakumari L, Vysotskaya TN, Duch DS, Hemmings HC Jr. Differential effects of anesthetic and nonanesthetic cyclobutanes on neuronal voltage-gated sodium channels. *Anesthesiology* 2000;92:529–541. [PubMed: 10691242]
226. Kendig JJ. Barbiturates: active form and site of action at node of Ranvier sodium channels. *J Pharmacol Exp Ther* 1981;218:175–181. [PubMed: 6264067]
227. Mohr JT, Gribble GW, Lin SS, Eckenhoff RG, Cantor RS. Anesthetic potency of two novel synthetic polyhydric alkanols longer than the n-alkanol cutoff: evidence for a bilayer-mediated mechanism of anesthesia? *J Med Chem* 2005;48:4172–4176. [PubMed: 15943489]
228. Curry S, Moss GWJ, Dickinson R, Lieb WR, Franks NP. Probing the molecular dimensions of general anaesthetic target sites in tadpoles (*Xenopus laevis*) and model systems using cycloalcohols. *Br J Pharmacol* 1991;102:167–173. [PubMed: 2043920]
229. Franks N, Lieb W. Partitioning of long-chain alcohols into lipid bilayers: Implications for mechanisms of general anesthesia. *Proc Natl Acad Sci USA* 1986;83:5116–5120. [PubMed: 3460084]
230. Cantor RS. The lateral pressure profile in membranes: A physical mechanism of general anesthesia. *Biochemistry* 1997;36:2339–2344. [PubMed: 9054538]
231. Roth R, Gillespie D, Nonner W, Eisenberg B. Bubbles, Gating, and Anesthetics in Ion Channels. *Biophys J*. 2008




Figure 1.

This graph indicates the fraction of MAC for each anesthetic of a pair that, in combination, produces immobility in response to noxious stimulation in rats. Each anesthetic pair was usually chosen because the two anesthetics differed in their capacities to inhibit or enhance the response of a specific channel (e.g., cyclopropane minimally enhances the response of gamma-aminobutyric acid (GABA)_A receptors to GABA, whereas halothane markedly enhances the response). The sum of the fractional contributions never was less than 0.9, an a priori value assigned as the boundary between additivity and synergism (i.e., no pair acted synergistically). The data are taken from Eger et al.¹⁶




Figure 2.

In rats, MAC for various anesthetics was determined in the presence of intrathecal infusions of strychnine [a glycine receptor blocker;¹²³ left panel] or picrotoxin [a gamma-aminobutyric acid (GABA)_A receptor blocker;¹¹² right panel]. For infusions that produced maximal increases in MAC, the increases correlated with the capacity of particular anesthetics to enhance the response of glycine receptors *in vitro* but did not correlate with the capacity of particular anesthetics to enhance the response of GABA_A receptors *in vitro* (the *in vitro* capacity is indicated for each abscissa). Such results are consistent with a role for glycine receptors as the mediators of the anesthetic effect of some (e.g., halothane – see open circles) but not other (e.g., cyclopropane – see open triangles) anesthetics. Data for isoflurane are shown

as closed diamonds, and data for isoflurane are shown as open circles. Such results also are not consistent with a role for GABA_A receptors as mediators of immobility.




Figure 3.

TREK-1 knockout mice have greater MAC values than their wild-type littermates. The upper graphs indicate the percentage increase in MAC that attends knockout.¹⁸⁰ The increases vary by nearly an order of magnitude (e.g., 7% for desflurane, 15% for sevoflurane, and 47% for halothane), but do not correlate with the capacity of these anesthetics to enhance the response of the TREK-1 channel *in vitro* (bottom graph).^{180,181} The failure of the correlation calls into question the relevance of TREK-1 channels as mediators of the immobility produced by inhaled anesthetics.