

Integrin $\alpha 3\beta 1$ -dependent β -catenin phosphorylation links epithelial Smad signaling to cell contacts

Young Kim, Matthias C. Kugler, Ying Wei, Kevin K. Kim, Xiaopeng Li, Alexis N. Brumwell, and Harold A. Chapman

Pulmonary and Critical Care Division, Department of Medicine, and Cardiovascular Research Institute, University of California, San Francisco, San Francisco, CA 94143

Injury-initiated epithelial to mesenchymal transition (EMT) depends on contextual signals from the extracellular matrix, suggesting a role for integrin signaling. Primary epithelial cells deficient in their prominent laminin receptor, $\alpha 3\beta 1$, were found to have a markedly blunted EMT response to TGF- $\beta 1$. A mechanism for this defect was explored in $\alpha 3$ -null cells reconstituted with wild-type (wt) $\alpha 3$ or point mutants unable to engage laminin 5 (G163A) or epithelial cadherin (E-cadherin; H245A). After TGF- $\beta 1$ stimulation, wt epithelial cells but not cells expressing the H245A mutant internalize complexes of

E-cadherin and TGF- $\beta 1$ receptors, generate phospho-Smad2 (p-Smad2)-pY654- β -catenin complexes, and up-regulate mesenchymal target genes. Although Smad2 phosphorylation is normal, p-Smad2-pY654- β -catenin complexes do not form in the absence of $\alpha 3$ or when $\alpha 3\beta 1$ is mainly engaged on laminin 5 or E-cadherin on adherens junctions, leading to attenuated EMT. These findings demonstrate that $\alpha 3\beta 1$ coordinates cross talk between β -catenin and Smad signaling pathways as a function of extracellular contact cues and thereby regulates responses to TGF- $\beta 1$ activation.

Introduction

Integrins, a group of heterodimeric surface molecules, provide a dynamic interface between the cell and ECM by regulating inside-out and outside-in signaling during embryonic development, wound repair, and tissue morphogenesis (Geiger et al., 2001; Hynes, 2002). Epithelial integrins play an important role in the process of differentiation and homeostasis of skin, kidney, and lung lining layers (Kreidberg et al., 1996). One integrin highly expressed in epithelial cells, $\alpha 3\beta 1$, mediates cell-matrix interactions with its major ligand laminin 5 (Ln5; Eble et al., 1998) and localizes to focal adhesion sites (Dogic et al., 1998). In addition, $\alpha 3\beta 1$ is also found along lateral cell borders in association with epithelial cadherin (E-cadherin) in adherens junctions (Nakamura et al., 1995; Chattopadhyay et al., 2003), where its colocalization is thought to contribute to assembly and maintenance of cell-cell contacts and normal barrier function (Wang et al., 1999; Lubman et al., 2000). Whether $\alpha 3\beta 1$ has a structural role in adherens junction maintenance or acts in some way to sense perturbations in cell-cell contacts is unknown.

Under certain conditions, epithelial cells undergo the process of epithelial to mesenchymal transition (EMT). EMT involves loss of epithelial characteristics by down-regulation of proteins such as E-cadherin and a shift toward a fibroblast-like phenotype with expression of mesenchymal proteins such as α smooth muscle actin (α -SMA), matrix metalloproteinases (MMPs), and enhanced motility (Savagner, 2001; Kalluri and Neilson, 2003). The most well-studied cytokine as an inducer of EMT is TGF- $\beta 1$ (Gotzmann et al., 2004; Zeisberg and Kalluri, 2004). TGF- $\beta 1$ binds to its receptor, through which it activates intracellular Smad-dependent and -independent signaling pathways and regulates gene transcription and cell shape (Derynck and Zhang, 2003).

TGF- $\beta 1$ signaling is regulated at multiple levels beginning with activation of latent TGF- $\beta 1$ itself, determinants of phosphorylation of receptor-associated Smads (i.e., Smads 2 and 3) by the TGF- $\beta 1$ receptor, and various factors that influence stability and translocation of phospho-Smads (p-Smads) to the nucleus. Smad-dependent transcription is further dependent on coactivators that stabilize Smad association with DNA, and these coregulators are both constitutive, such as p300, and cell

Y. Kim, M.C. Kugler, and Y. Wei contributed equally to this paper.

Correspondence to Harold A. Chapman: hal.chapman@ucsf.edu

Abbreviations used in this paper: α -SMA, α smooth muscle actin; AEC, alveolar epithelial cell; E-cadherin, epithelial cadherin; EMT, epithelial to mesenchymal transition; ERK, extracellular signal-regulated kinase; Fn, fibronectin; IP, immunoprecipitation; MDC, monodansylcadaverine; MMP, matrix metalloproteinase; p-Smad, phospho-Smad; shRNA, short hairpin RNA; wt, wild type.

© 2009 Kim et al. This article is distributed under the terms of an Attribution-Noncommercial-Share Alike-No Mirror Sites license for the first six months after the publication date [see <http://www.jcb.org/misc/terms.shtml>]. After six months it is available under a Creative Commons License [Attribution-Noncommercial-Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>].

type specific. As a consequence of this complexity, multiple signaling molecules are known to modulate TGF- β 1 signaling during EMT and tumorigenesis, including bone morphogenetic proteins and their receptors as well as Ras, MAPK, and various cytokines (Massague and Chen, 2000).

There is circumstantial evidence that integrins via their engagement with the ECM play a substantial role in TGF- β 1 activation and signaling. Integrin α V β 6 and binding to the ECM molecule fibronectin (Fn) is required for activation of latent TGF- β 1 (Fontana et al., 2005; Sheppard, 2005). Global inhibition of β 1-integrin matrix binding by blocking antibodies is reported to abrogate active TGF- β 1-mediated p38 MAPK activation and EMT in mammary cells (Bhowmick et al., 2001). Attachment of epithelial cells to various ECMs is known in some way to promote responses to active TGF- β 1 (Thannickal et al., 2003). Several prior studies implicate the β 1 integrin-associated protein, integrin-linked kinase in EMT, also implying an important role for β 1 integrins in the process (Lee et al., 2004; Oloumi et al., 2004). However, to date, there is no specific integrin α chain associated with β 1 that is known to be critically involved in EMT, and the molecular mechanisms by which ECM regulates EMT are undefined.

The impetus for this study was the observation that an immortalized α 3-null kidney epithelial cell line failed to up-regulate the expected mesenchymal markers after TGF- β 1 stimulation. We extended this observation by finding defective TGF- β 1 responses in primary lung epithelial cells selectively missing α 3 β 1. We have used the α 3-null cell line to explore the mechanisms by which α 3 β 1 influences cellular responses to TGF- β 1. These experiments reveal not only an unrecognized role for α 3 β 1 in EMT but also a molecular mechanism that can explain how cell contacts critically influence responses to TGF- β 1 signaling.

Results

Primary lung alveolar epithelial cells (AECs) deprived of α 3 β 1 show an impaired response to TGF- β 1

Primary AECs isolated from a conditional α 3-null mouse were exposed to adenovirus-Cre for 96 h to abrogate integrin α 3 expression. As a control, cells were infected in parallel with a GFP-expressing adenovirus. Cells were maintained on Matrigel/collagen because on this matrix the cells activate little TGF- β 1 and maintain an epithelial phenotype (Kim et al., 2006). At the end of 96 h, the primary cells were harvested and transferred to Fn-coated surfaces for an additional 4 d to allow for TGF- β 1 activation and initiation of EMT, as previously reported (Kim et al., 2006). This point is illustrated in Fig. 1 A (left), showing phosphorylation of Smad2 after 4 d on Fn and induction of α -SMA. Inhibition of TGF- β 1 receptor kinase activity with SB431542 blocked p-Smad2 formation and also attenuated up-regulation of α -SMA. Immunoblotting indicated a near-complete absence of α 3 in adenovirus-Cre-treated cells as compared with control cells bearing the unrecombined floxed α 3 allele (Fig. 1 A, right) and a marked decrease in TGF- β 1-induced expression of α -SMA and collagen I in α 3 β 1-null cells. The presence or

absence of α 3 had no effect on Fn-induced TGF- β 1 activation, as judged by equivalent p-Smad2 levels (Fig. 1 A, right). These findings were confirmed in experiments of primary AECs derived from mice with lung epithelial-specific loss of α 3 achieved by crossing α 3^{fl/fl} mice with mice carrying the human *SPC-rtTA* (surfactant protein C promoter–reverse tetracycline-controlled transactivator) and *tetO-Cre* (tetO-cytomegalovirus-Cre) transgenes (Kim et al., 2008). The phenotypes of these mice, termed FASC (floxed α 3 *SPC-rtTA tetO-Cre*) mice, are the subject of a separate study (Kim et al., 2008). AECs isolated from FASC mice also demonstrated a markedly impaired EMT response to Fn *ex vivo* (unpublished data). Thus, epithelial cell α 3 appears critical for up-regulation of mesenchymal markers in response to endogenous TGF- β 1 activation. We next undertook a series of experiments to explore the molecular mechanisms underlying the role of this integrin in TGF- β 1 signaling.

α 3 β 1 regulates kidney epithelial cell responses to TGF- β 1

To assess the specific role of α 3 β 1, we used immortalized kidney α 3^{-/-} epithelial cells (Kreidberg et al., 1996). The basal morphology of these cells on laminin (Matrigel), serum-coated surfaces, and Fn was found to be similar. Unlike primary AECs, this cell line did not activate Smads upon attachment to Fn and maintained a clustered phenotype, allowing us to separate determinants of the activation of TGF- β 1 from cellular responses to active TGF- β 1. We reintroduced into cells wild-type (wt) α 3 or α 3 head domain point mutants previously shown to block the interaction of α 3 β 1 with Ln5 (G163A) or its cis-acting ligand, uPAR (urokinase receptor; H245A; Zhang et al., 1999, 2003). Surface expression detected by FACS and total protein by Western blotting of both wt α 3 and the mutants were comparable with and similar to endogenous α 3 in mouse kidney epithelial cells (Fig. S1 A, available at <http://www.jcb.org/cgi/content/full/jcb.200806067/DC1>). We initially exposed α 3^{-/-} cells and cells reconstituted with the various integrins to active TGF- β 1 for 48 h and observed their phenotype (Fig. 1 B). TGF- β 1 induced striking declustering and scattering in wt α 3- and G163A-expressing cells, whereas α 3-null and H245A mutant cells appeared unaffected. Time-lapse microscopy indicated that wt cells began to scatter at \sim 18 h after exposure to active TGF- β 1, whereas H245A cells never underwent a scattering response at >48 h of observation (Videos 1 and 2). Further stimulation time up to 96 h did not alter the observed pattern (unpublished data). The G163A mutation mimicked the wt phenotype changes but responded to TGF- β 1 faster, suggesting that the capacity of the integrin to engage Ln5 may not be critical to TGF- β 1 signaling. The α 3^{-/-} cells showed membrane ruffling; however, declustering was more difficult to assess because these cells do not form extensive cell–cell contacts at baseline, as previously reported (Wang et al., 1999). Enhanced scattering was not observed (Fig. 1 B). Two independent clones each of α 3 wt, H245A, and G163A cells showed the same phenotypic response to TGF- β 1. Most subsequent experiments were performed on serum-coated surfaces because for cells exhibiting a morphological response to TGF- β 1, the responses appeared to be qualitatively similar on serum- or Fn-coated surfaces.





Figure 1. **TGF- β 1 responses correlate with α 3 β 1 association with E-cadherin but not Ln5.** (A) TGF- β 1 responses in primary AECs from wt (left) or α 3 conditional null mice (right). wt AECs were cultured on Fn to induce endogenous TGF- β 1 activation and were incubated with TGF- β 1 receptor kinase inhibitor SB431542 (SB) or DMSO (ctl) for 4 d. α 3 conditional AECs were initially cultured on Matrigel/collagen matrices and treated with either adeno-Cre (AdCre) virus or 20 plaque-forming U/cell adeno-GFP control virus. After 96 h, cells were replated on Fn for 4 d. Cell extracts were blotted for various proteins. SB431542 blocks Smad2 phosphorylation and the α -SMA response in wt AECs. Loss of α 3 does not affect Smad2 phosphorylation but blocks up-regulation of α -SMA and collagen I. (B) Phase photographs of α 3^{-/-}, α 3 wt, H245A, and G163A mutant cells at baseline and after 48-h stimulation with TGF- β 1. Videos showing phenotypic change and cell motility of α 3 wt and H245A cells after TGF- β 1 stimulation are provided (Videos 1 and 2, available at <http://www.jcb.org/cgi/content/full/jcb.200806067/DC1>). Bar, 50 μ m. (C) E-cadherin distribution in α 3 wt and H245A mutant cells without and with 48-h TGF- β 1 stimulation. Bar, 1 μ m. (D) E-cadherin coimmunoprecipitates with wt α 3- and G163A-expressing cells but does not coimmunoprecipitate with the H245A mutant. (E) E-cadherin- β -catenin association is not affected by α 3 β 1. The aforementioned experiments have been performed at least three times with similar results.

Figure 2. TGF- β 1 up-regulates the expression of MMP-9, PAI-1, and α -SMA in α 3 wt cells but not in H245A mutant cells. (A) TGF- β 1 induces expression of MMP-9, PAI-1, and α -SMA in α 3 wt and G163A mutant cells but not in H245A mutant cells. Serum-starved cells were stimulated with 4 ng/ml TGF- β 1 for 48 h. Supernatants were collected and concentrated 10 times. Cells were lysed in RIPA buffer and blotted for integrin α 3, E-cadherin, PAI-1, α -SMA, and β -actin. MMP-9 was detected by gelatin zymography. The aforementioned blots have been repeated at least three times with a similar pattern. (B) Time course of mRNA expression in α 3 wt and H245A cells stimulated with TGF- β 1. Serum-starved cells were stimulated with 4 ng/ml TGF- β 1 for different time periods. RNA was extracted from the cells and used for direct Taqman real-time PCR analysis. Data shown are representative of four independent experiments.



Note that E-cadherin was not down-regulated as judged by total protein in any of the cells after 48 h (Fig. 2 A), which is consistent with prior evidence that suppression of E-cadherin expression during EMT often requires 7–14 d (Thiery and Sleeman, 2006). However, the distribution of E-cadherin on the surface of epithelial cells was strikingly different among the cells at baseline and after 48-h TGF- β 1 stimulation. As previously reported, α 3-null cells have less surface E-cadherin even though total protein is equivalent to that of wt cells, indicating that surface expression is very unstable and most of the E-cadherin at steady state is contained in endosomal vesicles (Wang et al., 1999). This likely accounts for the scattered appearance of null cells in culture at baseline (Fig. 1 B). In contrast, cells expressing either wt or mutant α 3 integrins have comparable surface expression of E-cadherin and form extensive cell–cell contacts at baseline (Fig. 1 C). TGF- β 1 treatment for 48 h results in a marked shift of E-cadherin away from cell–cell contacts and into the necks of lamellipodia of wt (or G163A mutant) cells but had little or no apparent effect on H245A-expressing cells (Fig. 1 C).

The lack of change in E-cadherin distribution after TGF- β 1 stimulation in H245A-expressing cells raised the possibility that the integrin associates with and affects trafficking of E-cadherin. Chattopadhyay et al. (2003) previously reported that

α 3 β 1 formed complexes with CD151 and E-cadherin, although no role for this complex in growth factor signaling or trafficking of E-cadherin was investigated. Indeed, α 3 wt and G163A integrins were robustly coimmunoprecipitated with E-cadherin, whereas the H245A mutant was undetectable in E-cadherin immunoprecipitates (Fig. 1 D). It is noteworthy that β -catenin, known to associate with the cytoplasmic tails of E-cadherin, co-precipitated equally with E-cadherin among all of the cell lines examined (Fig. 1 E). Thus, there is a strong correlation between the capacity of α 3 β 1 to associate with E-cadherin and the cellular responses to TGF- β 1.

To further examine the obvious differences among the various cells after TGF- β 1 stimulation, we looked for changes in protein expression of known TGF- β 1 target genes. Western blot analysis and gel zymography confirmed that several such proteins could be reliably detected: plasminogen activator inhibitor-1 (PAI-1), MMP-9, and α -SMA. As expected, wt α 3- and G163A-bearing cells strongly up-regulated MMP-9, PAI-1, and α -SMA after 48-h TGF- β 1 stimulation, whereas H245A and α 3 $^{-/-}$ cells showed either partial or completely absent responses of MMP-9 and α -SMA (Fig. 2 A). PAI-1 was completely suppressed in H245A-expressing cells, but α 3-null cells continued to respond, indicating some difference in the




Figure 3. Identification of colocalized $\alpha 3\beta 1$, E-cadherin, and TGF- $\beta 1$ on cell surfaces of $\alpha 3$ wt and H245A mutant cells. Cells serum starved overnight were exposed to 300 μM MDC to inhibit clathrin-mediated endocytosis for 30 min and TGF- $\beta 1$ for 4 h. E-cadherin and $\alpha 3\beta 1$ were clustered with fluorescent secondary antibodies on live cells, and the cells were fixed and stained for TGF- $\beta 1$ (TRI). Representative confocal images with individual color and the merged images of $\alpha 3$ wt and H245A mutant cells are shown. Quantitative analysis of E-cadherin- $\alpha 3\beta 1$ -TGF- $\beta 1$ coclusters comparing multiple fields on each of six separate clusters for each cell type indicated significantly ($P < 0.001$) more colocalization on $\alpha 3$ wt cells (see Identification of a tripartite complex... for details). This experiment has been performed three times with similar results. Bar, 2 μm .

requirements for induction of PAI-1 compared with MMP-9 and α -SMA in these cells.

We compared protein levels and corresponding mRNA levels in $\alpha 3$ wt and H245A mutant cells at different time points after TGF- $\beta 1$ stimulation. $\alpha 3$ wt cell mRNAs for MMP-9, PAI-1, and α -SMA peak at 24 h of stimulation and then wane (Fig. 2 B). H245A follows a similar time course; however, expression levels are strongly decreased compared with the wt. These results are consistent with the time course of TGF- $\beta 1$ responses at the protein level (Fig. S1 B). We next turned to whether the differences in TGF- $\beta 1$ responses among the various cell lines could be explained by altered Smad phosphorylation and/or nuclear translocation.

Smad phosphorylation, nuclear translocation, and MAPK activation are $\alpha 3$ integrin independent

The striking differences in mRNA responses were not mirrored by changes in Smad phosphorylation (Fig. S2 A, available at <http://www.jcb.org/cgi/content/full/jcb.200806067/DC1>). Strong TGF- $\beta 1$ -dependent Smad2 phosphorylation and nuclear accumulation was seen at the 15-min time point as described by other groups (Nakao et al., 1997; Shen et al., 1998) and revealed no difference between $\alpha 3$ wt integrin and the mutants. Fractionation was confirmed by a lack of β -actin in the nuclear immunoblots. Separate experiments comparing wt and $\alpha 3$ -null cells also showed no differences in Smad2 phosphorylation (unpublished data). Smad7 protein was detectable in both wt and H245A mutant cells but was found to not be different either at baseline or after TGF- $\beta 1$ exposure (unpublished data). To further test this point, we transiently transfected $\alpha 3$ wt and H245A mutant cells with a luciferase construct containing 12 repeats of the Smad3/Smad4-binding element CAGA (Denmler et al., 1998) and measured transcriptional activity after TGF- $\beta 1$ stim-

ulation (Fig. S2 B). Both the $\alpha 3$ wt and H245A mutants respond to TGF- $\beta 1$ with an ~ 10 -fold increase in the CAGA promoter activity, indicating that Smad functionality in the nucleus is not impaired in the H245A mutant.

We next considered Smad-independent TGF- $\beta 1$ signaling pathways known to be influenced by integrins. TGF- $\beta 1$ affects the MAPK cascades via Ras to extracellular signal-regulated kinase (ERK) and via TGF- β -activated kinase 1 to JNK and p38, modulating TGF- $\beta 1$ -mediated transcription (Derynck and Zhang, 2003). Because MAPKs also integrate downstream signals derived from integrins, we asked whether the signaling cascades through ERK, JNK, and p38 were activated differentially among the cells expressing the various forms of $\alpha 3\beta 1$. Although ERK phosphorylation increased transiently after TGF- $\beta 1$ stimulation in all of the cell lines, inhibitors of MAPK kinase (PD98059), p38 (SB202190), and JNK (JNK-II inhibitor) over a wide concentration range did not affect the measured TGF- $\beta 1$ -mediated transcriptional responses in $\alpha 3$ wt or H245A mutant cells (unpublished data). However, these experiments did confirm that motility responses of the cells and transcriptional responses can be dissociated, as has been previously reported (Ozdamar et al., 2005; Wang et al., 2005). The 50–500-nM JNK-II inhibitor completely blocked $\alpha 3$ wt cell scattering after TGF- $\beta 1$ in a dose-dependent matter, but the induction of PAI-1, α -SMA, and MMP-9 was not affected by the JNK-II inhibitor (Fig. S3, available at <http://www.jcb.org/cgi/content/full/jcb.200806067/DC1>), suggesting that the TGF- $\beta 1$ signal divided up proximal of the JNK MAPK and that the integrin influence on TGF- $\beta 1$ signaling occurs proximally in a signaling cascade. We also tested an inhibitor of glycogen synthase kinase 3 β (GSK3 β ; TPK I inhibitor), which is the kinase known to influence β -catenin signaling and previously linked to $\beta 1$ -integrin signaling. The inhibitor did not affect induction of the mesenchymal markers in these cells (unpublished data). Thus, we could

not readily ascribe the marked functional differences in cellular responses to TGF- β 1 affected by α 3 β 1 to differences in MAPK or GSK3 β signaling.

Identification of a tripartite complex of E-cadherin, α 3 β 1, and TGF- β 1

Because cellular responses to TGF- β 1 strongly correlated with the capacity of integrin α 3 β 1 to associate with E-cadherin (Fig. 1), we considered the possibility that E-cadherin, α 3 β 1, and TGF- β 1 receptors all physically interact. Preliminary attempts to visualize E-cadherin–TGF- β 1 complexes on the cell surface indicated that little or no TGF- β 1 could be found enriched in adherens junctions. However, when E-cadherin (Fig. 3, red) was clustered on live cells expressing wt α 3, a fraction of α 3 (Fig. 3, blue) was found to cocluster (Fig. 3, purple) on the apical surfaces. The degree of coclustering appeared to depend on the degree of E-cadherin available for clustering, mainly around the edges of cellular islands and distinct from adherens junctions. Likewise, colocalization of E-cadherin, α 3 β 1, and TGF- β 1 (Fig. 3, white) were seen mainly at the periphery of the epithelial islands. In comparison, cells expressing the nonpermissive H245A mutant α 3 under the same conditions showed much less colocalization of the three receptors (Fig. 3, top). The mean white pixel intensity in multiple peripheral fields of six separate cellular islands for each cell type was \sim 10 times higher on wt as compared with H245A integrin-expressing cells ($P < 0.001$). It is noteworthy that detection of TGF- β 1 (Fig. 3, green) in surface coclusters with α 3 β 1 and E-cadherin was enhanced by a blockade of clathrin-dependent endocytosis and stimulation with TGF- β 1. TGF- β 1 is known to induce clathrin-mediated receptor turnover (Runyan et al., 2005), and the images in Fig. 3 were obtained in the presence of monodansylcadaverine (MDC), an inhibitor of this turnover pathway. Merged images without and with clathrin pathway blockers and TGF- β 1 stimulation are provided in Fig. S4 A (available at <http://www.jcb.org/cgi/content/full/jcb.200806067/DC1>).

To allow for further analysis of E-cadherin–TGF- β 1 complexes, α 3 wt epithelial or H245A mutant cells stably expressing a myc-tagged TGF- β 1 were established. E-cadherin was present in the myc–TGF- β 1 immunoprecipitates whether the integrin associates with E-cadherin or not (Fig. 4 A). Immunoprecipitation (IP) of myc–TGF- β 1 consistently coprecipitated wt α 3 integrin but failed to coprecipitate the nonpermissive H245A mutant (Fig. 4 B), suggesting that the integrin likely associates with TGF- β 1 indirectly through E-cadherin. Additional experiments revealed that $<10\%$ of the total E-cadherin associated with TGF- β 1 by co-IP (unpublished data), raising the possibility that TGF- β 1 only associates with a small subfraction of cell surface E-cadherin.

The extent of E-cadherin–TGF- β 1 coclustering was not obviously different at baseline between α 3 wt and H245A mutant cells. However, when clusters were visualized 24 h after TGF- β 1 stimulation, coclusters of E-cadherin and TGF- β 1 disappeared from the surface of α 3 wt cells, whereas these coclusters persisted on cells expressing the H245A mutant (Fig. 4 C and Fig. S4 B). Quantification of the total extent of yellow pixels as a fraction of the sum of both the yellow and red pixels in

multiple random fields indicated that this fraction dramatically decreased in α 3 wt cells (Fig. 4 D). The exact opposite was seen in H245A cells stimulated with TGF- β 1, in which this fraction actually increases, reflecting persistence of surface E-cadherin–TGF- β 1 clusters concurrent with loss of overall surface E-cadherin. Overall turnover of TGF- β 1 induced by TGF- β 1 was not impacted (Fig. 4 E), indicating that α 3 specifically determines the fate of TGF- β 1–E-cadherin complexes but not all TGF- β receptor turnover pathways. The observation that E-cadherin and TGF- β 1 receptors colocalize and appear to internalize together in response to TGF- β 1 raises the possibility that transcription factors known to be associated with these receptors, i.e., β -catenin and receptor-associated Smads, could also interact. We next investigated this possibility.

β -Catenin–p-Smad2 complex formation requires TGF- β 1, α 3 membrane complex, and endocytosis

As shown in Fig. 5 A, epithelial cells expressing wt α 3 formed clear β -catenin–p-Smad2 complexes 1 h after TGF- β 1 stimulation, whereas α 3-null or H245A mutant cells failed to support formation of this transcriptional complex. The appearance of β -catenin–p-Smad2 complexes did not result in canonical β -catenin signaling as judged by the lack of response of the TOPFlash T cell factor/catenin reporter to TGF- β 1 in transiently transfected α 3 wt cells (unpublished data), which is consistent with prior evidence that TOPFlash is not activated directly by TGF- β 1 stimulation in other cell types (Labbe et al., 2000). Also, we could not detect β -catenin in the nucleus by direct immunostaining at any time point after TGF- β 1 stimulation. However, the catenin is functionally involved in TGF- β 1 signaling, which is indicated by near-complete suppression of induction of α -SMA and PAI-1 protein in α 3 wt cells stably transfected with a dominant-negative version of β -catenin (Fig. 5 B, left). To further test this point, α 3-expressing cells were stably transfected with short hairpin RNA (shRNA) blocking β -catenin expression or a nonblocking shRNA, and the response to TGF- β 1 was again assessed. Compared with control, knockdown of β -catenin expression by $>50\%$ markedly attenuated the α -SMA response to TGF- β 1 without decreasing the p-Smad2 response (Fig. 5 B, right), confirming the importance of β -catenin in the cellular response to TGF- β 1 signaling.

The formation of β -catenin–p-Smad complexes after TGF- β 1 stimulation was found to require α 3 β 1-dependent endocytosis because both clathrin inhibitors MDC and chlorpromazine inhibited β -catenin–p-Smad complex formation in α 3 wt cells (Fig. S5 A, available at <http://www.jcb.org/cgi/content/full/jcb.200806067/DC1>). If β -catenin–p-Smad complexes are formed during or soon after receptor internalization, it is unclear how β -catenin would move from its physical interaction with E-cadherin to form transcriptionally active complexes with p-Smad2. An attractive possibility is tyrosine phosphorylation of β -catenin, especially tyrosine 654, which is known to lead to dissociation of β -catenin from E-cadherin (Piedra et al., 2001). Indeed, we found that Y-654 was phosphorylated within 60 min after active TGF- β 1 addition to the cultures. This phosphorylation absolutely required the presence of a permissive α 3 β 1.





Figure 4. Turnover of E-cadherin and TGF- β 1 complexes requires α 3 β 1 integrin association. (A) E-cadherin immunoprecipitates equally well with myc-TGF- β 1 in both α 3 wt cells and H245A mutant cells. Myc IP from α 3 wt cells without myc-TGF- β 1 expression (α 3 wt no myc) was used as a negative control. (B) α 3 integrin only immunoprecipitates with myc-TGF- β 1 in α 3 wt cells but not in H245A mutant cells. (C) Confocal microscopy analysis of the effect of TGF- β 1 on E-cadherin-TGF- β 1 coclusters in α 3 wt and H245A mutant cells. After overnight treatment with TGF- β 1, cell surface E-cadherin was clustered, fixed, and stained with anti-myc antibody. α 3 wt cells show decreased E-cadherin-TGF- β 1 coclusters after TGF- β 1 stimulation, whereas E-cadherin-TGF- β 1 coclusters persist in H245A mutant cells. Red, E-cadherin; green, TGF- β 1; blue, DAPI. (D) Quantification of E-cadherin-TGF- β 1 complex turnover. After TGF- β 1 stimulation, the fraction of E-cadherin-TGF- β 1 coclusters decreases in α 3 wt cells but increases in H245A mutant cells. (E) Immunoblotting of total myc-TGF- β 1 before and after TGF- β 1 stimulation overnight showing that the overall turnover of TGF- β 1 is not different between α 3 wt and H245A mutant cells. All of the aforementioned experiments have been performed at least three times with similar results. Error bars indicate standard deviation. Bar, 1 μ m.

Neither α 3-null cells nor cells bearing the H245A mutant phosphorylated β -catenin at Y-654 in response to TGF- β 1 stimulation (Fig. 5 C).

To examine whether endocytosis is important for β -catenin tyrosine phosphorylation, α 3 wt cells were serum

starved, exposed to an inhibitor of clathrin-mediated endocytosis (MDC), and stimulated with TGF- β 1. As shown in Fig. 5 D, pY654- β -catenin was absent in the inhibitor-treated cells, indicating that this phosphorylation requires endocytosis. Note that the blockade of endocytosis had no clear

Figure 5. β -Catenin-p-Smad2 complex formation requires wt $\alpha 3$ and is essential for TGF- $\beta 1$ -mediated up-regulation of mesenchymal genes. (A) β -Catenin effectively immunoprecipitates p-Smad2 in $\alpha 3$ wt cells but not in $\alpha 3^{-/-}$ or H245A mutant cells after TGF- $\beta 1$ stimulation. (B) Inhibition of β -catenin by expression of a dominant-negative β -catenin (dn β -cat; left) or an shRNA against β -catenin (sh β -cat; right) in $\alpha 3$ wt cells inhibits TGF- $\beta 1$ -induced α -SMA but not Smad2 phosphorylation compared with wt $\alpha 3$ cells transfected with vector alone, nonsilencing control shRNA (shctl), or left nontransfected. (C) Tyrosine phosphorylation of β -catenin and formation of the pY654- β -catenin-p-Smad2 complex is only detected in $\alpha 3$ wt cells treated with TGF- $\beta 1$ but not in $\alpha 3^{-/-}$ or H245A mutant cells. (D) Tyrosine phosphorylation of β -catenin (β -cat) and formation of the pY654- β -catenin-p-Smad2 complex in $\alpha 3$ wt cells requires endocytosis. $\alpha 3$ wt cells untreated or exposed to 300 μ M of the clathrin inhibitor MDC were stimulated with TGF- $\beta 1$ and analyzed for pY654- β -catenin-p-Smad2 complexes by co-IP. The supernatant from pY654- β -catenin IP was subsequently immunoprecipitated with total β -catenin antibody. Lane 3 shows p-Smad associated with pY654- β -catenin, and lane 5 shows the remainder of p-Smad2 on β -catenin after pY654- β -catenin depletion. GAPDH, glyceraldehyde 3-phosphate dehydrogenase. All of the aforementioned experiments have been performed at least three times with similar results.



effect on overall levels of p-Smad2 (Fig. 5 D, right) but markedly attenuated formation of complexes as judged by co-IP. Interestingly, re-IP of total β -catenin in the supernatant after pY654- β -catenin IP resulted in little p-Smad2 (Fig. 5 D), and pY654- β -catenin mAb coimmunoprecipitated similar amounts of p-Smad2 as antibodies to total β -catenin (not depicted), indicating that p-Smad2 preferentially associates with tyrosine-phosphorylated β -catenin, and the principal form of β -catenin in β -catenin-Smad complexes is Y-654 phosphorylated. These data imply that the internalization of colocalized E-cadherin and TGF- β R1 (Fig. 4) allows β -catenin to be tyrosine phosphorylated, dissociate from E-cadherin, and form complexes with p-Smad2.

Catenin-Smad complex formation is regulated by cell contacts

The finding of $\alpha 3\beta 1$ -dependent β -catenin-p-Smad2 complexes raised the intriguing possibility that formation of such complexes is regulated at the cell surface by determinants of an assembly of complexes containing $\alpha 3\beta 1$, E-cadherin, and TGF- β R1. Therefore, we asked whether limiting the availability of either $\alpha 3\beta 1$ or E-cadherin would influence the formation of β -catenin-p-Smad2 complexes and the subsequent transcriptional responses. To vary access of E-cadherin to TGF- β 1 receptors, we used two approaches. First, we varied the culture medium calcium concentration. Under high calcium conditions (~ 1.8 mM), $\alpha 3$ wt cells show a tight clustered phenotype (Fig. 6 A). Clustering





Figure 6. Effect of calcium on TGF- β 1-mediated signaling and β -catenin-p-Smad2 complex formation. (A) Phase photographs of α 3 wt cells cultured in either low (125 μ M) or high (1.8 mM) calcium media showing cells cultured in high calcium media form tight cell-cell contact, whereas cells cultured in low calcium media have loose cell-cell contact. Bar, 50 μ m. (B) Confocal microscopy of E-cadherin clusters in α 3 wt cells cultured under either low (125 μ M) or high (1.8 mM) calcium conditions. α 3 wt cells cultured in high calcium media show that E-cadherin clusters occur mainly in cells at the periphery of the colony, whereas α 3 wt cells cultured in low calcium media form E-cadherin clusters throughout the colony. The boxed areas are shown in higher magnifications in the bottom panels. Bar, 2 μ m. (C) Serum-starved α 3 wt and H245A mutant cells under high or low calcium conditions were stimulated with TGF- β 1 for 48 h. Supernatants were removed from the plates and concentrated for zymography. α 3 wt cells cultured in low calcium show higher MMP-9 levels than α 3 wt cells cultured in high calcium after TGF- β 1 stimulation. (D) Serum-starved α 3 wt cells in high or low calcium media were stimulated with TGF- β 1 for 48 h, and the lysates were blotted for α -SMA. α -SMA is only up-regulated in cells cultured in low calcium medium and not in high calcium condition. (E) Cells cultured under low calcium condition show increased β -catenin-p-Smad2 complex formation. α 3 wt cells cultured in either high or low calcium medium were treated with TGF- β 1, and the lysates were subjected to β -catenin IP followed by p-Smad2 immunoblotting. The β -catenin-p-Smad2 complex formation is only seen with cells cultured in low calcium condition and not in high calcium condition. All of the aforementioned experiments have been performed at least three times with similar results.

of E-cadherin on live cells is mainly confined to the periphery of well-compacted epithelial cell islands (Fig. 6 B), and there are relatively little β -catenin-p-Smad2 complexes formed after TGF- β 1 stimulation (Fig. 6 E). In contrast, under low calcium conditions (\sim 125 μ M Ca^{2+}), there is more extensive E-cadherin clustering (Fig. 6 B) and complex formation (Fig. 6 E). Again, there is a direct correlation between the degree of β -catenin-p-Smad complexes and the induction of MMP-9 (Fig. 6 C) and α -SMA (Fig. 6 D) after TGF- β 1 stimulation. Note that the complexes are observed within 60 min of exposure of the cells to TGF- β 1, whereas the protein responses are measured after 48 h. In a second approach, we varied the plating density of the epithelial cells 24 h before TGF- β 1 stimulation. At a high density in which there is extensive E-cadherin-dependent cell-cell contact, little or no β -catenin-p-Smad complexes were observed (Fig. S5 B) and little α -SMA was induced (not depicted). At low plating density, i.e., when

E-cadherin is unengaged, both transcriptional complexes and induction of α -SMA were robust, confirming the critical role of the extent of adherens junction formation as a regulator of cellular responses to TGF- β 1.

When α 3 wt cells were cultured on Ln5, sequestering α 3 β 1 to the basal surface, β -catenin-p-Smad2 complexes failed to form (Fig. 7 A), and the α -SMA transcriptional response to TGF- β 1 was again attenuated (Fig. 7 C, left). In contrast, on purified Fn, complex formation and α -SMA induction is robust. To further test the role of Ln5 engagement in regulating TGF- β 1 responses, we repeated the experiments using G163A mutant α 3-expressing cells that are unable to engage Ln5, and thus, Ln5 would be predicted to have less influence on the TGF- β 1 responses of these cells. Indeed, β -catenin-p-Smad2 complexes were readily observed when G163A cells were plated onto either Fn or Ln5 (Fig. 7 B), and the α -SMA induction was equivalent on either matrix (Fig. 7 C, right). Collectively, these data indicate that the α 3 β 1-E-cadherin-TGF- β 1

Figure 7. **Regulation of β -catenin-p-Smad2 complex formation by extracellular matrices.** (A) $\alpha 3\beta 1$ engagement on Ln5 limits the formation of β -catenin-p-Smad2 complex formation. Lysates of TGF- $\beta 1$ -stimulated $\alpha 3$ wt cells from either Fn- or Ln5-coated plates were subject to β -catenin IP followed by p-Smad2 immunoblotting. The β -catenin-p-Smad2 complex formation is only seen with cells plated on Fn and not with Ln5. (B) Ln5 does not limit formation of β -catenin-p-Smad2 complexes in cells unable to engage Ln5 through $\alpha 3\beta 1$ (G163A mutant cells). Lysates of TGF- $\beta 1$ -stimulated G163A mutant cells from either Fn- or Ln5-coated plates were subjected to β -catenin IP followed by p-Smad2 immunoblotting. The β -catenin-p-Smad2 complex formation is seen with cells plated on both Fn and Ln5. (C) $\alpha 3\beta 1$ engagement on Ln5 suppresses TGF- $\beta 1$ -induced α -SMA up-regulation. Up-regulation of α -SMA is suppressed in $\alpha 3$ wt cells plated on Ln5 but not in G163A mutant cells on Ln5. The aforementioned experiments have been performed three times with similar results.



membrane complexes can coordinate cues from cell-matrix and cell-cell interactions to determine the cellular response to TGF- $\beta 1$.

Discussion

The findings reported in this study identify a previously unrecognized role for the prominent epithelial integrin $\alpha 3\beta 1$ in TGF- $\beta 1$ signaling, providing evidence that the integrin functions as a sensor of cell contacts to regulate TGF- $\beta 1$ signaling. The essential function of $\alpha 3\beta 1$ appears to depend on the presence of surface complexes on epithelial cells, which are comprised of a subfraction of the surface pool of $\alpha 3\beta 1$, E-cadherin, and TGF- $\beta 1$ receptor, which critically influence the signaling response to TGF- $\beta 1$. The components of this complex by itself reveals the likely elements of its function, as modeled in Fig. 8. Surface complexes of E-cadherin and TGF- $\beta 1$ bring two key transcription factors involved in induction of EMT into spatial proximity: β -catenin and receptor Smads (Chilosi et al., 2003; Kalluri and Neilson, 2003; Yook et al., 2006). After TGF- $\beta 1$ stimulation, the surface complexes are internalized, and β -catenin is phosphorylated on Y-654. The formation of stable complexes between pY654- β -catenin and p-Smad2 and the appearance of these complexes strongly correlate with subsequent EMT-related protein expression (Figs. 5-7). Both internalization and β -catenin tyrosine phosphorylation require the third component

of the complex, $\alpha 3\beta 1$. The unique function of $\alpha 3\beta 1$ in this context appears to reside in its affinity for E-cadherin, without which E-cadherin-TGF- $\beta 1$ complex internalization is impaired, and this signaling system does not operate. Conceptually, these findings are similar to prior evidence in another system in which integrin engagement is found to regulate growth factor signaling at least in part by altering pathways of endocytosis that influence the quality and duration of receptor signaling (del Pozo et al., 2004). However, our findings also reveal a completely new point of intersection between β -catenin and Smad signaling pathways (Lei et al., 2004), and the linkage of this point of intersection to a dynamic interplay between adhesion receptors and their normal pericellular contacts provides new understanding for how the extracellular environment can regulate the epithelial cell response to TGF- $\beta 1$ (Masszi et al., 2004; Thiery and Sleeman, 2006).

TGF- $\beta 1$ has been previously reported to promote tyrosine phosphorylation of β -catenin (Tian and Phillips, 2002), although the specific site of phosphorylation and its functional significance in TGF- $\beta 1$ signaling has been unknown. Our finding of integrin-dependent tyrosine phosphorylation of Y654- β -catenin is important because phosphorylation of β -catenin at Y654 is known to promote both dissociation of β -catenin from E-cadherin and stabilization of β -catenin from ubiquitination and degradation (Brembeck et al., 2006). Therefore, independently of Wnt signaling, our findings indicate that TGF- $\beta 1$ can

promote a pathway of cross talk with β -catenin by generating stable pY654- β -catenin-Smad complexes. The data indicate that only a fraction of the β -catenin is phosphorylated, and presumably, this reflects, at least in part, the pool internalized with E-cadherin and TGF- β R1 after TGF- β 1 stimulation. However, internalization alone does not appear to be sufficient, as α 3-null cells, even though they display unstable surface E-cadherin, fail to phosphorylate β -catenin in response to TGF- β 1. We are uncertain of the tyrosine kinases responsible for Y654 phosphorylation. At least three different tyrosine kinases have been reported to phosphorylate this site: c-src (and possibly other src family members; Roura et al., 1999), Bcr-Abl (Coluccia et al., 2007), and hepatocyte growth factor receptor Met (Zeng et al., 2006). In addition, the TGF- β 1 receptor complex itself has recently been shown to have tyrosine kinase activity (Lee et al., 2007). As src family kinases are known to associate with the cytoplasmic tails of β 1 integrins (Hynes, 2002), the requirement for the integrin likely involves a src family kinase. However, the exact kinase and how this kinase activity is locally regulated by α 3 β 1 after TGF- β 1 stimulation remains to be defined in future work.

A recent study described defective TGF- β 1 signaling in α 3 β 1-deficient murine keratinocytes and attributed the defect to higher levels of the inhibitory Smad, Smad7 (Reynolds et al., 2008). Corresponding to the higher Smad7 levels was lower overall levels of TGF- β 1 receptors and lower Smad2/3 phosphorylation after TGF- β 1 stimulation, which is consistent with the known effects of Smad7 on the TGF- β 1 signaling pathway. However, neither in the kidney epithelial cell line studied here nor in α 3 β 1-deficient primary lung epithelial cells were we able to detect altered levels of Smad7. Consistent with our data, overall levels of TGF- β 1 receptors were not lower in α 3-deficient cells, and Smad2/3 phosphorylation was not lower after TGF- β 1 stimulation (Fig. 5). Together, these observations may imply that the impact of α 3 β 1 on TGF- β 1 signaling operates through different pathways in different cells but may also suggest that this integrin may critically influence TGF- β 1 signaling in many, if not all, epithelial cells.

We have previously reported that stable overexpression of uPAR in kidney epithelial cells promotes an EMT response via interactions of uPAR with α 3 β 1 (Zhang et al., 2003). The functionally important H245A α 3 point mutation used in this study was discovered in a screen of several integrin β propeller mutants for inhibitors of uPAR-dependent EMT. Cells expressing the H245A mutant were found to have disrupted uPAR- α 3 β 1 interactions and attenuated EMT compared with cells expressing wt α 3. However, basal epithelial cells express little or no uPAR, and uPAR is a known TGF- β 1 target gene (Yue et al., 2004). Therefore, we asked whether the response of these cells to TGF- β 1 involved uPAR. Surprisingly, H245A-expressing cells were found to have little or no response to active TGF- β 1, implying a critical role for α 3 β 1 in the early response of these cells even before induction of uPAR and leading to the series of experiments reported in this study. It is likely that uPAR, once induced by TGF- β 1, further promotes signaling amplifying EMT, as has been recently reported in other systems (Lester et al., 2007). However, findings reported in this study indicate




Figure 8. **Model of TGF- β 1 signaling.** The schematic illustration summarizes the influence of α 3 β 1 on TGF- β 1 signaling as a function of cell-cell and cell-matrix contacts. For details see the Discussion.

that, independently of uPAR, α 3 β 1 has a critical role in the initial TGF- β 1 signaling leading to EMT.

One limitation of our observations is that it is unclear how pY654- β -catenin-p-Smad2 complexes operate to promote mesenchymal gene responses to TGF- β 1. Although there is a strong correlation between formation of these complexes and initiation of EMT in both primary lung epithelial cells as well as the kidney epithelial cells primarily studied here, it is unclear what the critical promoter targets are for this complex. TGF- β 1 does not strongly activate the canonical Lef1/T cell factor reporter, and activation of the canonical Smad-binding CAGA reporters is not different between cells expressing wt and H245A mutant α 3 β 1 (Fig. S2), suggesting that a noncanonical signaling mechanism may be in play in this signaling pathway. Moreover, how can the pro-EMT signaling pathway involving p-Smad2 identified in this study be reconciled with recent findings that Smad2-null keratinocytes spontaneously develop EMT (Hoot et al., 2008), suggesting the possibility that p-Smad2 actually functions to suppress EMT? We note that total p-Smad2 levels after TGF- β 1 stimulation are strongly and comparably induced in both responding α 3 wt cells and nonresponsive α 3-null and H245A mutant-expressing cells (Fig. 5). It is well known that cell-specific coactivators and repressors critically regulate the repertoire of responses to Smad signaling (Derynck and Zhang, 2003). This is consistent with the view that p-Smad2 may function to suppress EMT unless and until pY654- β -catenin-p-Smad2 complexes form under the conditions revealed by our experiments. If so, the set of TGF- β 1 target genes

activated by pY654- β -catenin-p-Smad2 complexes and how this switches p-Smad2 from a suppressor to activator role in EMT are important areas for further investigation.

Materials and methods

Reagents

Recombinant active TGF- β 1 was obtained from R&D Systems. Inhibitors PD98095 (MEK1), SB202190 (p38), SB431542 (TGF- β), JNK-II (SP600125), GSK3 inhibitor VII (TPK I inhibitor), and polyclonal p-Smad2 antibody were purchased from EMD. Rat anti-E-cadherin mAb was obtained from Invitrogen. E-cadherin pAb, mouse anti-E-cadherin mAb, and β -catenin mAb were purchased from BD. MDC, chlorpromazine hydrochloride, myc pAb, α -SMA mAb, and β -actin mAb were obtained from Sigma-Aldrich. Sheep anti-mouse PAI-1 pAb was obtained from American Diagnostica Inc. TGF- β RI pAb, β -catenin pAb, and myc mAb were purchased from Cell Signaling Technology. α 3 pAb, pY654- β -catenin mAb, and fluorophore-labeled antibodies were purchased from Invitrogen. Secondary HRP-conjugated antibodies and Smad2/3 pAb were purchased from Santa Cruz Biotechnology, Inc. 804-G supernatant rich in Ln5 was provided by J.C. Jones (Northwestern University Medical School, Chicago, IL). Collagen I pAb was obtained from Abcam. Fn was obtained from Roche. Nuclear extraction kit and TOPFlash T cell factor reporter plasmid were obtained from Millipore.

Plasmid constructs and virus production

Myc-TGF- β RI (provided by Y. Henis, Tel Aviv University, Tel Aviv, Israel; Gilboa et al., 1998) was cloned into a retroviral expression vector by M. Wheelock and K. Johnson's laboratory (University of Nebraska Medical Center, Omaha, NE). Dominant-negative β -catenin (provided by P. McCrea, University of Texas MD Anderson Cancer Center, Houston, TX) missing the C-terminal Lef-binding sites and fused with the transcriptional repressor *en-gailed* was cloned into a retroviral expression vector (pWZL-blast). Retroviral supernatants were produced using Phoenix-E cells.

β -Catenin knockdown

Five different mouse β -catenin shRNAs in pLKO.1 lentiviral vector were obtained from Open Biosystems, and the lentiviral particles were packaged at the University of California, San Francisco core facility. Virus-infected α 3 wt cells were selected with puromycin, and knockdown efficiency of β -catenin was screened. Cells stably expressing active shRNA (sh β -cat: 5'-CCGGCCCA-AGCCTTAGTAAACATACTCGAGTTATGTTACTAAGGCTGGGTTTT-3') and cells expressing inactive control shRNA (shctl: 5'-CCGGCCATTGTTG-TGCAGTTGCTTCTCAGAAAGCAACTGCACAAACAATGGTTTT-3') were compared for TGF- β 1 response.

α 3-Null kidney epithelial cell culture

α 3-Null B12 cells expressing the wt α 3 or point mutants (H245A and G163A) were cultured in DME supplemented with 10% FBS, 50 μ g/ml zeocin, and penicillin-streptomycin as previously described (Zhang et al., 2003). All cells were sorted periodically to maintain surface levels of α 3 comparable with wt. Unless otherwise indicated, cells were seeded in DME 10% FBS for 24 h and switched to F-12 serum-free media (\sim 200 μ M Ca^{2+}) with 4 ng/ml TGF- β 1 for different time periods. Cells were analyzed by immunoblotting, IP, and Taqman real-time PCR as previously described (Zhang et al., 2003; Kim et al., 2006). For inhibitor experiments, the following concentrations were used: 100 nM JNK-II inhibitor, 20 μ M SB202190, 10 μ M PD98059, and 50 μ M TPK I inhibitor.

IP and Western blotting

For co-IP of E-cadherin and α 3 integrin, 1% Triton X-100 lysates were immunoprecipitated with E-cadherin mAb, and the immunoprecipitates were blotted for α 3 integrin and E-cadherin. For co-IP of myc-TGF- β RI and α 3 integrin or E-cadherin, 1% NP-40 lysates were immunoprecipitated with myc mAb for α 3 integrin co-IP or myc pAb for E-cadherin co-IP. The immunoprecipitates were blotted with α 3 integrin pAb or E-cadherin mAb. For co-IP of β -catenin and p-Smad2, serum-starved cells were treated with 2 ng/ml TGF- β 1 for 60 min, the RIPA (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1% Triton X-100, 1% sodium deoxycholate, and 0.1% SDS) lysates were immunoprecipitated with β -catenin mAb or pY654- β -catenin mAb, and the immunoprecipitates were blotted with p-Smad2 pAb and β -catenin pAb. In some cases, cells were pretreated with clathrin inhibitors, incubated in high Ca^{2+} -containing medium, or seeded on Fn or Ln5 surfaces before TGF- β 1 treatment and co-IP.

Time-lapse microscopy and bright field microscopy of α 3 wt and H245A mutant cells

Serum-starved cells in 60-mm dishes were treated with 4 ng/ml TGF- β 1 and maintained in a mini incubator (5% CO_2 at 37°C) under a microscope (TE300; Nikon), and time-lapse microscopy (10 \times magnification) was performed for 48 h (1 frame/10 min) using a digital imaging system and the Simple PCI software (Hamamatsu Photonics). Cells were filmed over the same times after plating to eliminate nonspecific differences. Bright field images of cells in culture were taken under a microscope (TE300; 20 \times magnification) using a camera (Spot; Diagnostic Instruments, Inc.) and processed using Photoshop (Adobe).

Immunofluorescence, coclustering, and confocal microscopy

Cells were seeded in chamber slides and cultured for 2–3 d until 50% confluent. For E-cadherin staining, cells treated with TGF- β 1 for 48 h were incubated with rat anti-E-cadherin antibody followed by secondary anti-rat IgG–Alexa Fluor 568 antibody. For coclustering, cells were first incubated with rat anti-E-cadherin or mouse anti- α 3 antibodies on ice for 30 min and washed with PBS. E-cadherin or α 3 was clustered at 37°C with Alexa Fluor 568-conjugated anti-rat or Alexa Fluor 350-conjugated anti-mouse secondary antibodies for 60 min. Washed cells were fixed in methanol, blocked with 10% goat serum PBS, and stained with anti-myc or anti-TGF- β RI primary antibodies and Alexa Fluor 488-conjugated anti-rabbit secondary antibody. IgG isotypes were used as negative controls. Confocal microscopy was performed on a laser-scanning microscope (LSM 510; Carl Zeiss, Inc.). Images were captured with a 63 \times oil immersion objective, analyzed with the Image Browser software (LSM; Carl Zeiss, Inc.), and processed with Photoshop. For quantitative analysis of E-cadherin-TGF- β RI coclusters, 16 random confocal microscopy fields were used for each treatment (α 3 wt and H245A mutant cells \pm TGF- β 1 treatment). Data are expressed as the fractional difference (\pm SD) between E-cadherin-TGF- β RI coclusters (total yellow pixels/total yellow + red pixels) in cells stimulated with TGF- β 1 minus that of unstimulated cells for both α 3 wt and H245A mutant cells. For quantitative analysis of E-cadherin- α 3-TGF- β RI coclusters, the exported confocal images were converted to black and white, and the fraction of white in each of at least six random fields along the edges of each of six colonies for each cell line (α 3 wt and H245A mutant cells) was measured using the ImageJ software (National Institutes of Health). Statistical analysis was conducted using a Student's two-tailed *t* test for independent groups.

Gelatin zymography

Supernatants from cells cultured in F-12 + 0.001% BSA treated with TGF- β 1 were collected and concentrated 10-fold. Protein concentration was normalized on cell number and volume. Lysates were prepared in 3 \times Laemmli SDS sample buffer, and 10% gelatin PAGE was performed (Invitrogen) and analyzed for MMP activity as described previously (Wei et al., 2007).

Mouse type II cell isolation and treatment with Cre recombinase

Isolation of type II alveolar cells from α 3 conditional knockout mice was performed as described previously (Corti et al., 1996) with recent modifications (Kim et al., 2006). Purified primary type II cells were cultured on Matrigel/collagen type I/small airway growth medium (70:5:25 vol/vol) as described previously (Rice and Leinwand, 2003). Cells were mixed with adeno-Cre virus (provided by L. Wu, University of California, Los Angeles, Los Angeles, CA) or adeno-GFP virus at 10–20 plaque-forming U/cell before seeding cells on Matrigel/collagen and were incubated with virus for 96 h. Cells were removed from the Matrigel/collagen and plated on Fn to activate endogenous TGF- β 1 signaling (Kim et al., 2006).

Online supplemental material

Videos 1 and 2 show TGF- β 1-induced cell motility and morphological changes in α 3 wt cells (Video 1) and H245A mutant cells (Video 2). Fig. S1 shows TGF- β 1-induced expression of MMP-9, PAI-1, and α -SMA in α 3 wt but not in H245A mutant cells. Equivalent Smad-dependent signaling in α 3 wt, H245A, and G163A mutant cells are shown in Fig. S2. Fig. S3 shows that inhibition of JNK MAPK blocks TGF- β 1-induced phenotype change but not expression of MMP-9 and α -SMA. Colocalization and turnover of α 3 β 1, E-cadherin, and TGF- β RI on cell surfaces of α 3 wt cells are identified in Fig. S4. Fig. S5 presents regulation of TGF- β 1-mediated β -catenin-p-Smad2 complex formation. Online supplemental material is available at <http://www.jcb.org/cgi/content/full/jcb.200806067/DC1>.

We thank Michael Galvez and Liliane Robillard for technical assistance, Dr. Lilly Wu for kind gifts of adenovirus-expressing Cre recombinase, Dr. J.C. Jones for the 804-G supernatant rich in Ln5, Drs. Y. Henis, M. Wheelock, and

K. Johnson for myc-TGF- β 1 in retroviral vector, and Dr. P. McCrea for dominant-negative β -catenin.

This work was supported by the National Institutes of Health (grant T32HL007185 to Y. Kim, grant K08HL085290 to K.K. Kim, and grant RO1 HL44712 to H.A. Chapman) and the Parker B. Francis Foundation (grant to K.K. Kim).

Submitted: 10 June 2008

Accepted: 19 December 2008

References

- Bhowmick, N.A., R. Zent, M. Ghiassi, M. McDonnell, and H.L. Moses. 2001. Integrin beta 1 signaling is necessary for transforming growth factor-beta activation of p38MAPK and epithelial plasticity. *J. Biol. Chem.* 276:46707–46713.
- Brembeck, F.H., M. Rosario, and W. Birchmeier. 2006. Balancing cell adhesion and Wnt signaling, the key role of beta-catenin. *Curr. Opin. Genet. Dev.* 16:51–59.
- Chattopadhyay, N., Z. Wang, L.K. Ashman, S.M. Brady-Kalnay, and J.A. Kreidberg. 2003. α 3 β 1 integrin-CD151, a component of the cadherin-catenin complex, regulates PTP μ expression and cell-cell adhesion. *J. Cell Biol.* 163:1351–1362.
- Chilosi, M., V. Poletti, A. Zamo, M. Lestani, L. Montagna, P. Piccoli, S. Pedron, M. Bertaso, A. Scarpa, B. Murer, et al. 2003. Aberrant Wnt/beta-catenin pathway activation in idiopathic pulmonary fibrosis. *Am. J. Pathol.* 162:1495–1502.
- Coluccia, A.M., A. Vacca, M. Dunach, L. Mogni, S. Redaelli, V.H. Bustos, D. Benati, L.A. Pinna, and C. Gambacorti-Passerini. 2007. Bcr-Abl stabilizes beta-catenin in chronic myeloid leukemia through its tyrosine phosphorylation. *EMBO J.* 26:1456–1466.
- Corti, M., A.R. Brody, and J.H. Harrison. 1996. Isolation and primary culture of murine alveolar type II cells. *Am. J. Respir. Cell Mol. Biol.* 14:309–315.
- del Pozo, M.A., N.B. Alderson, W.B. Kiesses, H.H. Chiang, R.G. Anderson, and M.A. Schwartz. 2004. Integrins regulate Rac targeting by internalization of membrane domains. *Science.* 303:839–842.
- Dennler, S., S. Itoh, D. Vivien, P. ten Dijke, S. Huet, and J.M. Gauthier. 1998. Direct binding of Smad3 and Smad4 to critical TGF beta-inducible elements in the promoter of human plasminogen activator inhibitor-type 1 gene. *EMBO J.* 17:3091–3100.
- Derynck, R., and Y.E. Zhang. 2003. Smad-dependent and Smad-independent pathways in TGF-beta family signalling. *Nature.* 425:577–584.
- Dogic, D., P. Rousselle, and M. Aumailley. 1998. Cell adhesion to laminin 1 or 5 induces isoform-specific clustering of integrins and other focal adhesion components. *J. Cell Sci.* 111:793–802.
- Eble, J.A., K.W. Wucherpennig, L. Gauthier, P. Dersch, E. Krukonis, R.R. Isberg, and M.E. Hemler. 1998. Recombinant soluble human alpha 3 beta 1 integrin: purification, processing, regulation, and specific binding to laminin-5 and invasion in a mutually exclusive manner. *Biochemistry.* 37:10945–10955.
- Fontana, L., Y. Chen, P. Prijatelj, T. Sakai, R. Fassler, L.Y. Sakai, and D.B. Rifkin. 2005. Fibronectin is required for integrin alphavbeta6-mediated activation of latent TGF-beta complexes containing LTBP-1. *FASEB J.* 19:1798–1808.
- Geiger, B., A. Bershadsky, R. Pankov, and K.M. Yamada. 2001. Transmembrane crosstalk between the extracellular matrix-cytoskeleton crosstalk. *Nat. Rev. Mol. Cell Biol.* 2:793–805.
- Gilboa, L., R.G. Wells, H.F. Lodish, and Y.I. Henis. 1998. Oligomeric structure of type I and type II transforming growth factor β receptors: homodimers form in the ER and persist at the plasma membrane. *J. Cell Biol.* 140:767–777.
- Gotzmann, J., M. Mikula, A. Eger, R. Schulte-Hermann, R. Foisner, H. Beug, and W. Mikulits. 2004. Molecular aspects of epithelial cell plasticity: implications for local tumor invasion and metastasis. *Mutat. Res.* 566:9–20.
- Hoot, K.E., J. Lighthall, G. Han, S.L. Lu, A. Li, W. Ju, M. Kulesz-Martin, E. Bottinger, and X.J. Wang. 2008. Keratinocyte-specific Smad2 ablation results in increased epithelial-mesenchymal transition during skin cancer formation and progression. *J. Clin. Invest.* 118:2722–2732.
- Hynes, R.O. 2002. Integrins: bidirectional, allosteric signaling machines. *Cell.* 110:673–687.
- Kalluri, R., and E.G. Neilson. 2003. Epithelial-mesenchymal transition and its implications for fibrosis. *J. Clin. Invest.* 112:1776–1784.
- Kim, K.K., M.C. Kugler, P.J. Wolters, L. Robillard, M.G. Galvez, A.N. Brumwell, D. Sheppard, and H.A. Chapman. 2006. Alveolar epithelial cell mesenchymal transition develops in vivo during pulmonary fibrosis and is regulated by the extracellular matrix. *Proc. Natl. Acad. Sci. USA.* 103:13180–13185.
- Kim, K.K., Y. Wei, C. Szekeres, M.C. Kugler, P.J. Wolters, M. Hill, J.A. Frank, A.N. Brumwell, S.E. Wheeler, J.A. Kreidberg, and H.A. Chapman. 2008. Epithelial cell α 3 β 1 integrin links β -catenin and Smad signaling to promote myofibroblast formation and pulmonary fibrosis. *J. Clin. Invest.* .
- Kreidberg, J.A., M.J. Donovan, S.L. Goldstein, H. Rennke, K. Shepherd, R.C. Jones, and R. Jaenisch. 1996. Alpha 3 beta 1 integrin has a crucial role in kidney and lung organogenesis. *Development.* 122:3537–3547.
- Labbe, E., A. Letamendia, and L. Attisano. 2000. Association of Smads with lymphoid enhancer binding factor 1/T cell-specific factor mediates cooperative signaling by the transforming growth factor-beta and wnt pathways. *Proc. Natl. Acad. Sci. USA.* 97:8358–8363.
- Lee, M.K., C. Pardoux, M.C. Hall, P.S. Lee, D. Warburton, J. Qing, S.M. Smith, and R. Derynck. 2007. TGF-beta activates Erk MAP kinase signalling through direct phosphorylation of ShcA. *EMBO J.* 26:3957–3967.
- Lee, Y.I., Y.J. Kwon, and C.K. Joo. 2004. Integrin-linked kinase function is required for transforming growth factor beta-mediated epithelial to mesenchymal transition. *Biochem. Biophys. Res. Commun.* 316:997–1001.
- Lei, S., A. Dubeykovskiy, A. Chakladar, L. Wojtkiewicz, and T.C. Wang. 2004. The murine gastrin promoter is synergistically activated by transforming growth factor-beta/Smad and Wnt signaling pathways. *J. Biol. Chem.* 279:42492–42502.
- Lester, R.D., M. Jo, V. Montel, S. Takimoto, and S.L. Gonias. 2007. uPAR induces epithelial-mesenchymal transition in hypoxic breast cancer cells. *J. Cell Biol.* 178:425–436.
- Lubman, R.L., X.L. Zhang, J. Zheng, L. Ocampo, M.Z. Lopez, S. Veeraraghavan, S.M. Zabski, S.I. Danto, and Z. Borok. 2000. Integrin alpha(3)-subunit expression modulates alveolar epithelial cell monolayer formation. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 279:L183–L193.
- Massague, J., and Y.G. Chen. 2000. Controlling TGF-beta signaling. *Genes Dev.* 14:627–644.
- Masszi, A., L. Fan, L. Rosivall, C.A. McCulloch, O.D. Rotstein, I. Mucsi, and A. Kapus. 2004. Integrity of cell-cell contacts is a critical regulator of TGF-beta 1-induced epithelial-to-myofibroblast transition: role for beta-catenin. *Am. J. Pathol.* 165:1955–1967.
- Nakamura, K., R. Iwamoto, and E. Mekada. 1995. Membrane-anchored heparin-binding EGF-like growth factor (HB-EGF) and diphtheria toxin receptor-associated protein (DRAP27)/CD9 form a complex with integrin α 3 β 1 at cell-cell contact sites. *J. Cell Biol.* 129:1691–1705.
- Nakao, A., E. Roijer, T. Imamura, S. Souchelnytskyi, G. Stenman, C.H. Heldin, and P. ten Dijke. 1997. Identification of Smad2, a human Mad-related protein in the transforming growth factor beta signaling pathway. *J. Biol. Chem.* 272:2896–2900.
- Oloumi, A., T. McPhee, and S. Dedhar. 2004. Regulation of E-cadherin expression and beta-catenin/Tcf transcriptional activity by the integrin-linked kinase. *Biochim. Biophys. Acta.* 1691:1–15.
- Ozdamar, B., R. Bose, M. Barrios-Rodiles, H.R. Wang, Y. Zhang, and J.L. Wrana. 2005. Regulation of the polarity protein Par6 by TGFbeta receptors controls epithelial cell plasticity. *Science.* 307:1603–1609.
- Piedra, J., D. Martinez, J. Castano, S. Miravet, M. Dunach, and A.G. de Herreros. 2001. Regulation of beta-catenin structure and activity by tyrosine phosphorylation. *J. Biol. Chem.* 276:20436–20443.
- Reynolds, L.E., F.J. Conti, R. Silva, S.D. Robinson, V. Iyer, R. Rudling, B. Cross, E. Nye, I.R. Hart, C.M. Dipersio, and K.M. Hodivala-Dilke. 2008. alpha3beta1 integrin-controlled Smad7 regulates reepithelialization during wound healing in mice. *J. Clin. Invest.* 118:965–974.
- Rice, N.A., and L.A. Leinwand. 2003. Skeletal myosin heavy chain function in cultured lung myofibroblasts. *J. Cell Biol.* 163:119–129.
- Roura, S., S. Miravet, J. Piedra, A. Garcia de Herreros, and M. Dunach. 1999. Regulation of E-cadherin/catenin association by tyrosine phosphorylation. *J. Biol. Chem.* 274:36734–36740.
- Runyan, C.E., H.W. Schnaper, and A.C. Poncelet. 2005. The role of internalization in transforming growth factor beta 1-induced Smad2 association with Smad anchor for receptor activation (SARA) and Smad2-dependent signaling in human mesangial cells. *J. Biol. Chem.* 280:8300–8308.
- Savagner, P. 2001. Leaving the neighborhood: molecular mechanisms involved during epithelial-mesenchymal transition. *Bioessays.* 23:912–923.
- Shen, X., P.P. Hu, N.T. Liberati, M.B. Datto, J.P. Frederick, and X.F. Wang. 1998. TGF-beta-induced phosphorylation of Smad3 regulates its interaction with coactivator p300/CREB-binding protein. *Mol. Biol. Cell.* 9:3309–3319.
- Sheppard, D. 2005. Integrin-mediated activation of latent transforming growth factor beta. *Cancer Metastasis Rev.* 24:395–402.
- Thannickal, V.J., D.Y. Lee, E.S. White, Z. Cui, J.M. Larios, R. Chacon, J.C. Horowitz, R.M. Day, and P.E. Thomas. 2003. Myofibroblast differen-

- tiation by transforming growth factor-beta1 is dependent on cell adhesion and integrin signaling via focal adhesion kinase. *J. Biol. Chem.* 278:12384–12389.
- Thiery, J.P., and J.P. Sleeman. 2006. Complex networks orchestrate epithelial-mesenchymal transitions. *Nat. Rev. Mol. Cell Biol.* 7:131–142.
- Tian, Y.C., and A.O. Phillips. 2002. Interaction between the transforming growth factor-beta type II receptor/Smad pathway and beta-catenin during transforming growth factor-beta1-mediated adherens junction disassembly. *Am. J. Pathol.* 160:1619–1628.
- Wang, S.E., F.Y. Wu, I. Shin, S. Qu, and C.L. Arteaga. 2005. Transforming growth factor {beta} (TGF-{beta})-Smad target gene protein tyrosine phosphatase receptor type kappa is required for TGF-{beta} function. *Mol. Cell. Biol.* 25:4703–4715.
- Wang, Z., J.M. Symons, S.L. Goldstein, A. McDonald, J.H. Miner, and J.A. Kreidberg. 1999. (Alpha)3(beta)1 integrin regulates epithelial cytoskeletal organization 3. *J. Cell Sci.* 112:2925–2935.
- Wei, Y., C.H. Tang, Y. Kim, L. Robillard, F. Zhang, M.C. Kugler, and H.A. Chapman. 2007. Urokinase receptors are required for alpha 5 beta 1 integrin-mediated signaling in tumor cells. *J. Biol. Chem.* 282:3929–3939.
- Yook, J.I., X.Y. Li, I. Ota, C. Hu, H.S. Kim, N.H. Kim, S.Y. Cha, J.K. Ryu, Y.J. Choi, J. Kim, et al. 2006. A Wnt-Axin2-GSK3beta cascade regulates Snail1 activity in breast cancer cells. *Nat. Cell Biol.* 8:1398–1406.
- Yue, J., B. Sun, G. Liu, and K.M. Mulder. 2004. Requirement of TGF-beta receptor-dependent activation of c-Jun N-terminal kinases (JNKs)/stress-activated protein kinases (Sapks) for TGF-beta up-regulation of the urokinase-type plasminogen activator receptor. *J. Cell. Physiol.* 199:284–292.
- Zeisberg, M., and R. Kalluri. 2004. The role of epithelial-to-mesenchymal transition in renal fibrosis. *J. Mol. Med.* 82:175–181.
- Zeng, G., U. Apte, A. Micsenyi, A. Bell, and S.P. Monga. 2006. Tyrosine residues 654 and 670 in beta-catenin are crucial in regulation of Met-beta-catenin interactions. *Exp. Cell Res.* 312:3620–3630.
- Zhang, F., C.C. Tom, M.C. Kugler, T.T. Ching, J.A. Kreidberg, Y. Wei, and H.A. Chapman. 2003. Distinct ligand binding sites in integrin $\alpha\beta 1$ regulate matrix adhesion and cell-cell contact. *J. Cell Biol.* 163:177–188.
- Zhang, X.P., W. Puzon-McLaughlin, A. Irie, N. Kovach, N.L. Prokopishyn, S. Laferte, K. Takeuchi, T. Tsuji, and Y. Takada. 1999. Alpha 3 beta 1 adhesion to laminin-5 and invasin: critical and differential role of integrin residues clustered at the boundary between alpha 3 N-terminal repeats 2 and 3. *Biochemistry.* 38:14424–14431.