



Published in final edited form as:

*Neuropsychopharmacology*. 2009 April ; 34(5): 1245–1254. doi:10.1038/npp.2008.171.

## **GABRG1 and GABRA2 as Independent Predictors for Alcoholism in Two Populations**

Mary-Anne Enoch, M.D.<sup>1</sup>, Colin A. Hodgkinson, PhD.<sup>1</sup>, Qiaoping Yuan, PhD.<sup>1</sup>, Bernard Albaugh, MSW.<sup>2</sup>, Matti Virkkunen, M.D.<sup>3</sup>, and David Goldman, M.D.<sup>1</sup>

<sup>1</sup> Laboratory of Neurogenetics, National Institute on Alcohol Abuse and Alcoholism, National Institutes of Health, Bethesda MD 20892 <sup>2</sup> Center for Human Behavior Studies, Inc, Weatherford, OK 73096 <sup>3</sup> Department of Psychiatry, University of Helsinki, Helsinki, Finland

### **Abstract**

The chromosome 4 cluster of GABA<sub>A</sub> receptor genes is predominantly expressed in the brain reward circuitry and this chromosomal region has been implicated in linkage scans for alcoholism. Variation in one chromosome 4 gene, *GABRA2*, has been robustly associated with alcohol use disorders (AUD) although no functional locus has been identified. Since HapMap data reveals moderate long-distance linkage disequilibrium across *GABRA2* and the adjacent gene, *GABRG1*, it is possible that the functional locus is in *GABRG1*. We genotyped 24 SNPs across *GABRG1* and *GABRA2* in two population isolates: 547 Finnish Caucasian men (266 alcoholics) and 311 community-derived Plains Indian men and women (181 alcoholics). In both the Plains Indians and the Caucasians: (a) the *GABRG1* haplotype block(s) did not extend to *GABRA2*; (b) *GABRG1* haplotypes and SNPs were significantly associated with AUD; (c) there was no association between *GABRA2* haplotypes and AUD; (d) there were several common ( $\geq 0.05$ ) haplotypes that spanned *GABRG1* and *GABRA2* (341 kb), three of which were present in both populations: one of these ancestral haplotypes was associated with AUD, the other two were more common in non-alcoholics; this association was determined by *GABRG1*; (e) in the Finns, three less common ( $< 0.05$ ) extended haplotypes showed an association with AUD that was determined by *GABRA2*. Our results suggest that there are likely to be independent, complex contributions from both *GABRG1* and *GABRA2* to alcoholism vulnerability.

### **Keywords**

Linkage disequilibrium; GABAA receptor; ancestral haplotypes; alcohol use disorders; anxiety; American Indians

### **INTRODUCTION**

A substantial body of evidence from preclinical studies has implicated GABA<sub>A</sub> receptors in the acute and chronic effects of ethanol including tolerance, dependence and withdrawal (reviewed in Enoch, 2008; Krystal *et al*, 2006). Rapid synaptic inhibition is mediated through GABA<sub>A</sub> receptors that are ligand-gated, chloride ion channels formed by

---

Corresponding author: Mary-Anne Enoch M.D., NIH/NIAAA/DICBR/LNG, 5625 Fishers Lane, Room 3S32, MSC 9412, Bethesda MD 20892, Tel: 301-496-2727, Fax: 301-480-2839, email: maenoch@niaaa.nih.gov.

#### **DISCLOSURE/CONFLICTS OF INTEREST**

All authors reported no biomedical financial interests or potential conflicts of interest.

pentameric complexes composed of subunits ( $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ ,  $\epsilon$ ,  $\pi$ ,  $\rho$ ), each of which have several isoforms (Barnard *et al*, 1998). Genes for the GABA<sub>A</sub> receptor subunit isoforms are clustered in several chromosomal regions. Expression of the various subunit isoforms varies across brain locations and during development. The mRNAs from the chromosome 4 cluster genes (*GABRA2*, *GABRA4*, *GABRB1*, *GABRG1*) predominate in rat embryo but these genes are generally down-regulated in the adult rat except in the hippocampus and in dopamine neurons in the substantia nigra and the ventral tegmental area (VTA) where they are highly expressed (Okada *et al*, 2004; Steiger and Russek, 2004; Wisden *et al*, 1992). This suggests that the chromosome 4 gene cluster may be implicated in addiction. Indeed, earlier genome-wide scans in American Indians and Caucasians have provided evidence for linkage of alcohol dependence to the location of the chromosome 4 GABA<sub>A</sub> gene cluster (Long *et al*, 1998; Reich *et al*, 1998; Zinn-Justin and Abel, 1999). Subsequently, several studies have found haplotype and SNP associations between *GABRA2* and alcoholism. All these studies, together with HapMap, have identified the same two *GABRA2* haplotype blocks, at least within Caucasians, American Indians and Asians. The significant association signals for alcoholism have been with two common yin-yang haplotypes within the haplotype block that extends downstream from intron 3 in the direction of the adjacent *GABRG1* gene (Agrawal *et al*, 2006; Covault *et al*, 2004; Edenberg *et al*, 2004; Enoch *et al*, 2006a; Fehr *et al*, 2006; Lappalainen *et al*, 2005; Soyka *et al*, 2008). However, other studies have found no *GABRA2* association with alcoholism (Covault *et al*, 2008 (Project MATCH dataset); Drgon *et al*, 2006; Matthews *et al*, 2007).

As yet, no functional *GABRA2* locus has been identified. Moreover, a recent study has shown significant haplotype and SNP association with alcoholism in a haplotype block that extends from the intergenic region between *GABRA2* and *GABRG1* up to *GABRG1* intron 3 in two large groups of U.S. Caucasians but not in African Americans (Covault *et al*, 2008). Thus it is possible that the apparent *GABRA2* association with alcoholism may be due to linkage disequilibrium (LD) with a functional variant in *GABRG1*. The aim of our study was therefore firstly to examine the association between *GABRG1* and alcoholism in Finnish Caucasians and Plains American Indians, two population isolates in which we had previously found a *GABRA2* association with alcoholism that was mediated by anxiety (Enoch *et al*, 2006a). Secondly, since HapMap (<http://www.hapmap.org/>) has identified long-distance LD across the two adjacent genes, we investigated whether there were long-distance haplotypes spanning *GABRA2* and *GABRG1* that were common to both population isolates, i.e. ancestral haplotypes. To this end we genotyped 24 SNPs spanning *GABRG1* from one intergenic region to the other and included two previously genotyped tag SNPs for the *GABRA2* risk haplotypes.

## PARTICIPANTS AND METHODS

### Plains American Indians

Volunteers (311 total: 180 women, 131 men) were recruited from a Plains Indian tribe living in rural Oklahoma. The mean (S.D.) ages of participants were; women: 44.0 (14.9) yrs; men: 41.7 (12.9) yrs. The study was approved by the Plains Indian Tribal Council. Proband were initially ascertained at random from the tribal register and the families of alcoholic probands were extended. Exclusion criteria included a history of brain trauma and neurological diseases, together with current use of psychotropic medications and evidence of alcohol intoxication or withdrawal at the time of testing. The study was carefully explained to participants and written informed consent was obtained according to a human research protocol approved by the human research committee of the National Institute on Alcohol Abuse and Alcoholism (NIAAA), National Institutes of Health (NIH). The protocol and consent forms were also formally approved by the Plains Indian Tribal Council.

Blind-rated DSM-III-R lifetime psychiatric diagnoses (American Psychiatric Association, 1987) were derived from the Schedule for Affective Disorders and Schizophrenia-Lifetime Version (SADS-L). A clinical social worker (B.A.), knowledgeable in the tribe's customs and culture, conducted the SADS-L interviews. The prevalence of lifetime alcohol use disorders (AUD) (nearly all alcohol dependence) in this sample was 73% in men and 47% in women (Enoch *et al*, 2006b) however 63% of the alcoholics had been abstinent for at least one year. The sample included 181 alcoholics (97 men, 84 women) and 130 non-alcoholics (34 men, 96 women).

### Finnish Caucasians

Finnish Caucasians are regarded as a population isolate. The sample from Helsinki, Finland, has been described in detail elsewhere (Lappalainen *et al*, 1998). In total, 547 men were genotyped: 266 alcoholics and 281 non-alcoholics. The sample was comprised of: 173 incarcerated alcoholic criminal offenders, 159 relatives (93 alcoholic, 66 non-alcoholic) and 215 population controls. Of the alcoholics, 217 had alcohol dependence and 49 had alcohol abuse. Mean ages were: alcoholics, 34.4 yrs (SD = 10.4), non-alcoholics: 33.1 yrs (SD = 12.0).

Written informed consent was obtained according to human research protocols approved by the human research committees of NIAAA and the National Institute of Mental Health, NIH, the Department of Psychiatry, University of Helsinki, and the University of Helsinki Central Hospital, Helsinki, Finland.

The Structured Clinical Interview for DSM-III-R (SCID) was administered by psychiatrists to both alcoholics and controls and blind-rated DSM-III-R psychiatric diagnoses were obtained. Individuals with major psychotic episodes were excluded.

### Genotyping

The *GABRG1* genomic region, including 5 kb upstream and 1 kb downstream, was retrieved from NCBI Human Genome Build 35.1. Genotype data from the most diverse population, Africans, were obtained from the HapMap Project Public Release #18 to construct the *GABRG1* haplotype structure using SNP-HAP (<http://www-gene.cimr.cam.ac.uk/clayton/software/snphap.txt>). To select minimum index SNPs that represent maximal haplotype structures information, a program based on a double classification tree search algorithm was used (Zhang *et al*, 2004). On this basis, 24 SNPs spanning *GABRG1* were selected (Figure 1). Genotyping was performed using SNPlex (Tobler *et al*, 2005). Allele-specific probes and optimized multiplexed assays for the 24 SNPs were designed by an automated multi-step pipeline (Applied Biosystems (ABI): <http://www.appliedbiosystems.com>). SNPlex was carried out on fragmented gDNA (88ng/well) according to the manufacturer's protocols. Samples were run on the 3730 DNA Analyzer (ABI) and data were analyzed using Gene Mapper v. 4.0 software (ABI). Genotype analysis was performed based on the SNPlex\_Rules\_3730 method following the factory default rules. Genotyping included 21% duplicates.

In order to determine the haplotype block structure across the 493 kb *GABRG1* and *GABRA2* region, 10 *GABRA2* SNPs from the Illumina Linkage IV-B panel, previously genotyped with the Illumina GoldenGate platform under contract at Illumina (Hodgkinson *et al*, 2008), were included in the analyses. In addition, two previously genotyped *GABRA2* SNPs (rs279863 and rs279858) (Enoch *et al*, 2006a) that tagged the two common *GABRA2* yin-yang haplotypes, were included. These two SNPs, particularly rs279858, have been most strongly associated with alcohol dependence in earlier studies. The inclusion of these two SNP allows for cross comparison with earlier studies. The genotyped *GABRA2* SNPs were

(5' to 3'): rs11503014; haplotype block 1: rs9291283, rs1442060; haplotype block 2: rs10805145, rs426463, rs279827, rs279843, rs279847, rs279858, rs279863, rs519270 and rs693547.

### Assessment of population stratification using ancestry informative markers (AIMS)

The Plains Indian and Finnish Caucasian samples were genotyped for 186 ancestry markers (Hodgkinson *et al*, 2008). These AIMS were also genotyped in 1051 individuals from the 51 worldwide populations represented in the HGDP-CEPH Human Genome Diversity Cell Line Panel (<http://www.cephb.fr/HGDP-CEPH-Panel>). PHASE *Structure 2.2* (<http://pritch.bsd.uchicago.edu/software.html>) was run, simultaneously using the AIMS genotypes from our two samples and the 51 CEPH populations to identify population substructure and compute individual ethnic factor scores. This ancestry assessment showed that within the Plains Indian sample there was on average 0.05 admixture with Caucasians (median value = 0.01) and 0.01 with African Americans. In the Finnish Caucasians, the only detectable admixture was with an Asian factor (mean = 0.07, median = 0.02). Since the ethnic factor scores were highly skewed (due to very low admixture) we used a median split in analyses. Ethnic factor scores were initially tested as covariates in the Plains Indian and Finnish Caucasian analyses but their inclusion had no significant effects on any analyses.

### Statistical Analyses

In both the Plains Indian and the Finnish Caucasian samples, the percentage of genetic identity shared between any two individuals through common descent was calculated for all possible pairs (related and unrelated) using S.A.G.E. (Case Western Reserve University). The average sharing of descent was 0.003 in the Plains Indians, which is less than the degree of relationship between third cousins. The average sharing of descent between any two Finnish Caucasian individuals was only 0.002.

Since the Plains Indians derive from one large pedigree we confirmed the significant p values for associations by performing 10,000 simulations using SIMPED (Leal *et al*, 2005). With this program genotypes can be generated conditional on user-specified frequencies and using the gene-drop method within pedigrees of virtually any size and complexity. Since most of the significant SNPs were in allelic identity, association analysis (t-test between the two homozygous genotypes) was computed for the simulated SNP rs12511372 and a null distribution for the t statistic was generated. This was subsequently used to determine the empirical p value (0.0004) which was close to the p value assuming independence (0.0003). Therefore we were able to undertake analyses that assume independence of individuals.

Haplotype frequencies were estimated using a Bayesian approach implemented with PHASE (Stephens and Donnelly, 2003). Haploview version 2.04 Software (Whitehead Institute for Biomedical Research, USA) was used to produce LD matrices with  $D'$  set to 0.80.

Many of the *GABRG1* SNPs were in allelic identity (Table 1). Twelve SNPs were selected to capture all the allelic variation across *GABRG1* and the intergenic regions. The two *GABRA2* SNPs that tag the *GABRA2* risk haplotypes were included. PHASE was employed to determine long-distance haplotypes across the two gene regions using the 14 SNPs indicated in Table 1.

## RESULTS

### *GABRG1* SNP associations

Allele frequencies in both samples were very similar across *GABRG1* however they began to differ in the intergenic region and differed markedly in *GABRA2* (Table 1). In both the

Plains Indians and the Finns, *GABRG1* SNPs extending from intron 1 to downstream from the 3' UTR region were significantly associated with AUD (Table 1). In the Finns, SNPs in the *GABRG1* promoter and the intergenic region between *GABRG1* and *GABRA2* were also associated with AUD. The allelic associations were in the same direction in both samples.

## ***GABRG1* haplotype associations**

### **Plains Indians**

There was one haplotype block extending from rs12511372 in intron 1 of *GABRG1* to rs1504501 distal to the gene, a distance of 155 kb (Figure 2). Twelve of the 16 SNPs within the haplotype block showed significant associations with AUD (Table 1). There were 7 haplotypes with  $\geq 0.01$  frequency that accounted for 0.93 of the haplotype diversity. From Figure 2 it can be seen that there are two cladistic groupings of haplotypes: H1 – H4 and H5 – H7. Haplotypes H1 – H4 were more common in alcoholics than non-alcoholics (0.54 vs 0.41). The other 3 haplotypes (H5 – H7) were more abundant in non-alcoholics compared with alcoholics (0.59 vs 0.46): (H1 – H4) vs (H5 – H7):  $\chi^2 = 8.8$ , 1 df,  $p = 0.003$ ; global analysis:  $\chi^2 = 21.4$ , 6 df,  $p = 0.002$ .

### **Finnish Caucasians**

There were two *GABRG1* haplotype blocks with a recombination region in intron 2 (Figure 3). Block 1 extended from the *GABRA2-GABRG1* intergenic SNP rs4695146 to the intron 2 SNP rs17536211, a distance of 83.2 kb. There were 9 haplotypes ( $\geq 0.01$  frequency) that accounted for 0.96 of the haplotype diversity. From Figure 3 it can be seen that haplotypes H1 and H2 differed by only one allele and were more abundant in alcoholics than non-alcoholics (0.45 vs 0.36). In contrast, haplotypes H8 and H9, differing by only one allele, were more common in non-alcoholics compared with alcoholics (0.49 vs 0.41): (H1 + H2) vs (H8 + H9):  $\chi^2 = 8.3$ , 1 df,  $p = 0.0039$ .

Block 2 extended from the intron 2 SNP rs13130508 to the intergenic SNP rs1394344, a distance of 113.7 kb. There were 6 haplotypes with a frequency of  $\geq 0.01$  that accounted for 0.98 of the haplotype diversity. From Figure 3 it can be seen that haplotypes could be cladistically arranged into two groups: H1 – H3, that were more abundant in alcoholics (0.52 vs 0.43) and H4 – H6 that were more common in non-alcoholics (0.57 vs 0.48): (H1 – H3) vs (H4 – H6):  $\chi^2 = 8.1$ , 1 df,  $p = 0.0045$ .

***GABRA2* SNP and haplotype associations**—Three SNPs, rs494270 (intergenic) and the *GABRA2* tag SNPs rs279863 and rs279858, were located within the *GABRA2* haplotype block 2 that has been associated with AUD in previous studies (Table 1). SNPs rs279863 and rs279858, which are in allelic identity, showed a significant genotypic association with AUD due to the fact that alcoholics had an excess of both homozygotes: 11 (0.44 vs 0.34) and 22 (0.18 vs 0.16), and a corresponding deficit of heterozygotes (0.38 vs 0.50) compared with non-alcoholics, respectively. There were no significant differences in allele frequencies.

The Plains Indians had three common haplotypes that accounted for 0.99 of the haplotype diversity: 122 (0.64), 211 (0.25) and 111 (0.10). The Finnish Caucasians had only two haplotypes that accounted for 0.93 of the haplotype diversity: 211 (0.57) and 122 (0.36). There was no difference in haplotype frequency distribution between alcoholics and non-alcoholics in the Plains Indians ( $p = 0.68$ , 2 df) and the Finns ( $p = 0.39$ , 1 df).

**Long-distance haplotypes across *GABRG1* – *GABRA2*: influence of *GABRG1* on association with AUD**—Figure 4 shows that in the Finnish Caucasians and Plains

American Indians there is a degree of LD extending from *GABRA2* block 2 to downstream of *GABRG1*. For example,  $D'$  between rs1504501 (in the intergenic region distal to *GABRG1*) and *GABRA2* block 2 SNPs was 0.6 in the Finns and 0.4 in the Plains Indians.

Fourteen SNPs (12 *GABRG1* and 2 *GABRA2*, identified in Table 1) were included in haplotype analyses across *GABRG1* and *GABRA2* block 2, a distance of more than 354 kb, in order to determine extended haplotypes.

### Finnish Caucasians

There were 13 long-distance haplotypes with  $\geq 0.01$  frequency that accounted for 0.89 of the haplotype diversity. Of these, five haplotypes had a frequency of  $\geq 0.05$  (H1: 0.30; H2: 0.07; H3: 0.11; H4: 0.16, H5: 0.07), accounting for 0.71 of the haplotype diversity in the total sample. For simplicity, analyses were conducted with the haplotypes  $\geq 0.05$  frequency. Analyses undertaken with these five haplotypes showed that H1 was more abundant in alcoholics, H3 – H5 were more common in non-alcoholics and H2 was neutral: H1 vs rest:  $\chi^2 = 7.3$ , 1 df,  $p = 0.007$  (Figure 5).

Figure 5 illustrates the relative contributions from *GABRG1* and *GABRA2* haplotypes to the extended haplotypes' association with AUD. H1: both the *GABRG1* block 1 and block 2 haplotypes are increased in alcoholics, therefore extended haplotype H1 is increased in alcoholics. H2: the *GABRG1* block 2 haplotype is increased in alcoholics, the block 1 haplotype is decreased in alcoholics therefore extended haplotype H2 is neutral to AUD diagnosis. Haplotypes H3 and H4 are less abundant in alcoholics: in both cases the *GABRG1* block 1 and block 2 haplotypes are decreased in alcoholics. Haplotype H5 is numerically increased in non-alcoholics: the *GABRG1* block 2 haplotype is neutral; the block 1 haplotype is more common in non-alcoholics. As discussed above, *GABRA2* haplotypes were not associated with AUD and, as seen in Figure 5, did not contribute to the extended haplotypes' association with AUD.

### Plains Indians

Nine haplotypes had a frequency of  $\geq 0.01$  and accounted for 0.90 of the haplotype diversity. Of these, six haplotypes had a frequency of  $\geq 0.05$ : H1: 0.16; H2: 0.09; H5: 0.18; H6: 0.22; H3: 0.06; H4: 0.09, accounting for 0.80 of haplotype diversity in the total sample. For simplicity, analyses were conducted with the haplotypes  $\geq 0.05$  frequency. As can be seen from Figure 5, there are two clastic groupings of haplotypes, (H1, H2, H5) and (H3, H4, H6). Haplotypes H1, H2, H5 were more abundant in alcoholics compared with non-alcoholics (0.58 vs 0.43); haplotypes H3, H4, H6 were more common in non-alcoholics (0.57 vs 0.42):  $\chi^2 = 10.4$ , 1 df,  $p = 0.0013$ .

Only one *GABRG1* haplotype contributes to H1, H2 and H5: this *GABRG1* haplotype is associated with AUD. The one *GABRG1* haplotype that contributes to H6, H3 and H4 is more abundant in non-alcoholics. In contrast, the *GABRA2* yin yang haplotypes 211 and 122 are found in the long distance haplotypes that are associated both with AUD and without AUD; likewise for the intergenic haplotypes 111 and 222.

**Ancestral Haplotypes**—The Finns and Plains Indians have in common three long-distance haplotypes: H1, H3 and H4 (see Figure 5). These ancestral haplotypes accounted for 0.57 of the long-distance haplotype diversity in Finns and 0.31 of diversity in the Plains. In both populations, H1 is associated with AUD and H3 and H4 are more abundant in non-alcoholics (Fig 5). Haplotypes H3 and H4 have very nearly opposite configuration to haplotype H1.

**Long-distance haplotypes across *GABRG1* – *GABRA2*: influence of *GABRA2* on association with AUD**—Two *GABRG1* SNPs that showed exceptionally high LD across the whole *GABRG1* –*GABRA2* region in both samples: rs13130508 ( $D' = 0.9 - 1.0$ ) and rs10517150 ( $D' = 0.9 - 1.0$ ), were included in the group of 14 SNPs (Table 1) selected to detect extended haplotypes.

In the Finns, the variant alleles of these two *GABRG1* SNPs were found on three extended haplotypes (total frequency = 0.06) that only differed in the *GABRA2* block 2 SNPs: haplotypes 22111221121122 (0.01); 22111221121211 (0.02); and 22111221121222 (0.03). These haplotypes differed in their association with AUD. Two haplotypes were more common in alcoholics than in non-alcoholics: 22111221121122 (0.29 vs 0.12) and 22111221121222 (0.57 vs 0.44). In contrast, 22111221121211 was more common in non-alcoholics compared with alcoholics (0.44 vs 0.14);  $\chi^2 = 6.8$ , 2 df,  $p = 0.034$ .

In the Plains Indians, the variant alleles of the *GABRG1* SNPs rs13130508 and rs10517150 were found on only one extended haplotype: 22111221222122 (frequency 0.04) that was not the same as any of the Finnish haplotypes.

## DISCUSSION

We found substantial, significant *GABRG1* haplotype and SNP associations with AUD in two independent populations. The ten SNPs that showed significant associations with AUD were in approximate allelic identity in both populations (non-alcoholics: Plains Indians,  $MAF = 0.40 - 0.46$ ; Finns,  $MAF = 0.35 - 0.50$ ) suggesting selection pressure. In the Plains Indians the significant SNP associations were located within the one haplotype block in which *GABRG1* (excluding the promoter region) is located. However in the Finns nearly all the SNPs were associated with AUD in a region that extended from the distal intergenic region, across *GABRG1* and up to the *GABRA2* haplotype block. This suggests that the functional locus may be found within a long-distance haplotype. Moreover, the examination of more population isolates might help to narrow down the genomic interval in which the functional variant(s) resides.

In contrast to our findings with *GABRG1* we did not find a *GABRA2* haplotype association with AUD in either sample. In a previous analysis in these two populations we had found a complex *GABRA2* association with AUD in men only that was mediated by trait anxiety (Enoch et al, 2006a). The observed findings in our current study suggest that the association signal coming from *GABRG1* is more robust than that from *GABRA2*.

The Finnish Caucasian and Plains Indian haplotype block structure and extended LD patterns across *GABRG1* and *GABRA2* were similar to that found in the HapMap European American and Asian populations and in a recent study of 13 SNPs covering the region between *GABRG1* intron 7 and *GABRA2* intron 3 (Ittiwut *et al*, 2007). It has been shown that LD between distant markers is due to the presence of extended haplotype superblocks in individuals with ancient chromosomes which have escaped historic recombination (Buzas *et al*, 2004). When we looked at extended haplotypes with frequencies  $\geq 0.05$ , we found three long-distance haplotypes that were common to both populations with total frequencies that were lower in the Plains Indians (0.31) than in the Finns: (0.57). The Plains Indians experienced a bottleneck in the recent genetic past therefore lower frequencies of ancestral haplotypes might be expected. The ancient superblock structure has been partially disrupted by recombination in the region between *GABRG1* and *GABRA2*, and also in the Caucasian *GABRG1* intron 2.

The most common of the three ancestral, extended haplotypes was associated with AUD in both populations; the other two ancestral haplotypes (identical except for the *GABRA2* contribution) were more abundant in non-alcoholics. Our results suggest that the signal for the extended haplotypes' association with AUD derives from *GABRG1* and is independent of *GABRA2* (Figure 5). However the differential impact of *GABRA2* haplotypes was observed in the less common (< 0.05 frequency) long distance haplotypes. We had noted that in both samples, two *GABRG1* SNPs: rs13130508 and rs10517150 (not associated with AUD), showed exceptionally high LD ( $D' = 0.9 - 1.0$ ) with *GABRA2* SNPs indicating the likelihood that they are located in ancient chromosomal regions. However, the variant alleles of these two SNPs were not embedded in the three common ancestral haplotypes described above. Instead, in the Finns they were found in three haplotypes that differed only in the *GABRA2* haplotype contribution and it was these *GABRA2* haplotypes that determined the haplotype association with AUD. These three ancestral haplotypes account for only 0.06 of the extended haplotype diversity and therefore they are unlikely to contribute significantly to the previously reported AUD associations with the two common yin yang *GABRA2* haplotypes in Caucasians (Agrawal *et al*, 2006; Covault *et al*, 2004; Edenberg *et al*, 2004; Enoch *et al*, 2006a; Fehr *et al*, 2006; Lappalainen *et al*, 2005; Soyka *et al*, 2008).

Edenberg *et al* (2004) genotyped six *GABRG1* SNPs in the family-based COGA dataset extending from the 5' region to intron 8. Their strongest association with alcohol dependence was with an intron 1 SNP rs1391175 ( $p = 0.05$ ). We genotyped two of these six SNPs: for the intron 1 SNP rs2221020 Edenberg *et al*. found a trend association with alcohol dependence ( $p = 0.07$ ) in contrast to our significant associations with AUD in both populations ( $p < 0.01$ ). For the intron 8 SNP rs1497570, Edenberg *et al*. found no association ( $p = 0.65$ ) in contrast to our significant results in both samples ( $p < 0.05$ ). On the other hand, as in our study Covault *et al* (2008) have shown significant haplotype and SNP association with alcoholism in a haplotype block that extends from *GABRG1* intron 2 to the intergenic region between *GABRG1* and *GABRA2* in two large groups of U.S. Caucasians (Covault *et al*, 2008). They genotyped 15 SNPs extending from *GABRA2* intron 3 to *GABRG1* intron 3 and found, as we did, that although the *GABRA2* SNPs were located in a distinct haplotype block from *GABRG1* SNPs, there was moderate LD between markers in the two blocks. Covault *et al* (2008) used only 4 SNPs to identify *GABRG1* haplotypes in contrast to our 11 SNPs, nevertheless both our studies have identified *GABRG1* haplotype associations with alcoholism. Our study expanded on this earlier result by demonstrating in two independent, ethnically diverse populations that the association between AUD and *GABRG1* extended across a region that included all of *GABRG1* and the downstream intergenic region.

As in our study, Covault *et al* (2008) found that in the extended haplotypes, the AUD risk was conferred by the *GABRG1* and not the *GABRA2* haplotype components. Analyses of a best-fit genetic model for markers in the two haplotype block regions led Covault *et al* (2008) to conclude that there may be a separate contribution to risk for AD by *GABRG1* and *GABRA2* and that their prior findings of a *GABRA2* association with alcoholism (Covault *et al*, 2004) might be partly due to LD with functional genetic variants in *GABRG1*.

As yet, there is no known functional locus in either *GABRA2* or *GABRG1*. Although *GABRA2* has four common isoforms (Tian *et al*, 2005) *GABRG1* has no known isoforms. There is a microRNA binding site in the *GABRG1* 3' UTR, however no SNPs overlap with that site. The fact that the chr 4 genes cluster together and that there is extensive LD across the cluster suggests the presence of distant enhancers and repressors. Using NCBI and HapMap we determined that the nearest gene, *GNPDA2* (glucosamine-6-phosphate deaminase 2), is at a distance of more than 1 Mb and several major haplotype blocks downstream from *GABRG1*. Moreover, long-distance LD does not extend downstream from



the *GABRG1* haplotype block. Therefore it seems likely that the functional locus is located in the, admittedly extensive, region covered by the extended haplotypes. We used HapMap haplotypes for the Caucasian, Asian and African samples to determine the two major ancestral *GABRG1* haplotypes (H1 and H3 + H4 as shown in Figure 5) and created the haplotype specific sequences from NCBI refGene's *GABRG1* transcript (NM\_173536.3). We ran Mfold 3.2 (Zuker *et al*, 1999) with the haplotype specific transcript sequences and computed ten different foldings for each haplotype. There were no significant differences in the folding energy between the two ancestral haplotypes ( $p = 0.187$ ).

*GABRG1* has until recently been in the shadow of its neighbor, *GABRA2*, and has not been considered a prime candidate gene for alcoholism. In the rat, *GABRG1* is expressed in only a few brain regions, primarily in the amygdala and all areas receiving innervation from the striatum including the substantia nigra (Pirker *et al*, 2000; Schwarzer *et al*, 2001). These are regions that are implicated in reward and addiction. The GABA<sub>A</sub> receptor subunit composition determines distinct pharmacological and electrophysiological properties. Studies have shown that  $\gamma_1$  subunits co-assemble with  $\alpha_2$  subunits in vivo largely as  $\alpha_2\beta_1\gamma_1$  receptors (Whiting, 2003) that are much less responsive to the effects of benzodiazepines than receptors with the much more common  $\gamma_2$  subunit (Wafford *et al*, 1993). In contrast, the anxiolytic effects of benzodiazepines and barbiturates appear to be mediated in part by  $\alpha_2$  subunits co-assembled with  $\gamma_2$  subunits (Dixon *et al*, 2008; Low *et al*, 2000). Thus it is possible that  $\alpha_2\beta_1\gamma_1$  receptors may be implicated in alcoholism vulnerability per se whereas variation in the more common receptors with co-assembled  $\alpha_2$  and  $\gamma_2$  subunits may predispose to alcoholism mediated by anxiety.

In conclusion, our similar results in two independent populations, Plains Indians and Finnish Caucasians, have identified a complex situation: (a) *GABRG1* haplotypes and SNPs are significantly associated with AUD; (b) *GABRA2* haplotypes are not associated with AUD except, as previously shown (Enoch *et al*, 2006a), when mediated by trait anxiety; (c) there are three common ancestral haplotypes that span *GABRG1* and *GABRA2*; their association with AUD is determined by *GABRG1* (d) in the Finns, the association of three less common ( $< 0.05$ ) long-distance haplotypes with AUD is determined by *GABRA2*. Our results suggest that there are likely to be independent contributions from both *GABRG1* and *GABRA2* to the risk for AUD.

## Acknowledgments


This research was supported by the Intramural Research Program of the National Institute on Alcohol Abuse and Alcoholism, NIH, and in part by the Office of Research on Minority Health.

## References


- Agrawal A, Edenberg HJ, Foroud T, Bierut LJ, Dunne G, Hinrichs AL, et al. Association of GABRA2 with drug dependence in the collaborative study of the genetics of alcoholism sample. *Behav Genet* 2006;36:640–650. [PubMed: 16622805]
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 3. American Psychiatric Association Press; Washington, DC: 1987.
- Barnard EA, Skolnick P, Olsen RW, Mohler H, Sieghart W, Biggio G, et al. International Union of Pharmacology. XV. Subtypes of gamma-aminobutyric acidA receptors: classification on the basis of subunit structure and receptor function. *Pharmacol Rev* 1998;50:291–313. [PubMed: 9647870]
- Buzas B, Belfer I, Hipp H, Lorincz I, Evans C, Phillips G, et al. Haplotype block and superblock structures of the alpha1-adrenergic receptor genes reveal echoes from the chromosomal past. *Mol Genet Genomics* 2004;272:519–529. [PubMed: 15503142]

- Covault J, Gelernter J, Hesselbrock V, Nellissery M, Kranzler HR. Allelic and haplotypic association of GABRA2 with alcohol dependence. *Am J Med Genet (Neuropsychiatr Genet)* 2004;129B:104–109.
- Covault J, Gelernter J, Jensen K, Anton R, Kranzler HR. Markers in the 5'-Region of GABRG1 Associate to Alcohol Dependence and are in Linkage Disequilibrium with Markers in the Adjacent GABRA2 Gene. *Neuropsychopharmacology* 2008;33:837–848. [PubMed: 17507911]
- Dixon CI, Rosahl TW, Stephens DN. Targeted deletion of the GABRA2 gene encoding alpha2-subunits of GABA(A) receptors facilitates performance of a conditioned emotional response, and abolishes anxiolytic effects of benzodiazepines and barbiturates. *Pharmacol Biochem Behav.* 2008 Jan 31; [Epub ahead of print].
- Drgon T, D'Addario C, Uhl GR. Linkage disequilibrium, haplotype and association studies of a chromosome 4 GABA receptor gene cluster: candidate gene variants for addictions. *Am J Med Genet B Neuropsychiatr Genet* 2006;141:854–860. [PubMed: 16894595]
- Edenberg HJ, Dick DM, Xuei X, Tian H, Almasy L, Bauer LO, et al. Variations in GABRA2, encoding the alpha 2 subunit of the GABA(A) receptor, are associated with alcohol dependence and with brain oscillations. *Am J Hum Genet* 2004;74:705–714. [PubMed: 15024690]
- Enoch MA, Schwartz L, Albaugh B, Virkkunen M, Goldman D. Dimensional anxiety mediates linkage of GABRA2 haplotypes with alcoholism. *Am J Med Genet B Neuropsychiatr Genet* 2006a; 141:599–607.
- Enoch M-A, Waheed J, Harris CR, Albaugh B, Goldman D. Sex Differences in the Influence of COMT Val158Met on Alcoholism and Smoking in Plains American Indians. *Alcohol Clin Exp Res* 2006b;30:399–406. [PubMed: 16499480]
- Enoch M-A. The role of GABA(A) receptors in the development of alcoholism. *Pharmacol Biochem Behav* 2008;90:95–104. [PubMed: 18440057]
- Fehr C, Sander T, Tadic A, Lenzen KP, Anghelescu I, Klawe C, et al. Confirmation of association of the GABRA2 gene with alcohol dependence by subtype-specific analysis. *Psychiatr Genet* 2006;16:9–17. [PubMed: 16395124]
- Hodgkinson CA, Yuan Q, Xu K, Shen PH, Heinz E, Lobos EA, et al. Addictions Biology: Haplotype-Based Analysis for 130 Candidate Genes on a Single Array. *Alcohol Alcohol.* 2008 May 12; [Epub ahead of print].
- Ittiwut C, Listman J, Mutirangura A, Malison R, Covault J, Kranzler HR, et al. Interpopulation linkage disequilibrium patterns of GABRA2 and GABRG1 genes at the GABA cluster locus on human chromosome 4. *Genomics* 2008;91:61–69. [PubMed: 17976953]
- Krystal JH, Staley J, Mason G, Petrakis IL, Kaufman J, Harris RA, et al. Gamma-aminobutyric acid type A receptors and alcoholism: intoxication, dependence, vulnerability, and treatment. *Arch Gen Psychiatry* 2006;63:957–968. [PubMed: 16952998]
- Lappalainen J, Long JC, Eggert M, Ozaki N, Robin RW, Brown GL, et al. Linkage of antisocial alcoholism to the serotonin 5-HT1B receptor gene in 2 populations. *Arch Gen Psychiatry* 1998;55:989–994. [PubMed: 9819067]
- Lappalainen J, Krupitsky E, Remizov M, Pchelina S, Taraskina A, Zvartau E, et al. Association between alcoholism and gamma-amino butyric acid alpha2 receptor subtype in a Russian population. *Alcohol Clin Exp Res* 2005;29:493–498. [PubMed: 15834213]
- Leal SM, Yan K, Muller-Myhsok B. SimPed: a simulation program to generate haplotype and genotype data for pedigree structures. *Hum Hered* 2005;60:119–122. [PubMed: 16224189]
- Long JC, Knowler WC, Hanson RL, Robin RW, Urbanek M, Moore E, et al. Evidence for genetic linkage to alcohol dependence on chromosomes 4 and 11 from an autosome-wide scan in an American Indian population. *Am J Med Genet* 1998;81:216–221. [PubMed: 9603607]
- Low K, Crestani F, Keist R, Benke D, Brunig I, Benson JA, et al. Molecular and neuronal substrate for the selective attenuation of anxiety. *Science* 2000;290:131–134. [PubMed: 11021797]
- Matthews AG, Hoffman EK, Zezza N, Stiffler S, Hill SY. The Role of the GABRA2 Polymorphism in Multiplex Alcohol Dependence Families With Minimal Comorbidity: Within-Family Association and Linkage Analyses. *J Stud Alcohol Drugs* 2007;68:625–633. [PubMed: 17690794]
- Okada H, Matsushita N, Kobayashi K, Kobayashi K. Identification of GABAA receptor subunit variants in midbrain dopaminergic neurons. *J Neurochem* 2004;89:7–14. [PubMed: 15030384]

- Pirker S, Schwarzer C, Wieselthaler A, Sieghart W, Sperk G. GABA(A) receptors: immunocytochemical distribution of 13 subunits in the adult rat brain. *Neuroscience* 2000;101:815–850. [PubMed: 11113332]
- Reich T, Edenberg HJ, Goate A, Williams JT, Rice JP, Van Eerdewegh P, et al. Genome-wide search for genes affecting the risk for alcohol dependence. *Am J Med Genet* 1998;81:207–215. [PubMed: 9603606]
- Schwarzer C, Berresheim U, Pirker S, Wieselthaler A, Fuchs K, Sieghart W, Sperk G. Distribution of the major gamma-aminobutyric acid(A) receptor subunits in the basal ganglia and associated limbic brain areas of the adult rat. *J Comp Neurol* 2001;433:526–549. [PubMed: 11304716]
- Soyka M, Preuss UW, Hesselbrock V, Zill P, Koller G, Bondy B. GABA-A2 receptor subunit gene (GABRA2) polymorphisms and risk for alcohol dependence. *J Psychiatr Res* 2008;42:184–191. [PubMed: 17207817]
- Steiger JL, Russek SJ. GABAA receptors: building the bridge between subunit mRNAs, their promoters, and cognate transcription factors. *Pharmacol Ther* 2004;101:259–281. [PubMed: 15031002]
- Stephens M, Donnelly P. A comparison of Bayesian methods for haplotype reconstruction from population genotype data. *Am J Hum Genet* 2003;73:1162–1169. [PubMed: 14574645]
- Tian H, Chen H-J, Cross TH, Edenberg HJ. Alternative splicing and promoter use in the human GABRA2 gene. *Brain Res Mol Brain Res* 2005;137:174–183. [PubMed: 15950776]
- Tobler A, Short S, Andersen M, Paner T, Briggs J, Lambert S, et al. The SNPlex genotyping system: a flexible and scalable platform for SNP genotyping. *J Biomol Tech* 2005;16:398–406. [PubMed: 16522862]
- Wafford KA, Bain CJ, Whiting PJ, Kemp JA. Functional comparison of the role of gamma subunits in recombinant human gamma-aminobutyric acidA/benzodiazepine receptors. *Mol Pharmacol* 1993;44:437–442. [PubMed: 8102787]
- Whiting PJ. GABA-A receptor subtypes in the brain: a paradigm for CNS drug discovery? *Drug Discov Today* 2003;8:445–450. [PubMed: 12801796]
- Wisden W, Laurie DJ, Monyer H, Seeburg PH. The distribution of 13 GABAA receptor subunit mRNAs in the rat brain. I Telencephalon, diencephalon, mesencephalon. *J Neurosci* 1992;12:1040–1062. [PubMed: 1312131]
- Zhang P, Sheng H, Uehara R. A double classification tree search algorithm for index SNP selection. *BMC Bioinformatics* 2004;5:89. [PubMed: 15238162]
- Zinn-Justin A, Abel L. Genome search for alcohol dependence using the weighted pairwise correlation linkage method: interesting findings on chromosome 4. *Genet Epidemiol* 1999;17(Suppl 1):S421–426. [PubMed: 10597473]
- Zuker, M.; Mathews, DH.; Turner, DH. Algorithms and Thermodynamics for RNA Secondary Structure Prediction: A Practical Guide. In: Barciszewski, J.; Clark, BFC., editors. *RNA Biochemistry and Biotechnology*. Kluwer Academic Publishers; Dordrecht, NL: 1999. p. 11–43. NATO ASI Series



**FIGURE 1. GABRG1 Gene Map and Haplotype Block Structure in HapMap Populations**  
 The positions of the genotyped SNPs are indicated. CDS = coding sequences, linked by dotted lines to the corresponding exons. The haplotype block structures of the three HapMap populations: CEU (Caucasian); YRI (African) and ASN (Chinese/Japanese) are provided. Conservation across 17 vertebrate species, from zebrafish to humans is indicated.




A

Haplotype Frequencies

		Non-Alc		Alc															
B	1	1	1	2	1	2	2	2	2	2	2	1	1	1	2	1	H1	0.17	0.25
	2	1	1	1	2	1	2	2	2	2	2	1	1	1	2	1	H2	0.01	0.03
	2	1	1	1	2	1	2	1	2	2	2	1	1	1	2	1	H3	0.04	0.02
	2	1	1	1	2	2	2	2	2	2	2	1	1	1	2	1	H4	0.19	0.24
	2	1	2	2	1	1	1	2	1	2	1	2	1	2	1	2	H5	0.07	0.03
	2	2	2	2	1	1	1	2	1	2	1	2	1	2	1	2	H6	0.49	0.37
	2	2	2	2	1	1	1	2	1	1	1	1	2	2	2	1	H7	0.03	0.06

**FIGURE 2. *GABRG1* Haplotype Block Structure and Haplotypes in Plains Indians**  
 A: The haplotype block extends from rs12511372 in intron 1 to rs1504501 distal to the gene (155 kb). The darker the color of squares, the greater the degree of LD ( $D'$ ). B: Cladistic groupings of haplotypes with freq  $\geq 0.01$  in total sample: increased frequency of group H1 – H4 in alcoholics; increased frequency of group H5 – H7 in non-alcoholics;  $p = 0.003$ . Alleles 1 and 2 are located on opposite DNA strands.



**FIGURE 3. GABRG1 Haplotype Block Structure and Haplotypes in Finnish Caucasians**  
 Haplotype block 1 extends 83 kb from the intergenic SNP rs4695146 to the intron 2 SNP rs17536211. Haplotype block 2 extends 114 kb from intron 2 SNP 13130508 to the downstream SNP rs1394344. The darker the color of squares, the greater the degree of LD (D').

Cladistic groupings of haplotypes with freq  $\geq 0.01$  in total sample: Block 2: increased frequency of group H1 – H3 in alcoholics; increased frequency of group H4 – H6 in non-alcoholics;  $p = 0.005$ . Block 1: increased frequency of group H1 – H2 in alcoholics; increased frequency of group H8 – H9 in non-alcoholics;  $p = 0.004$ . Alleles 1 and 2 are located on opposite DNA strands.

GABRG1

GABRA2






FIGURE 4. Long Distance Linkage Disequilibrium across *GABRG1* and *GABRA2*



**FIGURE 5. The Association of Long-distance Haplotypes across *GABRG1* and *GABRA2* with Alcohol Use Disorders**

See table 1 for the 14 SNPs used to derive these long-distance haplotypes that are at least 341 kb long. There are three haplotypes (H1, H3, H4) that are identical in both populations. H1 is associated with AUD in both populations. H3 and H4 are more abundant in non-alcoholics from both populations.

Upward arrow indicates increased haplotype frequency in alcoholics, downward arrow indicates decreased frequency in alcoholics, '=' indicates no association with AUD. Arrows and '=' on left indicate direction of *GABRG1* haplotypes' association with AUD. Arrows and '=' on right indicate the direction of *GABRG1-GABRA2* long-distance haplotypes' association with AUD.

Alleles 1 and 2 are located on opposite DNA strands.



TABLE 1

*GABRG1* and *GABRA2* SNP Associations with Alcohol Use Disorders

SNPs	Alleles 1 – 2	Location	Plains Indians, N = 311			Finnish Men, N = 547		
			Alcoholics Allele 1	Non- Alcoholics Frequency	Genotype P Value	Alcoholics Allele 1	Non- Alcoholics Frequency	Genotype P Value
<b><i>GABRG1</i></b>								
rs1504497	A – G	intergenic	0.46	0.54	0.056	0.58	0.65	<b>0.046</b>
rs4695138	A – T	intergenic	0.78	0.73	0.720	0.56	0.48	<b>0.022</b>
rs1504501	C – T	intergenic	0.25	0.17	0.054	0.25	0.21	0.060
rs1394344*	C – T	intergenic	0.56	0.47	<b>0.010</b>	0.61	0.54	<b>0.031</b>
rs1497568	A – G	3' UTR	0.53	0.4	<b>0.001</b>	0.52	0.44	<b>0.020</b>
rs6447493*	C – T	3' UTR	0.53	0.4	<b>0.001</b>	0.52	0.44	<b>0.012</b>
rs1497570*	A – C	intron 8	0.48	0.59	<b>0.002</b>	0.48	0.57	<b>0.018</b>
rs1566857	A – T	intron 7	0.76	0.81	0.052	0.96	0.99	–
rs10032631	A – G	intron 7	0.47	0.6	<b>0.002</b>	0.48	0.56	<b>0.028</b>
rs993677	C – T	intron 6	0.03	0.04	0.286	0.12	0.11	0.478
rs976156	C – T	exon 3	0.47	0.59	<b>0.001</b>	0.58	0.65	<b>0.030</b>
rs13130508*	C – T	intron 2	0.07	0.03	0.120	0.06	0.07	0.830
rs17536211	A – C	intron 2	0.98	0.97	–	0.81	0.76	0.137
rs1497575	A – G	intron 2	0.48	0.6	<b>0.0003</b>	0.55	0.63	<b>0.017</b>
rs6824361*	C – T	intron 2	0.59	0.44	<b>0.0002</b>	0.52	0.44	<b>0.035</b>
rs10517150*	A – G	intron 1	0.93	0.96	0.184	0.93	0.92	0.816
rs2221020	C – T	intron 1	0.52	0.4	<b>0.003</b>	0.45	0.36	<b>0.006</b>
rs6813633*	A – G	intron 1	0.41	0.55	<b>0.0003</b>	0.44	0.53	<b>0.014</b>
rs12511372*	A – G	intron 1	0.60	0.46	<b>0.0006</b>	0.59	0.50	<b>0.009</b>
rs17639557*	C – T	promoter	0.74	0.68	0.616	0.59	0.50	<b>0.019</b>
rs17536530	A – C	promoter	1	1	–	0.89	0.93	0.079
rs10938427*	A – G	intergenic	0.74	0.69	0.500	0.48	0.39	<b>0.007</b>
rs4695146*	A – G	intergenic	0.74	0.67	0.353	0.60	0.51	<b>0.013</b>
rs494270*	A – T	intergenic	0.73	0.72	0.970	0.37	0.42	<b>0.025</b>

SNPs	Alleles 1 – 2	Location	Plains Indians, N = 311			Finnish Men, N = 547		
			Alcoholics Allele 1	Non- Alcoholics Frequency	Genotype P Value	Alcoholics Allele 1	Non- Alcoholics Frequency	Genotype P Value
<i>GABRA2</i>								
rs279863*	A – C	intron 5	0.36	0.35	0.261	0.62	0.59	<b>0.021</b>
rs279858*	G – A	exon 5	0.36	0.36	0.106	0.63	0.59	<b>0.022</b>

\* SNPs used to determine ancestral haplotypes.

Alleles 1 and 2 are located on opposite DNA strands