

# A Signaling Pathway Linking Nitric Oxide Production to Heterotrimeric G Protein and Hydrogen Peroxide Regulates Extracellular Calmodulin Induction of Stomatal Closure in Arabidopsis<sup>1[W]</sup>

Jian-Hua Li<sup>2</sup>, Yin-Qian Liu<sup>2</sup>, Pin Lü, Hai-Fei Lin, Yang Bai, Xue-Chen Wang, and Yu-Ling Chen\*

Hebei Key Laboratory of Molecular and Cellular Biology, College of Life Science, Hebei Normal University, Shijiazhuang 050016, China (J.-H.L., Y.-Q.L., H.-F.L., Y.B., Y.-L.C.); School of Basic Medical Sciences, Hebei Medical University, Shijiazhuang 050017, China (P.L.); and National Laboratory of Plant Physiology and Biochemistry, College of Biological Sciences, China Agricultural University, Beijing 100094, China (X.-C.W.)

Extracellular calmodulin (ExtCaM) regulates stomatal movement by eliciting a cascade of intracellular signaling events including heterotrimeric G protein, hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), and Ca<sup>2+</sup>. However, the ExtCaM-mediated guard cell signaling pathway remains poorly understood. In this report, we show that Arabidopsis (*Arabidopsis thaliana*) NITRIC OXIDE ASSOCIATED1 (AtNOA1)-dependent nitric oxide (NO) accumulation plays a crucial role in ExtCaM-induced stomatal closure. ExtCaM triggered a significant increase in NO levels associated with stomatal closure in the wild type, but both effects were abolished in the *Atnoa1* mutant. Furthermore, we found that ExtCaM-mediated NO generation is regulated by GPA1, the G $\alpha$ -subunit of heterotrimeric G protein. The ExtCaM-dependent NO accumulation was nullified in *gpa1* knockout mutants but enhanced by overexpression of a constitutively active form of GPA1 (cG $\alpha$ ). In addition, cG $\alpha$  *Atnoa1* and *gpa1-2 Atnoa1* double mutants exhibited a similar response as did *Atnoa1*. The defect in *gpa1* was rescued by overexpression of *AtNOA1*. Finally, we demonstrated that G protein activation of NO production depends on H<sub>2</sub>O<sub>2</sub>. Reduced H<sub>2</sub>O<sub>2</sub> levels in guard cells blocked the stomatal response of cG $\alpha$  lines, whereas exogenously applied H<sub>2</sub>O<sub>2</sub> rescued the defect in ExtCaM-mediated stomatal closure in *gpa1* mutants. Moreover, the *atrbobD/F* mutant, which lacks the NADPH oxidase activity in guard cells, had impaired NO generation in response to ExtCaM, and H<sub>2</sub>O<sub>2</sub>-induced stomatal closure and NO accumulation were greatly impaired in *Atnoa1*. These findings have established a signaling pathway leading to ExtCaM-induced stomatal closure, which involves GPA1-dependent activation of H<sub>2</sub>O<sub>2</sub> production and subsequent AtNOA1-dependent NO accumulation.

Plant guard cells control opening and closure of the stomata in response to phytohormones (e.g. abscisic acid [ABA]) and various environmental signals such as light and temperature, thereby regulating gas exchange for photosynthesis and water status via transpiration (Schroeder et al., 2001). Cytosolic calcium ([Ca<sup>2+</sup>]<sub>i</sub>) has been shown to be a key second messenger that changes in response to multiple stimuli in guard cells (McAinsh et al., 1995; Grabov and Blatt, 1998; Wood et al., 2000). A large proportion of Ca<sup>2+</sup> is localized in extracellular space. It has been shown that external Ca<sup>2+</sup> concentration ([Ca<sup>2+</sup>]<sub>o</sub>) promotes sto-

matal closure and induces oscillation in [Ca<sup>2+</sup>]<sub>i</sub> in guard cells (MacRobbie, 1992; McAinsh et al., 1995; Allen et al., 2001). However, how the guard cells perceive [Ca<sup>2+</sup>]<sub>o</sub> concentration and convert [Ca<sup>2+</sup>]<sub>o</sub> changes into [Ca<sup>2+</sup>]<sub>i</sub> changes was not understood until a calcium-sensing receptor (CAS) in the plasma membrane of guard cells in Arabidopsis (*Arabidopsis thaliana*) was identified (Han et al., 2003). The external Ca<sup>2+</sup> (Ca<sup>2+</sup><sub>o</sub>)-induced [Ca<sup>2+</sup>]<sub>i</sub> increase is abolished in CAS antisense lines (Han et al., 2003). Both [Ca<sup>2+</sup>]<sub>o</sub> and [Ca<sup>2+</sup>]<sub>i</sub> show diurnal oscillation that is determined by stomatal conductance, whereas the amplitude of [Ca<sup>2+</sup>]<sub>i</sub> oscillation is reduced in CAS antisense lines (Tang et al., 2007). The reduced amplitude of [Ca<sup>2+</sup>]<sub>i</sub> diurnal oscillation in response to Ca<sup>2+</sup><sub>o</sub> treatment suggests the potential existence of other [Ca<sup>2+</sup>]<sub>o</sub> sensor(s) that may transmit [Ca<sup>2+</sup>]<sub>o</sub> information into the [Ca<sup>2+</sup>]<sub>i</sub> response in coordination with CAS. Extracellular calmodulin (ExtCaM) could be such an additional [Ca<sup>2+</sup>]<sub>o</sub> sensor.

Calmodulin is a well-known Ca<sup>2+</sup> sensor that is activated upon binding of Ca<sup>2+</sup>. It has been shown that calmodulin exists not only intracellularly but also extracellularly in many plant species (Biro et al., 1984; Sun et al., 1994, 1995; Cui et al., 2005). ExtCaM has been implicated in several important biological

<sup>1</sup> This work was supported by the National Science Foundation of China (grant no. 30670173 to Y.-L.C.), the Doctor Foundation of Hebei Province (grant no. B2003107 to Y.-L.C.), Hebei Normal University (grant no. B2002213 to Y.-L.C.), and the Science Foundation of Hebei Normal University (grant no. L2006Y06 to J.-H.L.).

<sup>2</sup> These authors contributed equally to the article.

\* Corresponding author; e-mail yulingchen@mail.hebtu.edu.cn.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors ([www.plantphysiol.org](http://www.plantphysiol.org)) is: Yu-Ling Chen (yulingchen@mail.hebtu.edu.cn).

<sup>[W]</sup> The online version of this article contains Web-only data.

[www.plantphysiol.org/cgi/doi/10.1104/pp.109.137067](http://www.plantphysiol.org/cgi/doi/10.1104/pp.109.137067)

functions, such as the promotion of cell proliferation, pollen germination, and tube growth (Sun et al., 1994, 1995; Ma and Sun, 1997; Ma et al., 1999; Cui et al., 2005; Shang et al., 2005). ExtCaM is found in the cell wall of guard cells in *Vicia faba* and in the epidermis of Arabidopsis by immunogold labeling/electron microscopy and western-blot analyses, respectively, and the endogenous CaM in the extracellular space has been shown to regulate stomatal movements (Chen et al., 2003; Xiao et al., 2004). Under natural conditions, once the activity of ExtCaM has been inhibited by its membrane-impermeable antagonist W7-agrose or CaM antibody, stomatal opening under light is enhanced and stomatal closure in darkness is inhibited in *V. faba* and Arabidopsis (Chen et al., 2003; Xiao et al., 2004).  $[Ca^{2+}]_i$  and cytosolic hydrogen peroxide ( $H_2O_2$ ) changes, two events involved in ExtCaM-regulated stomatal movement (Chen et al., 2004), are likely regulated by light/darkness (Chen and Gallie, 2004; Tang et al., 2007), suggesting that ExtCaM plays an important physiological role in the regulation of stomatal diurnal rhythm. Calmodulin-binding proteins have been found in the protoplast of suspension-cultured Arabidopsis cells, supporting the idea that ExtCaM functions as a peptide-signaling molecule (Cui et al., 2005). Furthermore, ExtCaM triggers  $[Ca^{2+}]_i$  elevation in guard cells of *V. faba* and Arabidopsis and in lily (*Lilium daviddi*) pollen (Chen et al., 2004; Xiao et al., 2004; Shang et al., 2005). These observations support the notion that ExtCaM could be a potential  $[Ca^{2+}]_o$  sensor for external calcium, and this external calcium sensing could subsequently regulate the  $[Ca^{2+}]_i$  level through a signaling cascade.

It is interesting that ExtCaM and ABA induce some parallel changes in second messengers in guard cell signaling. Our previous studies show that ExtCaM induces  $[Ca^{2+}]_i$  increase and  $H_2O_2$  generation through the  $G\alpha$ -subunit (GPA1) of a heterotrimeric G protein, and increased  $H_2O_2$  further elevates  $[Ca^{2+}]_i$  (Chen et al., 2004). G protein,  $Ca^{2+}$ , and  $H_2O_2$  are well-known second messengers in ABA-induced guard cell signaling (McAinsh et al., 1995; Grabov and Blatt, 1998; Pei et al., 2000; Wang et al., 2001; Zhang et al., 2001; Liu et al., 2007). However, the signaling cascade triggered by ExtCaM in guard cells is poorly understood. New ABA signaling components in guard cells could provide a clue in the study of the molecular mechanism of ExtCaM guard cell signaling.

Recently, nitric oxide (NO) has been shown to serve as an important signal molecule involved in many aspects of developmental processes, including floral transition, root growth, root gravitropism, adventitious root formation, xylogenesis, seed germination, and orientation of pollen tube growth (Beligni and Lamattina, 2000; Pagnussat et al., 2002; He et al., 2004; Prado et al., 2004; Gabaldón et al., 2005; Stohr and Stremmler, 2006). Increasing evidence points to a role for NO as an essential component in ABA signaling in guard cells (Garcia-Mata and Lamattina, 2001, 2002; Neill et al., 2002). It has been shown that nitrate


reductase (NR) reduces nitrite to NO, and the *nia1*, *nia2* NR-deficient mutant in Arabidopsis showed reduced ABA induction of stomatal closure (Desikan et al., 2002; Bright et al., 2006). Although animal nitric oxide synthase (NOS) activity has been detected in plants and inhibitors of mammalian NOS impair NO production in plants (Barroso et al., 1999; Corpas et al., 2001), the gene(s) encoding NOS in plants is still not clear. *AtNOS1* in Arabidopsis was initially reported to encode a protein containing NOS activity (Guo et al., 2003). However, recent studies have raised critical questions regarding the nature of *AtNOS1* and suggested that *AtNOS1* appears not to encode a NOS (Crawford et al., 2006; Zemojtel et al., 2006). However, the originally described *Atnos1* mutant is deficient in NO accumulation (Crawford et al., 2006). Consequently, *AtNOS1* was renamed *AtNOA1* (for NITRIC OXIDE ASSOCIATED1; Crawford et al., 2006). Therefore, the *Atnoa1* mutant provides a useful tool for dissecting the function of NO in plants. At present, the molecules that regulate NO generation in ABA-mediated guard cell signaling are not clear. Evidence suggests that  $H_2O_2$ , a second messenger important for the regulation of many developmental processes and stomatal movement (Pei et al., 2000; Zhang et al., 2001; Coelho et al., 2002; Demidchik et al., 2003; Kwak et al., 2003), regulates NO generation in guard cells (Lum et al., 2002; He et al., 2005; Bright et al., 2006).

Given the parallel signaling events induced by ABA and ExtCaM, we investigated whether NO is involved in the regulation of ExtCaM-induced stomatal closure in Arabidopsis and whether it is linked to G protein and  $H_2O_2$ , two key regulators of both ExtCaM and ABA regulation of stomatal movements. Using Arabidopsis mutants (e.g. *GPA1* null mutants, the NO-producing mutant *Atnoa1*, and the guard cell  $H_2O_2$  synthetic enzymatic mutant *atrbohD/F*) combined with pharmacological analysis, we present compelling evidence to establish a linear functional relationship between  $G\alpha$ ,  $H_2O_2$ , and NO in ExtCaM guard cell signaling.

## RESULTS

### AtNOA1-Dependent NO Accumulation Is Required for ExtCaM-Induced Stomatal Closure

To investigate a potential role for NO in ExtCaM-induced stomatal closure, we examined the effect of the NO scavenger 2-(4-carboxyphenyl)-4,4,5,5-tetramethylimidazole-1-oxyl 3-oxide potassium salt (cPTIO) on ExtCaM action. Both ExtCaM and the NO donor sodium nitroprusside (SNP) induced stomatal closure (Fig. 1A), as demonstrated previously (Garcia-Mata and Lamattina, 2001; Chen et al., 2004). cPTIO greatly reduced ExtCaM-induced stomatal closure compared with ExtCaM-treated stomata ( $P < 0.001$ ; Fig. 1A). We then monitored changes in NO levels in response to applied ExtCaM using the NO-specific



**Figure 1.** AtNOA1-dependent NO production regulates ExtCaM-induced stomatal closure. A, Open stomata in the wild type (WT) were incubated in MES buffer (control),  $10^{-8}$  M CaM,  $10^{-8}$  M  $\alpha$ -lactalbumin,  $60 \mu\text{M}$  SNP,  $200 \mu\text{M}$  cPTIO, or  $10^{-8}$  M CaM plus  $200 \mu\text{M}$  cPTIO. B, Open stomata in the wild type, *Atnoa1*, and rescued *Atnoa1* were incubated in MES buffer (control) or  $10^{-8}$  M CaM. Stomatal apertures were measured 2 h after treatment. C and D, ExtCaM induces a NO increase in guard cells in the wild type and rescued *Atnoa1* but fails in the *Atnoa1* mutant. Open stomata in each genotype were incubated in  $10 \mu\text{M}$  DAF2-DA buffer for 15 min and then incubated in MES buffer (control),  $10^{-8}$  M  $\alpha$ -lactalbumin,  $1 \text{ mM}$   $\text{CaCl}_2$ ,  $10^{-8}$  M CaM, or  $10^{-8}$  M CaM plus  $200 \mu\text{M}$  cPTIO for 30 min in darkness. Fluorescence images and intensities in guard cells were recorded and are shown in C and D, respectively. E, Open stomata in the wild type were incubated in  $10 \mu\text{M}$  4AF-DA, a negative control of DAF2-DA, for 15 min and then incubated in MES buffer or  $10^{-8}$  M CaM for 30 min in darkness, and fluorescence images in guard cells were recorded. Bars =  $10 \mu\text{m}$ .

fluorescent dye diaminofluorescein diacetate (DAF2-DA; Garcia-Mata and Lamattina, 2002; Guo et al., 2003). ExtCaM treatment induced a dramatic increase in DAF2-DA staining in guard cells compared with the untreated control ( $P < 0.001$ ), and cPTIO significantly reduced ExtCaM-induced NO generation in guard cells ( $P < 0.001$ ; Fig. 1, C and D). The same concentration of  $\alpha$ -lactalbumin, which contained an EF-hand like structure (Iyer and Qasba, 1999), had no effect on stomatal movement or NO levels in guard cells (Fig. 1, A, C, and D), indicating that induction of ExtCaM on stomatal closure and NO generation is CaM specific and is not induced by any other member of the EF-hand family of proteins. Application of calcium also caused NO synthesis, as did ExtCaM, supporting the possibility that ExtCaM transmits extracellular calcium changes into intracellular response (Fig. 1, C and D). Guard cells in wild-type (ecotype Columbia [Col-0]) epidermis were loaded with 4-aminofluorescein diacetate (4AF-DA), a negative control of DAF2-DA, and fluorescence intensity was examined when treated with ExtCaM. Fluorescence of 4AF in guard cells did

not increase upon ExtCaM treatment (Fig. 1E). These results support an essential role for NO in regulating ExtCaM-induced stomatal closure.

We next determined whether the *Atnoa1* mutant, which is defective in NO accumulation (Guo et al., 2003), exhibited different ExtCaM-induced guard cell responses. NO levels in guard cells and stomatal apertures in the *Atnoa1* mutant and rescued *Atnoa1* line were examined following ExtCaM treatment. ExtCaM-induced stomatal closure was impaired significantly in *Atnoa1* compared with the ExtCaM-treated wild type ( $P < 0.001$ ; Fig. 1B), consistent with the blocking effect of the NO scavenger cPTIO on ExtCaM-induced stomatal closure ( $P < 0.001$ ). These results were supported by NO changes: ExtCaM-induced NO accumulation was almost completely abolished in the *Atnoa1* mutant ( $P > 0.05$ ; Fig. 1, C and D). Both stomatal closure and NO accumulation upon ExtCaM treatment in the rescued *Atnoa1* line were similar to that of the wild type ( $P > 0.05$ ; Fig. 1, B–D). These data indicate that AtNOA1-mediated NO accumulation plays an essential role in ExtCaM-induced stomatal closure.

### G Protein $\alpha$ Subunit Modulates AtNOA1-Dependent NO Production in ExtCaM-Induced Stomatal Closure


The GPA1 G protein  $\alpha$  subunit ( $G\alpha$ ) has been shown to modulate ABA regulation of stomatal opening (Wang et al., 2001) and closure (Liu et al., 2007) and ExtCaM induction of stomatal closure (Chen et al., 2004). To investigate whether  $G\alpha$  also regulates ExtCaM induction of NO generation, we examined NO levels in guard cells of the *gpa1* null mutants and of transgenic lines expressing a constitutively active form of GPA1 (*cG $\alpha$* ; Okamoto et al., 2001; Wang et al., 2001). NO levels in the guard cells of two independent *cG $\alpha$*  lines were higher than that of the wild type (ecotype Wassilewskija [Ws]) in response to ExtCaM ( $P < 0.001$ ), but ExtCaM failed to induce NO accumulation in *gpa1-1* and *gpa1-2* mutants ( $P > 0.05$ ; Fig. 2, A and B; data from one line of *cG $\alpha$*  are shown). These results clearly indicate that  $G\alpha$  is required for ExtCaM induction of NO generation. These data are consistent with the faster response of *cG $\alpha$*  stomata to ExtCaM and the impaired stomatal closure of *gpa1* with ExtCaM treatment, as reported previously (Chen et al., 2004). Furthermore, the NO donor SNP induced stomatal closure in *gpa1-1* and *gpa1-2* mutants ( $P < 0.001$ ; Fig. 2C), further supporting that NO acts downstream of  $G\alpha$  during ExtCaM-induced stomatal closure.

To further test the role of  $G\alpha$  in the regulation of NO generation in ExtCaM guard cell signaling, we performed crosses between the *cG $\alpha$* 1 line (Okamoto et al., 2001) and the *Atnoa1* mutant and between *gpa1-2* and *Atnoa1* and examined stomatal responses and NO levels in guard cells in the double mutants. As shown in Figure 3, *cG $\alpha$* 1 *Atnoa1* stomata did not close when treated with ExtCaM ( $P > 0.05$ ; Fig. 3A) and ExtCaM failed to arouse NO production in guard cells of *cG $\alpha$* 1 *Atnoa1* ( $P > 0.05$ ; Fig. 3, B and C). The NO donor SNP rescued the defects of *cG $\alpha$* 1 *Atnoa1* in stomatal closure and NO generation, suggesting that the defects of *cG $\alpha$* 1 *Atnoa1* are due to a lack of NO production (Fig. 3).

At the same time, ExtCaM-induced stomatal response and NO level in the *gpa1-2 Atnoa1* double mutant were similar to those in both *gpa1-2* and *Atnoa1* mutants ( $P > 0.05$ ; Fig. 3), indicating that GPA1 and AtNOA1 act in the same pathway. Furthermore, we introduced the 35S: *AtNOA1* gene into the *gpa1-2* mutant. NO level and stomatal response following ExtCaM treatment were examined in three independent transgenic lines, and one set of representative data is shown. Stomatal closure and NO generation in guard cells in *gpa1-2 AtNOA1* were similar to those of the wild type when treated with ExtCaM ( $P > 0.05$ ; Fig. 3). Taken together, our results clearly demonstrate that the GPA1 G protein acts upstream of NO production to activate ExtCaM-induced stomatal closure.

### Essential Role of H<sub>2</sub>O<sub>2</sub> Generation in $G\alpha$ -Induced NO Production in ExtCaM Guard Cell Signaling

Pharmacological evidence supports a role for H<sub>2</sub>O<sub>2</sub> in the mediation of ExtCaM-induced stomatal closure




**Figure 2.** GPA1 modulates NO synthesis in ExtCaM guard cell signaling. A and B, Guard cells preloaded with 10  $\mu\text{M}$  DAF2-DA in the wild type (WT/WS), *gpa1* mutants, and the *cG $\alpha$*  line were incubated in MES buffer (control) or 10<sup>-8</sup> M CaM for 30 min in darkness. Fluorescence images and intensities in guard cells were recorded and are shown in A and B, respectively. Bar = 10  $\mu\text{m}$ . C, Open stomata in the wild type, *gpa1-1*, and *gpa1-2* were incubated in MES buffer (control) or 60  $\mu\text{M}$  SNP. Stomatal apertures were measured 2 h after treatment. WS, Ecotype Ws.


in *V. faba* (Chen et al., 2004). In this study, we vigorously tested the role of H<sub>2</sub>O<sub>2</sub> in Arabidopsis using both pharmacological and genetic approaches. Stomata of *atrbohD/F* failed to close upon ExtCaM treatment, resembling the response of the wild type (Col-0) pretreated with catalase (CAT), a H<sub>2</sub>O<sub>2</sub> scavenger ( $P > 0.05$ ; Fig. 4A). Furthermore, we measured H<sub>2</sub>O<sub>2</sub> changes using the fluorescent probe 2',7'-dichlorofluorescein diacetate (H<sub>2</sub>DCFDA). As shown in Figure 4, ExtCaM induced a dramatic increase in H<sub>2</sub>O<sub>2</sub> level in wild-type guard cells ( $P < 0.001$ ) but not in *atrbohD/F* guard cells,

while H<sub>2</sub>O<sub>2</sub> level in the control guard cells did not change significantly during the recording period (Fig. 4, B and C). These results demonstrate that ExtCaM-induced stomatal closure depends on its activation of H<sub>2</sub>O<sub>2</sub> production in guard cells.

We next sought to determine whether Gα and H<sub>2</sub>O<sub>2</sub> act in the same signaling pathway triggered by ExtCaM. We examined whether *gpa1* and *cGα* affect



**Figure 3.** The effect of GPA1 depends on AtNOA1-mediated NO production. **A**, Open stomata in the wild type (WT), *cGα1 Atnoa1* and *gpa1-2 Atnoa1* double mutants, and *gpa1-2 AtNOA1* lines were incubated in MES buffer (control), 10<sup>-8</sup> M CaM, or 60 μM SNP. Stomatal apertures were measured 2 h after treatment. **B** and **C**, Guard cells preloaded with 10 μM DAF2-DA in the wild type, *cGα1 Atnoa1* and *gpa1-2 Atnoa1* double mutants, and *gpa1-2 AtNOA1* lines were incubated in MES buffer (control), 10<sup>-8</sup> M CaM, or 60 μM SNP for 30 min in darkness. Fluorescence images and intensities in guard cells were recorded and are shown in **B** and **C**, respectively. Bar = 10 μm.




**Figure 4.** Impaired H<sub>2</sub>O<sub>2</sub> synthesis blocks ExtCaM-induced stomatal closure. **A**, Open stomata in the wild type (WT) and *atrbohD/F* were incubated in MES buffer (control), 10<sup>-8</sup> M CaM, or 10<sup>-8</sup> M CaM plus 100 units mL<sup>-1</sup> CAT. Stomatal apertures were measured 2 h after treatment. **B** and **C**, Fluorescence images (**B**) and a quantitative curve of fluorescence intensity (**C**) show that ExtCaM induces an H<sub>2</sub>O<sub>2</sub> increase in the wild type but fails in *atrbohD/F*. H<sub>2</sub>O<sub>2</sub> changes in guard cells preloaded with 50 μM H<sub>2</sub>DCFDA were measured with a confocal laser scanning microscope, and 10<sup>-8</sup> M CaM or MES buffer (control) was added at 0 s, as indicated in **C**. Both fluorescence intensities and images of the guard cells were recorded every 15 s. The experiments were repeated at least three times with seven to 10 cells each time, and one time set of data is presented. Bars = 10 μm.


H<sub>2</sub>O<sub>2</sub> generation in response to an ExtCaM stimulus. As shown in Figure 5, both diphenylene iodonium (DPI), an inhibitor of NADPH oxidases, and CAT blocked

the stomatal response to ExtCaM in *cGα* lines as well as in the wild type (Ws;  $P > 0.05$ ; Fig. 5A; data from one *cGα* line are shown). *cGα* lines have been reported to show faster stomatal closure in response to ExtCaM, although they exhibit the same final stomatal apertures as ExtCaM-treated wild-type controls (Chen et al., 2004). Therefore, the large final stomatal aperture in *cGα* lines under the treatment of ExtCaM together with DPI or CAT demonstrates that the effect of higher  $Gα$  activity depends on the generation of  $H_2O_2$ . Furthermore, exogenously applied  $H_2O_2$  induced stomatal closure in *gpa1* mutants ( $P < 0.001$ ; Fig. 5B), indicating that action of  $H_2O_2$  does not depend on the activity of  $Gα$ . These results suggest that  $H_2O_2$  acts downstream of  $Gα$ .

We next investigated whether  $H_2O_2$  is essential for NO generation by examining the NO level in the *atrobhD/F* double mutant upon treatment with ExtCaM. ExtCaM failed to induce NO generation in guard cells of *atrobhD/F* or the wild type (Col-0) incubated with CAT ( $P > 0.05$ ; Fig. 6, A and B). The NO donor SNP induced stomatal closure of *atrobhD/F* ( $P < 0.001$ ; Fig.




**Figure 5.**  $H_2O_2$  is required for action of *cGα* in ExtCaM guard cell signaling. A, Open stomata in the wild type (WT/WS) and *cGα* lines were incubated in MES buffer (control),  $10^{-8}$  M CaM,  $10^{-8}$  M CaM plus  $10 \mu M$  DPI, or  $10^{-8}$  M CaM plus 100 units  $mL^{-1}$  CAT. B, Open stomata in the wild type, *gpa1-1*, and *gpa1-2* were incubated in MES buffer (control) or  $5 \times 10^{-5}$  M  $H_2O_2$ . Stomatal apertures were measured 2 h after treatment. WS, Ecotype Ws.



**Figure 6.**  $H_2O_2$  generation is required for NO production in ExtCaM guard cell signaling. A and B, ExtCaM induces NO synthesis in the wild type (WT) but fails in *atrobhD/F* and the wild type incubated with CAT. Guard cells preloaded with  $10 \mu M$  DAF2-DA in the wild type and *atrobhD/F* were incubated in MES buffer (control),  $10^{-8}$  M CaM, or  $10^{-8}$  M CaM plus 100 units  $mL^{-1}$  CAT for 30 min in darkness. Fluorescence images and intensities in the guard cells were recorded and are shown in A and B, respectively. Bar =  $10 \mu m$ . C, Open stomata in the wild type and *atrobhD/F* were incubated in MES buffer (control) or  $60 \mu M$  SNP. Stomatal apertures were measured 2 h after treatment.

6C), supporting that NO acts downstream of  $H_2O_2$ . These results indicate that  $H_2O_2$  is essential for NO generation induced by ExtCaM. Based on these results and our previous finding that  $Gα$  is required for ExtCaM-induced  $H_2O_2$  generation in guard cells (Chen et al., 2004), we speculate that  $Gα$ -mediated NO production depends on  $H_2O_2$  generation induced by ExtCaM.



**Figure 7.** H<sub>2</sub>O<sub>2</sub>-induced stomatal closure depends on AtNOA1-mediated NO production. A, Open stomata in the wild type (WT), rescued *Atnoa1*, and *Atnoa1* were incubated in MES buffer (control),  $5 \times 10^{-5}$  M H<sub>2</sub>O<sub>2</sub>, or  $5 \times 10^{-5}$  M H<sub>2</sub>O<sub>2</sub> plus 200  $\mu$ M cPTIO. Stomatal apertures were measured 2 h after treatment. B, Guard cells preloaded with 10  $\mu$ M DAF2-DA in the wild type, rescued *Atnoa1*, and *Atnoa1* were incubated in MES buffer (control),  $5 \times 10^{-5}$  M H<sub>2</sub>O<sub>2</sub>, or  $5 \times 10^{-5}$  M H<sub>2</sub>O<sub>2</sub> plus 200  $\mu$ M cPTIO for 30 min in darkness, and fluorescence intensities in guard cells were recorded. ExtCaM-induced H<sub>2</sub>O<sub>2</sub> generation does not depend on AtNOA1-mediated NO production. C, Guard cells preloaded with 50  $\mu$ M H<sub>2</sub>DAFDA in the wild type and *Atnoa1* were incubated in MES buffer (control),  $10^{-8}$  M CaM, or  $10^{-8}$  M CaM plus 200

### AtNOA1-Dependent NO Accumulation Is Crucial for H<sub>2</sub>O<sub>2</sub>-Induced Stomatal Closure

Since interruption of H<sub>2</sub>O<sub>2</sub> generation blocked ExtCaM-induced NO generation, we further investigated the relationship between H<sub>2</sub>O<sub>2</sub> and NO. H<sub>2</sub>O<sub>2</sub> induced stomatal closure in the wild type (Col-0) control and the rescued *Atnoa1* line but not in the wild type incubated with cPTIO ( $P > 0.05$ ); meanwhile, H<sub>2</sub>O<sub>2</sub> failed to induce stomatal closure in the *Atnoa1* mutant ( $P > 0.05$ ; Fig. 7A). H<sub>2</sub>O<sub>2</sub> induced NO generation in guard cells of the wild type and the rescued *Atnoa1* line, and cPTIO reduced NO generation that was induced by H<sub>2</sub>O<sub>2</sub> ( $P < 0.001$ ); H<sub>2</sub>O<sub>2</sub> also stimulated NO production in *Atnoa1*, whereas the level was greatly lower than that in the H<sub>2</sub>O<sub>2</sub>-treated wild type ( $P < 0.001$ ; Fig. 7B). However, ExtCaM induced H<sub>2</sub>O<sub>2</sub> generation both in *Atnoa1* and in the wild type incubated with cPTIO ( $P > 0.05$ ; Fig. 7C). These results suggest that H<sub>2</sub>O<sub>2</sub>-induced stomatal closure mainly depends on AtNOA1-mediated NO generation.

### DISCUSSION

In this report, we have provided convincing evidence for a new signaling pathway controlling stomatal movement. Our combined genetic, pharmacological, and biochemical analyses show that ExtCaM stimulates AtNOA1-dependent production of NO, which acts as a second messenger to activate stomatal closure. Furthermore, we found that ExtCaM-triggered NO production is mediated by the activation of G $\alpha$  of the heterotrimeric G protein in Arabidopsis. We also showed that the modulation of NO production by G $\alpha$  requires NADPH oxidase-dependent H<sub>2</sub>O<sub>2</sub> generation. Therefore, together with previous findings (Chen et al., 2004; Joo et al., 2005; Bright et al., 2006), our results support an ExtCaM-activated guard cell signaling pathway that includes a cascade of ExtCaM, heterotrimeric G protein, NADPH oxidase, and AtNOA1-dependent NO generation.

### NO Plays an Important Role in ExtCaM Guard Cell Signaling

The role of NO in the regulation of stomatal closure has been well documented. The NO donor SNP was reported to induce stomatal closure in *V. faba*, *Salpiglossa organifolia*, and *Tradescantia* species and to reduce transpiration in wheat (*Triticum aestivum*; Garcia-Mata and Lamattina, 2001). A requirement of NO in ABA-mediated stomatal closure has also been found in pea (*Pisum sativum*) and Arabidopsis (Desikan et al., 2002; Neill et al., 2002; Bright et al., 2006). Our results in this report provide convincing evidence that NO plays an important role in the regulation of ExtCaM-induced

$\mu$ M cPTIO for 30 min in darkness, and fluorescence intensities in guard cells were recorded.

stomatal closure. The NO scavenger cPTIO reversed ExtCaM-induced stomatal closure (Fig. 1A), indicating that NO is required for ExtCaM-induced stomatal closure. The apertures of open stomata measured in this study are consistent with those in reports by Guo et al. (2003) and Liu et al. (2007) but somewhat larger than those reported by Desikan et al. (2002) and Bright et al. (2006). This discrepancy may be due to the different concentrations of KCl in the buffers used in these studies. The concentration of KCl in the MES/KCl buffer is 5 mM in the methods described by Desikan et al. (2002) and Bright et al. (2006) and 50 mM in the methods described by Liu et al. (2007). DAF2-DA is a NO-specific fluorescent dye. Using DAF2-DA dye, our results indicated that ExtCaM induced NO production in Arabidopsis (Fig. 1, C and D), further supporting the essential role of NO in ExtCaM signaling.

In animal systems, it has been reported that extracellular calcium induces NO production (Miles et al., 1998). In this study, we have checked NO levels after treatment with extracellular calcium, and the results indicated that NO levels in the guard cells of the wild type increased when treated with extracellular calcium (Fig. 1, C and D), suggesting that ExtCaM may act as an extracellular calcium sensor. We propose that G protein, H<sub>2</sub>O<sub>2</sub>, and NO play essential roles in transducing the extracellular calcium signal.


NR has been demonstrated to be an enzyme that mediates NO synthesis in plants. NR-mediated NO generation has also been suggested to regulate stomatal closure induced by ABA (Desikan et al., 2002; Bright et al., 2006). Although it has been shown that the NOS inhibitor nitro-L-Arg methyl ester suppresses stomatal closure and NO generation in guard cells of pea in response to ABA (Neill et al., 2002), the gene(s) encoding NOS in plants is not clear. AtNOA1 was initially isolated as a NOS in Arabidopsis (Guo et al., 2003), whereas the AtNOA1 protein showed no NOS activity in recent studies (Crawford et al., 2006; Zemojtel et al., 2006). Interestingly, AtNOA1 contains a GTP-binding domain and has been shown to be a member of the circularly permuted GTPase family for RNA/ribosome binding and to be involved in ribosome assembly (Moreau et al., 2008). Although it is not clear how AtNOA1 affects NO accumulation indirectly, the *Atnoa1* mutant shows impaired stomatal closure and less NO accumulation in response to ABA and salt stress (Guo et al., 2003; Zhao et al., 2007). In this study, we showed that ExtCaM-induced NO accumulation requires AtNOA1: ExtCaM failed to induce stomatal closure and NO generation in the guard cells in *Atnoa1*, consistent with the blocking effect of cPTIO on ExtCaM-induced stomatal closure and NO production (Fig. 1). Thus, our pharmacological and genetic results suggest that NO is an essential molecule in ExtCaM-induced stomatal closure and that NO accumulation induced by ExtCaM depends on AtNOA1.

AtNOA1 was found to be targeted to mitochondria in Arabidopsis root (Guo and Crawford, 2005). How-

ever, recent evidence shows that AtNOA1 localizes in chloroplast (Flores-Pérez et al., 2008; Gas et al., 2009). Organelle localization of AtNOA1 suggests that activation of this protein requires signal transduction from extracellular stimuli to intracellular organelle. In this study, we show that AtNOA1-dependent NO accumulation in response to ExtCaM requires the activation of G protein and the accumulation of H<sub>2</sub>O<sub>2</sub>.

#### Modulation of NO Generation by G Protein in ExtCaM-Induced Stomatal Closure

Involvement of heterotrimeric G proteins in guard cell signaling has been supported by genetic evidence from null mutants for GPA1 and the G protein-coupled receptors GCR1 and GCR2 (Wang et al., 2001; Pandey and Assmann, 2004; Liu et al., 2007). Guard cells in *gpa1* mutants were insensitive to ABA inhibition of inward K<sup>+</sup> currents and pH independent of ABA activation of anion channels (Wang et al., 2001) and induction of stomatal closure (Liu et al., 2007). It is speculated that G protein regulates ion channels directly in guard cells, whereas modulation of radical oxygen species (ROS) by G protein has been found both in ExtCaM-induced stomatal closure and in ozone stress responses (Chen et al., 2004; Joo et al., 2005). In addition, sphingosine-1-phosphate has been found to be a signaling molecule involved in ABA regulation of guard cell turgor (Ng et al., 2001), and the G $\alpha$  subunit acts downstream of sphingosine-1-phosphate during this process (Coursol et al., 2003). Those reports indicate that G $\alpha$  may play a central mediating role in guard cell signaling. ExtCaM and ABA likely act independently to regulate the G protein pathway, although this has not been experimentally demonstrated. However, G $\alpha$  regulation of NO generation in guard cells has not been reported. In mammalian cells, heterotrimeric G protein regulates endothelial NOS activity by increasing cellular levels of endothelial NOS (Andreeva et al., 2006). If such G protein-dependent regulation of NO generation exists in Arabidopsis, NO



**Figure 8.** Model showing the possible signaling pathway for ExtCaM-induced stomatal closure. ExtCaM activates G protein to induce H<sub>2</sub>O<sub>2</sub> generation by NADPH oxidases, which increases NO production depending on AtNOA1; NO subsequently arouses stomatal closure.



production will be decreased in *gpa1* mutants and increased in *cGα* lines. Our results were consistent with this speculation. *gpa1* guard cells generated less NO and *cGα* lines generated higher NO levels when treated with ExtCaM (Fig. 2, A and B). Furthermore, the similarity of *cGα Atnoa1* and *gpa1-2 Atnoa1* double mutants to the *Atnoa1* mutant in response to ExtCaM, the rescue of *gpa1* mutants by *AtNOA1* in stomatal closure and NO generation in response to ExtCaM (Fig. 3), and the enhancement of stomatal closure by SNP in *gpa1* mutants (Fig. 2C) suggest that *Gα* action depends on *AtNOA1*-mediated NO generation. These results suggest that *Gα* acts upstream of NO generation in ExtCaM-mediated guard cell signaling.

### ***Gα* Regulation of NO Production in ExtCaM-Induced Stomatal Closure Depends on H<sub>2</sub>O<sub>2</sub> Generation**

ROS has been shown to be an important second messenger in ABA-induced stomatal movement in both Arabidopsis and *V. faba* (Pei et al., 2000; Zhang et al., 2001). Pharmacological results show that H<sub>2</sub>O<sub>2</sub> is also involved in ExtCaM-induced stomatal closure and acts downstream of *Gα* in *V. faba* (Chen et al., 2004). In this report, we confirmed the essential role of ROS in the ExtCaM-mediated guard cell response using the *atrbohD/F* double mutant in Arabidopsis. The *atrbohD/F* mutant showed impaired ROS increases and stomatal closure when treated with ExtCaM (Fig. 4), providing genetic evidence for the involvement of ROS in ExtCaM guard cell signaling. Furthermore, ExtCaM-triggered stomatal closure in *cGα* lines was blocked by CAT or DPI, and the defect in stomatal closure in *gpa1* was rescued by H<sub>2</sub>O<sub>2</sub> (Fig. 5). These results indicate that H<sub>2</sub>O<sub>2</sub> acts downstream of *Gα* in ExtCaM guard cell signaling.

Reports from several groups imply that H<sub>2</sub>O<sub>2</sub> induces NO generation in guard cells of *Phaseolus aureus*, *V. faba*, and Arabidopsis (Lum et al., 2002; He et al., 2005; Bright et al., 2006). It has been shown that antioxidant ascorbate and CAT partially reverse SNP-induced H<sub>2</sub>DCF fluorescence and stomatal closure in *V. faba* (He et al., 2005). However, Bright et al. (2006) reported that SNP-induced stomatal closure is not affected by CAT in Arabidopsis, implying that guard cell responses differ between Arabidopsis and *V. faba*. In this study, we showed that ExtCaM-induced H<sub>2</sub>O<sub>2</sub> synthesis is essential for NO production, since ExtCaM-induced NO production was greatly impaired in the guard cells of *atrbohD/F* and CAT blocked the ExtCaM-induced NO production in the guard cells of the wild type (Fig. 6, A and B). Nitro-L-Arg methyl ester inhibits H<sub>2</sub>O<sub>2</sub>-mediated NO generation in guard cells of *P. aureus* and *V. faba*, implying that NOS activity is related to H<sub>2</sub>O<sub>2</sub>-mediated NO generation (Lum et al., 2002; He et al., 2005). However, using Arabidopsis *Atnoa1* and *nia1, nia2* mutants, Bright et al. (2006) reported that it is NR, not *AtNOA1*, that is responsible for the stomatal closure and NO generation in guard cells in response to H<sub>2</sub>O<sub>2</sub>, although Desikan et al.

(2002) showed that stomatal closure in the *nia1, nia2* double mutant was induced by H<sub>2</sub>O<sub>2</sub> as in the wild type. Our data indicate that H<sub>2</sub>O<sub>2</sub>-induced stomatal closure was greatly impaired by pretreatment with cPTIO. Meanwhile, H<sub>2</sub>O<sub>2</sub>-induced stomatal closure and NO generation were significantly reduced in *Atnoa1* (Fig. 7, A and B). These results indicate that H<sub>2</sub>O<sub>2</sub>-induced stomatal closure and NO production depend mainly on *AtNOA1*, which contradicts the findings of Bright et al. (2006) but is somewhat supported by Desikan et al. (2002). The discrepancy in the *AtNOA1* or NR source of H<sub>2</sub>O<sub>2</sub> induction of NO generation may be due to the nitrogen nutrition conditions in the soil, because NR is a nitrate-dependent enzyme (Raven, 2003).

In this study, we present pharmacological, biochemical, and genetic evidence to support the essential role played by *AtNOA1*-dependent NO generation in ExtCaM-induced stomatal closure. We also show that a signaling cascade of heterotrimeric G protein, NADPH oxidase-dependent H<sub>2</sub>O<sub>2</sub> generation, and *AtNOA1*-dependent NO production regulates ExtCaM-mediated guard cell responses. Both ExtCaM and H<sub>2</sub>O<sub>2</sub> raise the [Ca<sup>2+</sup>]<sub>i</sub> level by promoting Ca<sup>2+</sup><sub>o</sub> influx through Ca<sup>2+</sup> channels in the plasma membrane (Pei et al., 2000; Xiao et al., 2004; Shang et al., 2005). In addition, ExtCaM receptor-like binding protein has been found in the plasma membrane (Cui et al., 2005). Based on these observations, we propose a working model for ExtCaM action: ExtCaM is activated by Ca<sup>2+</sup><sub>o</sub> binding; active ExtCaM converts Ca<sup>2+</sup><sub>o</sub> status to intracellular events through its receptor-like binding protein in the plasma membrane. ExtCaM could elevate the [Ca<sup>2+</sup>]<sub>i</sub> level by at least two means: (1) by promoting Ca<sup>2+</sup><sub>o</sub> influx directly, or (2) by stimulating Ca<sup>2+</sup> channels in the plasma membrane via increased H<sub>2</sub>O<sub>2</sub>. [Ca<sup>2+</sup>]<sub>i</sub> increases induced by ExtCaM might trigger Ca<sup>2+</sup>-dependent NO accumulation, and subsequently NO further releases Ca<sup>2+</sup> from intracellular Ca<sup>2+</sup> stores (Sokolovski et al., 2005). ExtCaM could act as another Ca<sup>2+</sup><sub>o</sub> sensor together with CAS to regulate [Ca<sup>2+</sup>]<sub>i</sub> oscillation under natural conditions. A similar pathway appears to be responsible for ABA regulation of stomatal closure: ABA activates G protein (Liu et al., 2007) to arouse H<sub>2</sub>O<sub>2</sub> generation by NADPH oxidases (Kwak et al., 2003), which subsequently induces NO levels by raising the [Ca<sup>2+</sup>]<sub>i</sub> level, and increased NO may cause [Ca<sup>2+</sup>]<sub>i</sub> elevation (Desikan et al., 2004). Therefore, a signaling cascade involving G protein, H<sub>2</sub>O<sub>2</sub>, NO, and calcium may integrate both ABA and ExtCaM to regulate stomatal movement (Fig. 8).

## **MATERIALS AND METHODS**

### **Plant Materials and Growth Conditions**

Seeds of the wild type and various mutants of Arabidopsis (*Arabidopsis thaliana*) were sown in potting mix and grown in a greenhouse under a 16-h-light/8-h-dark cycle, a photon flux density of 0.30 mmol m<sup>-2</sup> s<sup>-1</sup>, and a day/night temperature cycle of 18°C/22°C for 3 to 4 weeks. Seeds of *cGα* lines

(background Ws) overexpressing a constitutively active form of *GPA1* and *gpa1-1* and *gpa1-2* mutants (background Ws) were obtained from Dr. L.G. Ma (National Institute of Biological Sciences). *cGα* plants were grown in the presence of 70 nM dexamethasone (Sigma) according to the method described by Okamoto et al. (2001). *atrbohD/F* double mutant seeds (background Col-0) were obtained from Dr. M.A. Torres (University of North Carolina at Chapel Hill). Seeds of *Atnoa1* mutant and the rescued *Atnoa1* line (background Col-0) and the construct *35S:AtNOA1* were obtained from Dr. N.M. Crawford (University of California at San Diego). The *cGα1 Atnoa1* and *gpa1-2 Atnoa1* double mutants and the *gpa1-2* mutant for transformation were back-crossed three times with Col-0 to remove the Ws background of *cGα1* or *gpa1-2*. Genotypes of all mutants were confirmed by PCR analysis. Recombinant AtCaM2 protein was obtained from Dr. S.J. Cui (Hebei Normal University; Cui et al., 2005).

## Transformation of Arabidopsis and PCR Analysis of Mutants

*AtNOA1* cDNA was cut from pBIN-JIT (Guo et al., 2003) and ligated into binary vector pCAMBIA-1300 with hygromycin selection. Transformation of pCAMBIA-1300 harboring *35S:AtNOA1* into the *gpa1-2* mutant was performed according to the flower-dip method (Clough and Bent, 1998). Hygromycin-resistant seedlings were transferred to soil for examination of the NO level and stomatal response to ExtCaM. Data from three independent transgenic lines were obtained, and representative results are shown. Primers for PCR analysis of the mutants are shown in the Supplemental Materials and Methods S1.

## Stomatal Bioassay

Stomatal assays were performed essentially as described by Chen et al. (2004). Abaxial epidermal strips were peeled gently and incubated in MES buffer (10 mM MES-Tris, 50 mM KCl, and 0.1 mM CaCl<sub>2</sub>, pH 6.1) for 90 min under light to open the stomata. To study the effects of CaM,  $\alpha$ -lactalbumin, SNP, and H<sub>2</sub>O<sub>2</sub> on stomatal closure, the epidermal strips with open stomata were transferred to and incubated in MES buffer containing 10<sup>-8</sup> M CaM, 10<sup>-8</sup> M  $\alpha$ -lactalbumin, 60  $\mu$ M SNP, or 5  $\times$  10<sup>-5</sup> M H<sub>2</sub>O<sub>2</sub> for 2 h. To investigate the effect of cPTIO, CAT, or DPI on CaM-induced stomatal closure, the strips with open stomata were transferred to and incubated in MES buffer containing 10<sup>-8</sup> M CaM plus 200  $\mu$ M cPTIO, 100 units mL<sup>-1</sup> CAT, or 10  $\mu$ M DPI for 2 h. To investigate the effect of cPTIO on the effect of H<sub>2</sub>O<sub>2</sub>-induced stomatal closure, the strips with open stomata were transferred to and incubated in MES buffer containing 5  $\times$  10<sup>-5</sup> M H<sub>2</sub>O<sub>2</sub> plus 200  $\mu$ M cPTIO for 2 h. Stomatal apertures were measured with a microscope. Fifty stomata were randomly selected for three independent repeats. The data are presented as means  $\pm$  SE (*n* = 150).

## Fluorescence Microscopy

NO was visualized using the specific fluorescent NO dye DAF2-DA (Sigma). Abaxial epidermal strips with open stomata were loaded with 10  $\mu$ M DAF2-DA for 15 min, followed by a wash step. The strips were subsequently incubated in MES buffer alone, 10<sup>-8</sup> M CaM, 10<sup>-8</sup> M  $\alpha$ -lactalbumin, 1 mM CaCl<sub>2</sub>, or 5  $\times$  10<sup>-5</sup> M H<sub>2</sub>O<sub>2</sub> in MES buffer (or coincubated with other reagents as indicated in the figure legends) for 30 min. All images were acquired with a fluorescence microscope (Nikon; ELLIPE TE2000-U) with the following settings: excitation = 488 nm, emission = 515 nm; the fluorescence intensities were analyzed using MetaMorph. Data are presented as mean pixel intensities. Fifty guard cells were recorded for each treatment for three independent repeats.

## Confocal Laser Scanning Microscopy

H<sub>2</sub>O<sub>2</sub> measurement in guard cells was carried out using the method described previously (Chen et al., 2004). Abaxial epidermal strips with open stomata were incubated in 50  $\mu$ M H<sub>2</sub>DCFDA (Molecular Probes; D399) for 15 min at room temperature and then washed three times. The strips were then incubated in MES buffer alone or 10<sup>-8</sup> M CaM in MES buffer (or coincubated with 200  $\mu$ M cPTIO) for 30 min. The H<sub>2</sub>O<sub>2</sub> fluorescence in guard cells was measured using a confocal laser scanning microscope (Bio-Rad; CLSM 1024) with the following settings: excitation = 488 nm, emission = 535 nm. For real-time recording of H<sub>2</sub>DCF fluorescence intensity, images were recorded every

15 s. When the fluorescence stabilized around 60 s after scanning, CaM solution was added directly to the buffer in which the strips were placed, and we treated this agent addition point as zero time. Fluorescence changes were recorded, and relative fluorescence intensity was calculated by subtracting the basal signal at different time points as indicated in the figure legends. The experiments were repeated at least three times with seven to 10 cells each time, and one set of data is presented to illustrate the changes in fluorescence intensity.

Sequence data from this article can be found in the GenBank/EMBL data libraries under accession numbers At2g26300 (AtGPA1), At3g47450 (AtNOA1), At5g47910 (AtrbohD), and At1g64060 (AtrbohF).

## Supplemental Data

The following materials are available in the online version of this article.

**Supplemental Materials and Methods S1.** Primers for PCR analysis of mutants.

## ACKNOWLEDGMENTS

We thank Dr. L.G. Ma, Dr. M.A. Torres, and Dr. N.M. Crawford for providing the seeds and constructs. We also thank Dr. S.J. Cui for providing purified AtCaM2 protein.

Received February 15, 2009; accepted March 19, 2009; published March 25, 2009.

## LITERATURE CITED

- Allen GJ, Chu SP, Harrington CL, Schumacher K, Hoffmann T, Tang YY, Grill E, Schroeder JI (2001) A defined range of guard cell calcium oscillation parameters encodes stomatal movements. *Nature* **411**: 1053–1057
- Andreeva AV, Vaiskunaite R, Kutuzov MA, Profirovic J, Skidgel RA, Voyno-Yasenetskaya T (2006) Novel mechanisms of G protein-dependent regulation of endothelial nitric-oxide synthase. *Mol Pharmacol* **69**: 975–982
- Barroso JB, Corpas FJ, Carreras A, Sandalio LM, Valderrama R, Palma JM, Lupiáñez JA, del Río LA (1999) Localization of nitric-oxide synthase in plant peroxisomes. *J Biol Chem* **274**: 36729–36733
- Beligni MV, Lamattina L (2000) Nitric oxide stimulates seed germination and de-etiolation, and inhibits hypocotyl elongation, three light inducible responses in plants. *Planta* **210**: 215–221
- Biro RL, Sun DY, Serlin BS, Terry ME, Datta N, Sopory SK, Roux SJ (1984) Characterization of oat calmodulin and radioimmunoassay of its sub-cellular distribution. *Plant Physiol* **75**: 382–386
- Bright J, Desikan R, Hancock JT, Weir IS, Neill SJ (2006) ABA-induced NO generation and stomatal closure in Arabidopsis are dependent on H<sub>2</sub>O<sub>2</sub> synthesis. *Plant J* **45**: 113–122
- Chen YL, Huang R, Xiao YM, Lü P, Chen J, Wang XC (2004) Extracellular calmodulin-induced stomatal closure is mediated by heterotrimeric G protein and H<sub>2</sub>O<sub>2</sub>. *Plant Physiol* **136**: 4096–4103
- Chen YL, Zhang XQ, Chen J, Wang XC (2003) Existence of extracellular calmodulin in the lower epidermis of the leaves of *Vicia faba* and its role in regulating stomatal movements. *Acta Bot Sin* **45**: 40–46
- Chen Z, Gallie DR (2004) The ascorbic acid redox state controls guard cell signaling and stomatal movement. *Plant Cell* **16**: 1143–1162
- Clough SJ, Bent AF (1998) Floral dip: A simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. *Plant J* **16**: 735–743
- Coelho SM, Taylor AR, Ryan KP, Sousa-Pinto I, Brown MT, Brownlee C (2002) Spatiotemporal patterning of reactive oxygen production and Ca<sup>2+</sup> wave propagation in *Fucus* rhizoid cells. *Plant Cell* **14**: 2369–2381
- Corpas FJ, Barroso JB, del Río LA (2001) Peroxisomes as a source of reactive oxygen species and nitric oxide signal molecules in plant cells. *Trends Plant Sci* **6**: 145–150
- Coursol S, Fan LM, Le Stunff H, Spiegel S, Gilroy S, Assmann SM (2003)

- Sphingolipid signalling in *Arabidopsis* guard cells involves heterotrimeric G proteins. *Nature* **423**: 651–654
- Crawford NM, Galli M, Tischner R, Heimer YM, Okamoto M, Mack A (2006) Response to Zemojtel et al.: Plant nitric oxide synthase: back to square one. *Trends Plant Sci* **11**: 526–527
- Cui S, Guo X, Chang F, Cui Y, Ma L, Sun Y, Sun D (2005) Apoplastic calmodulin receptor-like binding proteins in suspension-cultured cells of *Arabidopsis thaliana*. *J Biol Chem* **280**: 31420–31427
- Demidchik V, Shabala SN, Coutts KB, Tester MA, Davies JM (2003) Free oxygen radicals regulate plasma membrane  $\text{Ca}^{2+}$ - and  $\text{K}^{+}$ - permeable channels in plant root cells. *J Cell Sci* **116**: 81–88
- Desikan R, Cheung M, Bright J, Hancock JT, Neill SJ (2004) ABA, hydrogen peroxide and nitric oxide signalling in stomatal guard cells. *J Exp Bot* **55**: 205–212
- Desikan R, Griffiths R, Hancock JT, Neill S (2002) A new role for an old enzyme: nitrate reductase-mediated nitric oxide generation is required for abscisic acid-induced stomatal closure in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* **99**: 16314–16318
- Flores-Pérez U, Sauret-Güeto S, Gas E, Jarvis P, Rodríguez-Concepción M (2008) A mutant impaired in the production of plastome-encoded biosynthetic enzymes in *Arabidopsis* plastids. *Plant Cell* **20**: 1303–1315
- Gabalón C, Ros-Gómez LV, Pedreño MA, Ros-Barceló A (2005) Nitric oxide production by the differentiating xylem of *Zinnia elegans*. *New Phytol* **165**: 121–130
- García-Mata C, Lamattina L (2001) Nitric oxide induces stomatal closure and enhances the plant adaptive responses against drought stress. *Plant Physiol* **126**: 1196–1204
- García-Mata C, Lamattina L (2002) Nitric oxide and abscisic acid cross talk in guard cells. *Plant Physiol* **128**: 790–792
- Gas E, Flores-Pérez U, Sauret-Güeto S, Rodríguez-Concepción M (2009) Hunting for plant nitric oxide synthase provides new evidence of a central role for plastids in nitric oxide metabolism. *Plant Cell* **21**: 18–23
- Grabov A, Blatt MR (1998) Membrane voltage initiates  $\text{Ca}^{2+}$  waves and potentiates  $\text{Ca}^{2+}$  increases with abscisic acid in stomatal guard cells. *Proc Natl Acad Sci USA* **95**: 4778–4783
- Guo FQ, Crawford NM (2005) *Arabidopsis* nitric oxide synthase1 is targeted to mitochondria and protects against oxidative damage and dark-induced senescence. *Plant Cell* **17**: 3436–3450
- Guo FQ, Okamoto M, Crawford NM (2003) Identification of a plant nitric oxide synthase gene involved in hormonal signaling. *Science* **302**: 100–103
- Han S, Tang R, Anderson LK, Woerner TE, Pei ZM (2003) A cell surface receptor mediates extracellular  $\text{Ca}^{2+}$  sensing in guard cells. *Nature* **425**: 196–200
- He JM, Xu H, She XP, Song XG, Zhao WM (2005) The role and interrelationship of hydrogen peroxide and nitric oxide in the UV-B-induced stomatal closure in broad bean. *Funct Plant Biol* **32**: 237–247
- He Y, Tang RH, Hao Y, Stevens RD, Cook CW, Ahn SM, Jing L, Yang Z, Chen L, Guo F, et al (2004) Nitric oxide represses the *Arabidopsis* floral transition. *Science* **305**: 1968–1971
- Iyer LK, Qasba PK (1999) Molecular dynamics simulation of  $\alpha$ -lactalbumin and calcium binding c-type lysozyme. *Protein Eng* **12**: 129–139
- Joo JH, Wang S, Chen JG, Jones AM, Fedoroff NV (2005) Different signaling and cell death roles of heterotrimeric G protein  $\alpha$  and  $\beta$  subunits in the *Arabidopsis* oxidative stress response to ozone. *Plant Cell* **17**: 957–970
- Kwak JM, Mori IC, Pei ZM, Leonhardt N, Torres MA, Dangl JL, Bloom RE, Bodde S, Jones JDC, Schroeder JI (2003) NADPH oxidase *AtrbohD* and *AtrbohF* genes function in ROS-dependent ABA signaling in *Arabidopsis*. *EMBO J* **22**: 2623–2633
- Liu X, Yue Y, Li B, Nie Y, Li W, Wu WH, Ma L (2007) A G protein-coupled receptor is a plasma membrane receptor for the plant hormone abscisic acid. *Science* **315**: 1712–1716
- Lum HK, Butt YK, Lo SC (2002) Hydrogen peroxide induces a rapid production of nitric oxide in mung bean (*Phaseolus aureus*). *Nitric Oxide* **6**: 205–213
- Ma LG, Sun DY (1997) The effects of extracellular calmodulin on initiation of *Hippeastrum rutilum* pollen germination and tube growth. *Planta* **202**: 336–340
- Ma LG, Xu XD, Cui SJ, Sun DY (1999) The presence of a heterotrimeric G protein and its role in signal transduction of extracellular calmodulin in pollen germination and tube growth. *Plant Cell* **11**: 1351–1364
- MacRobbie E (1992) Calcium and ABA-induced stomatal closure. *Philos Trans R Soc Lond B Biol Sci* **338**: 5–18
- McAinsh MR, Webb AAR, Taylor JE, Hetherington AM (1995) Stimulus-induced oscillations in guard cell cytosolic free calcium. *Plant Cell* **7**: 1207–1219
- Miles PR, Bowman L, Rengasamy A, Huffman L (1998) Constitutive nitric oxide production by rat alveolar macrophages. *Am J Physiol Lung Cell Mol Physiol* **274**: L360–L368
- Moreau M, Lee GI, Wang Y, Crane BR, Klessig DF (2008) AtNOS/A1 is a functional *Arabidopsis thaliana* cGTPase and not a nitric oxide synthase. *J Biol Chem* **283**: 32957–32967
- Neill SJ, Desikan R, Clarke A, Hancock JT (2002) Nitric oxide is a novel component of abscisic acid signalling in stomatal guard cells. *Plant Physiol* **128**: 13–16
- Ng CK, Carr K, McAinsh MR, Powell B, Hetherington AM (2001) Drought-induced guard cell signal transduction involves sphingosine-1-phosphate. *Nature* **410**: 596–599
- Okamoto H, Matsui M, Deng XW (2001) Overexpression of the heterotrimeric G-protein  $\alpha$ -subunit enhances phytochrome mediated inhibition of hypocotyl elongation in *Arabidopsis*. *Plant Cell* **13**: 1639–1652
- Pagnussat GC, Simontacchi M, Puntarulo S, Lamattina L (2002) Nitric oxide is required for root organogenesis. *Plant Physiol* **129**: 954–956
- Pandey S, Assmann SM (2004) The *Arabidopsis* putative G protein coupled receptor GCR1 interacts with the G protein  $\alpha$  subunit GPA1 and regulates abscisic acid signaling. *Plant Cell* **16**: 1616–1632
- Pei ZM, Murata Y, Benning G, Thomine S, Klüsenner B, Allen GJ, Grill E, Schroeder JI (2000) Calcium channels activated by hydrogen peroxide mediate abscisic acid signaling in guard cells. *Nature* **406**: 731–734
- Prado AM, Porterfield DM, Feijó JA (2004) Nitric oxide is involved in growth regulation and re-orientation of pollen tubes. *Development* **131**: 2707–2714
- Raven JA (2003) Can plants rely on nitrate? *Trends Plant Sci* **8**: 314–315; author reply 315–316
- Schroeder JI, Allen GJ, Hugouvieux V, Kwak JM, Waner D (2001) Guard cell signal transduction. *Annu Rev Plant Physiol Plant Mol Biol* **52**: 627–658
- Shang ZL, Ma LG, Zhang HL, He RR, Wang XC, Cui SJ, Sun DY (2005)  $\text{Ca}^{2+}$  influx into lily pollen grains through a hyperpolarization-activated  $\text{Ca}^{2+}$ -permeable channel which can be regulated by extracellular CaM. *Plant Cell Physiol* **46**: 598–608
- Sokolovski S, Hills A, Gay R, Garcia-Mata C, Lamattina L, Blatt MR (2005) Protein phosphorylation is a prerequisite for intracellular  $\text{Ca}^{2+}$  release and ion channel control by nitric oxide and abscisic acid in guard cells. *Plant J* **43**: 520–529
- Stohr C, Stremmlau S (2006) Formation and possible roles of nitric oxide in plant roots. *J Exp Bot* **57**: 463–470
- Sun DY, Bian YQ, Zhao BH, Zhao LY, Yu XM, Duan SJ (1995) The effect of extracellular calmodulin on cell wall regeneration of protoplasts and cell division. *Plant Cell Physiol* **36**: 133–138
- Sun DY, Li HB, Cheng G (1994) Extracellular calmodulin accelerates the proliferation of suspension-cultured cells of *Angelica dahurica*. *Plant Sci* **99**: 1–8
- Tang RH, Han S, Zheng H, Cook CW, Choi CS, Woerner TE, Jackson EB, Pei ZM (2007) Coupling diurnal cytosolic  $\text{Ca}^{2+}$  oscillations to the CAS-IP<sub>3</sub> pathway in *Arabidopsis*. *Science* **315**: 1423–1426
- Wang XQ, Ullah H, Jones AM, Assmann SM (2001) G protein regulation of ion channels and abscisic acid signaling in *Arabidopsis* guard cells. *Science* **292**: 2070–2072
- Wood NT, Allan AC, Haley A, Viry-Moussaïd M, Trewavas AJ (2000) The characterization of differential calcium signaling in tobacco guard cells. *Plant J* **24**: 335–344
- Xiao Y, Chen YL, Huang R, Chen J, Wang XC (2004) Depolymerization of actin cytoskeleton is involved in stomatal closure induced by extracellular calmodulin in *Arabidopsis*. *Sci China C Life Sci* **47**: 454–460
- Zemojtel T, Fröhlich A, Palmieri MC, Kolanczyk M, Mikula I, Wyrwicz LS, Wanker EE, Mundlos S, Vingron M, Martasek P, et al (2006) Plant nitric oxide synthase: a never-ending story? *Trends Plant Sci* **11**: 524–525
- Zhang X, Zhang L, Dong F, Gao J, Galbraith DW, Song CP (2001) Hydrogen peroxide is involved in abscisic acid-induced stomatal closure in *Vicia faba*. *Plant Physiol* **126**: 1438–1448
- Zhao MG, Tian QY, Zhang WH (2007) Nitric oxide synthase-dependent nitric oxide production is associated with salt tolerance in *Arabidopsis*. *Plant Physiol* **144**: 206–217