

Research article

Open Access

Acclimatory responses of the *Daphnia pulex* proteome to environmental changes. II. Chronic exposure to different temperatures (10 and 20°C) mainly affects protein metabolism

Susanne Schwerin¹, Bettina Zeis¹, Tobias Lamkemeyer², Rüdiger J Paul¹, Marita Koch¹, Johannes Madlung², Claudia Fladerer² and Ralph Pirow*¹

Address: ¹Institute of Zoophysiology, University of Münster, Münster, Germany and ²Proteom Centrum Tübingen, Interfaculty Institute for Cell Biology, University of Tübingen, Tübingen, Germany

Email: Susanne Schwerin - susanne.schwerin@freenet.de; Bettina Zeis - zeis@uni-muenster.de; Tobias Lamkemeyer - tobias.lamkemeyer@uni-tuebingen.de; Rüdiger J Paul - paulr@uni-muenster.de; Marita Koch - marita@uni-muenster.de; Johannes Madlung - johannes.madlung@uni-tuebingen.de; Claudia Fladerer - claudia.fladerer@uni-tuebingen.de; Ralph Pirow* - pirow@uni-muenster.de

* Corresponding author

Published: 21 April 2009

Received: 15 February 2008

BMC Physiology 2009, 9:8 doi:10.1186/1472-6793-9-8

Accepted: 21 April 2009

This article is available from: <http://www.biomedcentral.com/1472-6793/9/8>

© 2009 Schwerin et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Temperature affects essentially every aspect of the biology of poikilothermic animals including the energy and mass budgets, activity, growth, and reproduction. While thermal effects in ecologically important groups such as daphnids have been intensively studied at the ecosystem level and at least partly at the organismic level, much less is known about the molecular mechanisms underlying the acclimation to different temperatures. By using 2D gel electrophoresis and mass spectrometry, the present study identified the major elements of the temperature-induced subset of the proteome from differently acclimated *Daphnia pulex*.

Results: Specific sets of proteins were found to be differentially expressed in 10°C or 20°C acclimated *D. pulex*. Most cold-repressed proteins comprised secretory enzymes which are involved in protein digestion (trypsins, chymotrypsins, astacin, carboxypeptidases). The cold-induced sets of proteins included several vitellogenin and actin isoforms (cytoplasmic and muscle-specific), and an AAA+ ATPase. Carbohydrate-modifying enzymes were constitutively expressed or down-regulated in the cold.

Conclusion: Specific sets of cold-repressed and cold-induced proteins in *D. pulex* can be related to changes in the cellular demand for amino acids or to the compensatory control of physiological processes. The increase of proteolytic enzyme concentration and the decrease of vitellogenin, actin and total protein concentration between 10°C and 20°C acclimated animals reflect the increased amino-acids demand and the reduced protein reserves in the animal's body. Conversely, the increase of actin concentration in cold-acclimated animals may contribute to a compensatory mechanism which ensures the relative constancy of muscular performance. The sheer number of peptidase genes (serine-peptidase-like: > 200, astacin-like: 36, carboxypeptidase-like: 30) in the *D. pulex* genome suggests large-scaled gene family expansions that might reflect specific adaptations to the lifestyle of a planktonic filter feeder in a highly variable aquatic environment.



Background

Planktonic crustaceans of the genus *Daphnia* experience pronounced variations in ambient parameters such as oxygen concentration and temperature in the field and show plastic adaptive responses to these environmental changes. Differential regulation of gene expression provides specific sets of proteins for the maintenance of cellular function under altered ambient conditions. The recent release of the *Daphnia pulex* genome sequence [1,2] offers the opportunity to relate proteomic adjustments to the differentially regulated genes.

Temperature affects the performance of poikilothermic animals at all levels of biological organization ranging from biochemical reactions *via* physiological processes to organismic properties such as fecundity and reproductive success. Acute changes in water temperature, for example, have a strong effect on systemic parameters such as heart and ventilation rate of *Daphnia* spp. (e.g. [3]). However, such physiological perturbations can be damped by acclimatory processes. Previous studies [3-6] have shown that the metabolic rates, heart and ventilation rates, and muscular performances of several *Daphnia* species at 10°C and 20°C are not as different as expected from the Q₁₀ rule, provided the animals have the chance to acclimate to the temperature at which they were tested. Such a type of compensatory control (metabolic cold adaptation) is primarily based on adjustments in enzyme concentration [7]. Nevertheless, a more or less reduced metabolic rate in the cold decreases the nutritive requirements [8] and causes also a retardation in somatic growth and development [9-11]. To mechanistically explain the role of temperature acclimation for the control of physiological processes, it is essential to know the adjustments which occur at the proteomic level.

The present study analyzed the protein expression patterns of 10°C and 20°C acclimated animals of *Daphnia pulex* under normoxic conditions. Two-dimensional gel electrophoresis and mass spectrometry were employed to identify the major elements of the temperature-induced subset of the proteome. Based on their putative functions, the probable physiological role of these sets of proteins are discussed.

Results

Two-dimensional gels were prepared from total soluble proteins extracted from 10°C or 20°C acclimated cultures of *Daphnia pulex* kept under normoxia (oxygen partial pressure: 20 kPa). A total of 224 spots were detected in representative fusion images for each acclimation condition (Figure 1A, B; encircled spots). The dual-channel representation of both fusion gels revealed a large set of cold-induced proteins of low molecular weight ($M_r < 40$ kDa) in the lower right diagonal half of the gel (Figure 1C; red-

colored spots). Proteins of reduced expression in the cold were mainly confined to the low-pI range (pI = 4–5) in the upper left diagonal half of the gel (green-colored spots).

A total number of 17 spots comprising cold-repressed proteins (36–41, Figure 1A) and cold-induced proteins (spots 42–52, Figure 1B) were successfully identified by mass spectrometry (Tables 1, see Table 2 for corresponding protein IDs and gen models). Additionally included into the inter-gel comparison was a set of spots (1, 19–22, 28–32, 34–35), the identity of which was already known from a previous study [12]. These spots showed either constitutive or temperature-dependent expressions.

It is conspicuous, that a separation of cold-induced and cold-repressed proteins by M_r /pI leads to protein groups of similar classification. Almost all of the identified proteins with a reduced expression in the cold (expression reduction by 40–80%) were secretory enzymes involved in protein digestion (spots 31–32 and 36–41, Table 1). These include three trypsins (TRY4B, TRY5F, TRY5L), two chymotrypsins (CHY1A, CHY1C), one astacin (ACN2), and two carboxypeptidases (CPA1A, CPA1B). All these proteins are synthesized as pro-enzymes (zymogens), which are activated by the removal of an N-terminal propeptide (3–11 kDa). Owing to the similarities in their M_r /pI values, these proteins were multiply identified among the analysed spots. In addition, the multiple occurrence of TRY5F and CHY1C in spots with assigned M_r values of 25 and ≈ 30 kDa may be explained by the possible co-presence of pro-enzymes and enzymes. The only non-proteolytic proteins identified among these spots were two secretory proteins (ERNA, ERNB) carrying the characteristic domain of the EndoU/XendoU family of endoribonucleases [13,14]. The spot region 28, which was excised and analyzed in a previous study [12], contained a mixture of enzymes (including peptidases of the family M2 and M13), which made an expression evaluation impossible.

Most dominant among the identified cold-induced proteins were the vitellogenins (VTGs) and actins. These proteins showed a 4–7-fold induction and were detected in ten spots (43–52). The multiple detection of these proteins and the large discrepancies between the experimental (15–40 kDa) and predicted M_r values (actins: 42 kDa, VTGs: 190–220 kDa) indicate that the main share of the cold-induced protein spots in the lower right diagonal half of the gel (Figure 1C; red-colored spots) were proteolytic cleavage fragments. However, it is important to note that VTG cleavage fragments of 65–155 kDa may naturally occur in developing *Daphnia* embryos (see discussion). The tryptic peptides used for the identification of VTGs covered a large part of the VTG sequences including the superoxide dismutase-like domain (SOD), the large-lipid-




Figure 1
2D protein gels from *Daphnia pulex* acclimated at 20°C (A) and 10°C (B). Gel images represent fusion (average) images from a set of three (A) or two (B) biological replicates. Consensus spots used for comparison in (D) are encircled. Blue and red numbers indicate cold-repressed and cold-induced protein spots that were picked from the 2D gels for mass-spectrometric analysis. Black numbers indicate previously identified proteins [12]. (C) Dual-channel representation of the gel images shown in (A) and (B). Protein spots of similar expression intensity appear in yellow. Green indicates that spots are much stronger or unique on the gel from 20°C-acclimated animals, whereas red means that spots are much stronger or unique in the gel from 10°C-acclimated *D. pulex*. (D) Scatter plot showing the comparison of expression levels in the two fusion images (V_{rel} : relative spot volume).

transfer module (Vit-N), and the von Willebrand-factor type-D domain (VWD) (Figure 2). None of the tryptic peptides could be allocated to the domain of unknown function (DUF1943) and to the interdomain regions. Based on the high sequence coverage by tryptic fragment analysis, two vitellogenins (VTG1, VTG4) could be identified (Figure 2, lower part). Although the present study did not yield any tryptic peptides for the N-terminal SOD-like

domain of VTG2, the presence of VTG2 among the analyzed spots cannot be excluded because of the very large sequence identity of VTG2 and VTG1 (98% identity when excluding the SOD-like domain).

Actins were detected in four spots (44, 47, 48, 51). The tryptic peptides used for the identification of actins (Figure 3) covered only the C-terminal half of the 42-kDa pro-

Table 1: Identified proteins from *Daphnia pulex* acclimated to 10°C or 20°C

Spot no.	Specificity N10: N20	Matched peptide sequences	Sequence coverage ^{a)}	Mascot score ^{b)}	M _r gel/ M _r predicted	pI gel/pI predicted	SP Length	PP Length	Putative function (symbolic name)
Proteolytic enzymes									
36	0.3*	1. VVAGEHSLR 2. SVDVPVDDDDTC NR	8.9%	149	30/26–29	4.4/4.4–4.8	15	27	Trypsin (TRYSL)
		1. LTAAEEPTRVEIR 2. IRNDVALIK	7.5%	80	30/25–30	4.4/4.5–5.3	18	48	Chymotrypsin (CHYIA)
37	0.2*	1. GVTDLTIFR 2. VVAGEHSLR 3. VVAGEHSLRTDSG LEQNR	9.8%	159	29/26–29	4.4/4.4–4.8	15	27	Trypsin (TRY5F)
		1. VVAGEHSLR 2. SVDVPVDDDDTC NR	8.9%	149	29/26–29	4.4/4.4–4.8	15	27	Trypsin (TRYSL)
38	0.5*	1. GLADADIIVFK 2. LIWVGQYNR 3. YRDELAGK	10.7%	123	29/30	4.5/4.5	19		Endoribonuclease-like protein (ERNA)
		1. GLADADIIVFK 2. LIWVGQYNR 3. YRDELAGK	8.0%	123	29/39	4.5/4.6	20		Endoribonuclease-like protein (ERNB)
		1. VVAGEHSLR 2. SVDVPVDDDDTC NR	8.9%	149	29/26–29	4.5/4.4–4.8	15	27	Trypsin (TRYSL)
		1. GVTDLTIFR 2. VVAGEHSLR	6.5%	80	29/26–29	4.5/4.4–4.8	15	27	Trypsin (TRY5F)

Table 1: Identified proteins from *Daphnia pulex* acclimated to 10°C or 20°C (Continued)

39	0.4*	1. VVAGEHSLR	8.9%	149	29/26–29	4.6/4.4–4.8	15	27	Trypsin (TRY5L)	
		2. SVDVPVDDDDTC NR								
		1. GVTDLTIFR	6.5%	120	29/26–29	4.6/4.4–4.8	15	27	Trypsin (TRY5F)	
		2. VVAGEHSLR								
40	0.6*	1. TTEEYYSVQK	6.5%	112	25/23–27	4.5/4.7–4.6	?	39 ?	Astacin (ACN2)	
		2. TGGGCYSYIGR								
		1. GVTDLTIFR	6.5%	109	25/26–29	4.5/4.4–4.8	15	27	Trypsin (TRY5F)	
		2. VVAGEHSLR								
41	0.4*	1. LTAAEEPTR	9.1%	141	25/25–30	4.7/4.4–5.0	18	47	Chymotrypsin (CHYIC)	
		2. LTAAEEPTRVEVR								
		3. IINDVALIK								
28	1.2	see [12]							Peptidase M13 Peptidase M2 Carboxylesterase, type B Sphingomyelin phosphodiesterase Sphingomyelin phosphodiesterase	
31	0.6	see [12]			30/34–45	4.8/4.9–4.8	16	92	Carboxypeptidase A (CPAIA) Carboxypeptidase A (CPAIB) Trypsin (TRY5F)	
					30/35–46	4.8/5.1–4.9	16	93		
					29/26–29	4.4/4.4–4.8	15	27		
32	0.3	see [12]			23/24–27	5.0/5.2–5.4	17	24	Trypsin (TRY4B)	
Egg yolk proteins & precursors										
43	7.3*	see Figure 2			/190–220	/6.4–6.7		17–20		Vitellogenin (VTG1 , VTG2 , VTG4)
44	7.1*	see Figure 3	15.7%	271	25/42	5.8/5.3	17	17		Actin Vitellogenin (VTG1)
		see Figure 2	2.8%		25/220	5.8/6.7				

Table 1: Identified proteins from *Daphnia pulex* acclimated to 10°C or 20°C (Continued)

45	5.9*	see Figure 2	2.2%	132	21/190	5.2/6.4	20	Vitellogenin (VTG4)
46	5.2*	see Figure 2	2.9%	361	21/220	5.9/6.7	17	Vitellogenin (VTG1)
47	4.9*	see Figure 3 see Figure 2	9.6% 2.0%		25/42 25/220	5.6/5.3 5.6/6.7	17	Actin Vitellogenin (VTG1 , VTG2)
		1. EDQMDYLEEK 2. LLVEKER 3. YSVDEELNK	3.6%		25/83	5.6/4.7		HSP90
49	4.4*	see Figure 2	2.2%	???	21/190–220	4.8/6.4–6.7	17–20	Vitellogenin (VTG1 , VTG2 , VTG4)
50	4.2*	see Figure 2	2.2%	163	20/220	5.7/6.7	17	Vitellogenin (VTG1)
52	3.7	see Figure 2	3.0%	344	18/220	5.1/6.7	17	Vitellogenin (VTG1 , VTG2)
Cytoskeleton & muscle proteins								
48	4.5*	see Figure 2 see Figure 3	2.2%	???	27/42	5.2/5.3		Actin Vitellogenin (VTG4)
		1. EQLDEESEAK 2. AEELEDAKR 3. ATVLANQMEK			27/220	5.2/5.9		Myosin heavy chain (MHC-1)
		1. LTTDPAFLEK 2. NAAAVHEIR 3. GDLGIEIPPEK			27/??	5.2/?		Pyruvate kinase
51	3.6*	see Figure 3	2.2%	???	36/42	5.7/5.3		Actin
ATPase								

Table 1: Identified proteins from *Daphnia pulex* acclimated to 10°C or 20°C (Continued)

42	9.8*	2. GNEDLSTAILK 3. MDELQLFK 4. GDIFIVR 5. KQLALIK 6. EMVELPLR	5.1%	214	16/89	5.3/5.0	AAA+ ATPase
Carbohydrate-modifying enzymes							
35	1.2	see [12]					α -Amylase (AMY)
34	1.0	see [12]					Exo- β -1,3-Glucanase (EXG5)
1	0.4	see [12]					Cellubiohydrolase (CEL7A)
29	0.3	see [12]					Endo- β -1,4-Glucanase (CEL9A) Paramyosin (PMY)
30	0.6	see [12]					Endo- β -1,4-Mannanase (MANS5A) β -1,3-Glucan-binding protein
19	0.6	see [12]					Enolase (ENO)

Identification was based on 2D gel electrophoresis and nano-HPLC-ESI-MS/MS analysis of trypsin-digested proteins matched against the "Frozen Gene Catalog" of the *D. pulex* protein database [2]. The compiled information includes the spot number (Figure 1A, B), the 10-to-20°C expression ratio, the number and sequences of matched peptides, the sequence coverage, the Mascot score as a statistical measure of identification probability, the experimental and theoretical molecular weight (M_r) and isoelectric point (pI) of the mature protein (without signal peptide), the predicted length of the N-terminal signal peptide (SP) in secretory proteins, as well as the putative function and symbolic name of the protein. The length of the putative pro-peptide (PP) is additionally provided for proteolytic enzymes that are secreted as inactive precursors (zymogens). The predicted M_r and pI values of zymogens and the mature enzymes are given as value ranges. The amino acid sequences of the identified proteins were derived from the gene models listed in Table 2. ^apercentage of predicted protein sequence covered by matched peptides. ^bProbability-based MOWSE score: $-10^* \log(P)$, where P is the probability that the observed match is a random event. Scores >38 indicate identity or extensive homology ($p < 0.05$). Protein scores are derived from ions scores as a non-probabilistic basis for ranking protein hits. The Mascot-score calculation was performed using whole-protein sequence (including the N-terminal signal peptide in case of extracellular proteins). * $p < 0.05$ (t-Test).

Table 2: List of referred proteins and gene models

Putative function	Symbol	Model name	Protein ID	Reference ID
Trypsin	TRY1	PIR_PASA_GEN_1500076	347826	301879
Trypsin	TRY2	PIR_PASA_GEN_5300037	347779	306771
Trypsin	TRY3	PIR_PASA_GEN_6100026	347764	307264
Trypsin	TRY4A	PIR_fgenesH_pg.C_scaffold_23000179	347804	102943
Trypsin	TRY4B	PIR_estExt_fgenesH_kg.C_230008	347242	230885
Trypsin	TRY5A	PIR_PASA_GEN_4200081	347777	305924
Trypsin	TRY5B	PIR_NCBI_GNO_4200123	347775	321745
Trypsin	TRY5C	PIR_NCBI_GNO_4200124	347774	106429
Trypsin	TRY5D	PIR_estExt_fgenesH_kg.C_420021	347772	231151
Trypsin	TRY5E	PIR_NCBI_GNO_4200126	347773	321748
Trypsin	TRY5F	PIR_SNAP_00016212	347771	231152
Trypsin	TRY5G	PIR_NCBI_GNO_4200130	347770	248155
Trypsin	TRY5H	PIR_PASA_GEN_4200082	347769	305925
Trypsin	TRY5I	PIR_PASA_GEN_4200034	347768	305886
Trypsin	TRY5J	PIR_PASA_GEN_4200035	347765	305887
Trypsin	TRY5K	PIR_estExt_fgenesH_kg.C_850001	347782	231482
Trypsin	TRY5L	PIR_NCBI_GNO_8500013	347781	59836
Trypsin	TRY5M	PIR_NCBI_GNO_24500018	347780	65745
Chymotrypsin	CHY1A	PIR_PASA_GEN_2900126	347760	304512
Chymotrypsin	CHY1B	PIR_NCBI_GNO_2900206	347750	319507
Chymotrypsin	CHY1C	PIR_NCBI_GNO_2900207	347751	52244
Chymotrypsin	CHY1D	PIR_PASA_GEN_2900062	347749	52244
Chymotrypsin	CHY1E	PIR_PASA_GEN_2900063	347752	304463
Chymotrypsin	CHY1F	PIR_NCBI_GNO_2900210	347753	26258
Chymotrypsin	CHY1G	PIR_PASA_GEN_2900130	347754	304515
Chymotrypsin	CHY1H	PIR_estExt_fgenesH_kg.C_290019	347757	231027
Endoribonuclease	ERNA	PIR_PASA_GEN_12200001	347694	301221
Endoribonuclease	ERNB	PIR_PASA_GEN_6000032	347697	307196


Table 2: List of referred proteins and gene models (Continued)

Astacin	ACN2	PIR_NCBI_GNO_18200007	347841	93694
Peptidase M13		estExt_GenewiseIPlus.C_750105	200882	200882
Peptidase M2		PASA_GEN_6000071	307230	307230
Carboxylesterase, type B		PASA_GEN_25200006	304160	304160
Sphingomyelin phosphodiesterase		PASA_GEN_2900053	304453	304453
Sphingomyelin phosphodiesterase		PASA_GEN_13800028	301526	301526
Carboxypeptidase A	CPA1A	estExt_GenewiseIPlus.C_150058	195011	195011
Carboxypeptidase A	CPA1B	NCBI_GNO_1500041	315693	315693
Vitellogenin	VTG1	BDE_estExt_GenewiseI.C_9580001	299677	219769
Vitellogenin	VTG2	PIR_estExt_fgeneshI_pg.C_9580001	347667	229959
Vitellogenin	VTG2	PIR_estExt_fgeneshI_pg.C_470029	347678	226075
Vitellogenin	VTG3	estExt_fgeneshI_pg.C_470021	226068	226068
Vitellogenin	VTG4	PASA_GEN_8300068	308693	308693
Actin	ACT1A	PIR_PASA_GEN_0400163	347740	305550
Actin	ACT1B	estExt_fgeneshI_pg.C_1190006	228751	301040
Actin	ACT1C	PIR_PASA_GEN_0500107	347742	306442
Actin	ACT1D	PIR_fgeneshI_pm.C_scaffold_66000006	347743	129328
Actin	ACT2A	PIR_PASA_GEN_0100278	347739	300012
Actin	ACT2B	PIR_estExt_GenewiseIPlus.C_20413	347736	190689
Actin	ACT2C	PIR_e_gwI.2.692.1	347703	40361
Myosin	MHC-I	PIR_7_PIR_NCBI_GNO_0600448	347733	192727
Myosin	PMY	PIR_estExt_GenewiseI.C_2380001	347700	219409
HSP90		PASA_GEN_17300027	302452	302452
Pyruvate kinase		NCBI_GNO_29900007	334106	334106
AAA+ ATPase		PASA_GEN_8000045	308570	308570
α -Amylase	AMY	FRA_PASA_GEN_2100059	347603	303445
Exo- β -1,3-glucanase	EXG5	PIR_PASA_GEN_1000289	347606	300436
Cellubiohydrolase	CEL7A	PIR_PASA_GEN_1000209	347598	300366
Endo- β -1,4-glucanase	CEL9A	PIR_estExt_fgeneshI_kg.C_70001	347602	230437
β -1,3-glucan-binding protein		PASA_GEN_0200102	303036	303036

Table 2: List of referred proteins and gene models (Continued)

Endo-β-1,4-mannanase	MAN5A	PIR_PASA_GEN_8600009	347627	308762
Enolase	ENO	PIR_PASA_GEN_1500033	347595	301844

Putative functions and symbolic names of identified proteins are given in relation to gene model names and protein identification numbers of those loci which were referred to in the present study. DappuDraft protein IDs in bold type indicate manually curated gene models that may differ from those contained in the 'Filtered Models v1.1' set (released by the Joint Genome Institute in July 2007). The Reference DappuDraft gene ID can be used to retrieve the corresponding models from the Filtered Models set.



Domain	Tryptic peptide	VTG				Spot number									
		1	2	3	4	43	44	45	46	47	48	49	50	52	
SOD	GVIYGLEPGR	•					•	•							•
	SVSYIGNIK	•					•	•							•
	TYSGVPSR	•					•	•							
	AVVIHQLPDDR	•						•							
Vit-N	FNTDKIEQR	•					•	•							
	LTLENAAPTR	•	•				•	•	•				•	•	
	LTLENAAPTRK	•	•					•							
	SAAGMYTLK				•	•									
	QONAVIELQSR				•	•									
	VIVQSIDENTIVVK				•	•									
	QFNEDVMEK	•	•			•				•					
	YSEVDPGQEVK	•	•			•				•					
	VETVSIQK	•	•							•					
	LPVEAVEAEPK	•	•							•			•	•	
	CNLVNGAGCENK	•	•										•	•	
	SPVAVSTLHK				•							•	•		
	ETPDSEKDNIER				•								•		
	NTSDKINMLR	•	•										•		
	ALWQEVK				•								•		
	EWILQGK	•	•										•		
ELVESQQITGAK				•								•	•		
ITTALSLSR	•	•			•							•			
EFESSQSSVDR				•								•			
VWD	DAASLTGGVSESER				•				•						
	TSEIAVTAHQSGGQK				•				•						
	YKFAVLSR	•	•											•	
	VVTVLLNK	•	•											•	
	IVTVIYGK				•				•				•		
	EKIEIFPAQQK	•	•											•	
NAENEVLAVIK	•	•											•		

Figure 2
Assignment of protein spots to the vitellogenins of *Daphnia pulex*. *Daphnia* vitellogenins (VTGs) are generally composed of an N-terminal large-lipid-transfer-module (Vit-N), a domain of unknown function (DUF1943), and a C-terminal von Willebrand-factor type-D domain (VWD). Of the multiple VTGs of *D. pulex*, only four are shown in respect to their domain composition (top). Note that VTG1, VTG2 and VTG3 additionally contain a superoxide dismutase-like domain (SOD) at the N-terminus. Interdomain regions are shown in gray, the signal peptide in black. Conserved residues of the SOD for Cu²⁺ and Zn²⁺ binding are indicated by blue (histidines) and green characters (histidines, aspartic acid), respectively. Potential N-linked glycosylation sites are indicated by plus signs. 'ARRI' indicates primary cleavage sites between two arginine residues. The lower part lists the tryptic peptides in the order of their appearance in the VTG sequences and in the analyzed spots.

Tryptic peptide	ACT						Spot No.				
	1A	1B	1C	1D	2A	2B	2C	44	47	48	51
DLTDYLMK	•	•	•		•	•	•	•	•	•	•
GYSFTTTAER	•	•	•		•	•	•	•	•	•	•
EIVRDIK	•	•	•	•	•	•	•			•	•
CDVDIRK	•	•	•		•			•	•	•	•
EITALAPSTMK	•	•	•	•				•	•	•	•
EITALAPSTIK					•	•	•	•		•	•
IKIIPPER	•	•	•	•	•	•	•	•			
IIAPPERK	•	•	•		•	•	•	•		•	
QEYDESGPSIVHR	•	•	•	•				•			
QEYDESGPGIVHR					•	•		•			

Figure 3
Assignment of protein spots to the actin sequences of *Daphnia pulex*. The *D. pulex* genome contains seven actin genes which code for cytoplasmic (ACT1A-D) and muscle-specific isoforms (ACT2A-C). The tryptic peptides identified in mass spectrometry are listed in the order of their appearance in the sequence of gene products and gel spots.

teins, suggesting that the N-terminal half was proteolytically cleaved during the preparation of whole-animal extracts. Proteolytic cleavage is additionally indicated by the discrepancy between experimental (25–36 kDa) and predicted M_r values (42 kDa). Owing to the high sequence identity ($\approx 97\%$), it was impossible to discriminate the expression of cytoplasmic isoforms (ACT1A-D) and muscle-specific isoforms (ACT2A-C). The lower number of tryptic-peptide assignments and the complete lack of EST evidences for ACT1D and ACT2C, however, suggests that these two actins were probably not expressed.

A ten-fold up-regulation in the cold was found for an AAA+ adenosine triphosphatase (ATPase associated with diverse cellular activities; [15]), a fragment of which was detected in spot 42. Additional identifications comprised proteolytic cleavage fragments of a molecular chaperone (HSP90, spot 47), the heavy chain of myosin (MHC-1) and a pyruvate kinase (both in spot 48). Since the latter three proteins were co-identified with actins and VTGs in the same spots, it was impossible to assess their induction states.

Among the remaining identifications was a group of carbohydrate-modifying enzymes with a constitutive or reduced expression in the cold. Constitutive expressions showed the exo- β -1,3-glucanase EXG5 (spot 34) and the α -amylase AMY (spot 35). The cold-repressed proteins included a cellulohydrolase (CEL7A, spot 1), an endo- β -1,4-glucanase (CEL9A, co-localized with paramyosin in spot 19), an endo- β -1,4-mannanase (MAN5A, co-local-

ized with a β -1,3-glucan-binding protein in spot 30), and the enolase ENO (spot 19).

Discussion

As a companion study to a previous investigation of acclimatory adjustments of the *Daphnia pulex* proteome to hypoxia [12], the effects of two different acclimation temperatures (10°C and 20°C) on the *Daphnia pulex* proteome were analyzed by 2D gel electrophoresis and mass spectrometry. Temperature acclimation mostly affected the expression of sets of proteins different from those identified under variable oxygen conditions. Several proteins constitutively expressed or subjected to hypoxic induction were also detected in the 2D gels presented here. The specific sets of proteins up- or down-regulated in the cold (10°C) were identified here for the first time.

Cold-induced protein sets I: Egg yolk proteins and precursors

The most dominant group among the cold-induced proteins in *D. pulex* were the vitellogenins (Table 1). Vitellogenin (VTG) is a precursor of the yolk protein vitellin. It is a lipoglycoprotein that is employed as a vehicle to provide the developing embryo with proteins, lipids, carbohydrates, and other essential resources. In many oviparous animals such as insects and non-mammalian vertebrates, VTG is synthesized in extraovarian tissues (e.g. fat body or liver) and is then transported *via* the blood/hemolymph to the developing oocytes [16,17]. An exception are the decapod crustaceans which show, in addition to the extraovarian synthesis in the hepatopancreas, an intraovarian synthesis of yolk proteins [18]. Since the VTGs of the branchiopod crustacean *Daphnia* spp. are more closely related to insect VTGs than to the yolk protein precursors of decapods [19-21], it is reasonable to postulate a vitellogenetic tissue that is homologous to the VTG-synthesizing fat body of insects. Although there are some cytological indications for an endogenous synthesis of yolk proteins in amphigonic oocytes [22], the main site of VTG synthesis in *Daphnia* appears to be the highly polyploid fat cells, which exhibit periodic variations in lipid and glycogen content, cell size and ultrastructure in relation to the parthenogenetic reproduction cycle [22-24].

The screening of the *D. pulex* genome database suggests 19 loci with VTG-like coding sequences. Two gene products, VTG1 and VTG4, were identified in the present study (Figure 2). The additional expression of VTG2, however, which shares a high sequence similarity with VTG1, cannot be excluded. VTG1 and VTG2 are homologous to the vitellogenins DmagVTG1 and DmagVTG2 of *D. magna* [25]. As in *D. magna*, the VTG1 and VTG2 genes are arranged in a tandem array in a back-to-back orientation, which might enable a coordinated hormonal regulation of their transcription [25]. DmagVTG1 and (probably)

DmagVTG2 are the most abundant polypeptides in *D. magna* parthenogenetic eggs at initial stages of development [19]. At least one of the primary cleavage sites is present in VTG1 and VTG2 of *D. pulex* (Figure 2, top: 'ARRI'). Given the high sequence identity (88–90%) between the corresponding VTGs of both *Daphnia* species, it is likely that primary cleavage fragments of similar size occur in the developing eggs of *D. pulex* as well. However, none of these primary cleavage fragments could be detected in full length (65–155 kDa) among the analyzed spots, which contained only smaller VTG fragments of 18–27 kDa, possibly as a consequence of a residual proteolytic activity during the preparation of whole-animal extracts. Alternatively, smaller-than-expected fragments may have arisen prior to extract preparation by an advanced cleavage of yolk material during embryonic development.

The 4–7-fold up-regulation of VTGs in 10°C acclimated *D. pulex* (Table 1) was an unexpected finding. About 50–100 adult daphnids were randomly sampled irrespective of their reproductive states for single protein extractions. The protein extracts consequently contained contributions from parthenogenetic eggs and embryos in the brood chamber as well as from maternal tissues. A greater share of vitellogenin in the protein extracts from 10°C acclimated animals may therefore result from a greater amount of eggs in the ovaries and the brood pouch or from an increased vitellogenin concentration in the synthesizing tissues, ovaries, eggs and embryos. An inspection of both acclimation groups did not reveal any differences in clutch size or in the share of animals carrying eggs and embryos. Previous findings on the impact of temperature on clutch sizes in *Daphnia* are ambiguous: there were reports on lowered [9], unchanged [11] or increased [26] clutch sizes in *D. magna* at lower temperatures. In this study, the protein concentration in the extracts was quantified and the extracts were appropriately diluted to guarantee the application of identical amounts of protein (142 µg protein) per 2D gel. Compared to the extracts from 20°C acclimated animals, the extracts from 10°C acclimated animals had a 50% higher protein concentration. The slower growth and development of *D. pulex* in the cold may possibly result in a higher concentration of whole-body protein with the VTGs particularly contributing to this effect.

A striking feature of VTG1-VTG3 is the presence of an N-terminal superoxide dismutase (SOD)-like domain (Figure 2), which is related to the Cu/Zn SODs of prokaryotes [25]. The catalytic activity of this class of SODs depends on Zn²⁺ and Cu²⁺ ions, which are coordinated by six histidine residues and one aspartic residue [27]. These residues are still present in VTG3. VTG1 and VTG2 have lost all Zn-binding residues and one of the four histidine residues

involved in Cu²⁺ binding. Functional studies on the purified yolk-protein complexes of *D. magna* revealed some residual SOD activity per constituent VTG chain (≈ 1%, in comparison to the activity of a bovine Cu/Zn SOD) [19]. Because of the great number of VTG loci in the *D. pulex* genome and the presence of an apparently intact SOD-like domain in VTG3 (for which EST evidence is available), it is difficult to analyze any (residual) detoxifying capacity of VTG1 and VTG2. Future experimental studies will be necessary to evaluate the suggested implications of the SOD-like domains of the *Daphnia* VTGs in superoxide detoxification [19] and copper binding/transportation [25].

Cold-induced protein sets II: Cytoskeleton and muscle proteins

Actins were the second large set of proteins up-regulated in the cold (Table 1). Although actins were often co-identified with VTGs during the proteomic analysis, the identification of only actin in spot 51 indicates the manifold induction of these proteins. Actin is a highly conserved protein. As major building block of the cytoskeleton and the thin filaments of myofibrils, it is involved in many important cell functions including cell motility, muscle contraction and intracellular transport. Actin generally occurs in multiple isoforms which are expressed in a tissue- and development-specific manner [28]. Compared to the genomes of human, mouse, and fly, which contain six actin loci [29], seven actin loci are present in the genome of *D. pulex* (Figure 3). Four of the predicted amino acid sequences (ACT1A, ACT1B, ACT1C, ACT1D) of *D. pulex* are related to cytoplasmic actin isoforms (5C, 42A) of *Drosophila melanogaster* [28,30]. The other three *D. pulex* sequences (ACT2A, ACT2B, ACT2C) are similar to the muscle-specific actin isoforms (57B, 79B, 87E, 88F) of *Drosophila*. The ACT2C gene is very likely a pseudogene as it lacks about 50% of the actin sequence information. Among the putative cytoplasmic actins of *D. pulex*, ACT1D possesses less conserved sequence characteristics. The complete lack of EST support for ACT1D and ACT2C suggests that only three cytoplasmic and two muscle-specific actin isoforms are expressed in *D. pulex*. Because of high sequence identity, a discrimination between these isoforms was not possible in the present study.

Two additional muscle proteins, the heavy chain(s) of muscle myosin (e.g. MHC-1) and a paramyosin (PMY), were identified on the 2D gels (Table 1). These proteins were detected in separate spots together with other proteins, which make an assessment of induction state difficult. The MHC gene of *D. pulex* deserves special attention as it shares interesting features with the MHC gene of *Drosophila melanogaster* (Figure 4) [58]. In contrast to many complex organisms with physiologically distinct muscle types, in which MHC isoforms are encoded by multiple




Figure 5
Trypsin-like proteins of *Daphnia pulex*. (A) Derived amino-acid sequence and domain structure of three trypsin genes (TRY4B, TRY5F, and TRY5L) from *D. pulex*. Predicted domain characteristics include the N-terminal signal peptide (white frame), the propeptide (blue), the chymotrypsin-like domain (red), the conserved disulfide bridges (connected cysteine residues), the catalytic triade (red characters), and substrate-specificity residues (blue characters). Residues numbering was taken from bovine chymotrypsinogen [59]. (B) Phylogenetic tree for selected trypsin-like sequences based on a multiple-sequence alignment of the trypsin-like domain including three adjacent propeptide residues (see Additional file 1). Proteins detected in the present study are labeled in red. The tree was constructed using the neighbor-joining algorithm and was rooted with chymotrypsin sequences. Bootstrap analysis was performed with 100 replicates (bootstrap values <80 are omitted). Abbreviations and NCBI accession numbers: TRY1-TRY5M, *Daphnia pulex*; TAFi, trypsin from *Aplysina fistularis* (AAO12215); TPC, trypsin from *Paralithodes camtschaticus* (AAL67442); TPV, trypsin from *Litopenaeus vannamei* (CAA75311); TPL, trypsin from *Pacifastacus leniusculus* (CAA10915); PSS, plasminogen activator from *Scolopendra subspinipes* (AAD00320); TLS1 and TLS7, trypsin from *Lepeophtheirus salmonis* (CAH61270, AAP55755); TAAe, trypsin from *Aedes aegypti* (P29787); TAS, trypsin from *Anopheles stephensi* (AAB66878); TNV, trypsin from *Nasonia vitripennis* (XP_001599779); TDM, trypsin from *Drosophila melanogaster* (P04814); TTC, trypsin from *Tribolium castaneum* (XP_967332); TBT, trypsin precursor from *Bos taurus* (Q29463); TSS, trypsin-I precursor from *Salmo salar* (P35031); ChPO, chymotrypsinogen 2 from *Paralichthys olivaceus* (Q9W7Q3); ChBT, chymotrypsinogen A from *Bos taurus* (P00766).

which is a pocket adjacent to Ser195 [36]. The S1-site residues of trypsin are Asp189, Gly216 and Gly226 [37]. All three residues are present in the detected trypsins of *D. pulex* (Figure 5A). Multiple-sequence alignment (Additional Files 1, 2) and phylogenetic-tree analysis of serine peptidase sequences from the *D. pulex* genome database revealed many other trypsin-like proteins. Two of them (TRY5F, TRY5L) together with 11 other sequences from *D. pulex* form a monophyletic cluster (Figure 5B). In CHY1A and CHY1C, the primary specificity residues comprise Ser189, Gly216 and Ala226 (Figure 6B). The first two residues are the same as in bovine chymotrypsin [37]. At the third position, Ala226 replaces the typical Gly226. These two residues are similar in shape and electrostatic

character, suggesting that substrate specificity is not significantly altered by this replacement. CHY1A and CHY1C together with six additional chymotrypsin-like proteins from *D. pulex* form a monophyletic cluster (Figure 6C). *D. pulex* chymotrypsins are closely related to the C-type brachyurins (MEROPS classification: S01.122), which include the decapod chymotrypsins and collagenolytic proteases [38-42]. C-type brachyurins are characterized by a broad substrate specificity [41]. Among the *D. pulex* chymotrypsins, an even enlarged range of substrate specificity may be assumed because of the sporadic replacements of Ser189 and Gly226 by residues of different electrostatic properties (Figure 6B).




Figure 6

Chymotrypsin-like proteins of *Daphnia pulex*. (A) Derived amino-acid sequence and domain structure of the CHY1A gene from *D. pulex*. Predicted domain characteristics include the N-terminal signal peptide (white frame), the propeptide (blue), the chymotrypsin-like domain (red), the conserved disulfide bridges (connected cysteine residues), the catalytic triade (red characters), and substrate-specificity residues (blue characters). (B) Sequence alignment of chymotrypsin-like enzymes showing the substrate recognition site with the primary specificity (S1) residues at 189, 216 and 226 (numbering system of bovine chymotrypsinogen; [59]). The shape (tiny, small) and electrostatic character (non-polar, polar, charged) of S1 residues is indicated by color shading. (C) Phylogenetic tree for selected chymotrypsin-like sequences based on a multiple-sequence alignment of the chymotrypsin-like domain including four adjacent propeptide residues (see Additional file 2). Proteins detected in the present study are labeled in red (CHY1A and CHY1C). The tree was constructed using the neighbor-joining algorithm and was rooted with trypsin sequences. Bootstrap analysis was performed with 100 replicates (bootstrap values <80 are omitted). Abbreviations and NCBI accession numbers: CHY1A-H, *Daphnia pulex*; CPC, collagenolytic protease from *Paralithodes camtschaticus* (AAL67441); CUP, collagenolytic protease from *Celuca pugilator* (P00771); ChPV, chymotrypsin BII from *Litopenaeus vannamei* (CAA71673); CES, protease from *Euphausia superba* [39]; ChPH, protease from *Pediculus humanus corporis* (AAV68346); CHL, collagenase precursor from *Hypoderma lineatum* (P08897); JoDM, Jonah 43E from *Drosophila melanogaster* (NP_724699); ChPC, chymotrypsin precursor from *Phaedon cochleariae* (O97398); ChAG, protease from *Anopheles gambiae* (AGAP005663-PA); ChCF, protease from *Chlamys farreri* (ABB89132); ChAM, chymotrypsinogen from *Arenicola marina* (CAA64472); ChRP, serine peptidase 2 from *Radix peregra* (ABL67951); ChBT, chymotrypsinogen A from *Bos taurus* (P00766); ChPO, chymotrypsinogen 2 from *Paralichthys olivaceus* (Q9W7Q3); TBT, trypsin precursor from *Bos taurus* (Q29463); TSS, trypsin-I precursor from *Salmo salar* (P35031).

The MS analysis could identify and assign only those tryptic peptides which were specific for mature proteolytic enzymes. No support was obtained for the N-terminal signal peptides, which direct the nascent proteins to the secretory pathway, and for the pro-peptides, which shield the active sites in the immature trypsinogens or chymotrypsinogens (Figure 5 and 6). Therefore, it can be assumed that the proteases originated from the gut lumen, which (in *D. magna*) contain the major share of proteases [43]. During the preparation of whole-animal extracts for the present study, intestinal proteins such as proteases are included along with those from other tissues. The pres-

ence of high amounts of proteases causes methodical problems [12], resulting in a contribution of proteolytic fragments to the observed protein spots. On the other hand, the high concentration of proteases being present in the whole-animal extracts documents a high digestive capacity for nutritional protein resources which increases with acclimation temperature. The marked induction of proteases between 10°C and 20°C acclimated animals probably reflects a higher rate of protein turnover at the higher temperature. Between identical ambient and acclimation temperatures of 10 and 20°C, the oxygen consumption rate of *D. magna* increased by 30% [4] and that

of *D. pulex* by 60% (unpublished results). Accordingly, the observed induction of proteolytic capacity by a factor of 2–5 (Table 1: trypsin, chymotrypsin) may reflect at least in parts the temperature effect on metabolic rate in acclimated *D. pulex*. In addition, higher needs for proteins may arise at higher temperatures due to modifications in the allocation and/or requirement of nutrient resources (e.g. enlarged protein needs for growth and reproduction). Previous reports on the impact of temperature on clutch sizes in *Daphnia* were ambiguous; however, a reduction of vitellogenin and protein concentration was detected in this study between 10°C and 20°C acclimation (see Discussion above). At 20°C acclimation (in comparison to 10°C acclimation), the higher growth rate (and possibly a higher reproduction rate) of *D. pulex* and/or a faster passage of nutrients through the digestive tract with possibly incomplete nutrient digestion and reduced assimilation efficiency goes hand in hand with a reduced concentration of total protein and vitellogenin in the animals. These relationships at least indicate higher demands for proteins at 20°C acclimation, which may explain the induction of intestinal proteases.

Miscellaneous proteins

Among the miscellaneous proteins with an unambiguous (one spot-one protein) identification were several carbohydrate-modifying enzymes, which were either down-regulated in the cold (cellubiohydrolase, enolase) or remained constitutively expressed (α -amylase, exo- β -1,3-glucanase), and an AAA+ ATPase, which was strongly up-regulated under cold conditions. AAA+ ATPases are molecular machines that are involved in a variety of cellular functions including vesicle transport, organelle assembly, membrane dynamics and protein unfolding [15]. They contribute to the non-destructive recycling of proteins, play an important role in protein quality control (e.g. chaperone function), and can act as microtubule motor proteins or microtubule-severing enzymes [15].

Conclusion

Major sets of proteins (egg yolk proteins and precursors, cytoskeleton and muscle proteins, proteolytic enzymes) were differentially expressed in 10°C and 20°C acclimated *D. pulex*. Compared to 10°C, the acclimation to 20°C was associated with a decrease of vitellogenins, actins and even total protein concentration, as well as with an increase of proteases. The increase of proteolytic enzymes probably reflects a higher cellular demand for amino acids, which may result from higher growth and reproduction rates and/or from a lower efficiency of intestinal protein digestion/assimilation. The decrease of protein reserves (vitellogenins, actins or total protein) also indicates an increasing bottle-neck in the amino acid supply of cells. Conversely, the acclimation to cold conditions induced an increase in protein concentration which

may be related to metabolic cold adaptation, a phenomenon for which multiple physiological support exists. Metabolic cold adaptation is a compensatory mechanism which ensures a relative constancy of metabolic rate and muscular performance. Particularly, the increase of actins in the cold maybe related to a compensatory control of muscular proteins to establish a relative constancy of muscular activity and performance.

Methods

Acclimation conditions

Water fleas, *Daphnia pulex*, were raised in the laboratory as described previously [12]. The animals were acclimated at least for three weeks (mostly months) to 10°C or 20°C at normoxic conditions (100% air saturation; oxygen partial pressure: 20 kPa), which was obtained by mild aeration using an aquarium pump. To guarantee an adequate nutrient supply at each acclimation temperature, animals were fed with green algae (*Desmodesmus subspicatus*) *ad libitum* (>1 mg C L⁻¹) every second day. Only adult females were used for protein extraction.

Proteomics

Protein extraction, two-dimensional gel electrophoresis and statistical analysis of protein expression were carried out as described previously [12]. Spots showing a sufficient size and staining intensity (relative spot volume, $V_{rel} > 0.1\%$) and differential expression between 10°C or 20°C acclimation, were excised from representative gels and subjected to in-gel digestion using trypsin and mass spectrometric analysis (nano-HPLC-ESI-MS/MS) [12]. Ratios of relative spot volumes at both temperatures were considered as induction factors. Several spots of high but constitutive expression were also included in the analysis.

Identification and characterization of proteins

Proteins were identified by correlating the ESI-MS/MS spectra with the "Frozen Gene Catalog" of the *D. pulex* v1.1 gene builds (July, 2007) [2] using the MOWSE-algorithm as implemented in the MS search engine MASCOT (Matrix Science Ltd., London, UK) [44]. The "Frozen Gene Catalog" contains all manual curations as of July 3, 2007 as well as automatically annotated models chosen from the "Filtered Models" v1.1 set. "Filtered Models" is the filtered set of models representing the best gene model for each locus. The putative function of identified proteins was inferred by sequence homology either from the automated blastp search provided by Joint Genome Institute [2] or from a manual blastp search provided by NCBI. Derived protein sequences were checked for the presence of N-terminal signal sequences using the SignalP V3.0 server [45-47]. The theoretical molecular weight (M_r) and isoelectric point (pI) of mature proteins (without N-terminal signal peptide) was calculated using the ExPASy Proteomics tool "Compute pI/MW" [48-50]. Characteristic

domains of protein families were identified using the conserved domain database (CDD) and search engine v2.13 at NCBI [51,52]. Putative N-glycosylation sites in vitellogenins were predicted using the NetNGlyc 1.0 Server [53].

Sequence alignments and phylogenetic analysis

Multiple-sequence alignments were performed using the T-Coffee algorithm [54-56]. Phylogenetic trees were constructed using the neighbor-joining algorithm [57] and a bootstrap analysis with 100 replicates.

Abbreviations

M_r : molecular weight; pI: isoelectric point; V_{rel} : relative spot volume.

Authors' contributions

SS and MK were involved in the culturing of animals and performed the protein extraction as well as the 2D-PAGE. 2D-gel image analysis was carried out by TL, BZ and RP. JM and CF were responsible for mass spectrometry and protein identification. RP and SS retrieved the information contained in Table 1. Figures were designed by RP. The annotation and manual curation of identified genes, the sequence alignments and phylogenetic analysis were performed by RP. BZ, RJP, RP and SS conceived and coordinated the study, and prepared the manuscript.

Additional material

Additional File 1

Multiple sequence alignment of trypsin-like sequences. *Multiple sequence alignment of trypsin-like sequences. The multiple-sequence alignment was performed using the T-Coffee algorithm [54]. NCBI accession numbers for the symbolic sequence names are listed in the Figure legend 5.*
Click here for file
[http://www.biomedcentral.com/content/supplementary/1472-6793-9-8-S1.pdf]

Additional File 2

Multiple sequence alignment of chymotrypsin-like sequences. *Multiple sequence alignment of chymotrypsin-like sequences. The multiple-sequence alignment was performed using the T-Coffee algorithm [54]. NCBI accession numbers for the symbolic sequence names are listed in the Figure legend 6.*
Click here for file
[http://www.biomedcentral.com/content/supplementary/1472-6793-9-8-S2.pdf]

Acknowledgements

The authors would like to thank Simone König from the Integrated Functional Genomics (University of Münster), and Marco Matthes (at that time at the Technical University Dresden) for providing the animals used for the clonal culture. We are also grateful for support from DECODON GmbH (BioTechnikum Greifswald). The Proteom Centrum Tübingen is supported

by the Ministerium für Wissenschaft und Kunst, Landesregierung Baden-Württemberg.

The sequencing and portions of the analyses were performed at the DOE Joint Genome Institute under the auspices of the U.S. Department of Energy's Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Livermore National Laboratory under Contract No. W-7405-Eng-48, Lawrence Berkeley National Laboratory under Contract No. DE-AC02-05CH11231, Los Alamos National Laboratory under Contract No. W-7405-ENG-36 and in collaboration with the Daphnia Genomics Consortium, DGC <http://daphnia.cgb.indiana.edu>. Additional analyses were performed by wFleaBase, developed at the Genome Informatics Lab of Indiana University with support to Don Gilbert from the National Science Foundation and the National Institutes of Health. Coordination infrastructure for the DGC is provided by The Center for Genomics and Bioinformatics at Indiana University, which is supported in part by the METACyt Initiative of Indiana University, funded in part through a major grant from the Lilly Endowment, Inc. Our work benefits from, and contributes to the Daphnia Genomics Consortium.

References

1. wFleaBase: **Daphnia waterflea genome database.** [<http://wFleaBase.org>].
2. JGI: **Joint Genome Institute.** [<http://www.jgi.doe.gov/Daphnia/>].
3. Pinkhaus O, Schwerin S, Pirow R, Zeis B, Buchen I, Gigengack U, Koch M, Horn W, Paul RJ: **Temporal environmental change, clonal physiology and the genetic structure of a Daphnia assemblage (D. galeata-hyalina hybrid species complex).** *Freshwater Biol* 2007, **52**:1537-1554.
4. Lamkemeyer T, Zeis B, Paul RJ: **Temperature acclimation influences temperature-related behaviour as well as oxygen-transport physiology and biochemistry in the water flea Daphnia magna.** *Can J Zool* 2003, **81**:237-249.
5. Paul RJ, Lamkemeyer T, Maurer J, Pinkhaus O, Pirow R, Seidl M, Zeis B: **Thermal acclimation in the microcrustacean Daphnia: a survey of behavioural, physiological and biochemical mechanisms.** *J Therm Biol* 2004, **29**:655-662.
6. Zeis B, Maurer J, Pinkhaus O, Bongartz E, Paul RJ: **A swimming activity assay shows that the thermal tolerance of Daphnia magna is influenced by temperature acclimation.** *Can J Zool* 2004, **82**:1605-1613.
7. Hochachka PV, Somero GN: *Biochemical adaptation: Mechanism and process in physiological evolution* Oxford: Oxford University Press; 2002.
8. Lampert W: **Feeding and nutrition in Daphnia.** In *Memorie dell'Istituto Italiano di Idrobiologia, Daphnia Volume 45*. Edited by: Peters RH, DeBernardi R. Pallanza: Istituto Italiano di Idrobiologia; 1987:143-192.
9. Goss LB, Bunting DL: **Daphnia development and reproduction – responses to temperature.** *J Therm Biol* 1983, **8**:375-380.
10. Lampert W, Trubetskova I: **Juvenile growth rate as a measure of fitness in Daphnia.** *Funct Ecol* 1996, **10**:631-635.
11. Giebelhausen B, Lampert W: **Temperature reaction norms of Daphnia magna: the effect of food concentration.** *Freshwater Biol* 2001, **46**:281-289.
12. Zeis B, Lamkemeyer T, Paul RJ, Nunes F, Schwerin S, Koch M, Schütz W, Madlung J, Fladerer C, Pirow R: **Acclimatory responses of the Daphnia pulex proteome to environmental changes. I. Chronic exposure to hypoxia affects the oxygen transport system and carbohydrate metabolism.** *BMC Physiol* 2009, **9**:7.
13. Renzi F, Caffarelli E, Laneve P, Bozzoni I, Brunori M, Vallone B: **The structure of the endoribonuclease XendoU: From small nuclear RNA processing to severe acute respiratory syndrome coronavirus replication.** *Proc Natl Acad Sci USA* 2006, **103**:12365-12370.
14. Worrall JAR, Luisi BF: **Information available at cut rates: structure and mechanism of ribonucleases.** *Curr Opin Struct Biol* 2007, **17**:128-137.

15. Roehl White S, Lauring B: **AAA+ ATPases: Achieving diversity of function with conserved machinery.** *Traffic* 2007, **8**:1657-1667.
16. Sappington TW, Raikhel AS: **Molecular characteristics of insect vitellogenins and vitellogenin receptors.** *Insect Biochem Mol Biol* 1998, **28**:277-300.
17. Romano M, Rosanova P, Anteo C, Limatola E: **Vertebrate yolk proteins: a review.** *Mol Reprod Dev* 2004, **69**:109-116.
18. Avarre J-C, Michelis R, Tietz A, Lubzens E: **Relationship between vitellogenin and vitellin in a marine shrimp (*Penaeus semisulcatus*) and molecular characterization of vitellogenin complementary DNAs.** *Biol Reprod* 2003, **69**:355-364.
19. Kato Y, Tokishita S, Ohta T, Yamagata H: **A vitellogenin chain containing a superoxide dismutase-like domain is the major component of yolk proteins in cladoceran crustacean *Daphnia magna*.** *Gene* 2004, **334**:157-165.
20. Smolenaars MMW, Madsen O, Rodenburg KW, Horst DJ Van der: **Molecular diversity and evolution of the large lipid transfer protein superfamily.** *J Lipid Res* 2006, **48**:489-502.
21. Avarre J-C, Lubzens E, Babin PJ: **Apolipoprotein, formerly vitellogenin, is the major egg yolk precursor in decapod crustaceans and is homologous to insect apolipoprotein III/I and vertebrate apolipoprotein B.** *BMC Evol Biol* 2007, **7**:3.
22. Zaffagnini F, Zeni C: **Considerations on some cytological and ultrastructural observations on fat cells in *Daphnia* (Crustacea, Cladocera).** *Boll Zool* 1986, **53**:33-39.
23. Jäger G: **Ueber den Fettkörper von *Daphnia magna*.** *Z Zellforsch* 1935, **22**:89-131.
24. Sterba G: **Zytologische Untersuchungen an grosskernigen Fettzellen von *Daphnia pulex* unter besonderer Berücksichtigung des Mitochondrien-Formwechsels.** *Z Zellforsch* 1956, **44**:456-487.
25. Tokishita S, Kato Y, Kobayashi T, Nakamura S, Ohta T, Yamagata H: **Organization and repression by juvenile hormone of a vitellogenin gene cluster in the crustacean, *Daphnia magna*.** *Biochem Biophys Res Commun* 2006, **345**:362-370.
26. McKee D, Ebert D: **The interactive effects of temperature, food level and maternal phenotype on offspring size in *Daphnia magna*.** *Oecologia* 1996, **107**:189-196.
27. Bordo D, Djinic K, Bolognesi M: **Conserved patterns in the Cu, Zn superoxide dismutase family.** *J Mol Biol* 1994, **238**:366-386.
28. Röper K, Mao Y, Brown NH: **Contribution of sequence variation in *Drosophila* actins to their incorporation into actin-based structures in vivo.** *J Cell Sci* 2005, **118**:3937-3948.
29. Hooper SI, Thuma JB: **Invertebrate muscles: muscle specific genes and peptides.** *Physiol Rev* 2005, **85**:1001-1060.
30. Fyrberg EA, Fyrberg CC, Biggs JR, Saville D, Beall CJ, Ketchum A: **Functional nonequivalence of *Drosophila* actin isoforms.** *Biochem Genet* 1998, **36**:271-287.
31. George E, Ober MB, Emerson CP: **Functional domains of the *Drosophila melanogaster* muscle myosin heavy-chain gene are encoded by alternatively spliced exons.** *Mol Cell Biol* 1989, **9**:2957-2974.
32. Swank DM, Wells L, Kronert WA, Morrill GE, Bernstein SI: **Determining structure/function relationships for sarcomeric myosin heavy chain by genetic and transgenic manipulation of *Drosophila*.** *Microsc Res Techniq* 2000, **50**:430-442.
33. Rawlings ND, Morton FR, Barrett AJ: **MEROPS: the peptidase database.** *Nucleic Acids Res* 2006, **34**:D270-D272.
34. Krem MM, Rose T, Di Cera E: **Sequence determinants of function and evolution in serine proteases.** *Trends Cardiovasc Med* 2000, **10**:171-176.
35. Colbourne JK, Eads BD, Shaw J, Bohuski E, Bauer DJ, Andrews J: **Sampling *Daphnia*'s expressed genes: preservation, expansion and invention of crustacean genes with reference to insect.** *BMC Genomics* 2007, **8**:217.
36. Hedstrom L: **Serine protease mechanism and specificity.** *Chem Rev* 2002, **102**:4501-4523.
37. Perona JJ, Craik CS: **Structural basis of substrate specificity in the serine proteases.** *Protein Sci* 1995, **4**:337-360.
38. Benjamin DC, Kristjánsson S, Gudmundsdóttir Á: **Increasing the thermal stability of euphauserase. A cold-active and multi-functional serine protease from antarctic krill.** *Eur J Biochem* 2001, **268**:131.
39. Kristjánsson S, Gudmundsdóttir Á: **Propeptide dependent activation of the antarctic krill euphauserase precursor produced in yeast.** *Eur J Biochem* 2000, **267**:2632-2639.
40. Gudmundsdóttir Á: **Cold-adapted and mesophilic brachyurins.** *Biol Chem* 2002, **383**:1125-1131.
41. Rudenskaya GN: **Brachyurins, serine collagenolytic enzymes from crabs.** *Rus J Bioorg Chem* 2003, **29**:101-111.
42. Rudenskaya GN, Kisilitsin YA, Rebrikov DV: **Collagenolytic serine protease PC and trypsin PC from king crab *Paralithodes camtschaticus*: cDNA cloning and primary structure of the enzymes.** *BMC Struct Biol* 2004, **4**:2.
43. von Elert E, Agrawal MK, Gebauer C, Jaensch H, Bauer U, Zitt A: **Protease activity in gut of *Daphnia magna*: evidence for trypsin and chymotrypsin enzymes.** *Comp Biochem Physiol B Biochem Mol Biol* 2004, **137**:287-296.
44. Perkins DN, Pappin DJC, Creasy DM, Cottrell JS: **Probability-based protein identification by searching sequence data bases using mass spectrometric data.** *Electrophoresis* 1999, **20**:3551-3567.
45. **SignalP 3.0** [<http://www.cbs.dtu.dk/services/SignalP/>]
46. Bendtsen JD, Nielsen H, von Heijne G, Brunak S: **Improved prediction of signal peptides: SignalP 3.0.** *J Mol Biol* 2004, **340**:783-795.
47. Emanuelsson O, Brunak S, von Heijne G, Nielsen H: **Locating proteins in the cell using TargetP, SignalP, and related tools.** *Nature Protocols* 2007, **2**:953-971.
48. **Compute pI/Mw tool** [http://www.expasy.ch/tools/pi_tool.html]
49. Bjellqvist B, Basse B, Olsen E, Celis JE: **Reference points for comparisons of two-dimensional maps of proteins from different human cell types defined in a pH scale where isoelectric points correlate with polypeptide compositions.** *Electrophoresis* 1994, **15**:529-539.
50. Gasteiger E, Hoogland C, Duvaud S, Wilkins MR, Appel RD, Bairoch A: **Protein Identification and Analysis Tools on the ExPASy Server.** In *The Proteomics Protocols Handbook* Edited by: Walker JM. Totowa: Humana Press; 2005:571-607.
51. **CDD: a Conserved Domain Database and Search Service.** [<http://www.ncbi.nlm.nih.gov/Structure/cdd/cdd.shtml>].
52. Marchler-Bauer A, Anderson JB, Derbyshire MK, DeWeese-Scott C, Gonzales NR, Gwadz M, Hao L, He S, Hurwitz DJ, Jackson JD, Ke Z, Krylov D, Lanczycki CI, Liebert CA, Liu C, Lu F, Lu S, Marchler GH, Mullokandov M, Song JS, Thanki N, Yamashita RA, Yin JJ, Zhang D, Bryant SH: **CDD: a conserved domain database for interactive domain family analysis.** *Nucleic Acids Res* 2006, **35**:D237-D240.
53. **NetNGlyc 1.0 Server** [<http://www.cbs.dtu.dk/services/NetNGlyc/>]
54. **T-Coffee** [<http://www.tcoffee.org/>]
55. Notredame C, Higgins DG, Heringa J: **T-Coffee: a novel method for fast and accurate multiple sequence alignment.** *J Mol Biol* 2000, **302**:205-217.
56. Poirrot O, O'Toole E, Notredame C: **Tcoffee@igs: a web server for computing, evaluating and combining multiple sequence alignments.** *Nucleic Acids Res* 2003, **31**:3503-3506.
57. Saitou N, Nei M: **The neighbor-joining method – a new method for reconstructing phylogenetic trees.** *Mol Biol Evol* 1987, **4**:406-425.
58. Polyák E, Standiford DM, Yakopson V, Emerson CP, Franzini-Armstrong C: **Contribution of myosin rod protein to structural organization of adult and embryonic muscles in *Drosophila*.** *J Mol Biol* 2003, **331**:1077-1091.
59. Greer J: **Comparative modeling methods: Application to the family of the mammalian serine proteases.** *Proteins Struct Funct Genet* 1990, **7**:317-334.