

# Ancient Egyptian herbal wines

Patrick E. McGovern<sup>a,1</sup>, Armen Mirzozian<sup>b</sup>, and Gretchen R. Hall<sup>a</sup>

<sup>a</sup>Museum Applied Science Center for Archaeology, University of Pennsylvania Museum of Archaeology and Anthropology, Philadelphia, PA 19104; and <sup>b</sup>Scientific Services Division, Alcohol and Tobacco Tax and Trade Bureau, U.S. Treasury, Beltsville, MD 20705

Edited by Ofer Bar-Yosef, Harvard University, Cambridge, MA, and approved February 23, 2009 (received for review November 17, 2008)

**Chemical analyses of ancient organics absorbed into pottery jars from the beginning of advanced ancient Egyptian culture, ca. 3150 B.C., and continuing for millennia have revealed that a range of natural products—specifically, herbs and tree resins—were dispersed by grape wine. These findings provide chemical evidence for ancient Egyptian organic medicinal remedies, previously only ambiguously documented in medical papyri dating back to ca. 1850 B.C. They illustrate how humans around the world, probably for millions of years, have exploited their natural environments for effective plant remedies, whose active compounds have recently begun to be isolated by modern analytical techniques.**

ancient medicine | biomolecular archaeology | herbs | Middle East | wine

**B**efore the rise of modern medicine and likely extending back into the Paleolithic period, humans treated disease and physical ailments by experimenting with natural products derived from plants, other animals, and minerals (1). Fruit-bearing trees, which appeared around 100 million years ago (Mya), offered unparalleled access to sugar and ethanol. The latter had already established themselves as prime energy sources in the animal kingdom. The sweet liquid that oozes out of ruptured ripened fruit provides the ideal conditions of water and nutrients for yeast on their surfaces to multiply and convert the sugar into alcohol (2).

A close symbiosis developed between plants and animals over time, in which the plants provided nourishment and other benefits to the animals, which, in turn, pollinated the plants' flowers, dispersed their seeds, and carried out other functions. The smaller molars and canines of *Proconsul* and other early hominid fossils as early as 24 Mya (3) were well adapted to consuming soft fleshy foods like fruit. These dentitions are broadly comparable to those of modern apes, including gibbons, orangutans, and lowland gorillas, which obtain most of their calories from fruit. Modern chimpanzees, whose genome is the closest to *Homo sapiens*, consume over 90% plants, of which more than 75% is fruit. It can be concluded that early hominids and their descendants favored ripe, often fermented, fruit for millions of years.

In the warm tropical climate of sub-Saharan Africa, where the human species emerged, sweet fruit slurries can achieve an alcoholic content of 5% or more (2). If early hominids were primarily fruit eaters, at least up until about 1–2 Mya, when they began focusing more on tubers and animal fat and protein, they can be expected to have adapted biologically. One result, among many, is that about 10% of the enzymes in the human liver, including alcohol dehydrogenase, function to generate energy from alcohol (4). The genetic underpinnings of the presumed early human penchant for alcoholic fruit compotes, according to the so-called “drunken monkey hypothesis” (5), has been partly borne out by the diet of Malaysian tree shrews (6). These nocturnal animals, which belong to a family believed to be ancestral to all living primates from more than 55 Mya ago, spend their nights feasting year-round on a frothy strongly scented palm “wine,” with an alcoholic content as high as 3.8%.

Plant fruits, exudates (including resins and nectar), and other structures, (such as flowers, roots, and leaves), also contain many additional compounds with anesthetic, antimicrobial, and psychotropic properties that early humans likely exploited (1, 7). Although some of these compounds might have been intended to protect the plants from predation, they could also have benefited host animals. By trial and error, humans might well have made use of some of

these properties by preparing “medicinal wines” and external salves in which the plant products were dissolved or decocted in alcoholic media.

At present, the earliest biomolecular archaeological evidence for plant additives in fermented beverages dates from the early Neolithic period in China (8) and the Middle East (9), when the first plants and animals were taken into domestication and provided the basis for complex society and permanent settlement. Possibilities for extending the evidential base back into the Paleolithic period are limited by the lack of containers, which were probably made from perishable materials, such as wood, leather, or woven textiles. Under the right environmental conditions, however, we can expect future discoveries. For example, at Monte Verde in Chile, around 13,000 B.P., the bog-like conditions of one of the first human settlements in the Americas resulted in the preservation of an incredible diversity of marsh, dune, mountain, and sea plants, which were likely exploited for their alcoholic, medicinal, and nutritional benefits (10).

The dry climate of Egypt has similarly contributed to excellent preservation of ancient organic materials, in addition to providing very detailed literary and botanical evidence for medicinal wines from one of the most long-standing ancient traditions (11, 12). We deliberately chose samples from 2 ancient Egyptian jars that illuminate the earliest and latest stages of Egyptian winemaking history and applied highly sensitive chemical techniques to obtain biomolecular information about what the vessels originally contained. Because we had already analyzed both samples by less precise methods, the latest results provide a means to test our previous results and conclusions; at the same time, they shed additional light on the vessels' contents.

**Archaeological Samples Analyzed.** The earlier sample comes from a multichambered tomb (U-j) at Abydos on the middle Nile River in Upper Egypt (2, 13). Dated to the Naqada IIIa2 period (ca. 3150 B.C.) by radiocarbon determinations, the tomb was built in the desert and belongs to one of the first rulers of the country, Scorpion I of Dynasty 0, at the beginning of Egyptian dynastic history. Its contents were exceptionally rich and included some 700 jars of foreign type that were stacked high in 3 chambers.

In 1994, a sample of yellowish flaky residue from jar no. 156 (Fig. 1) was analyzed by FTIR spectrometry, HPLC, and a Feigl spot test for tartaric acid/tartrate (14). The residue represented the agglomerated remains of materials on the surface of a liquid that had gradually evaporated and left a ring or tide-line on the interior of the jar (see Fig. S1). The 3 independent methods gave results that pointed strongly to the presence of tartrate, a principal biomarker for wine and other grape products in the Middle East. The HPLC data supported the interpretation that a tree resin—very likely terebinth—had been added to the vessel's contents.

The identification of tartrate, as the calcium salt, was based on


Author contributions: P.E.M., A.M., and G.R.H. designed research; A.M. and G.R.H. performed research; A.M. contributed new reagents/analytical tools; P.E.M., A.M., and G.R.H. analyzed data; and P.E.M., A.M., and G.R.H. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

<sup>1</sup>To whom correspondence should be addressed. E-mail: mcgovern@sas.upenn.edu.

This article contains supporting information online at [www.pnas.org/cgi/content/full/0811578106/DCSupplemental](http://www.pnas.org/cgi/content/full/0811578106/DCSupplemental).



**Fig. 1.** Jar no. 115 [Hartung U, ed (2001) *Umm el-Qaab II: Importkeramik aus dem Friedhof U in Abydos (Umm el-Qaab) und die Beziehungen ägyptens zu Vorderasien im 4. Jahrtausend v. Chr.* (P. von Zabern, Mainz, Germany): cat. no. 156, pls. 25 and 79:156] from chamber 10 of the Scorpion I tomb. Swirling “tiger-stripes” in red paint and the form of this narrow-mouthed jar are paralleled by southern Levantine vessels of the same period (Early Bronze I, ca. 3300–3000 B.C.); the manufacture of the vessel in this region was borne out by instrumental neutron activation analysis (INAA). The jar contained a yellowish flaky residue, which was analyzed by multiple chemical techniques, and several grape pips. Height: 31.7 cm. Drawing courtesy of German Archaeological Institute in Cairo.

the 3 methods independently agreeing with one another. If a single test result had proven negative, that would have invalidated the identification of calcium tartrate by the other methods. Although mixtures of compounds can be equivocal for IR and HPLC UV spectra, these were deconvoluted by statistical methods and scrutinized for the presence or absence of key absorptions. If a known absorption for a compound was lacking, that compound was excluded as a possibility. The IR and UV spectra of the Abydos sample were also searched for “matches” against large databases of relevant natural products and processed organic materials, synthetic compounds, modern wine samples, and “ancient wine reference samples.” The latter were residues from ancient vessels that likely originally contained wine, based on strong archaeological criteria or exterior inscriptions that recorded their contents.

Once calcium tartrate was identified to a high level of probability in the residue, archaeological and enological considerations come into play in determining whether the intended grape product was wine and not grape juice, syrup, or vinegar. A syrup, produced by heating grape juice and concentrating it down, was unlikely, because its viscosity would have left a uniform coating of residue on the inside of the vessel. Instead, the residue was confined to the tide-line and the bottom of the vessel, where precipitates from a liquid accumulate. Minimally, the jar had contained grape juice.

Any grape juice, however, will naturally ferment to wine in several days in the warm climate of the Middle East. The identification of a 256-bp DNA segment of a larger 840-bp fragment belonging to a ribosomal sequence of the principal wine yeast, *Saccharomyces cerevisiae*, bore out this interpretation (15). Intentional fermentation to vinegar was unlikely because of the precautions that were taken to protect the liquid in this jar from oxygen by sealing its mouth and adding a tree resin with antioxidant properties.

Grape seeds from jar no. 156 provided further corroboration that the jars originally held a grape wine. Additionally, and bearing on the issue of additives to the wine, some jars (not including jar no. 156) yielded a single whole fig, which had been preserved by desiccation. It had been sliced, perhaps to assure greater surface contact for enhancing the wine’s sweetness, taste, and other properties and to provide sufficient yeast for starting and sustaining the fermentation.



**Fig. 2.** Wine amphora from tomb 217 of cemetery 4 at Gebel Adda (Egypt), dated to early Byzantine times (the Ballana period of Lower Nubia, fourth to early sixth century A.D.). Height: 67.3 cm. With permission of the Royal Ontario Museum; photograph courtesy of W. Pratt (museum no. 973.24.1217).

The second sample, a residue deposited on or intentionally applied to the interior of an amphora (Fig. 2) from Lower Nubia in southern Egypt, dates more than 3,500 years later than the Abydos jar and represents a terminal phase of Egyptian winemaking before the Islamic conquest. Belonging to the fourth to early sixth century A.D. (Ballana period), the amphora was recovered from a tomb at Gebel Adda (16). Based on its form and an inscription on its shoulder, it has been identified as a wine jar (N.B. Millet, personal communication, 1989). Numerous such vessels littered the ground around taverns in Nubian villages of the period, demonstrating how wine had gone from being an elite beverage in Pharaonic times to becoming a beverage of commoners, who were also buried with it.

In our 1990 pioneering study of ancient Near Eastern wine (16), the Gebel Adda residue provided an ancient wine reference sample for detecting tartaric acid/tartrate and other wine constituents. As a known wine residue from antiquity, which had undergone aging processes, its composition served to assess whether other less definitive vessels, such as Abydos jar no. 156, once held grape wine. Modern reference samples of tartaric acid/tartrate and other wine constituents provided additional controls.

As expected, the Gebel Adda residue tested positive for tartaric acid/tartrate according to the 3 methods. Its FTIR spectrometry spectrum also showed the characteristic and very specific absorptions for tartaric acid and a tartrate salt by location, shape, relative intensity, and multiplicity. These results were directly comparable to the residue in Abydos jar no. 156, except that the latter’s FTIR spectrometry spectrum lacked the sharp intense carbonyl peak of the acid, with a shoulder, at  $1,720\text{--}1,740\text{ cm}^{-1}$ . In other words, although the tartaric acid in the Abydos jar had been converted wholly to the calcium salt, the younger Gebel Adda residue still retained the acid.

Another important inference could be drawn from the FTIR spectrometry spectra: other unidentified hydrocarbon-rich com-



samples gave peaks for tartaric acid at a retention time of 0.75 min, which matched a 0.1- $\mu\text{g}/\text{mL}$  (0.1 ppm) standard of tartaric acid. The results of analyses on similar extracts of an ancient pottery blank and aqueous controls, run before and after each analysis, were negative (data not shown). An ancient bowl rim from a site in Jordan (Tell el-Fukhar, Late Bronze II, ca. 1400–1200 B.C., no. 763) served as the pottery blank. It had been made of local clay according to instrumental neutron activation analysis (INAA) (20) and had a very low probability of having originally contained wine or another grape product.

### Headspace Solid Phase Microextraction and Thermal Desorption GC/MS Analyses

Having confirmed tartaric acid/tartrate by LC/MS/MS in the Abydos and Gebel Adda samples, we then analyzed them by headspace solid phase microextraction (SPME) and thermal desorption (TD) GC/MS. The goal was to discover whether the Egyptian wines contained any plant additives, which had been implied by our earlier analyses.

Headspace SPME and TD GC/MS have great utility in biomolecular archaeology because of their high sensitivity and selectivity for volatile compounds of interest (8, 21). SPME extraction efficiency and sensitivity are considerably increased by dissolving or suspending the solid sample in an aqueous saline solution. Moreover, the methods require only milligram quantities of valuable archaeological samples, and analyses can be performed rapidly without extraction in an organic solvent.

Compounds were identified in the Abydos and Gebel Adda samples (Table S1) by retention time and/or matches to a mass spectral library of more than 160,000 samples (NIST05). Illustrative SPME experimental data are provided in Figs. S2 and S3.

Compounds detected in the ancient pottery blank are assumed to derive either from ancient and/or modern “background contaminants,” attributable to groundwater percolation or sample handling (e.g., plasticizers and antioxidants from plastic, including compounds in the phthalate family and 3,5-di-tert-butyl-4-hydroxybenzaldehyde). Possibly, some of the low-boiling compounds up to hexenal were also contaminants; however, it is more likely that they were preserved within the ionic clay structure. Any ancient ethanol would have been metabolized by microorganisms.

A thorough search of the chemical literature [using SciFinder Scholar of the Chemical Abstract Services, American Chemical Society; Dr. Duke’s Phytochemical and Ethnobotanical Databases (<http://www.ars-grin.gov/duke/>) of the U.S. Department of Agriculture, Agricultural Research Service; the chemical database of the Amber Research Laboratory of Vassar College (<http://cima.ng-london.org.uk/ar1/>); and other bioinformatics tools (22)] enabled several groups of probable ancient compounds to be distinguished in the Abydos and Gebel Adda samples.

As might be anticipated for samples that tested positive for tartaric acid/tartrate, constituents of modern grape wine (23) are very well represented in one or both of the ancient samples, including alcohols, acids, esters, aldehydes, fatty acid derivatives, and terpenoids. Although benzaldehyde, 2-ethyl-1-hexanol, nonanal, and ethyl palmitate occur in wine, they might also be contaminants. Excellent preservation of the more ancient Abydos sample, in particular, is implied by these findings.

For the Abydos sample, 3 herbs—savory (*Satureja*), *Artemisia seibeni* (a member of the wormwood family), and blue tansy (*Tanacetum annuum*)—could account for the combined presence of 8 terpenoid compounds (labeled 2 in Table S1): linalool, camphor, borneol, L-menthol, alpha-terpineol, carvone, thymol, and geranyl acetone. The same compounds, except for geranyl acetone, occur in an additional 7 herbal genera, including balm (*Melissa*), senna (*Cassia*), coriander (*Coriandrum*), germander (*Teucrium*), mint (*Mentha*), sage (*Salvia*), and thyme (*Thymus/Thymbra*).

It may be significant that only *Satureja*, *Cassia*, and *Salvia* are possibly native to Egypt. Because *A. seibeni* and *T. annuum* are probably native to Iran and Morocco, respectively, they are unlikely to have been traded or transplanted to Egypt at this early date. Of the remaining 7 plants, all are arguably native to the southern Levant (24) (for coriander, see ref. 25). Today, they grow in the vicinity of the areas where the Abydos wine was made according to the INAA results (26), viz., the Jordan Valley, the uplands to its east and west, and the Mediterranean littoral near Gaza.

For the Gebel Adda sample, only rosemary (*Rosmarinus officinalis*), a member of the mint family (Lamiaceae or Labiatae), could explain fenchone, camphor, borneol, cuminaldehyde, and vanillin (labeled 3 in Table S1). Rosemary, which can tolerate relatively dry conditions, grows both in the southern Levant and Egypt (27).

As yet, unique biomarkers for the herbs that might have been added to the Abydos and Gebel Adda wines have not been delimited. Moreover, many of the compounds that we identified are found in other plants and herbs of the eastern Mediterranean region. For example, camphor, borneol, carvone, and thymol are reported in yarrow (*Achillea*); the same compounds, with the exception of borneol, in wild fennel (*Foeniculum vulgare*); linalool and thymol in marjoram; and only borneol in oregano. Two herbs of Egyptian origin should also be noted: *Ambrosia maritima* contains camphor and carvone and *Conyza dioscorides* has camphor and linalool. Linalool, alpha-terpineol, and geranyl acetone also occur in grape wine, and camphor and borneol are found in pine resin, in addition to rosemary and many other herbs. Thus, although many different permutations of the detected compounds deriving from various natural products might be proposed, the most straightforward and simplest explanation is that one or more of the herbs, which can account for the greatest number of relatively rare compounds in the Abydos and Gebel Adda samples, were the likely additives to the wines.

Both ancient samples also contained compounds found in pine resin (labeled 4 in Table S1). Our previous HPLC results for the Abydos sample had already pointed to a tree resin additive, which we had tentatively identified as terebinth. However, a diterpenoid compound specific to pine—retene, an oxidative product of abietic acid (17)—was observed by TD GC/MS. Conventional GC/MS analysis of the Abydos sample also revealed the presence of additional oxidative products of the acid, viz., dehydroabietic acid and 7-oxodehydroabietic acid. Triterpenoids characteristic of terebinth resin (28) were absent.

The Gebel Adda sample had not been analyzed for tree resins in our initial study. In addition to retene, methyl dehydroabietate was detected by SPME and TD GC/MS. Methyl dehydroabietate (Fig. S4) points to the pine resin having been processed by heat to the tar, which is consistent with the entire interior of the amphora having been lined with a dark coating as a sealant.

### Discussion and Conclusions

By employing LC/MS/MS to analyze the Abydos and Gebel Adda samples, 2 Egyptian samples separated in time by over 3,500 years, we have confirmed our earlier methodology for identifying tartaric acid/tartrate, the key biomarker for wine and grape products. By demonstrating that 3 independent techniques—FTIR spectrometry, HPLC, and a Feigl spot test—attested to the presence of tartaric acid/tartrate in the ancient samples and by making statistical comparisons with large IR and UV databases of ancient wine reference samples for wine and modern standards, we were reasonably confident that our results would hold up. Moreover, archaeological considerations—admittedly at a lower level of probability—supported a “wine hypothesis” as best accounting for the available evidence.

Certainly, a wine hypothesis stands or falls on whether tartaric acid/tartrate can be detected in the ancient residues. As analytical instrumentation improves, chemists and archaeologists alike

should continually test previous results, cull out any “false-positives,” and generate data as they are made available by excavation or analytical techniques. Our analyses of the Abydos and Gebel Adda samples provide an illustrative example of how this can be accomplished in the constantly evolving field of biomolecular archaeology.

The reanalysis of the Abydos and Gebel Adda samples by SPME and TD GC/MS had another welcome unintended consequence in addition to confirming the presence of tartaric acid/tartrate. These sensitive versatile techniques enabled us to make a start on the chemical detection of ancient Egyptian herbal medicines. Much work remains to be done in refining and substantiating such findings, but ancient wine and other alcoholic beverages are known to be an excellent means to dissolve and administer herbal concoctions externally and internally (29). Indeed, before modern synthetic medicines became available, alcoholic beverages were the universal palliative.

Chemical analysis opens a door onto early Egyptian pharmacology by providing contemporaneous data of plant additives in ancient alcoholic beverages. In particular, adding a tree resin to wine, principally to protect against wine disease as already noted, was one of the most popular and widespread practices throughout the ancient world. Resinated wines were still being made in the Middle Ages according to the extensive agricultural and medical compilations based on classical writings collectively known as the *Geoponica* (e.g., ref. 30). An appreciation for the medicinal value of tree resins was not restricted to the Middle East and Mediterranean. In the Yellow River valley of China, probable resins of China fir, in the *Elemi* family of fragrant trees, and other native species were added to fermented beverages made from rice, millet, and fruits as early as 7000 B.C., according to multiple analyses by FTIR spectrometry, GC/MS, LC/MS, isotope analysis, and spot tests (8).

Until now, textual sources, in particular a series of medical papyri, have provided the primary evidence for the ancient Egyptian *materia medica*, which was renowned in the ancient world (11, 12, 31). The majority of the papyri belong to the New Kingdom, including the longest one, the 108-page Ebers papyrus, dated to ca. 1550 B.C. Several papyri have been dated as early as the mid-12th Dynasty, ca. 1850 B.C., and the Egyptian word for “physician” (*swmw*), which involved diagnosing disease and often included treatment with herbal remedies, occurs as early as Dynasty 3, ca. 2650 B.C. (12). A later tradition, which is not independently supported, states that Djer, the second pharaoh of Dynasty 1, was also a *swmw* (12); this is an intriguing possibility that may relate to the chemical findings from tomb U-j, because Djer ruled shortly after Scorpion I, ca. 3100 B.C., and his tomb at Abydos is close to U-j.

The prescriptions in the papyri, numbering more than 1,000, present a detailed picture of the ancient Egyptian pharmacopeia, even if more than 80% of some 160 hieroglyphic plant names defy translation. Many vegetables and fruits, including garlic, onion, celery, *Cyperus* grass tubers, watermelon, fig, moringa, persea, and zizyphus, for example, figure prominently as ingredients in the formulations; however, by far the most numerous are alcoholic beverages (wine and beer), tree resins (e.g., terebinth, pine, frankincense, myrrh, fir), and herbs of all kinds (e.g., bryony, coriander, cumin, mandrake, dill, aloe, wormwood). These plants and their exudates are described as being macerated; mixed together; steeped as a decoction or infusion in wine or vinegar, beer, honey, milk, oil, and/or water; strained; and administered for specific ailments (e.g., laxatives, emollients, expectorants, anthelmintics, analgesics, diuretics, aphrodisiacs). Many of the ingredients are still part of the herbal medical tradition of modern Egypt.

Among the most probable herbal additives to the Abydos wine, only coriander is known by its ancient Egyptian name (*š3w*). Eight baskets (half a liter) of coriander mericarp in the tomb of Tutankhamun underline its importance in ancient Egyptian culture and medicine (32). Coriander is explicitly listed in several medical prescriptions (11). Thus, stomach problems called for drinking a

beer mixed with this herb, bryony, flax, and dates. For the treatment of blood in the stool, it was to be grated and mixed with chaste-tree and another unidentified fruit, infused into beer, strained, and drunk. For treating herpes, an external salve was prepared from coriander seeds, myrrh, and fermented honey.

The other herbal additive possibilities for the Abydos wine—balm, senna, germander, mint, sage, savory, and thyme—are yet to be certainly identified by their Egyptian names, although more intensive linguistic study promises their elucidation. For example, quite possibly, *k3y* in a recipe for *kyphi* (11), a well-known temple fumigant and beverage additive, is to be translated as “mint.” According to inscriptions in the late first millennium B.C. temples at Edfu and Philae, *kyphi* was prepared by grinding and sieving equal amounts of sweet flag (*Acorus calamus*), aromatic rush (*Andropogon schoenanthus*), terebinth resin, cassia, mint, and possibly *aspalathus*. This powder was then combined with separately prepared concoctions of wine with juniper berries, *Cyperus* and other plants, raisins and wine, and frankincense and honey. The addition of finely ground myrrh completed the recipe.

This literary attestation of an ancient Egyptian resinated herbal wine has only been partly borne out by our analysis of the Abydos sample, because, apart from the wine itself and possibly mint, the other plant components of *kyphi* were not chemically attested. We are not arguing that *kyphi*, per se, had been developed and used as early as 3150 B.C. Rather, our contention is that plant additives, including various herbs and tree resins, were already being dispensed via alcoholic beverages millennia earlier than the Edfu and Philae temple inscriptions.

The most probable herbal additives to the Abydos wine, on current evidence, share another important feature in common: nearly all were domesticated or cultivated in the southern Levant in advance of their introduction into Egypt (33). This circumstance is in accord with the wine in the Abydos jars having most likely been made in this region. Beginning ca. 3000 B.C., as the domesticated grapevine was transplanted to the Nile Delta, one may reasonably hypothesize that some southern Levantine herbs accompanied or soon followed it into the gardens and fields of the country. These developments considerably expanded the Egyptian pharmacopeia.

Other researchers have begun to report botanical and chemical evidence for herbal concoctions in alcoholic beverages. From about the same time as the Abydos wine, native rosemary and mint, together with thyme, were added to a fermented emmer wheat-barley beverage at Genó in Spain (34). Wild rosemary was also an ingredient, along with bog myrtle, yarrow, and other herbs, in gruit, the principal bittering agent in early medieval European beer (35).

Other archaeochemical and archaeobotanical evidence for rosemary is of special interest, because only this herb had a high probability of having been added to the Gebel Adda resinated wine. Unfortunately, the ancient Egyptian word for the herb is unknown; thus, its place in Egyptian medicine cannot be tracked. It is known that rosemary was a popular food flavorant in Roman and Byzantine times (36) when the Gebel Adda resinated wine was produced. Moreover, it contains numerous antioxidant compounds (e.g., rosmarinic acid, carnosol) that have potentially wide-ranging medical benefits (37–38).

Although much remains to be discovered about ancient Egyptian herbal wines, our chemical findings from Abydos and Gebel Adda, together with the results from Malkata, attest to their great antiquity and importance from the country’s initial unification under the pharaohs and continuing for millennia.

Further refinements in analytical techniques will undoubtedly reveal other important compounds in ancient wines. For example, the Gebel Adda sample was recently analyzed by Fourier-transform ion cyclotron resonance/MS. Signals consistent with the structure of resveratrol, the well-known antioxidant that has anticancer effects and has been shown to extend life in many organisms, were detected. Follow-up confirmatory studies by ultraperformance

liquid chromatography/MS are now being carried out (P. Schmitt-Kopplin, R. Gougeon, personal communication, 2009).

**ACKNOWLEDGMENTS.** We thank Abdul Mabud, Jeffrey Ammann, and Vanessa Kinton of the Tax and Trade Bureau's Scientific Services Division for contributing technical expertise and ongoing support. Günter Dreyer of the German Archaeological Institute in Cairo advised on the archaeology of the Abydos tomb. W. Christian Petersen assisted in the GC/MS analyses at the Scientific Research and Analysis Laboratory of Winterthur Museum and Country Estate and consulted on chemical matters, together with Olga Jáuregui of the University of Barcelona and

Theodore Davidson of the Museum Applied Science Center for Archaeology, Naomi F. Miller of the Museum Applied Science Center for Archaeology and Mark Nesbitt of the Royal Botanic Gardens, Kew, aided in the interpretation of the archaeobotanical data. Recent work using Fourier-transform ion cyclotron resonance/MS and ultraperformance liquid chromatography/MS is part of a collaboration with Philippe Schmitt-Kopplin of the Helmholtz Zentrum München and Régis Gougeon of the Institut Universitaire de la Vigne et du Vin-Jules Guyot, Dijon. Detailed information on the extraction methods for the LC/MS/MS analyses and on the experimental conditions for the SPME, TD, and liquid-injection GC/MS analyses is provided in *SI Text*. A grant from the Abramson Cancer Center of the University of Pennsylvania provided partial financial assistance.

1. Johns T (1990) *With Bitter Herbs They Shall Eat It: Chemical Ecology and the Origins of Human Diet and Medicine* (Univ of Arizona Press, Tucson, AZ).
2. McGovern PE (2007) *Ancient Wine: The Search for the Origins of Viniculture* (Princeton Univ Press, Princeton).
3. Dudley R (2004) Ethanol, fruit ripening, and the historical origins of human alcoholism in primate frugivory. *Integr Comp Biol* 44:315–323.
4. Dudley R (2000) Evolutionary origins of human alcoholism in primate frugivory. *Q Rev Biol* 75:3–15.
5. Stephens D, Dudley R (2004) The drunken monkey hypothesis. *Nat Hist* 113:40–44.
6. Wiens F, et al. (2008) Chronic intake of fermented floral nectar by wild treeshrews. *Proc Natl Acad Sci USA* 105:10426–10431.
7. Nesse RM, Berridge KC (1997) Psychoactive drug use in evolutionary perspective. *Science* 278:63–67.
8. McGovern PE, et al. (2004) Fermented beverages of pre- and proto-historic China. *Proc Natl Acad Sci USA* 101:17593–17598.
9. McGovern PE, Voigt MM, Glusker DL, Exner LJ (1996) Neolithic resinated wine. *Nature* 381:480–481.
10. Dillehay TD, et al. (2008) Monte Verde: seaweed, food, medicine, and the peopling of South America. *Science* 320:784–786.
11. Manniche L (1989) *An Ancient Egyptian Herbal* (Univ of Texas Press, Austin, TX), pp 57–58, 94, 120, 144, 159–162.
12. Nunn JF (1996) *Ancient Egyptian Medicine* (Univ. of Oklahoma Press, Norman, OK), pp 24–42, 121–122, 124, 131–132, 136–162, 211–212 (Appendix B).
13. Dreyer G (1998) *Umm el-Qaab I: Das prädynastische Königsgrab U-j und seine frühen Schriftzeugnisse*, Archäologische Veröffentlichungen 86 (P von Zabern, Mainz, Germany).
14. McGovern PE, Glusker DL, Exner LJ (2001) The organic contents of the Tomb U-j Syro-Palestinian type jars: resinated wine flavored with fig. *Umm el-Qaab II: Importkeramik aus dem Friedhof U in Abydos (Umm el-Qaab) und die Beziehungen ägyptens zu Vorderasien im 4. Jahrtausend v. Chr.*, ed Hartung U (P von Zabern, Mainz, Germany), pp 399–403.
15. Cavalieri D, McGovern PE, Hartl DL, Mortimer R, Polsinelli M (2003) Evidence for *S. cerevisiae* fermentation in ancient wine. *J Mol Evol* 57:5226–5232.
16. McGovern PE, Badler V, Michel RH (1990) Drink and be merry!: Infrared spectroscopy and ancient Near Eastern wine. *Organic Contents of Ancient Vessels: Materials Analysis and Archaeological Investigation*, MASCA Research Papers in Science and Archaeology, eds Biers WR, McGovern PE (MASCA, University of Pennsylvania Museum, University of Pennsylvania, Philadelphia), Vol 7, pp 25–36.
17. Beck CW, Stout E, Lee KC, Chase AA, DeRosa N (2008) Analysis of organic remains in the fabric of Minoan and Mycenaean pottery sherds by gas chromatography-mass spectrometry. *Archaeology Meets Science: Biomolecular Investigations in Bronze Age Greece*, eds Tzedakis Y, Martlew H, Jones MK (Oxbow Books, Oxford), pp 12–47.
18. McGovern PE (1997) Wine of Egypt's golden age: An archaeochemical perspective. *J Egypt Archaeol* 83:69–108.
19. Guasch-Jané MR, Ibern-Gómez M, Andrés-Lacueva C, Jáuregui O, Lamuela-Raventós RM (2004) Liquid chromatography with mass spectrometry in tandem mode applied for the identification of wine markers in residues from ancient Egyptian vessels. *Anal Chem* 76:1672–1677.
20. McGovern PE (1997) A ceramic sequence for northern Jordan: An archaeological and chemical perspective. *Studies in the History and Archaeology of Jordan VI*, eds Bisheh G, Zaghoul M, Kehrberg I (Department of Antiquities, Amman, Egypt), pp 421–425.
21. Hamm S, Bleton J, Connan J, Tchapl A (2005) A chemical investigation by headspace SPME and GC-MS of volatile and semi-volatile terpenes in various olibanum samples. *Phytochemistry* 66:1499–1514.
22. Seidel A, ed (2004) *Kirk-Othmer Encyclopedia of Chemical Technology* (Wiley-Interscience, Hoboken, NJ). Available at <http://mrw.interscience.wiley.com/emrw/9780471238966/home/>.
23. Clarke RJ, Bakker J (2004) *Wine Flavour Chemistry* (Blackwell, Oxford).
24. Feinbrun-Dothan N (1978) *Flora Palaestina*, Part 3: *Ericaceae to Compositae*, ed Zohary M (Israel Academy of Sciences and Humanities, Jerusalem).
25. Zohary D, Hopf M (2000) *Domestication of Plants in the Old World: The Origin and Spread of Cultivated Plants in West Asia, Europe, and the Nile Valley* (Oxford Univ Press, Oxford), pp 205–206.
26. McGovern PE (2001) The origins of the tomb U-j Syro-Palestinian type jars as determined by neutron activation analysis. *Umm el-Qaab II: Importkeramik aus dem Friedhof U in Abydos (Umm el-Qaab) und die Beziehungen ägyptens zu Vorderasien im 4. Jahrtausend v. Chr.*, ed Hartung U (P von Zabern, Mainz, Germany), pp 407–416.
27. Davis, PH (1982) *Flora of Turkey and the East Aegean Islands* (Edinburgh Univ Press, Edinburgh), Vol 7.
28. Mills JS, White R (1994) *The Organic Chemistry of Museum Objects* (Butterworth-Heinemann, Oxford).
29. Majno G (1975) *The Healing Hand: Man and Wound in the Ancient World* (Harvard Univ Press, Cambridge, MA).
30. Owen T, trans (1805–1806) *Geoponika: Agricultural Pursuits* (printed for the author, London).
31. Grapow H (1954–1973) *Grundriss der Medizin der alten Ägypter* (Akademie, Berlin).
32. Germer R (1989) *Die Pflanzenmaterialien aus dem Grab des Tutanchamun* (Gerstenberg, Hildesheim, Germany).
33. Germer R (1993) Ancient Egyptian pharmaceutical plants and the eastern Mediterranean. *The Healing Past: Pharmaceuticals in the Biblical and Rabbinic World*, eds Jacob I and W (E.J. Brill, Leiden, Germany), pp 69–80.
34. Juan-Tresserras J (1998) La cerveza prehistórica: investigaciones arqueobotánicas y experimentales. *Genó: un poblado del Bronce Final en el Bajo Segre (Lleida)*, eds Maya JL, Cuesta F, López Cachero J (University of Barcelona, Barcelona), pp 239–252.
35. Nelson M (2005) *The Barbarian's Beverage: A History of Beer in Ancient Europe* (Routledge, London).
36. Dalby A (1996) *Siren Feasts: A History of Food and Gastronomy in Greece* (Routledge, London), p 190, n. 21.
37. Lai PK, Roy J (2004) Antimicrobial and chemopreventive properties of herbs and spices. *Curr Med Chem* 11:1451–1460.
38. Aggarwal MM, Shishodia S (2006) Molecular targets of dietary agents for prevention and therapy of cancer. *Biochem Pharmacol* 71:1397–1421.