

# An alternate pathway to long-chain polyunsaturates: the *FADS2* gene product $\Delta 8$ -desaturates 20:2n-6 and 20:3n-3

Woo Jung Park, Kumar S. D. Kothapalli, Peter Lawrence, Cynthia Tyburczy, and J. Thomas Brenna<sup>1</sup>

Division of Nutritional Sciences, Cornell University, Ithaca, NY 14853

**Abstract** The mammalian  $\Delta 6$ -desaturase coded by fatty acid desaturase 2 (*FADS2*; HSA11q12-q13.1) catalyzes the first and rate-limiting step for the biosynthesis of long-chain polyunsaturated fatty acids. *FADS2* is known to act on at least five substrates, and we hypothesized that the *FADS2* gene product would have  $\Delta 8$ -desaturase activity. *Saccharomyces cerevisiae* transformed with a *FADS2* construct from baboon neonate liver cDNA gained the function to desaturate 11,14-eicosadienoic acid (20:2n-6) and 11,14,17-eicosatrienoic acid (20:3n-3) to yield 20:3n-6 and 20:4n-3, respectively. Competition experiments indicate that  $\Delta 8$ -desaturation favors activity toward 20:3n-3 over 20:2n-6 by 3-fold. Similar experiments show that  $\Delta 6$ -desaturase activity is favored over  $\Delta 8$ -desaturase activity by 7-fold and 23-fold for n-6 (18:2n-6 vs 20:2n-6) and n-3 (18:3n-3 vs 20:3n-3), respectively. In mammals, 20:3n-6 is the immediate precursor of prostaglandin E1 and thromboxane B1. 20:3n-6 and 20:4n-3 are also immediate precursors of long-chain polyunsaturated fatty acids arachidonic acid and eicosapentaenoic acid, respectively. **■** These findings provide unequivocal molecular evidence for a novel alternative biosynthetic route to long-chain polyunsaturated fatty acids in mammals from substrates previously considered to be dead-end products.—Park, W. J., K. S. D. Kothapalli, P. Lawrence, C. Tyburczy, and J. T. Brenna. An alternate pathway to long-chain polyunsaturates: the *FADS2* gene product  $\Delta 8$ -desaturates 20:2n-6 and 20:3n-3. *J. Lipid Res.* 2009. 50: 1195–1202.

**Supplementary key words** polyunsaturated fatty acid biosynthesis • dihomo- $\gamma$ -linolenic acid • eicosanoid precursor biosynthesis

Long-chain polyunsaturated fatty acids (LCPUFAs) are ubiquitous in mammalian tissue, achieving highest concentrations in the membranes of neural and other excitable tissue (1). LCPUFA of the n-3 and n-6 families, especially eicosapentaenoic acid (EPA; 20:5n-3), docosahexaenoic acid (22:6n-3), and arachidonic acid (20:4n-6), are bioactive components of membrane phospholipids and serve as sub-

strates for signaling molecules (2). The degree of unsaturation of the membranes is determined by the action of enzymes involved in fatty acid biosynthesis and metabolism (3). Most organisms synthesize unsaturated fatty acids, but the pathways are specific to cell types and species.

Fatty acid desaturases are enzymes that catalyze the introduction of *cis* double bonds at specific positions in a fatty acid chain (4). Desaturases in plants and lower animal species can introduce double bonds near the methyl end. Eukaryotic cells of higher animals, fungi, and dinoflagellates express membrane-bound acyl-CoA front-end desaturases (5, 6) catalyzing double bond introduction into the  $\Delta 6$ ,  $\Delta 5$ ,  $\Delta 8$ , and  $\Delta 4$  positions. Mammalian front-end desaturases operate on diet-derived PUFA to synthesize LCPUFA, which can also be derived from the diet but possibly not in sufficient quantities to optimize health (7).

The front-end desaturases are remarkable for their structural similarity and functional diversity. They all contain the N-terminal cytochrome b5 domain (HPGG) as electron donor and three histidine motifs, HXXXH, HXXXH, and QXXHH, conserved from human to microalgae (8). Molecular cloning and isolation of a  $\Delta 5$ -desaturase from *Caenorhabditis elegans* (9) and *Mortierella alpina* (10) and a  $\Delta 6$ -desaturase from *C. elegans* (11), *M. alpina* (12), rat (13), and mouse (14) have all been reported. The human fatty acid desaturase (*FADS*) gene cluster at 11q12-q13.1 encodes two desaturases with known function,  $\Delta 5$ -desaturase (*FADS1*) and  $\Delta 6$ -desaturase (*FADS2*) (14, 15), as well as a third putative desaturase gene (*FADS3*) (16), which thus far has no known substrate despite high homology to *FADS1* and *FADS2*.

**Figure 1** shows the common n-3 and n-6 LCPUFA pathways mediated by  $\Delta 6$  and  $\Delta 5$  desaturases. The  $\Delta 6$ -desaturase (*FADS2*) is known to operate on both 18:3n-3 and 18:2n-6, resulting in the synthesis of 6,9,12,15-18:4 and 6,9,12-18:3 ( $\gamma$ -linolenic acid), respectively. This step is rate limiting

Abbreviations: EPA, eicosapentaenoic acid; FADS, fatty acid desaturase; FAME, fatty acid methyl ester; GC-CACI-MS/MS, gas chromatography-covalent adduct chemical ionization tandem mass spectrometry; LCPUFA, long-chain polyunsaturated fatty acids; wt, wild type.

<sup>1</sup> To whom correspondence should be addressed.

e-mail: jtb4@cornell.edu

This work was supported by National Institutes of Health Grant GM071534.


Manuscript received 5 December 2008 and in revised form 21 January 2009.

Published, JLR Papers in Press, February 6, 2009.

DOI 10.1194/jlr.M800630-JLR200

Copyright © 2009 by the American Society for Biochemistry and Molecular Biology, Inc.

This article is available online at <http://www.jlr.org>



**Fig. 1.** Pathways for LCPUFA biosynthesis. The conventional pathway consists of alternating desaturation and elongation leading to LCPUFA.  $\Delta 8$ -Desaturation of 20:2n-6 and 20:3n-3 would yield 20:3n-6 and 20:4n-3, intermediates in the conventional pathway to 20:4n-6 and 20:5n-3, as well as immediate eicosanoid precursors.

and is followed by elongation to 8,11,14,17-20:4 and 8,11,14-20:3 (dihomo- $\gamma$ -linolenic acid). A rapid  $\Delta 5$ -desaturation (*FADS1*) on these PUFA produces EPA and arachidonic acid. EPA can be further elongated and desaturated to yield docosaheptaenoic acid by the pathway shown, which is accepted as mammalian pathway, or via a  $\Delta 4$ -desaturase as demonstrated in *Thraustochytrium* (17).

The operation of an alternative pathway via C20 fatty acids using a  $\Delta 8$ -desaturase reported in unicellular organisms (18–21) has been verified by molecular cloning and functional characterization studies in *Euglena gracilis* (22), *Acanthamoeba castellanii* (23), and *Perkinsus marinus* (24). There are many reports of  $\Delta 8$ -desaturation activity in mammalian cells (25, 26), in rat and human testes (27, 28), and in mouse liver (29), though it has not been verified by molecular cloning, and the existence of  $\Delta 8$ -desaturation in rat microsomes has been questioned (30). The putative substrate of the  $\Delta 8$ -desaturase, 11,14-eicosadienoic acid (20:2 n-6), is found in human plasma and red cells as well as other tissues, and its concentration has recently been associated with human genetic variation in the *FADS* gene cluster (31, 32).

The mammalian  $\Delta 6$ -desaturase coded by *FADS2* uses at least five substrates, 18:2n-6, 18:3n-3, 24:6n-3, 24:5n-3 (33, 34), and 16:0.  $\Delta 6$ -desaturase in the sebaceous glands catalyzes desaturation of 16:0 to 16:1n-10 (sapienate), the most abundant fatty acid in human sebum, showing that substrate specificity is influenced by the cellular environment in which it is expressed (35). We hypothesized that the primate *FADS2* gene product would have  $\Delta 8$ -desaturase activity and cloned baboon *FADS2* into *Saccharomyces cerevisiae*, an organism with no native PUFA biosynthetic capability, to test for gain of  $\Delta 8$ -desaturase activity. Here, we report unambiguous evidence of the existence of  $\Delta 8$ -desaturation in primates, suggesting alternative pathway for LCPUFA biosynthesis.

## RNA isolation and cDNA synthesis

Total RNA from 30 mg neonate baboon liver tissue homogenate was extracted using the RNeasy Mini kit (Qiagen, Valencia, CA). The yield of total RNA was assessed by 260 nm UV absorption. The quality of RNA was analyzed by 260/280 nm ratios of the samples and by agarose gel electrophoresis to verify RNA integrity. One microgram total RNA was reverse transcribed into first-strand cDNA using the iScript cDNA synthesis kit (Bio-Rad, Hercules, CA). The resulting cDNA was used as template for RT-PCR reactions.

## Cloning of baboon *FADS2* and sequence analysis


To identify baboon *FADS2* cDNA sequence, primers were generated using human cDNA sequences for *FADS2* (GenBank accession number NM\_004265). PCR primers, *FADS2* forward (5'-ATGGGGAAGGGAGGGAACCAGGGCGA-3') and *FADS2* reverse (5'-TCATTGTGAAGGTAGGCGTCCAGCCA-3') were ordered from Integrated DNA Technologies (Coralville, IA) and were amplified with baboon liver cDNA as template and high-fidelity Taq polymerase (Roche Diagnostics) using Eppendorf gradient thermal cycler. Cycling conditions were as follows: initial denaturation at 95°C for 5 min followed by 40 cycles of denaturation at 95°C for 30 s, annealing at 72°C for 45 s, and extension at 72°C for 1 min, with a final extension at 72°C for 5 min. PCR product was separated by electrophoresis on 2% agarose gel stained with ethidium bromide and band of appropriate size was obtained. The PCR product was gel purified and cloned in pGEM T-Easy vector (Promega) and sequenced using T7 forward and SP6 reverse universal primers at Cornell University Life Sciences Core Laboratories Center using the Applied Biosystems automated 3730 DNA analyzer. We successfully cloned the baboon *FADS2* protein coding region (GenBank accession number EU780003). The pGEM T-Easy vector with *FADS2* was named pTFADS2.

## Transformation into yeast (*S. cerevisiae*)

The entire coding regions of baboon *FADS2* was amplified from pTFADS2 with primers *FADS2*-KOZAK forward (5'-CCCAAGCTTACCATGGGGAAGGGAGGGAACCAGGGCGA-3') including the *Hind*III site and *FADS2*-KOZAK reverse (5'-CCGCTCGAGTCATTTGTGAAGGTAGGCGTCCAGCCA-3') including *Xho*I site. The high-fidelity Taq polymerase (Roche Diagnostics) was used to minimize potential PCR errors. The amplified PCR product containing baboon *FADS2* was gel purified, restriction digested, and inserted into *Hind*III and *Xho*I sites behind the GAL1 promoter of pYES2 vector (Invitrogen) to yield the plasmid pYFADS2. The constructed plasmid of pYFADS2 was transformed into *S. cerevisiae* (strain INVSc1 from Invitrogen) using *S. c.* Easy Comp™ Transformation Kit (Invitrogen), and the transformants were verified by DNA sequencing.

## Expression of baboon *FADS2*

For functional expression characterization, transformed yeast strains with pYES2 (empty vector) as a negative control and pYFADS2 were grown for 24 h in *S. cerevisiae* minimal media without uracil. As another negative control, wild *S. cerevisiae* (INVSc1) was cultured in *S. cerevisiae* minimal medium with uracil. Expression of the transgene was induced when OD<sub>600</sub> reached 0.4. At that time, appropriate fatty acids, 1 mM linoleic acid (18:2n-6),  $\alpha$ -linolenic acid (18:3n-3), eicosadienoic acid (20:2n-6), and eicosatrienoic acid (20:3n-3), were added in the presence of 1% tert-butylhydroquinone (Sigma-Aldrich) to the cultures and were grown at 30°C with constant shaking. The samples were collected after 48 h for fatty acid analysis. All treatments were performed in duplicate.



**Fig. 2.** Data showing *FADS2* action on 20:2n-6. A: Reconstructed ion chromatograms of FAME derived from 48 h incubation of 20:2n-6 with wt *S. cerevisiae*; B: *S. cerevisiae* transformed with empty vector; and C: *S. cerevisiae* transformed with *FADS2* showing novel product. D: CACI-MS1 spectrum of novel product in C showing diagnostic ions for a 20:3 FAME. E: CACI-MS/MS spectrum of the  $[M+54]^+$  showing diagnostic ions characteristic of 20:3. Masses of fragments shown in the structural inset indicate the diagnostic ions characteristic of 20:3n-6 ( $m/z$  190 and 262).


### Fatty acid analysis

The yeast cells were harvested by centrifugation at 4,000 rpm for 5 min. The cell pellets were washed twice with tergitol-Nonidet P-40 and finally twice with distilled water. Fatty acid methyl esters (FAMES) were prepared using modified one-step lipid extraction method of Garces and Mancha (36). FAMES were structurally identified by gas chromatography-covalent adduct chemical ionization tandem mass spectrometry (GC-CACI-MS/MS) (37–39) and quantitatively analyzed by GC-flame ionization detection. An equal

weight FAME mixture was used to verify response factors on a daily basis (40). For competition experiments, GC analyses were performed in triplicate.

### Materials (Chemicals)

Fatty acids (18:2n-6, 18:3n-3, 20:2n-6, and 20:3n-3) were purchased from Nu-Chek Prep (MN). Uracil dropout SD-U medium and supplement contents including amino acids were obtained from Clontech, TaKaRa Bio. Uracil and tergitol-Nonidet P-40 was from Sigma-



**Fig. 3.** Data showing *FADS2* action on 20:3n-3. Panels are analogous to those in Fig. 2 and demonstrate synthesis of 20:4n-3.

Aldrich. pGEM-T Easy Vector II system was purchased from Promega. The pYES2 vector, INVSc1 strain, *S. c.* Easy Comp™ Transformation Kit, and restriction enzymes (*Hind*III and *Xho*I) were obtained from Invitrogen. Total RNA was isolated by using RNeasy Mini kit from Qiagen. The cDNA synthesis kit was purchased from Bio-Rad.

## RESULTS

### Baboon *FADS2*

The sequenced PCR product (Baboon *FADS2*; GenBank EU780003) revealed an open reading frame of 1,335 bp, en-

coding a protein of 444 amino acids and a stop codon. It shares a 60% homology with baboon *FADS1* (EF531577) and 62% homology with the putative baboon *FADS3* (EU780002), including HPGG characteristic of a cytochrome b5 domain and three conserved histidine motifs, HXXXXH, HXXXHH, and QXXXHH. Analysis and comparison of amino acid sequence of baboon *FADS2* showed 97% identity and 99% similarity with human *FADS2* (AAH09011), and 64% identity and 79% similarity with the bifunctional zebrafish desaturase (AAG25710). Baboon *FADS2* also shares homology with  $\Delta 8$ -desaturases from unicellular organisms [27%

identity and 43% similarity with *E. gracilis* (AAD45877), 27% identity and 40% similarity with *P. marinus* (ABF58684), and 23% identity and 38% similarity with *A. castellanii* (CAO00489)]. Analysis of the baboon *FADS2* secondary structure by SOSUI software (41) predicted three transmembrane regions, whereas baboon *FADS1* and the putative baboon *FADS3* had four transmembrane regions (data not shown).


The transformed yeast grown on minimal media were supplemented with various fatty acids, incubated at 30°C and harvested after 48 h. Wild-type (wt) *S. cerevisiae* and *S. cerevisiae* containing empty pYES2 vector were used as controls for every replicate. GC-CACI-MS chromatograms of FAME are presented in **Figs. 2 and 3** for 20:2n-6 and 20:3n-3 incubations, respectively. A–C of both figures correspond to wt *S. cerevisiae*, *S. cerevisiae* containing empty pYES2 vector, and *S. cerevisiae* containing *FADS2*, respectively. *S. cerevisiae* wt and *S. cerevisiae* with empty pYES2 vector have no activity toward 20:3n-6 and 20:4n-3, as expected. Fig. 2C shows a new product appearing upon incubation with 20:2n-6. Panel D is the MS-1 spectrum showing peaks at *m/z* 374, 321, 289, and 271, corresponding to the  $[M+54]^+$ ,  $[MH]^+$ ,  $[MH-32]^+$ , and  $[MH-32-18]^+$  ions, respectively, characteristic of a 20:3 FAME. Panel E is the collisional dissociation spectrum of  $[M+54]^+$ , yielding ions at *m/z* 262 and 190 corresponding, respectively, to the  $\alpha$  and  $\omega$  diagnostic ions for 20:3n-6 and positively identifying this product. Similarly, Fig. 3D displays MS1 ions characteristic of 20:4 (*m/z* 372, 319, 287, and 269), and collisional dissociation yields diagnostic ions *m/z* 302 and 188 in MS/MS, positively identifying 20:4n-3.

An alternative hypothesis to  $\Delta 8$ -desaturation of 20:2n-6 (20:2n-6  $\rightarrow$  20:3n-6) is  $\beta$ -oxidation followed by entry into the normal pathway (20:2n-6  $\rightarrow$  18:2n-6  $\rightarrow$  18:3n-6  $\rightarrow$  20:3n-6). The wt *S. cerevisiae* contain peroxisomes and mitochondria

and thus have native  $\beta$ -oxidation activity, so this pathway is plausible. **Figure 4A** presents the relative product distribution with 18:2 as a substrate. No  $\Delta 6$ -desaturation is observed for the controls (wt or vector only), while the cells transformed with *FADS2* accumulate 18:3n-6, and  $\sim 8\%$  of this product is further elongated to 20:3n-6 (18:2n-6  $\rightarrow$  18:3n-6  $\rightarrow$  20:3n-6). There is also some elongation of 18:2n-6 to 20:2n-6.

The putative intermediate of a  $\beta$ -oxidation-mediated alternative pathway, 18:3n-6, was detected in only one of two trials with 20:2n-6 used as a substrate. Fig. 4B presents relative product distribution with 20:2n-6 for that trial. There is indeed some  $\beta$ -oxidation to 18:2n-6 in all treatments. However,  $\Delta 6$ -desaturation of this product to 18:3n-6 is detected in only the *FADS2* cells and is  $\sim 7\%$  of the 18:2n-6 product. The  $\Delta 8$ -desaturation product, 20:3n-6, is  $>10$ -fold greater in abundance than 18:3n-6, its putative intermediate in the alternative pathway. Considering the data of Fig. 4A, the conversion of 18:3n-6  $\rightarrow$  20:3n-6 can only account for a negligible fraction of the 20:3n-6. We conclude that  $\Delta 8$ -desaturation by the *FADS2* product mediates direct conversion of 20:2n-6 to 20:3n-6.

Once establishing that *FADS2*-transformed *S. cerevisiae* gained the ability to  $\Delta 8$ -desaturate 20-carbon PUFA, the relative activity was tested in competition experiments.  $\Delta 8$ -desaturase activity toward n-3 and n-6 fatty acids was investigated by supplementing media with a 1:1 mixture of 20:2n-6 and 20:3n-3 as substrates. Substrates and products were analyzed quantitatively by GC-flame ionization detection after confirmation of structure by GC-CACI-MS and GC-CACI-MS/MS. **Table 1** shows that 20:2n-6  $\rightarrow$  20:3n-6 was  $0.9 \pm 0.16\%$ , and 20:3n-3  $\rightarrow$  20:4n-3 was  $2.8 \pm 0.7\%$  over a 48 h incubation, yielding a relative conversion efficiency of  $\sim 3.1$ -fold, favoring the n-3 PUFA.



**Fig. 4.** Analysis of putative intermediates. A: Peak areas of n-6 PUFA, normalized to 20:0, for wt, vector-only, and *FADS2* transformed cells treated with 18:2n-6. A small amount of chain shortening is detected (16:2n-6) in wt and *FADS2* cells. The *FADS2* cells uniquely show accumulation of the  $\Delta 6$ -desaturated product 18:3n-6, the elongated product 20:2n-6, and a small amount of 20:3n-6, the chain elongated product of 18:3n-6. B: All 20:2n-6 treated cells show the chain-shortened product 18:2n-6, and only the *FADS2* cells show a small amount of the elongated product 18:3n-6. The product of  $\Delta 8$ -desaturase activity on 20:2n-6, 20:3n-6 is present only in the *FADS2* cells at about 10-fold higher concentration than 18:3n-6.

TABLE 1. Competition between n-6 and n-3 fatty acids

Substrate Mixture	Reactions	Conversion of Substrate (%)	
		Mean $\pm$ SD	Ratio
20:2n-6+ 20:3n-3	$\Delta 8$ -desaturase 20:2n-6 $\rightarrow$ 20:3n-6	0.90 $\pm$ 0.16	3.1
	$\Delta 8$ -desaturase 20:3n-3 $\rightarrow$ 20:4n-3	2.8 $\pm$ 0.7	

In a second competition experiment, 1:1 mixtures of 18:2n-6+20:2n-6 or 18:3n-3+20:3n-3 were added to media to test the relative  $\Delta 6$ -desaturase and  $\Delta 8$ -desaturase activities. **Table 2** shows that conversions 18:2n-6  $\rightarrow$  18:3n-6 and 20:2 n-6  $\rightarrow$  20:3n-6 were 12.2  $\pm$  0.16% and 1.7  $\pm$  0.54%, respectively, yielding a relative activity of 7.2-fold favoring  $\Delta 6$ -desaturase. The relative conversions 18:3n-3  $\rightarrow$  18:4n-3 and 20:3n-3  $\rightarrow$  20:4n-3 were 24.1  $\pm$  2.4% and 1.03  $\pm$  0.31%, respectively, yielding the substantially greater conversion ratio of 23.

## DISCUSSION

More than 50 years ago, Thomasson (42) showed that the n-6 PUFA were active in supporting growth of water-deprived young rats raised on a diet with saturates as their only source of fat. The order of relative vitamin F activity was 20:4 > 18:2 = 20:3 > 20:2 (131:100:100:43), with relative activity of <10 for all other fatty acids tested, including n-3s. Within two decades, the pathway from 18:2 to 20:4 by sequential  $\Delta 6$ -desaturation-elongation- $\Delta 5$ -desaturation (18:2  $\rightarrow$  18:3  $\rightarrow$  20:3  $\rightarrow$  20:4) emerged as the major route of biosynthesis. The alternative, elongation- $\Delta 8$ -desaturation- $\Delta 5$ -desaturation (18:2  $\rightarrow$  20:2  $\rightarrow$  20:3  $\rightarrow$  20:4), has been of interest over the years because of the appearance of intermediates in mammalian tissue, as has a third alternative, elongation- $\Delta 5$ -desaturation- $\Delta 8$ -desaturation (18:2  $\rightarrow$  20:2  $\rightarrow$  5,11,14-20:3  $\rightarrow$  20:4) because 20:2 is converted to 5,11,14-20:3 (sciadonic acid) by the action of  $\Delta 5$ -desaturase. Experiments involving only activity measurements in mammalian tissue or cells that definitively establish or rule out participation of a  $\Delta 8$ -desaturase are difficult to design because of the low concentration of the intermediates, indicating that molecular techniques capable of isolating a particular biochemical activity are required (30). Indeed, the existence of  $\Delta 8$ -desaturation as an alternative pathway to LCPUFA has been reported periodically, and  $\Delta 8$ -desaturase activity has been found by some and not by others (30). Presently, 20:2n-6 and 20:3n-3 are widely considered dead-end products, in part because their conversion to LCPUFA has not been unequivocally established (43). However, recent studies have associated 20:2n-2 with *FADS2* polymorphisms and/or fatty acid compositions in humans (32, 44), including

patients with cardiovascular disease (31) and type 2 diabetes mellitus (45). In addition, apolipoprotein D knockout mice, a model for psychiatric disorders, show increased CNS 20:2n-6 and 18:2n-6 compared with wild-type mice (46).

Several fatty acids of chain length 16, 18, and 24 carbons are substrates for the *FADS2* gene product. Apart from 18:2n-6 and 18:3n-3, it is also known to  $\Delta 6$ -desaturate 24:5n-3 and 24:4n-6, as required for the coupled microsomal-peroxisomal pathway for 22:6n-3 and 22:5n-6 biosynthesis (33). The *FADS2* gene product catalyzes the  $\Delta 6$ -desaturation of 16:0 (palmitic acid) to 16:1n-10 (*cis*-6-16:1, sapienic acid) when expressed natively in human skin sebocytes (35), and when COS-7 cells were transfected with rat *FADS2* they acquired the ability to  $\Delta 6$ -desaturate 16:0 (34). However, there are no previous reports of *FADS2* gene product activity toward 20 carbon PUFA.

The alternative synthetic pathways to arachidonic acid from 20:2n-6 are either by sequential action of a  $\Delta 8$ -desaturase and a  $\Delta 5$ -desaturase or vice versa. Initial  $\Delta 8$ -desaturation yields the eicosanoid precursor 20:3n-6, whereas initial  $\Delta 5$ -desaturase activity yields 5,11,14-20:3. There are numerous reports showing that 11,14-20:2 is  $\Delta 5$ -desaturated to 5,11,14-20:3 (30, 47–50). However, no clear evidence for the conversion of 5,11,14-20:3 to 20:4n-6 has been found (30, 47, 49). Sprecher and coworkers have studied the desaturation of 11,14-20:2 with isotope labeling in vitro and in vivo, and consistently find that rat liver does  $\Delta 5$ -desaturate it to 5,11,14-20:3, but they find no evidence of  $\Delta 8$ -desaturation activity on this product (47, 49). Fourteen day feeding of 5,11,14-20:3 led to the accumulation of this PUFA in liver phosphoglycerides where it decreased 20:4n-6, while feeding of 11,14-20:2 did not alter 20:4n-6 levels (30). The production of 5,11,14-20:3 and 5,11,14,17-20:4 in human leukemia K562 cells has been reported in which the  $\Delta 5$ -desaturase is the only active desaturase operating because of the lack of  $\Delta 6$ -desaturase activity in these cells (50). Consistent with this report, we recently found significant amounts of 7,11,14-20:3, 7,11,14,17-20:4, and 9,13,16,19-22:4 in the liver lipids of chow-fed *FADS2* null mice, all of which may be synthesized by action of  $\Delta 5$ -desaturase, coded by *FADS1*, on 18:2n-6 or 18:3n-3, followed by prompt elongation (C. Stroud, P. Lawrence, J. T. Brenna, and M. Nakamura, unpublished observations). These data support the hypothesis that the  $\Delta 5$ -desaturase acts on PUFA only when its preferred


TABLE 2. Competition between  $\Delta 6$ - and  $\Delta 8$ -desaturase activities

Substrate Mixture	Reactions	Conversion of Substrate (%)	
		Mean $\pm$ SD	Ratio
18:2n-6+ 20:2n-6	$\Delta 6$ -desaturase 18:2n-6 $\rightarrow$ 18:3n-6	12.2 $\pm$ 0.16	7.2
	$\Delta 8$ -desaturase 20:2n-6 $\rightarrow$ 20:3n-6	1.7 $\pm$ 0.54	
18:3n-3+ 20:3n-3	$\Delta 6$ -desaturase 18:3n-3 $\rightarrow$ 18:4n-3	24.1 $\pm$ 2.4	23
	$\Delta 8$ -desaturase 20:3n-3 $\rightarrow$ 20:4n-3	1.03 $\pm$ 0.31	

substrate is not available, which may well imply that its products found in experimental studies are not relevant in vivo under normal conditions.

Reports of  $\Delta 8$ -desaturase activity in rodent and human testes have appeared (27, 28), and the most recent study shows stable isotope labeling best explained by direct conversion of 11,14-20:2 to 20:4n-6 via  $\Delta 8$ -desaturation, albeit as a minor pathway (30), but there are no existing molecular data to implicate a specific gene responsible for coding for vertebrate  $\Delta 8$ -desaturase activity.  $\Delta 8$ -Desaturation has been shown unequivocally in unicellular organisms where the gene has been cloned and is active when expressed in *Arabidopsis thaliana* (51). A  $\Delta 8$ -desaturase was first reported in the single cell protist *E. gracilis* (22) and later, along with a  $\Delta 9$ PUFA-elongase, in *Isochrysis galbana* (21) as well as the free living amoeba *A. castellanii* (23). The present report is the first to show that a vertebrate gene product introduces a double bond at the  $\Delta 8$  position, demonstrating an alternative pathway to LCPUFA biosynthesis.

The competition experiments provide insight as to whether  $\Delta 8$ -desaturase activity of the *FADS2* protein product can be important in vivo. The synthesis of 20:4n-3 dominates by 3.1-fold over 20:3n-6, consistent with long-established observations for  $\Delta 6$ -desaturase preference for 18:3n-3 over 18:2n-6 (52–54). These observations are also consistent with in vitro work showing that the biosynthesis of n-6 PUFA is strongly suppressed by <2% of calories of 18:3n-3, whereas nearly 10 times as much 18:2n-6 is required to equally suppress n-3 PUFA biosynthesis (55), indicating that the affinity of the biosynthetic apparatus favors n-3 PUFA. Our competition experiments (Table 2) also establish that the *FADS2* gene product exhibits both  $\Delta 6$ -desaturase and  $\Delta 8$ -desaturase activities when both substrates are available. As expected, the *FADS2* gene product showed higher  $\Delta 6$ -desaturase activity by acting on the 18:2n-6 substrate to generate 7-fold more product than for the 20:2n-6. The relative action  $\Delta 6/\Delta 8$  activity toward the n-3 was much greater, at 23-fold, indicating that the conventional pathway would be strongly favored when both substrates are available.

In conclusion, baboon *FADS2* gene cloned into *S. cerevisiae* causes gain of  $\Delta 8$ -desaturase activity, in addition to coding for  $\Delta 6$ -desaturase activity.  $\Delta 8$ -Desaturase activity on 20:2n-6 leads directly to 20:3n-6, the immediate precursor of PGE1 and of 20:4n-6. All available evidence indicates that  $\Delta 8$ -desaturation is a minor pathway, but further study may show that it becomes important when there is high demand for eicosanoid synthesis, such as in inflammation or vasodilation, particularly in situations in which specialized tissues require 20:3n-6 as a precursor to prostaglandins E1 and F1 $\alpha$ , hydroxy-eicosatrienoic acids, or thromboxane B1. This alternative pathway to the eicosanoid precursors may explain data suggesting that 20:2n-6 levels are related to human health. 

## REFERENCES

1. Brenna, J. T., and G. Y. Diau. 2007. The influence of dietary docosahexaenoic acid and arachidonic acid on central nervous system polyunsaturated fatty acid composition. *Prostaglandins Leukot. Essent. Fatty Acids*. **77**: 247–250.
2. Kinsella, J. E., B. Lokesh, S. Broughton, and J. Whelan. 1990. Dietary polyunsaturated fatty acids and eicosanoids: potential effects on the modulation of inflammatory and immune cells: an overview. *Nutrition*. **6**: 24–44 (discussion 59–62).
3. Voss, A., M. Reinhart, S. Sankarappa, and H. Sprecher. 1991. The metabolism of 7,10,13,16,19-docosapentaenoic acid to 4,7,10,13,16,19-docosahexaenoic acid in rat liver is independent of a 4-desaturase. *J. Biol. Chem.* **266**: 19995–20000.
4. Los, D. A., and N. Murata. 1998. Structure and expression of fatty acid desaturases. *Biochim. Biophys. Acta*. **1394**: 3–15.
5. Tocher, D. R., M. J. Leaver, and P. A. Hodgson. 1998. Recent advances in the biochemistry and molecular biology of fatty acyl desaturases. *Prog. Lipid Res.* **37**: 73–117.
6. Nakamura, M. T., and T. Y. Nara. 2004. Structure, function, and dietary regulation of delta6, delta5, and delta9 desaturases. *Annu. Rev. Nutr.* **24**: 345–376.
7. Salem, N., Jr., B. Wegher, P. Mena, and R. Uauy. 1996. Arachidonic and docosahexaenoic acids are biosynthesized from their 18-carbon precursors in human infants. *Proc. Natl. Acad. Sci. USA*. **93**: 49–54.
8. Sperling, P., P. Ternes, T. K. Zank, and E. Heinz. 2003. The evolution of desaturases. *Prostaglandins Leukot. Essent. Fatty Acids*. **68**: 73–95.
9. Michaelson, L. V., J. A. Napier, M. Lewis, G. Griffiths, C. M. Lazarus, and A. K. Stobart. 1998. Functional identification of a fatty acid delta5 desaturase gene from *Caenorhabditis elegans*. *FEBS Lett.* **439**: 215–218.
10. Michaelson, L. V., C. M. Lazarus, G. Griffiths, J. A. Napier, and A. K. Stobart. 1998. Isolation of a Delta5-fatty acid desaturase gene from *Mortierella alpina*. *J. Biol. Chem.* **273**: 19055–19059.
11. Napier, J. A., S. J. Hey, D. J. Lacey, and P. R. Shewry. 1998. Identification of a *Caenorhabditis elegans* Delta6-fatty-acid-desaturase by heterologous expression in *Saccharomyces cerevisiae*. *Biochem. J.* **330**: 611–614.
12. Huang, Y. S., S. Chaudhary, J. M. Thurmond, E. G. Bobik, Jr., L. Yuan, G. M. Chan, S. J. Kirchner, P. Mukerji, and D. S. Knutzon. 1999. Cloning of delta12- and delta6-desaturases from *Mortierella alpina* and recombinant production of gamma-linolenic acid in *Saccharomyces cerevisiae*. *Lipids*. **34**: 649–659.
13. Aki, T., Y. Shimada, K. Inagaki, H. Higashimoto, S. Kawamoto, S. Shigeta, K. Ono, and O. Suzuki. 1999. Molecular cloning and functional characterization of rat delta-6 fatty acid desaturase. *Biochem. Biophys. Res. Commun.* **255**: 575–579.
14. Cho, H. P., M. T. Nakamura, and S. D. Clarke. 1999. Cloning, expression, and nutritional regulation of the mammalian Delta-6 desaturase. *J. Biol. Chem.* **274**: 471–477.
15. Cho, H. P., M. Nakamura, and S. D. Clarke. 1999. Cloning, expression, and fatty acid regulation of the human delta-5 desaturase. *J. Biol. Chem.* **274**: 37335–37339.
16. Marquardt, A., H. Stohr, K. White, and B. H. Weber. 2000. cDNA cloning, genomic structure, and chromosomal localization of three members of the human fatty acid desaturase family. *Genomics*. **66**: 175–183.
17. Qiu, X., H. Hong, and S. L. MacKenzie. 2001. Identification of a Delta 4 fatty acid desaturase from *Thraustochytrium* sp. involved in the biosynthesis of docosahexanoic acid by heterologous expression in *Saccharomyces cerevisiae* and *Brassica juncea*. *J. Biol. Chem.* **276**: 31561–31566.
18. Korn, E. D. 1964. The polyunsaturated 20-carbon and 22-carbon fatty acids of *Euglena*. *Biochem. Biophys. Res. Commun.* **14**: 1–6.
19. Korn, E. D. 1964. Biosynthesis of unsaturated fatty acids in *Acanthamoeba* Sp. *J. Biol. Chem.* **239**: 396–400.
20. Lees, A. M., and E. D. Korn. 1966. Metabolism of unsaturated fatty acids in protozoa. *Biochemistry*. **5**: 1475–1481.
21. Qi, B., F. Beaudoin, T. Fraser, A. K. Stobart, J. A. Napier, and C. M. Lazarus. 2002. Identification of a cDNA encoding a novel C18-Delta (9) polyunsaturated fatty acid-specific elongating activity from the docosahexaenoic acid (DHA)-producing microalga, *Isochrysis galbana*. *FEBS Lett.* **510**: 159–165.
22. Wallis, J. G., and J. Browse. 1999. The Delta8-desaturase of *Euglena gracilis*: an alternate pathway for synthesis of 20-carbon polyunsaturated fatty acids. *Arch. Biochem. Biophys.* **365**: 307–316.
23. Sayanova, O., R. Haslam, B. Qi, C. M. Lazarus, and J. A. Napier. 2006. The alternative pathway C20 Delta8-desaturase from the non-photosynthetic organism *Acanthamoeba castellanii* is an atypical cytochrome b5-fusion desaturase. *FEBS Lett.* **580**: 1946–1952.
24. Chu, F. L., E. D. Lund, E. Harvey, and R. Adlof. 2004. Arachidonic acid synthetic pathways of the oyster protozoan parasite, *Perkinsus marinus*: evidence for usage of a delta-8 pathway. *Mol. Biochem. Parasitol.* **133**: 45–51.

25. Nakazawa, I., J. F. Mead, and R. H. Yonemoto. 1976. In vitro activity of the fatty acyl desaturases of human cancerous and noncancerous tissues. *Lipids*. **11**: 79–82.
26. Bardon, S., M. T. Le, and J. M. Alessandri. 1996. Metabolic conversion and growth effects of n-6 and n-3 polyunsaturated fatty acids in the T47D breast cancer cell line. *Cancer Lett.* **99**: 51–58.
27. Albert, D. H., and J. G. Coniglio. 1977. Metabolism of eicosa-11, 14-dienoic acid in rat testes. Evidence for delta8-desaturase activity. *Biochim. Biophys. Acta.* **489**: 390–396.
28. Albert, D. H., R. K. Rhamy, and J. G. Coniglio. 1979. Desaturation of eicosa-11,14-dienoic acid in human testes. *Lipids*. **14**: 498–500.
29. Schenck, P. A., H. Rakoff, and E. A. Emken. 1996. Delta 8 desaturation in vivo of deuterated eicosatrienoic acid by mouse liver. *Lipids*. **31**: 593–600.
30. Chen, Q., F. Q. Yin, and H. Sprecher. 2000. The questionable role of a microsomal delta8 acyl-coA-dependent desaturase in the biosynthesis of polyunsaturated fatty acids. *Lipids*. **35**: 871–879.
31. Malerba, G., L. Schaeffer, L. Xumerle, N. Klopp, E. Trabetti, M. Biscuola, U. Cavallari, R. Galavotti, N. Martinelli, P. Guarini, et al. 2008. SNPs of the FADS gene cluster are associated with polyunsaturated fatty acids in a cohort of patients with cardiovascular disease. *Lipids*. **43**: 289–299.
32. Schaeffer, L., H. Gohlke, M. Muller, I. M. Heid, L. J. Palmer, I. Kompauer, H. Demmelmair, T. Illig, B. Koletzko, and J. Heinrich. 2006. Common genetic variants of the FADS1 FADS2 gene cluster and their reconstructed haplotypes are associated with the fatty acid composition in phospholipids. *Hum. Mol. Genet.* **15**: 1745–1756.
33. D'Andrea, S., H. Guillou, S. Jan, D. Catheline, J. N. Thibault, M. Bouriel, V. Rioux, and P. Legrand. 2002. The same rat Delta6-desaturase not only acts on 18- but also on 24-carbon fatty acids in very-long-chain polyunsaturated fatty acid biosynthesis. *Biochem. J.* **364**: 49–55.
34. Guillou, H., V. Rioux, D. Catheline, J. N. Thibault, M. Bouriel, S. Jan, S. D'Andrea, and P. Legrand. 2003. Conversion of hexadecanoic acid to hexadecenoic acid by rat Delta 6-desaturase. *J. Lipid Res.* **44**: 450–454.
35. Ge, L., J. S. Gordon, C. Hsuan, K. Stenn, and S. M. Prouty. 2003. Identification of the delta-6 desaturase of human sebaceous glands: expression and enzyme activity. *J. Invest. Dermatol.* **120**: 707–714.
36. Garces, R., and M. Mancha. 1993. One-step lipid extraction and fatty acid methyl esters preparation from fresh plant tissues. *Anal. Biochem.* **211**: 139–143.
37. Van Pelt, C. K., and J. T. Brenna. 1999. Acetonitrile chemical ionization tandem mass spectrometry to locate double bonds in polyunsaturated fatty acid methyl esters. *Anal. Chem.* **71**: 1981–1989.
38. Lawrence, P., and J. T. Brenna. 2006. Acetonitrile covalent adduct chemical ionization mass spectrometry for double bond localization in non-methylene-interrupted polyene fatty acid methyl esters. *Anal. Chem.* **78**: 1312–1317.
39. Michaud, A. L., G. Y. Diau, R. Abril, and J. T. Brenna. 2002. Double bond localization in minor homoallylic fatty acid methyl esters using acetonitrile chemical ionization tandem mass spectrometry. *Anal. Biochem.* **307**: 348–360.
40. Diau, G. Y., A. T. Hsieh, E. A. Sarkadi-Nagy, V. Wijendran, P. W. Nathanielsz, and J. T. Brenna. 2005. The influence of long chain polyunsaturate supplementation on docosahexaenoic acid and arachidonic acid in baboon neonate central nervous system. *BMC Med.* **3**: 11.
41. Hirokawa, T., S. Boon-Chieng, and S. Mitaku. 1998. SOSUI: classification and secondary structure prediction system for membrane proteins. *Bioinformatics.* **14**: 378–379.
42. Thomasson, H. J. 1953. Biological standardization of essential fatty acids; a new method. *Int. Z. Vitaminforsch.* **25 (Beih)**: 62–82.
43. Stubhaug, L., D. R. Tocher, J. G. Bell, J. R. Dick, and B. E. Torstensen. 2005. Fatty acid metabolism in Atlantic salmon (*Salmo salar*L.) hepatocytes and influence of dietary vegetable oil. *Biochim. Biophys. Acta.* **1734**: 277–288.
44. Rzehak, P., J. Heinrich, N. Klopp, L. Schaeffer, S. Hoff, G. Wolfram, T. Illig, and J. Linseisen. 2009. Evidence for an association between genetic variants of the fatty acid desaturase 1 fatty acid desaturase 2 (FADS1 FADS2) gene cluster and the fatty acid composition of erythrocyte membranes. *Br. J. Nutr.* **101**: 20–26.
45. Kusunoki, M., K. Tsutsumi, M. Nakayama, T. Kurokawa, T. Nakamura, H. Ogawa, Y. Fukuzawa, M. Morishita, T. Koide, and T. Miyata. 2007. Relationship between serum concentrations of saturated fatty acids and unsaturated fatty acids and the homeostasis model insulin resistance index in Japanese patients with type 2 diabetes mellitus. *J. Med. Invest.* **54**: 243–247.
46. Thomas, E. A., and J. K. Yao. 2007. Clozapine specifically alters the arachidonic acid pathway in mice lacking apolipoprotein D. *Schizophr. Res.* **89**: 147–153.
47. Sprecher, H., and C. J. Lee. 1975. The absence of an 8-desaturases in rat liver: a reevaluation of optional pathways for the metabolism of linoleic and linolenic acids. *Biochim. Biophys. Acta.* **388**: 113–125.
48. Takagi, T. 1965. The dehydrogenation of all-cis-5,11,14-eicosatrienoic acid to arachidonic acid. *Bull. Chem. Soc. Jpn.* **38**: 2055–2057.
49. Ullman, D., and H. Sprecher. 1971. An in vitro and in vivo study of the conversion of eicosa-11,14-dienoic acid to eicosa-5,11,14-trienoic acid and of the conversion of eicosa-11-enoic acid to eicosa-5,11-dienoic acid in the rat. *Biochim. Biophys. Acta.* **248**: 186–197.
50. Naval, J., M. J. Martinez-Lorenzo, I. Marzo, P. Desportes, and A. Pineiro. 1993. Alternative route for the biosynthesis of polyunsaturated fatty acids in K562 cells. *Biochem. J.* **291**: 841–845.
51. Qi, B., T. Fraser, S. Mugford, G. Dobson, O. Sayanova, J. Butler, J. A. Napier, A. K. Stobart, and C. M. Lazarus. 2004. Production of very long chain polyunsaturated omega-3 and omega-6 fatty acids in plants. *Nat. Biotechnol.* **22**: 739–745.
52. Christiansen, K., Y. Marcel, M. V. Gan, H. Mohrhauer, and R. T. Holman. 1968. Chain elongation of alpha- and gamma-linolenic acids and the effect of other fatty acids on their conversion in vitro. *J. Biol. Chem.* **243**: 2969–2974.
53. Brenner, R. R., and R. O. Peluffo. 1966. Effect of saturated and unsaturated fatty acids on the desaturation in vitro of palmitic, stearic, oleic, linoleic, and linolenic acids. *J. Biol. Chem.* **241**: 5213–5219.
54. Cook, H. W., D. M. Byers, F. B. Palmer, M. W. Spence, H. Rakoff, S. M. Duval, and E. A. Emken. 1991. Alternate pathways in the desaturation and chain elongation of linolenic acid, 18:3(n-3), in cultured glioma cells. *J. Lipid Res.* **32**: 1265–1273.
55. Holman, R. T. 1998. The slow discovery of the importance of omega 3 essential fatty acids in human health. *J. Nutr.* **128**: 427S–433S.