

Acetylcholine-Stimulated [^3H]GABA Release from Mouse Brain Synaptosomes is Modulated by $\alpha 4\beta 2$ and $\alpha 4\alpha 5\beta 2$ Nicotinic Receptor Subtypes

Tristan D. McClure-Begley, Nathan M. King, Allan C. Collins, Jerry A. Stitzel, Jeanne M. Wehner, and Christopher M. Butt¹

Institute for Behavioral Genetics (T.D.M.B., N.M.K., A.C.C., J.A.S., J.M.W., C.M.B.) and Department of Integrative Physiology (T.D.M.B., J.A.S.) University of Colorado, Boulder, Boulder, Colorado

Received October 29, 2008; accepted January 12, 2009

ABSTRACT

Nicotinic acetylcholine receptor (nAChR) agonists stimulate the release of GABA from GABAergic nerve terminals, but the nAChR subtypes that mediate this effect have not been elucidated. The studies reported here used synaptosomes derived from the cortex, hippocampus, striatum, and thalamus of wild-type and $\alpha 4$ -, $\alpha 5$ -, $\alpha 7$ -, $\beta 2$ -, and $\beta 4$ -null mutant mice to identify nAChR subtypes involved in acetylcholine (ACh)-evoked GABA release. Null mutation of genes encoding the $\alpha 4$ or $\beta 2$ subunits resulted in complete loss of ACh-stimulated [^3H]GABA release in all four brain regions. In contrast, $\alpha 5$ gene deletion exerted a small but significant decrease in maximal ACh-evoked [^3H]GABA release in hippocampus and striatum, with a more profound effect in cortex. Acetylcholine-stimulated [^3H]GABA release from thalamic synaptosomes was not significantly affected by $\alpha 5$ gene

deletion. No effect was detected in the four brain regions examined in $\alpha 7$ - or $\beta 4$ -null mutant mice. Further analysis of ACh-evoked [^3H]GABA release revealed biphasic concentration-response relationships in the four brain regions examined from all wild-type animals and in $\alpha 5$ null mutant mice. Moreover, a selective reduction in the maximum response of the high-affinity component was apparent in $\alpha 5$ -null mutant mice. The results demonstrate that $\alpha 4\beta 2$ -type nAChRs are critical for ACh-stimulated [^3H]GABA release from all four brain regions examined. In addition, the results suggest that $\alpha 5$ -containing receptors on GABAergic nerve terminals comprise a fraction of the high ACh-sensitivity component of the concentration-response curve and contribute directly to the ability of nicotinic agonists to evoke GABA release in these regions.

The nicotinic acetylcholine receptor (nAChR) arguably represents one of the most evolutionarily conserved and well characterized neurotransmitter receptors, and the majority of high-resolution structural data have been accumulated from studies of the *torpedo*- and muscle-type nAChR (Millar and Gotti, 2009). It is generally believed that the structures of neuronal nAChRs are very similar to the muscle-type nAChR subunits because the amino acid sequences of neuronal nAChR subunits are similar to the muscle-type subunits (Lindstrom et al., 1998). Assuming that the neuronal nAChRs are pentameric assemblies that resemble the peripheral-type

nAChR, the fact that mammalian neurons express mRNAs for nine nAChR genes (designated $\alpha 2$ – $\alpha 7$ and $\beta 2$ – $\beta 4$) suggests that many different nAChR subtypes might be expressed in the central nervous system.

The importance of identifying the sites of expression and subunit compositions of those nAChRs that are actually expressed in brain is demonstrated by observations that the biophysical and pharmacological properties of nAChRs are affected when different subunit combinations are examined in heterologous expression systems (Zwart and Vijverberg, 1998; Moroni et al., 2006; Kuryatov et al., 2008). Of particular importance is the observation that $\alpha 4\alpha 5\beta 2$ nAChRs are more cation-permeable than other high-affinity heteromeric nAChRs, and the calcium-permeability of these $\alpha 4\alpha 5\beta 2$ nAChRs is exceeded only by $\alpha 7$ -type nAChRs and *N*-methyl-D-aspartate glutamate receptors (Kuryatov et al., 2008). Indeed, previous studies performed in our laboratory (Brown et al., 2007) showed that deletion of the $\alpha 5$ subunit decreased maximal agonist-evoked $^{86}\text{Rb}^+$ efflux in several brain re-

This work was supported by the National Institutes of Health National Institute on Alcohol Abuse and Alcoholism [Grants AA13108, AA13465]; the National Institutes of Health National Cancer Institute [Grant CA089392]; and the National Institutes of Health National Institute on Drug Abuse [Grant DA015663].

¹ Current affiliation: Martek Biosciences, Neuroscience Research, Discovery, Boulder, Colorado.

Article, publication date, and citation information can be found at <http://molpharm.aspetjournals.org>.
doi:10.1124/mol.108.052274.

ABBREVIATIONS: nAChR, nicotinic acetylcholine receptor; NO-711, 1-[2-[[[(diphenylmethylene)imino]oxy]ethyl]-1,2,5,6-tetrahydro-3-pyridinecarboxylic acid hydrochloride; ANOVA, analysis of variance; ACh, acetylcholine; DH β E, dihydro- β -erythroidine.

gions without any loss of total receptor binding. These results suggest that the $\alpha 5$ subunit exerts a distinct and neuron phenotype-specific effect on sequelae of nAChR activation, such as neurotransmitter release.

The development of genetically engineered (gene knockout or null mutant) mice has provided a vital tool that has facilitated the identification and characterization of native neuronal nAChRs. For example, the findings that $\alpha 4$ and $\beta 2$ mRNAs are found together in many brain regions (Marks et al., 1992, Whiteaker et al., 2006) that also express high-affinity [^3H]nicotine binding sites (Marks and Collins, 1982; Clarke et al., 1985) led to the suggestion that those nAChRs that bind nicotinic agonists with high affinity are composed of $\alpha 4$ and $\beta 2$ subunits. Definitive proof that the $\alpha 4$ and $\beta 2$ subunits are required to form these binding sites was provided by studies that showed the absence of high-affinity [^3H]nicotine binding in brain tissue derived from $\alpha 4$ - (Marubio et al., 1999) and $\beta 2$ - (Picciotto et al., 1995) null mutant mice. Likewise, the findings that ^{125}I - α -bungarotoxin binding is eliminated in $\alpha 7$ -null mutant mice helped establish that $\alpha 7$ -type receptors bind ^{125}I - α -bungarotoxin with high affinity (Orr-Urtreger et al., 1997). More recent studies using null mutant mice have demonstrated that the $\alpha 6$ (Champ-tieux et al., 2001), $\beta 2$ (Salminen et al., 2005), and $\beta 3$ (Cui et al., 2003) subunits are required to form the ^{125}I - α -conotoxin MII binding sites expressed in catecholaminergic neurons. In addition, studies from our laboratory (Marks et al., 2007) have used null mutants to identify the subunit compositions of seven different nAChRs that can be measured with the aid of high- and low-affinity [^3H]epibatidine binding.

Many nAChRs are expressed on presynaptic nerve terminals, where they modulate the release of acetylcholine (ACh), GABA, glutamate, serotonin, and dopamine (Wonnacott, 1997). We (Salminen et al., 2004) used $\alpha 4$ -, $\alpha 5$ -, $\alpha 7$ -, $\beta 2$ -, $\beta 3$ -, and $\beta 4$ -null mutant mice, along with an assay that measures [^3H]dopamine release from striatal synaptosomes, to identify the subunit compositions of the native receptors that are expressed in dopaminergic nerve terminals. Our results indicate that four different nAChR subtypes ($\alpha 4\beta 2$, $\alpha 4\alpha 5\beta 2$, $\alpha 4\alpha 6\beta 2\beta 3$, and $\alpha 6\beta 2\beta 3$) are expressed in dopaminergic nerve terminals derived from mouse striatum. An identical conclusion was drawn by Gotti et al. (2005) using immunological methodologies.

The studies reported here used synaptosomes obtained from wild-type and $\alpha 4$ -, $\alpha 5$ -, $\alpha 7$ -, $\beta 2$ -, and $\beta 4$ -null mutant mice in an attempt to identify the role of these subunits in modulating GABA release. ACh-stimulated [^3H]GABA release was studied using synaptosomes prepared from the hippocampus, striatum, cortex, and thalamus. Immunological (Gahring et al., 2004) and reverse-transcription polymerase chain reaction (Klink et al., 2001) data have demonstrated that these subunits may be found in GABAergic neurons in several mouse brain regions. The results indicate that ACh-evoked [^3H]GABA release in all four brain regions requires $\alpha 4$ and $\beta 2$ subunits and that the $\alpha 5$ subunit contributes to the formation of functional receptors in three of the four brain regions examined, but to differing degrees.

Materials and Methods

Materials. [^3H]GABA (33.4 Ci/mmol) was purchased from Perkin-Elmer Life and Analytical Sciences (Waltham, MA). Sucrose and

HEPES were obtained from Roche Diagnostics (Indianapolis, IN). The following compounds were products of Sigma Chemical Co. (St. Louis, MO): ACh, atropine sulfate, NO-711, aminooxyacetic acid, GABA, sodium chloride, potassium chloride, calcium chloride, magnesium sulfate, potassium dihydrogen phosphate, D-glucose, and diisopropylfluorophosphate. Econosafe scintillation cocktail was purchased from Research Products International Corp. (Mt. Prospect, IL), and OptiPhase Supermix scintillation cocktail was obtained from PerkinElmer.

Mice. Male and female mice carrying gene deletions (knockouts) for *Chrna4*, *Chrna5*, *Chrna7*, *Chrn2*, and *Chrn4*, which encode $\alpha 4$, $\alpha 5$, $\alpha 7$, $\beta 2$, and $\beta 4$ nAChR subunits, respectively, were used in these studies. All nAChR subunit-null mutant mice were bred onto the C57BL/6 strain for the indicated number of generations: $\alpha 4$ (Ross et al., 2000), 5 generations; $\alpha 5$ (Salas et al., 2003), 8 generations; $\alpha 7$ (Orr-Urtreger et al., 1997), 10 generations; $\beta 2$ (Picciotto et al., 1995), 10 generations; and $\beta 4$ (Xu et al., 1999), 10 generations. All of the animals were produced by heterozygous (+/-) matings maintained at the Institute for Behavioral Genetics (University of Colorado, Boulder, CO).

The mice were weaned and separated by gender at 25 days of age and were housed in groups of five to a cage. The animals were maintained on a 12-h light/dark cycle (lights on from 7:00 AM to 7:00 PM), were given unlimited access to food (Teklad Rodent Diet) and water and were used when they were 60 to 120 days old. All animal care and experimental procedures were performed in accordance with the National Institutes of Health's *Guide for Care and Use of Laboratory Animals* and were approved by the University of Colorado's Animal Care and Use Committee.

Genotyping. All genotyping was performed from tail clip samples (~1 cm). QIAGEN (Valencia, CA) DNEasy Tissue Kits were used to extract DNA from the tail clippings. The $\alpha 4$, $\alpha 5$, $\alpha 7$, $\beta 2$, and $\beta 4$ genotypes were determined by polymerase chain reaction with oligonucleotide probes specific for the *Chrna4*, *Chrna5*, *Chrna7*, *Chrn2*, and *Chrn4* sequences, respectively, as described in Salminen et al. (2004). The gene products were electrophoresed on 1.5% agarose gels and stained with ethidium bromide. Two independent observers then scored the genotypes.

Synaptosome Preparation. Each mouse was sacrificed by cervical dislocation, and the brain was removed from the skull. The cortex, hippocampus, striatum, and thalamus were then dissected on ice. Crude synaptosomes were prepared from each dissected brain region by hand homogenization in 4 volumes of 0.5 ml of ice-cold




Fig. 1. Representative data of ACh-stimulated [^3H]GABA release from hippocampal synaptosomes. A total of 23 10-s fractions were taken from each synaptosomal sample, and the radioactivity in each fraction was measured in a scintillation spectrometer. Thus, these data are presented as cpm. The synaptosomes were stimulated with a range of ACh concentrations (0.03–1000 μM) for 12 s in the middle of each run (hash marks beneath each trace). The fractions before and after the stimulation were then used to determine the baseline [^3H]GABA release, which was subsequently used to determine the size of the response (see *Materials and Methods* for details). The size of the responses increased in a concentration-dependent manner.

(4°C) Percoll medium I (320 mM sucrose and 5 mM HEPES, pH 7.5) with a glass Teflon homogenizer. The homogenates were then centrifuged at 12,000*g* for 20 min at 4°C. The pellets of each homogenate were resuspended in uptake buffer (128 mM NaCl, 2.4 mM KCl, 3.2 CaCl₂, 1.2 mM KH₂PO₄, 1.2 mM MgSO₄ · 7H₂O, 25 mM HEPES, and 10 mM glucose, pH 7.5). The volume of perfusion buffer used for resuspending the synaptosomes was 0.8 ml for each brain region.

[³H]GABA Release. These experiments used a protocol adapted from that developed by Lu et al. (1998) with modifications as specified below. Synaptosomal preparations were incubated for 10 min at 37°C in uptake buffer containing 1.25 mM aminooxyacetic acid, a GABA transaminase inhibitor. [³H]GABA (final concentration, 0.3 μM), unlabeled GABA (final concentration, 0.25 μM), and diisopropylfluorophosphate (final concentration, 10 μM), an irreversible cholinesterase inhibitor, were then added to the suspension, and the suspension was incubated for another 10 min. Aliquots (80 μl) were collected with gentle suction onto 6-mm diameter A/E glass-fiber filters (Gelman Science, Ann Arbor, MI) on the perfusion apparatus and then perfused with buffer containing 1 g/l bovine serum albumin (1.8 ml/min) for 10 min before fraction collection was started. ACh (0.03–1000 μM) was added to the perfusate for 12 s in the middle of each fraction collection run. The amount of radioactivity in each

fraction was then determined with scintillation spectrometry (45% counting efficiency). Atropine (1 μM) was included in the perfusion buffer to block any possible muscarinic responses. NO-711 (100 nM), a potent (*K_d* ~6 nM) and selective antagonist of the neuronal GABA transporter (Borden, 1996), was also added to the perfusion buffer to prevent calcium-independent [³H]GABA release due to reversal of the GABA transporter by large concentrations of ACh (data not shown).

Data Analysis. The amount of ACh-stimulated [³H]GABA release was determined in 12-s fractions. The data were normalized using Sigma Plot 5.0 (Systat Software, Inc., San Jose, CA) as described by Grady et al. (1992). In short, the fractions recorded in the absence of agonist are fit to a single-phase exponential decay equation to obtain a measure of baseline release. The stimulated release was divided by the calculated baseline release during the time of stimulation to calculate “response units,” where one unit is defined as twice basal release. The responses at each time interval of stimulation were then summed to yield the total response for a given agonist exposure. For comparisons between genotypes, the mean value of a near-maximal concentration of ACh (30 μM) for each brain region was used as a further normalizing factor, and all were responses plotted as a percentage of this control value. Concentration-effect curves for [³H]-




Fig. 2. Effect of $\alpha 4$ nAChR subunit gene deletion on ACh-evoked [³H]GABA release. ChRNA4 gene deletion resulted in a total loss of ACh-stimulated [³H]GABA release from synaptosomes made from the cortex (A), hippocampus (B), striatum (C), and thalamus (D). Animals lacking the $\alpha 4$ nAChR subunit did not exhibit a concentration-dependent response to ACh in any of the four brain regions; *n* = 2 to 8 for each data point in each graph. Error bars represent S.E.M.

GABA release were plotted and fit with the four-parameter hyperbolic equation $f = VS/(K + S) + vS/(k + S)$, where S is agonist concentration, maximum response for high and low-sensitivity components are V and v , respectively, and half-maximal agonists concentrations for high- and low-sensitivity components are represented by K and k , respectively. Curve-fits were performed using SigmaPlot (SigmaPlot DOS or SigmaPlot 2001), and an F test was conducted on each data set to ensure that the four-parameter equation was statistically preferred. To obtain concentration-response parameters, initial curve-fits were calculated from mean concentration-response curves, and the theoretical values were applied to each individual experiment to generate a series of individual experiment curve-fits. Significant differences ($p < 0.05$) between two-group data sets were determined with univariate analysis of variance (SPSS 16.2). Both raw data values and calculated curve-fit parameters were analyzed with ANOVA to assess genotype differences that may be evident across a concentration-response curve but less apparent when calculated from theoretical curve-fit parameters. When EC_{50} values were compared, their log values were used. This was done because $\log(EC_{50})$ values are normally distributed, but EC_{50} values are not (Hancock et al., 1988).

Results

Concentration-Dependent Nature of ACh-Stimulated [3H]GABA Release. Figure 1 presents a typical data trace. The data are presented as cpm of radioactivity obtained when hippocampal synaptosomes loaded with [3H]GABA were perfused with increasing concentrations of ACh. As higher ACh concentrations were used, more [3H]GABA release (units above baseline) was observed. Similar results were obtained with synaptosomes obtained from the other brain regions. These data are consistent with those reported by Lu et al. (1998).

Effects of $\alpha4$, $\alpha5$, and $\beta2$ Gene Deletion on ACh-Stimulated [3H]GABA Release. Figure 2 illustrates that null mutation of *Chrna4*, the gene encoding the $\alpha4$ subunit, resulted in a complete loss of ACh-stimulated GABA release from synaptosomes prepared from all four of the brain regions. Acetylcholine produced a concentration-dependent increase in GABA release in all four of the brain regions, but the maximal responses of the brain regions differed. High concentrations (0.3 and 1 mM) often produced a marked decrease in release from that seen at the determined maximally efficacious concentration (i.e., the concentration effects were of the inverted "U" type). These effects were most likely caused by agonist "channel block," in which the agonist accumulates in the ion channel and prevents cation flux, even though the receptor is in its active conformation (Zwart and Vijverberg, 1998). In all cases, a minimum of three to four log units of ACh was required to progress from minimal to maximal release. This suggests that more than one receptor subtype may be mediating this response. Therefore, the data were analyzed to determine whether they fit one- or two-site models. The ACh concentration-effect curves in wild-type animals were best fit by a two-site model in all brain regions examined (F test, $p < 0.05$). The calculated curve-fit parameters [R_{max} (maximal release) and EC_{50} (concentration that elicited half-maximal release)] for the cortex, hippocampus, striatum, cortex, and thalamus are presented in Table 1.

Figure 3 shows that null mutation of *Chrn2*, the gene encoding the $\beta2$ subunit, also resulted in a total elimination of ACh-stimulated [3H]GABA release. The effects of $\beta2$ knockout were statistically significant in all four brain regions. The ACh concentration-effect curves for all four brain regions in wild-

TABLE 1

Effects of gene deletion on high- and low-sensitivity components of ACh-evoked [3H]GABA release

R_{max} and EC_{50} values were calculated from concentration-response curves as described under *Materials and Methods*. Each value represents the mean \pm S.E.M. as compiled from four to eight individual concentration-effect curves for each genotype. The two components of the concentration-response curve are designated as high-ACh sensitivity (HS) and low-ACh sensitivity (LS).

Genotype	R_{max} (HS)	log EC_{50} (HS)	R_{max} (LS)	log EC_{50} (LS)
		μM		μM
Cortex				
$\alpha4(+/+)$	100.31 \pm 11.80	-0.17 \pm 0.098	39.78 \pm 25.21	1.40 \pm 0.480
$\alpha4(-/-)$				
$\beta2(+/+)$	80.58 \pm 29.56	-0.44 \pm 0.381	15.06 \pm 15.05	1.35 \pm 0.66
$\beta2(-/-)$				
$\alpha5(+/+)$	84.45 \pm 13.16	-0.32 \pm 0.171	50.10 \pm 10.36	0.93 \pm 0.45
$\alpha5(-/-)$	40.15 \pm 6.48	-0.66 \pm 0.14	56.37 \pm 17.85	1.31 \pm 0.52
Hippocampus				
$\alpha4(+/+)$	81.28 \pm 16.13	-0.34 \pm 0.199	53.03 \pm 16.72	2.14 \pm 0.28
$\alpha4(-/-)$				
$\beta2(+/+)$	77.98 \pm 6.46	-0.44 \pm 0.13	105.28 \pm 39.65	2.20 \pm 0.23
$\beta2(-/-)$				
$\alpha5(+/+)$	69.40 \pm 10.92	-3.42 \pm 2.70	61.24 \pm 7.11	1.27 \pm 0.53
$\alpha5(-/-)$	51.30 \pm 14.31	-1.98 \pm 1.43	51.23 \pm 9.84	1.20 \pm 0.20
Striatum				
$\alpha4(+/+)$	42.83 \pm 4.31	-1.12 \pm 0.15	64.98 \pm 24.01	1.00 \pm 0.55
$\alpha4(-/-)$				
$\beta2(+/+)$	48.10 \pm 8.03	-1.67 \pm 0.62	60.35 \pm 9.02	0.60 \pm 0.21
$\beta2(-/-)$				
$\alpha5(+/+)$	51.79 \pm 6.96	-0.73 \pm 0.18	63.17 \pm 12.12	0.998 \pm 0.12
$\alpha5(-/-)$	51.18 \pm 9.37	-0.46 \pm 0.18	41.61 \pm 13.68	0.82 \pm 0.47
Thalamus				
$\alpha4(+/+)$	54.30 \pm 10.31	-2.59 \pm 2.10	62.98 \pm 25.91	1.16 \pm 0.62
$\alpha4(-/-)$				
$\beta2(+/+)$	40.58 \pm 8.24	-2.37 \pm 1.29	75.72 \pm 17.67	0.94 \pm 0.44
$\beta2(-/-)$				
$\alpha5(+/+)$	41.54 \pm 5.23	-0.33 \pm 0.20	82.71 \pm 22.09	1.46 \pm 0.21
$\alpha5(-/-)$	47.17 \pm 12.00	-2.75 \pm 2.13	47.10 \pm 17.77	1.00 \pm 0.40

type animals were best fit by a two-site model (F test, $p < 0.05$). The calculated curve-fit parameters for the hippocampus, striatum, cortex, and thalamus are presented in Table 1.

Figure 4 demonstrates that deletion of *Chrna5*, the gene encoding the $\alpha 5$ subunit, caused robust decreases in ACh-stimulated [3 H]GABA release in the cortex ($F_{1,118} = 24.95$; $p < 0.001$). Deletion of the $\alpha 5$ subunit also produced a modest but significant decrease in [3 H]GABA release in the striatum and hippocampus ($F_{1,152} = 7.67$, $p < 0.007$; $F_{1,132} = 17.12$, $p < 0.001$, respectively). No observable effect of $\alpha 5$ gene deletion was observed in thalamic synaptosomes ($F_{1,90} = 0.76$, $p = 0.390$). The ACh concentration-effect curves for all four brain regions in both wild-type and knockout animals were best fit by a two-site model (F test, $p < 0.05$). The calculated curve-fit parameters for the cortex, thalamus, striatum, and hippocampus are presented in Table 1. As noted in the table, $\alpha 5$ gene deletion resulted in no change in EC_{50} values in the cortex, hippocampus, or striatum. The R_{max} for the high-affinity com-

ponent was significantly reduced in cortex only. The calculated R_{max} values for hippocampus and striatum were not significantly different from control, but the ANOVAs performed on the entire concentration-response curves did detect a significant effect of genotype, with a trend toward a decrease in the high-affinity component. Deletion of the $\alpha 5$ subunit did not produce any significant changes in the maximal response or EC_{50} values for the low-affinity component.

High- and low-sensitivity $\alpha 4\beta 2^*$ -type nAChRs differ in their sensitivity to antagonism by the competitive inhibitor dihydro- β -erythroidine (DH β E) (Marks et al., 1999). At low concentrations ($\leq 2 \mu M$), only the high-sensitivity fraction is blocked, allowing for a pharmacological isolation of the receptor responses observed. In synaptosomes from $\alpha 5$ -null mutant mice, DH β E ($2 \mu M$) blocked the high-sensitivity component of ACh-evoked [3 H]GABA; removal of the $\alpha 5$ gene did not alter the low-sensitivity $\alpha 4\beta 2$ responses to ACh (Fig. 5).

Deletion of either the $\beta 4$ or $\alpha 7$ subunit did not result in any




Fig. 3. Effect of $\beta 2$ nAChR subunit gene deletion on ACh-evoked [3 H]GABA release. ACh-stimulated [3 H]GABA release was not detected in synaptosomes made from the cortex (A), hippocampus (B), striatum (C), or thalamus (D) of *Chrnb2*-null mutant mice. As with $\alpha 4$ -null mutants, animals lacking the $\beta 2$ nAChR subunit did not exhibit a concentration-dependent GABA release response to ACh in any of the four brain regions studied; $n = 1$ to 3 for each data point in each graph. Error bars represent the S.E.M.

measurable changes in [^3H]GABA release evoked by ACh (30 μM) in any of the four brain regions screened (Fig. 6, A and B, respectively). Thus, no further experimentation proceeded with these animals.

Discussion

Null mutation of either *Chrna4* or *Chrn2* totally eliminated ACh-stimulated [^3H]GABA release from every brain region studied. These null mutations also eliminate [^3H]nicotine binding (Picciotto et al., 1995; Marubio et al., 1999) and cytosine-sensitive [^3H]epibatidine binding (Marks et al., 2007) throughout the brain. We interpreted the binding results to indicate that $\alpha 4\beta 2^*$ -type receptors are the major nAChR subtype that binds these ligands with high affinity. By analogy, we conclude that $\alpha 4\beta 2^*$ -type nAChRs modulate GABA release from the presynaptic nerve terminals in the four brain regions that were studied.

The International Union of Basic and Clinical Pharmacol-

ogy receptor nomenclature committee (Lukas et al., 1999) recommended that nAChRs should be named by including the Greek letters for known subunits along with an asterisk (*) to designate the potential contribution of other subunits. We found that *Chrna5* gene deletion resulted in decreased maximal ACh-evoked [^3H]GABA release from cortical, striatal, and hippocampal synaptosomes with differing degrees of effect. This outcome suggests that $\alpha 4\alpha 5\beta 2$ -type nAChRs play regionally distinct modulatory roles in ACh-evoked GABA release. *Chrna7* and *Chrn4* gene deletion did not affect ACh-evoked [^3H]GABA release. These observations indicate that receptors which include these subunits are not significant contributors to [^3H]GABA release from presynaptic nerve terminals in the four brain regions that were studied.

The finding that both $\alpha 4$ and $\beta 2$ gene deletion eliminated [^3H]GABA release from synaptosomes prepared from all four brain regions examined is not surprising given that the mRNAs for both of these subunits are expressed in high concen-




Fig. 4. Effect of $\alpha 5$ nAChR subunit gene deletion on ACh-evoked [^3H]GABA. *Chrna5* gene deletion generally resulted in a decrease of ACh-stimulated [^3H]GABA release from synaptosomes made from the cortex (A), hippocampus (B), striatum (C), and thalamus (D). The decreased response in the $\alpha 5$ -null mutants was readily observable in the cortex, diminished in the striatum and hippocampus, and was not significant in the thalamus; $n = 2$ to 11 for each data point in each graph. Error bars represent the S.E.M.

trations in these brain regions (Marks et al., 1992). Likewise, it is not surprising that *Chrna5* deletion resulted in a change in cortical [^3H]GABA release given that single-cell reverse-transcription polymerase chain reaction analyses detected $\alpha 5$ mRNA expression in some but not all cortical GABAergic neurons (Porter et al., 1999). It should be noted, however, that in situ hybridization measures mRNA that is expressed primarily in cell bodies.

Chrna5 gene deletion had a small effect on ACh-stimulated [^3H]GABA release from striatal and hippocampal synaptosomes. The in situ hybridization experiments indicate that $\alpha 4$, $\alpha 5$, and $\beta 2$ mRNAs are all expressed in both of these brain regions. However, the expression pattern for $\alpha 5$ mRNA is limited compared with those observed for $\alpha 4$ and $\beta 2$ mRNA. These findings may explain why we obtained evidence that suggests both $\alpha 4\beta 2$ and $\alpha 4\alpha 5\beta 2$ nAChRs are expressed on GABAergic nerve terminals in three of the four brain regions examined. This correlation between *Chrna5* mRNA expression and $\alpha 5$ -null mutation effect on ACh-evoked [^3H]GABA release could also explain the lack of $\alpha 5$

subunit removal on [^3H]GABA release measured from thalamic synaptosomes, because there is no appreciable $\alpha 5$ mRNA expressed in that region. Therefore, regional effects of *Chrna5* gene deletion could reflect preferential loss of activity in local GABAergic interneuron populations.

Many studies have used electrophysiological methods to study nicotinic modulation of GABA neuron activity, and many of these studies have focused on the hippocampus (Alkondon and Albuquerque, 1993; Alkondon et al., 1997; Kawai et al., 2002), but nicotinic modulation of GABAergic neurons has also been studied in the cortex (Alkondon et al., 2000), thalamus (Léna and Changeux, 1997), and dopamine-rich regions such as the midbrain (Klink et al., 2001), the ventral tegmental area (Mansvelder and McGehee, 2000), and the nucleus accumbens (de Rover et al., 2002). Many of these studies found that pretreatment with the nAChR antagonist DH β E blocked cholinergic activation of GABAergic neurons, leading to the conclusion that $\alpha 4\beta 2$ -type nAChRs influence the activity of GABAergic neurons. Electrophysiological analyses of the potential roles of nAChRs containing




Fig. 5. Effect of $\alpha 5$ nAChR subunit gene deletion on DH β E-resistant nAChR function measured with ACh-evoked [^3H]GABA release. ACh-stimulated [^3H]GABA release in the presence of 2 μM DH β E from cortex (A), hippocampus (B), striatum (C), and thalamus (D) shows no effect of *Chrna5* gene deletion ($p > 0.05$). Error bars represent S.E.M..

the $\alpha 5$ subunit in modulating GABAergic function have not been published. Our results suggest that such studies will yield different results depending on the brain region studied.

Many of the electrophysiological studies also demonstrated a role for $\alpha 7$ -containing nAChRs in the modulation of somatodendritic activities of GABAergic neurons. The lack of an effect of $\alpha 7$ subunit gene deletion on [^3H]GABA release from synaptosomal preparations suggests that $\alpha 7$ -containing nAChRs present on GABAergic neurons are expressed preferentially in somatodendritic locations, whereas the dominant presynaptic nAChR expressed by GABAergic neurons is the $\alpha 4\beta 2^*$ type.

It is exceedingly likely that our synaptosomal preparation contains a heterogeneous population of nerve terminals, with contributions to the overall pool being provided by many potentially distinct subregions. We found that null mutation of *Chrna5* significantly decreases the calculated R_{max} for ACh in cortex. The R_{max} for ACh-stimulated GABA release was not significantly altered in striatum and hippocampus, but the ANOVA, which is more sensitive because it analyzes effects over the entire concentration-effect curve, did detect a significant effect of gene deletion. Thus, deleting the $\alpha 5$ subunit produces a more subtle effect in striatum and hippocampus. No differences in agonist potency were observed in any brain region studied. In agreement with a previously published functional assessment of *Chrna5* gene deletion using $^{86}\text{Rb}^+$ efflux, it seems that the loss of the $\alpha 5$ subunit results in a preferential decrease in the maximal response of high ACh-sensitivity nAChRs with no subsequent loss of total receptor number (Brown et al., 2007). In addition, isolation of the low-sensitivity $\alpha 4\beta 2$ functional contribution by examining ACh-evoked [^3H]GABA release in the presence of $2\ \mu\text{M}$ DH β E (Marks et al., 1999) showed no differences between $\alpha 5$ wild-type and null-mutant animals in any brain region examined. These results are consistent with studies comparing the functional properties of $\alpha 4\beta 2$ and $\alpha 4\alpha 5\beta 2$ nAChRs in heterologous expression systems (Kuryatov et al., 2008) and suggest that the $\alpha 4\alpha 5\beta 2$ -type nAChR expressed on GABAergic nerve terminals is uniquely capable of influencing neurotransmitter release. This enhanced functionality may be due to distinct biophysical properties, such as the enhanced ion

(including Ca^{2+}) conductance of $\alpha 4\alpha 5\beta 2$ nAChRs compared with high-affinity nAChRs containing $\alpha 4$ and $\beta 2$ subunits only.

The biphasic responses for $\alpha 4\beta 2^*$ nAChRs have been described previously in $^{86}\text{Rb}^+$ efflux studies (Marks et al., 1999; Brown et al., 2007) and in electrophysiological measurements of heterologous expression systems (Zwart and Vijverberg, 1998; Moroni et al., 2006). It has been suggested that the high- and low-affinity components represent receptors with different stoichiometries: $(\alpha 4)_2(\beta 2)_3$ (high affinity), and $(\alpha 4)_3(\beta 2)_2$ (low affinity) (Zwart and Vijverberg, 1998; Marks et al., 2007). It is possible that the loss of the $\alpha 5$ subunit results in the compensatory formation of receptors with the stoichiometry $(\alpha 4)_2(\beta 2)_3$ that have an equally high affinity for ACh but are distinctly less efficacious at eliciting neurotransmitter release than the $\alpha 4\alpha 5\beta 2$ receptor. This event could explain the selective reduction in the high-sensitivity component of ACh-evoked [^3H]GABA release without any significant decrease in low-sensitivity $\alpha 4\beta 2$ nAChR function or loss of receptor number as a whole, which was measured by receptor binding assays in $\alpha 5$ -null mutant animals (Brown et al., 2007).

Conclusions

The $\alpha 4$ and $\beta 2$ nAChR subunits are clearly necessary for nicotinic agonist-stimulated GABA release in all four of the brain regions examined. *Chrna5* gene deletion does not result in a total loss of ACh-stimulated GABA release in any brain region. However, it is frequently included along with $\alpha 4$ and $\beta 2$ subunits, and the absence of the $\alpha 5$ subunit decreases overall nAChR-mediated [^3H]GABA release that is restricted to an effect on the high-affinity component of the functional profile. These findings suggest that $\alpha 4\alpha 5\beta 2$ nAChRs are important modulators of [^3H]GABA release from striatal, hippocampal, and cortical synaptosomes. Furthermore, $\alpha 5$ -containing nAChRs are distinct in their regional distribution and functionality. Thus, nAChRs containing the $\alpha 5$ subunit may serve as therapeutic targets with some selectivity for distinct neuronal pathways.




Fig. 6. Effects of $\beta 4$ and $\alpha 7$ subunit null mutation on $30\ \mu\text{M}$ ACh-evoked [^3H]GABA release from cortex, hippocampus, striatum, and thalamus. No statistically significant effect of gene deletion was observed.

References

- Alkondon M and Albuquerque EX (1993) Diversity of nicotinic acetylcholine receptors in rat hippocampal neurons. I. Pharmacological and functional evidence for distinct structural subtypes. *J Pharmacol Exp Ther* **265**:1455–1473.
- Alkondon M, Pereira EF, Barbosa CT, and Albuquerque EX (1997) Neuronal nicotinic acetylcholine receptor activation modulates gamma-aminobutyric acid release from CA1 neurons of rat hippocampal slices. *J Pharmacol Exp Ther* **283**:1396–1411.
- Alkondon M, Pereira EF, Eisenberg HM, and Albuquerque EX (2000) Nicotinic receptor activation in human cerebral cortical interneurons: a mechanism for inhibition and disinhibition of neuronal networks. *J Neurosci* **20**:66–75.
- Borden LA (1996) GABA transporter heterogeneity: pharmacology and cellular localization. *Neurochem Int* **29**:335–356.
- Brown RW, Collins AC, Lindstrom JM, and Whiteaker P (2007) Nicotinic alpha5 subunit deletion locally reduces high-affinity agonist activation without altering nicotinic receptor numbers. *J Neurochem* **103**:204–215.
- Champtiaux N, Gotti C, Cordero-Erausquin M, David DJ, Przybyski C, Lena C, Clementi F, Moretti M, Rossi FM, Le Novère N, et al. (2003) Subunit composition of functional nicotinic receptors in dopaminergic neurons investigated with knockout mice. *J Neurosci* **23**:7820–7829.
- Clarke PB, Schwartz RD, Paul SM, Pert CB, and Pert A (1985) Nicotinic binding in rat brain: autoradiographic comparison of [³H]acetylcholine, [³H]nicotine, and [¹²⁵I]-alpha-bungarotoxin. *J Neurosci* **5**:1307–1315.
- Cui C, Booker TK, Allen RS, Grady SR, Whiteaker P, Marks MJ, Salminen O, Tritto T, Butt CM, Allen WR, et al. (2003) The beta3 nicotinic receptor subunit: a component of alpha-conotoxin MII-binding nicotinic acetylcholine receptors that modulate dopamine release and related behaviors. *J Neurosci* **23**:11045–11053.
- de Rover M, Lodder JC, Kits KS, Schoffelmier AN, and Brussaard AB (2002) Cholinergic modulation of nucleus accumbens medium spiny neurons. *Eur J Neurosci* **16**:2279–2290.
- Gahring LC, PERSIYANOV K, Dunn D, Weiss R, Meyer EL, and Rogers SW (2004) Mouse strain-specific nicotinic acetylcholine receptor expression by inhibitory interneurons and astrocytes in the dorsal hippocampus. *J Comp Neurol* **468**:334–346.
- Gotti C, Moretti M, Clementi F, Riganti L, McIntosh JM, Collins AC, Marks MJ, and Whiteaker P (2005) Expression of nigrostriatal $\alpha 6$ -containing nicotinic acetylcholine receptors is selectively reduced, but not eliminated, by $\beta 3$ subunit gene deletion. *Mol Pharmacol* **67**:2007–2015.
- Grady S, Marks MJ, Wonnacott S, and Collins AC (1992) Characterization of nicotinic receptor-mediated [³H]dopamine release from synaptosomes prepared from mouse striatum. *J Neurochem* **59**:848–856.
- Hancock AA, Bush EN, Stanisic D, Kyncl JJ, and Lin CT (1988) Data normalization before statistical analysis: keeping the horse before the cart. *Trends Pharmacol Sci* **9**:29–32.
- Kawai H, Zago W, and Berg DK (2002) Nicotinic alpha7 receptor clusters on hippocampal GABAergic neurons: regulation by synaptic activity and neurotrophins. *J Neurosci* **22**:7903–7912.
- Klink R, de Kerchove d'Exaerde A, Zoli M, and Changeux JP (2001) Molecular and physiological diversity of nicotinic acetylcholine receptors in the midbrain dopaminergic nuclei. *J Neurosci* **21**:1452–1463.
- Kuryatov A, Onksen J, and Lindstrom J (2008) Roles of accessory subunits in $\alpha 4\beta 2$ nicotinic receptors. *Mol Pharmacol* **74**:132–143.
- Léna C and Changeux JP (1997) Role of Ca²⁺ ions in nicotinic facilitation of GABA release in mouse thalamus. *J Neurosci* **17**:576–585.
- Lindstrom J, Peng X, Kuryatov A, Lee E, Anand R, Gerzanich V, Wang F, Wells G, and Nelson M (1998) Molecular and antigenic structure of nicotinic acetylcholine receptors. *Ann NY Acad Sci* **841**:71–86.
- Lu Y, Grady S, Marks MJ, Picciotto M, Changeux JP, and Collins AC (1998) Pharmacological characterization of nicotinic receptor-stimulated GABA release from mouse brain synaptosomes. *J Pharmacol Exp Ther* **287**:648–657.
- Lukas RJ, Changeux JP, Le Novère N, Albuquerque EX, Balfour DJ, Berg DK, Bertrand D, Chiappinelli VA, Clarke PB, Collins AC, et al. (1999) International Union of Pharmacology. XX. Current status of the nomenclature for nicotinic acetylcholine receptors and their subunits. *Pharmacol Rev* **51**:397–401.
- Mansvelder HD and McGehee DS (2000) Long-term potentiation of excitatory inputs to brain reward areas by nicotine. *Neuron* **27**:349–357.
- Marks MJ and Collins AC (1982) Characterization of nicotine binding in mouse brain and comparison with the binding of α -bungarotoxin and quinuclidinyl benzilate. *Mol Pharmacol* **22**:554–564.
- Marks MJ, Meinerz NM, Drago J, and Collins AC (2007) Gene targeting demonstrates that alpha4 nicotinic acetylcholine receptor subunits contribute to expression of diverse [³H]epibatidine binding sites and components of biphasic 86Rb⁺ efflux with high and low sensitivity to stimulation by acetylcholine. *Neuropharmacology* **53**:390–405.
- Marks MJ, Pauly JR, Gross SD, Deneris ES, Hermans-Borgmeyer I, Heinemann SF, and Collins AC (1992) Nicotine binding and nicotinic receptor subunit RNA after chronic nicotine treatment. *J Neurosci* **12**:2765–2784.
- Marks MJ, Whiteaker P, Calcaterra J, Stitzel JA, Bullock AE, Grady SR, Picciotto MR, Changeux JP, and Collins AC (1999) Two pharmacologically distinct components of nicotinic receptor-mediated rubidium efflux in mouse brain require the $\beta 2$ subunit. *J Pharmacol Exp Ther* **289**:1090–1103.
- Marubio LM, del Mar Arroyo-Jimenez M, Cordero-Erausquin M, Léna C, Le Novère N, de Kerchove d'Exaerde A, Huchet M, Damaj MI, and Changeux JP (1999) Reduced antinociception in mice lacking neuronal nicotinic receptor subunits. *Nature* **398**:805–810.
- Millar NS and Gotti C (2009) Diversity of vertebrate nicotinic acetylcholine receptors. *Neuropharmacology* **56**:237–246.
- Moroni M, Zwart R, Sher E, Cassels BK, and Bermudez I (2006) alpha4beta2 nicotinic receptors with high and low acetylcholine sensitivity: pharmacology, stoichiometry, and sensitivity to long-term exposure to nicotine. *Mol Pharmacol* **70**:755–768.
- Orr-Urtreger A, Göldner FM, Saeki M, Lorenzo I, Goldberg L, De Biasi M, Dani JA, Patrick JW, and Beaudet AL (1997) Mice deficient in the alpha7 neuronal nicotinic acetylcholine receptor lack alpha-bungarotoxin binding sites and hippocampal fast nicotinic currents. *J Neurosci* **17**:9165–9171.
- Picciotto MR, Zoli M, Léna C, Bessis A, Lallemand Y, Le Novère N, Vincent P, Pich EM, Brûlet P, and Changeux JP (1995) Abnormal avoidance learning in mice lacking functional high-affinity nicotine receptor in the brain. *Nature* **374**:65–67.
- Porter JT, Cauli B, Tsuzuki K, Lambolez B, Rossier J, and Audinat E (1999) Selective excitation of subtypes of neocortical interneurons by nicotinic receptors. *J Neurosci* **19**:5228–5235.
- Ross SA, Wong JY, Clifford JJ, Kinsella A, Massalas JS, Horne MK, Scheffer IE, Kola I, Waddington JL, Berkovic SF, et al. (2000) Phenotypic characterization of an alpha 4 neuronal nicotinic acetylcholine receptor subunit knock-out mouse. *J Neurosci* **20**:6431–6441.
- Salas R, Orr-Urtreger A, Broide RS, Beaudet A, Paylor R, and De Biasi M (2003) The nicotinic acetylcholine receptor subunit $\alpha 5$ mediates short-term effects of nicotine in vivo. *Mol Pharmacol* **63**:1059–1066.
- Salminen O, Murphy KL, McIntosh JM, Drago J, Marks MJ, Collins AC, and Grady SR (2004) Subunit composition and pharmacology of two classes of striatal presynaptic nicotinic acetylcholine receptors mediating dopamine release in mice. *Mol Pharmacol* **65**:1526–1535.
- Salminen O, Whiteaker P, Grady SR, Collins AC, McIntosh JM, and Marks MJ (2005) The subunit composition and pharmacology of alpha-Conotoxin MII-binding nicotinic acetylcholine receptors studied by a novel membrane-binding assay. *Neuropharmacology* **48**:696–705.
- Whiteaker P, Cooper JF, Salminen O, Marks MJ, McClure-Begley TD, Brown RW, Collins AC, and Lindstrom JM (2006) Immunolabeling demonstrates the interdependence of mouse brain alpha4 and beta2 nicotinic acetylcholine receptor subunit expression. *J Comp Neurol* **499**:1016–1038.
- Wonnacott S (1997) Presynaptic nicotinic ACh receptors. *Trends Neurosci* **20**:92–98.
- Xu W, Orr-Urtreger A, Nigro F, Gelber S, Sutcliffe CB, Armstrong D, Patrick JW, Role LW, Beaudet AL, and De Biasi M (1999) Multiorgan autonomic dysfunction in mice lacking the beta2 and the beta4 subunits of neuronal nicotinic acetylcholine receptors. *J Neurosci* **19**:9298–9305.
- Zwart R and Vijverberg HP (1998) Four pharmacologically distinct subtypes of alpha4beta2 nicotinic acetylcholine receptor expressed in *Xenopus laevis* oocytes. *Mol Pharmacol* **54**:1124–1131.

Address correspondence to: Dr. Allan C. Collins, Institute for Behavioral Genetics, University of Colorado, Boulder, UCB 447, Boulder, CO 80309-0447. E-mail: al.collins@colorado.edu