

Research article

Open Access

Association of IL-10 and IL-10R β gene polymorphisms with graft-versus-host disease after haematopoietic stem cell transplantation from an HLA-identical sibling donor

Jyrki Sivula^{†1}, Hannu Turpeinen^{†1}, Liisa Volin² and Jukka Partanen^{*1}

Address: ¹Research and Development, Finnish Red Cross Blood Service, Helsinki, Finland and ²Department of Medicine, Helsinki University Central Hospital, Helsinki, Finland

Email: Jyrki Sivula - jyrki.sivula@helsinki.fi; Hannu Turpeinen - hannu.turpeinen@veripalvelu.fi; Liisa Volin - liisa.volin@hus.fi; Jukka Partanen* - jukka.partanen@veripalvelu.fi

* Corresponding author †Equal contributors

Published: 4 May 2009

Received: 23 June 2008

BMC Immunology 2009, 10:24 doi:10.1186/1471-2172-10-24

Accepted: 4 May 2009

This article is available from: <http://www.biomedcentral.com/1471-2172/10/24>

© 2009 Sivula et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Extensive allelic matching in the human leukocyte antigen (HLA) genes is regarded as a prerequisite for good clinical success of allogeneic haematopoietic stem cell transplantation (HSCT). Also other genetic factors can be assumed to play a role in preventing and controlling the complications associated with allogeneic HSCT, in particular graft-versus-host disease (GvHD). Interleukin-10 (IL-10) and its receptor (IL-10R), key regulators of the immune response, are among these candidates. We studied the association of IL-10 and IL-10R β gene polymorphisms with the occurrence of GvHD in 309 HLA-identical sibling donor and recipient pairs.

Results: The difference in genotypic IL-10 production between patient and donor in combination with patient IL-10R β A/A genotype predisposed strongly to acute GvHD (OR = 7.15, p = 0.00023). On the other hand, a combination of same genotypic IL-10 production with patient IL-10R β A/A genotype protected from chronic GvHD (OR = 0.407, p = 0.0097).

Conclusion: Our results suggest that IL-10 and IL-10R β genes have a synergistic effect on the risk of GvHD.

Background

Extensive allelic matching in the human leukocyte antigen (HLA) genes between recipient and donor is regarded as a prerequisite for good clinical success of allogeneic haematopoietic stem cell transplantation (HSCT) [1]. Graft-versus-host disease (GvHD) can not be totally avoided even with fully HLA-matched sibling donors or by using current GvHD prophylaxis, indicating complex and multifactorial nature of GvHD. This has lead into search of novel genetic factors, in addition to the HLA genes, to prevent and predict the occurrence of GvHD. Genes for killer

cell immunoglobulin like receptors (KIR), KIR ligands and minor histocompatibility antigens as well as several cytokines and cytokine receptors have obtained a considerable interest [2-4].

Cytokine interleukin 10 (IL-10) is a good functional candidate for GvHD-related gene [5,6]. Previous studies have shown that patient or donor IL-10 genotypes separately associate with both acute and chronic GvHD [6-9]. IL-10 is produced by a variety of different cells, of both haematopoietic and non-haematopoietic lineages [10]. IL-10 is

usually regarded as a potent suppressor of the immune responses and hence it is thought to be useful in preventing GvHD. However, it has also been shown to have some immunostimulatory effects [11-13]. Interleukin 10 receptor beta (IL-10R β) is also expressed on several cell types, and its role varies depending on the cell type it is expressed on [14]. In haematopoietic cell lineages IL-10R β functions as a part of IL-10 receptor together with IL-10 receptor alpha (IL-10R α). IL-10R α , the IL-10 specific part of the receptor complex, is only expressed on haematopoietic cells [5]. Thus in haematopoietic cells IL-10R β is mediating the IL-10 signal. In many non-haematopoietic cell lineages IL-10R β functions as a part of a receptor for other cytokines [14].

To address the role of IL-10 and IL-10R β in HSCT we here report the effects of polymorphisms in these genes on the incidence of acute and chronic GvHD in 309 Finnish HSCT recipients who had an HLA identical sibling donor. In particular, we addressed the questions of interplay of patient- and donor-specific factors of IL-10 and IL-10R β genes.

Results

Altogether 309 allogeneic HSCT recipients and their HLA-identical sibling donors were included in the study. The demographic and other background information is given in Table 1. Genetic variation in IL-10 and IL-10R β genes was studied and genetic association with the occurrence of GvHD in the patients was tested.

1. Genotype and haplotype association of IL-10 and IL-10R β gene polymorphisms with GvHD

The distributions of genotypes and haplotypes in patients and donors were studied for their association with the occurrence of acute (none versus grade III-IV) and chronic (none versus extensive) GvHD (Additional files 1 and 2).

In the patients, the genotype rs1800872 A/A in IL-10 gene predisposed to grade III-IV acute GvHD ($p = 0.031$, OR = 3.83) when compared to other genotypes (A/C and C/C). As a result of strong linkage disequilibrium between the IL-10 gene markers, the same association was also detected for the rs1800872 - rs1800896 haplotypes (Additional files 1 and 2). In the IL-10R β gene, patient's A/A genotype predisposed to acute GvHD ($p = 0.0035$, OR = 3.88) and A/G genotype showed protection from it ($p = 0.017$, OR = 0.30).

When the patients and donors were stratified based on their IL-10 and IL-10R β genotypes together, this approach did not suggest any novel genetic association.

In the patients, none of the genotypes of IL-10 or IL-10R β showed statistically significant association with chronic

Table 1: Sample demographics

Patient age, years mean (range)	47 (17-66)
Donor age, years mean (range)	45 (11-68)
Diagnosis, frequency	
Acute myeloid leukaemia	77
Multiple myeloma	54
Acute lymphoid leukaemia	48
Chronic lymphoid leukaemia	45
Myelodysplastic syndrome	23
Chronic myeloid leukaemia	13
Non-Hodgkin lymphoma	13
Other	36
Disease status	
Good prognosis	148
Bad prognosis	142
Gender, patient-donor	
male-male	82
male-female	67
female-male	76
female-female	63
Graft origin	
Peripheral blood	128
Bone marrow	162
Conditioning	
Myeloablative	233
Other	57
GvHD prevention	
Cyclosporine, Methotrexate and Methylprednisolone	207
Cyclosporine and Mycophenolate Mofetil	46
Cyclosporine and Methotrexate	37
Acute GvHD	
No	197
grade III-IV	29
Chronic GvHD	
No	117
Extensive	72

GvHD. In the donors, no genotypes or haplotypes showed statistically significant association with either chronic or acute GvHD (Additional files 1 and 2).

Previously, Lin et al. [7] described that a certain combination of patient IL-10 and donor IL-10R β polymorphisms protected very strongly from acute GvHD. Hence, we carried out similar analysis (Table 2). We also found that none of the three sibling pairs in our data set with IL-10 rs1800872 A/A (patient) - IL-10R β rs28341676 G/G (donor) genotypes actually resulted in acute GvHD, but no statistical significance or similar trend of decreasing acute GvHD frequency as Lin et al. observed toward this genotype combination could be seen.

Table 2: Incidence of grade III-IV acute GvHD according to patient IL-10 (-592) rs1800872 and donor IL-10Rβ(+238) rs28341676 genotypes/all cases (percent)

	Patient IL-10 rs1800872		
	A/A	A/C	C/C
Donor IL-10Rβ rs28341676			
A/A	3/8 (38%)	5/53 (9%)	9/90 (10%)
A/G	2/5 (40%)	4/32 (13%)	4/70 (6%)
G/G	0/3 (0%)	0/12 (0%)	1/24 (4%)

2. Associations with predicted IL-10 production levels

Genotype classes with different predicted production levels were tested for their association with the occurrence of GvHD but no statistically significant results could be found. However, patient intermediate genotypic production level was borderline protective from acute GvHD ($p = 0.073$, $OR = 0.431$).

We furthermore subdivided the study group based on their IL-10Rβ genotype and whether the donor – recipient pair had the same or different production level of IL-10 (Additional file 3). Those patients with the IL-10Rβ A/A genotype who received the graft from a donor with a different predicted IL-10 level to that of the recipient, had a very high risk for acute GvHD ($p = 0.000023$, $OR = 7.15$). Similarly, when the IL-10Rβ A/A homozygous patients received a graft with a same IL-10 production level, they were protected from chronic GvHD ($p = 0.0097$, $OR = 0.407$; Additional file 3).

3. Multivariate analysis

Logistic regression multivariate analysis was then performed to test if the statistically significant results observed in univariate analyses would hold, when tested in the context of other known factors influencing GvHD; disease status (good prognosis versus bad prognosis), gender match (male versus female donor), graft origin (bone marrow versus peripheral blood), pre-transplantation conditioning (myeloablative versus other) and aGvHD as a risk factor for cGvHD. All statistically significant results of univariate IL-10 and IL-10Rβ remained significant in multivariable analyses. Results of these analyses are shown in Table 3. We also performed multivariable analyses to see if the studied IL-10 and IL-10Rβ polymorphisms correlated with the other factors used in these analysis. These results showed no correlation between disease status, gender match, graft origin and pre-transplantation conditioning, and the IL-10 and IL-10Rβ polymorphisms.

Discussion

In the present study we analysed the role of IL-10 and IL-10Rβ SNPs on the incidence of acute and chronic GvHD after HSCT between HLA-identical siblings. Previously

published studies on interaction of IL-10 and HSCT have found association between IL-10 polymorphism and GvHD outcome [6,8,9,15-18]. In these studies the results predominantly rely on polymorphisms of either recipient or donor. Here we addressed the question also by studying the association of the SNPs with GvHD in biologically more meaningful context; that is analysing simultaneously both donor and patient genotypes.

In our analysis patient IL-10 rs1800872 genotype A/A was associated with worse acute GvHD outcome (Additional file 1 and Table 3). Similar result was also observed with rs1800896 and rs1800872 genotype combination AA/AA. Also patient IL-10Rβ rs28341676 genotype affected the risk of developing GvHD. Patient IL-10Rβ rs28341676 A/A predisposed to and A/G protected from grades III-IV acute GvHD. Donor IL-10 rs1800896 and rs1800872 or IL-10Rβ rs28341676 polymorphisms did not have any clear effect on the GvHD outcome (Additional file 2).

Because the cells of the graft are mainly haematopoietic cells, the effect of IL-10Rβ polymorphism on the donor side is restricted to recognition of IL-10. Since most of the patient's original haematopoietic cells are eliminated, the effect of IL-10Rβ on patient side is largely restricted to the signalling of other cytokines where the IL-10Rβ also functions as a part of the receptor complex. Cytokines known to date to utilise IL-10Rβ are IL-10, IL-22, IL-26, IL-28A, IL-28B and IL-29 [14]. Since donor IL-10Rβ had no significance on GvHD predisposition, the effects of IL-10Rβ polymorphism would seem to rely largely on functions of these other cytokines utilizing IL-10Rβ in their signalling.

When analysing both the patient and donor IL-10 genotypes the results showed a trend for lower risk of developing GvHD when both had similar IL-10 production levels estimated from genotypes (Additional file 3). Similar results were also seen by Bertinetto et al [18]. In their study the presence of IL-10 rs180087 G allele in both patient and donor was associated with a trend for lower risk of developing GvHD. The rationale for this is the fact that IL-10 is a strong regulator of immune system and it might be that either higher or lower production level of the protein as compared to original one would not be

Table 3: Multivariable logistic regression analysis of acute and chronic GvHD

Acute GvHD	B	S.E.	p
Disease status	0.92	0.47	0.053
Gender match	-0.29	0.50	0.56
Graft origin	0.34	0.50	0.49
Conditioning	0.055	0.60	0.93
Patient IL10 rs1800872 A/A	1.30	0.60	0.031
Constant	-2.65	0.53	

Acute GvHD	B	S.E.	p
Disease status	1.26	0.53	0.017
Gender match	-0.32	0.50	0.53
Graft origin	0.13	0.49	0.80
Conditioning	-0.11	0.61	0.85
Patient IL10R β rs28341676 A/A	1.33	0.46	0.0041
Constant	-3.13	0.58	

Acute GvHD	B	S.E.	p
Disease status	1.26	0.53	0.017
Gender match	-0.39	0.56	0.49
Graft origin	0.09	0.52	0.86
Conditioning	-0.52	0.67	0.44
IL-10 production level different and patient IL-10R β A/A	2.16	0.49	0.000088
Constant	-3.02	0.56	

Chronic GvHD	B	S.E.	p
Disease status	0.38	0.50	0.50
Gender match	-1.20	0.55	0.028
Graft origin	-1.58	0.48	0.0010
Conditioning	1.10	0.65	0.087
aGvHD	2.64	0.95	0.0052
IL-10 production level same and patient IL-10R β A/A	-1.18	0.50	0.017
Constant	0.58	0.43	

optimal for the patient's immune system. The effect of patient IL-10R β was also amplified when analyzed together with IL-10 production level. IL-10R β A/A strongly predisposed to acute GvHD when patient and donor IL-10 production level differed, but protected from chronic GvHD when patient and donor had similar IL-10 production level (Additional file 3 and Table 3).

We were not able to reproduce most of the significant results of Lin et al [7] with our study material. In our results the donor IL-10R β had no significant effect on the GvHD even in combination with IL-10 polymorphisms. Also, patient IL-10 rs1800872 A/A predisposed to acute GvHD (Additional file 1 p = 0.031, OR = 3.83). However, as Lin et al also observed, no acute GvHD cases were present among those three pairs with the combination of patient IL-10 rs1800872 A/A and donor IL-10R β rs28341676 G/G (Table 2). Compared to our results, in Lin et al study the frequency of IL-10 rs1800872 A allele was higher. This might in part explain the striking differ-

ence between our results. The rs1800872 A allele is often associated with lower risk of developing GvHD, but there are also studies which show similar association as we found [6,18]. It is possible that the rs1800872 is not it self affecting to the risk of GvHD, but is in linkage disequilibrium with the actual effector. In different populations this linkage might vary and cause the discrepancy in the results.

In analyses of IL-10R β rs28341676 the results differed between acute and chronic GvHD. Patient IL-10R β A/A predisposed to acute GvHD and seemed to protect from chronic GvHD. The different effect of IL-10R β polymorphism between acute GvHD and chronic GvHD and overall weaker results in chronic GvHD can be understood by the immunologic differences between acute and chronic GvHD. During the period of which acute GvHD is observed, the immune system is just starting to reconstitute anew after the HSCT. Also, many of the effector cells had been developed within the donor. In many cell line-

ages their numbers start rising only well after the usual 100 days time limit of acute GvHD diagnosis. More cells of donor origin are developing within the recipient during chronic GvHD. The immune system is more mature and complex than in the newly transplanted patient [19]. Notably, amounts of regulatory cells rise only relatively late in immune reconstitution [20]. The full reconstitution of immune system after transplantation may take years. Hence it is not surprising that IL-10 and IL-10R β polymorphisms have less effect in chronic GvHD than in acute GvHD. This could also explain the opposing effect of IL-10R β polymorphism in acute GvHD and chronic GvHD.

We are fully aware of the possible problems in analyzing all the transplantations as a single group. Grafting peripheral blood stem cells instead of bone marrow cells predisposes to higher incidence of chronic GvHD. Also pre-treatment conditioning, fully myeloablative versus reduced intensity conditioning, highly affects the complications after HSCT. To address this issue we performed logistic regression multivariate analysis to test the results of univariate analyses in the context of known GvHD affecting factors; disease status, gender match, graft origin, pre-transplantation conditioning and aGvHD as a risk factor for cGvHD. All the statistically significant results from univariate analyses remained significant in the multivariate analyses (Table 3). The p-values in this study were not corrected for multiple comparisons. These results need to be validated in other cohorts and further studies are also needed to clarify the relationships between IL-10 and IL-10R β .

Conclusion

We did observe synergistic effect between IL-10 and IL-10R β in predisposition to GvHD. The effect of IL-10R β polymorphism was strengthened when it was analysed together with IL-10 polymorphisms. In our samples similarity of IL-10 production level was associated with lower incidence of both acute and chronic GvHD. Different effect of IL-10R β genotype in acute GvHD and chronic GvHD could be explained by the developing immune system, arise of new regulatory cells and passing cytokine storm associated with the transplantation.

Methods

Patients

Altogether 309 adult allogeneic HSCT recipients and their HLA-identical sibling donors were included in this retrospective study. All transplantations were performed in a single centre (Department of Medicine, Helsinki University Central Hospital, Helsinki, Finland) between years 1993 and 2005. Clinical data were collected on diagnosis, age, gender match, transplantation protocol, date of transplantation and acute GvHD and chronic GvHD manifestation. GvHD was diagnosed and graded according to

standard criteria [21]. Patient and donor descriptive demographics are shown in Table 1.

This study was approved by the Ethical Review Board of the Helsinki University Hospital, Helsinki, Finland.

IL-10 and IL-10R β genotyping

Two single nucleotide polymorphisms (SNPs) in IL-10 and one in IL-10R β were genotyped. The IL-10 SNPs at locations -1082 (rs1800896) and -592 (rs1800872) and the IL-10R β SNP at location +238 (rs28341676) were typed using restriction fragment length polymorphism method as described earlier [7,22]. EcoNI recognition site primers and restriction enzyme was used for IL-10.

In the Finnish population, the selected two markers for IL-10 tag completely the often used three-marker haplotype (-1082 A/G (rs1800896), -819 T/C (rs1800871), and -592 A/C (rs1800872)) for the gene [23].

Statistical analysis

To analyse the effect of IL-10 and IL-10R β polymorphisms on the GvHD outcome we used different combinations of patient and donor genotyping data. We (i) analysed IL-10 and IL-10R β genotypes and haplotypes separately in patients and donors, and (ii) combined the genotyping data from each patient-donor pair. We also (iii) grouped the IL-10 genotypes into three production level groups: high, intermediate and low producing. These groups are based on the effect of IL-10 polymorphisms on the level of protein production as published earlier [24,25]: IL-10 rs1800896 (SNP1) and rs1800872 (SNP2) genotype combination (SNP1SNP2/SNP1SNP2) GC/GC belongs to high producing, GC/AC and GC/AA to intermediate producing, and AC/AC, AC/AA and AA/AA to low producing group.

Analyses of different IL-10 and IL-10R β combinations were done in relation to the acute GvHD and chronic GvHD incidence using χ^2 or Fisher's exact tests. Recipients with acute GvHD of grade III-IV and their donors formed the positive group which was compared to patients with no acute GvHD and their donors. Chronic GvHD positive group patients had extensive chronic GvHD and their controls had no chronic GvHD. To test the statistically significant results of univariate analyses in the context of other risk factors we constructed a multivariable logistic regression model adjusted for disease status, gender match, graft origin and pre-transplantation conditioning. We also included aGvHD as a risk factor for cGvHD. The SPSS for Windows v15.0 software (SPSS Inc., Chicago, IL, USA) was used for the analyses. All p-values are considered as two-sided and uncorrected for multiple testing. P-values under 0.05 were regarded to indicate statistical significance.

Abbreviations

GvHD: Graft versus host disease; aGvHD: Acute graft versus host disease; cGvHD: Chronic graft versus host disease; HLA: Human leukocyte antigen; HSCT: Haematopoietic stem cell transplantation; IL-10: Interleukin-10; IL-10R: Interleukin-10 receptor; IL-10R β : Interleukin-10 receptor beta subunit; IL-10R α : Interleukin-10 receptor alpha subunit; KIR: Killer cell immunoglobulin like receptor.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

LV clinical expertise, JS and HT genetic typing and data analysis. All authors contributed in the study hypothesis and study design and all authors contributed to the preparation of the manuscript. All authors read and approved the final manuscript. JS and HT share an equal contribution to the article.

Authors' information

JS, HT and JP: Research and Development, Finnish Red Cross Blood Service, Helsinki, Finland

LV: Department of Medicine, Helsinki University Central Hospital, Helsinki, Finland

Address for correspondence: Dr Jukka Partanen, Research and Development, Finnish Red Cross Blood Service, Kivihaantie 7, 00310 Helsinki, Finland. jukka.partanen@bts.redcross.fi

Additional material

Additional file 1

Table s1. Distribution of IL-10 (-1082) rs1800896 and (-592) rs1800872, and IL-10R β (+238) rs28341676 genotypes and their association with the occurrence of acute and chronic GvHD in the patients
Click here for file
[\[http://www.biomedcentral.com/content/supplementary/1471-2172-10-24-S1.doc\]](http://www.biomedcentral.com/content/supplementary/1471-2172-10-24-S1.doc)

Additional file 2

Table s2. Distribution of IL-10 (-1082) rs1800896 and (-592) rs1800872, and IL-10R β (+238) rs28341676 genotypes and their association with the occurrence of acute and chronic GvHD in the patients donors
Click here for file
[\[http://www.biomedcentral.com/content/supplementary/1471-2172-10-24-S2.doc\]](http://www.biomedcentral.com/content/supplementary/1471-2172-10-24-S2.doc)

Additional file 3

Table s3. Incidence of GvHD according to patient IL-10R β (+238) rs28341676 and patient and donor IL-10 genotypic production levels
Click here for file
[\[http://www.biomedcentral.com/content/supplementary/1471-2172-10-24-S3.doc\]](http://www.biomedcentral.com/content/supplementary/1471-2172-10-24-S3.doc)

Acknowledgements

This research was supported by grants from the Sigrid Juselius Foundation, the Academy of Finland and Government EVO Fund. We would like to thank Sisko Lehmonen and the staff of Tissue Typing Laboratory of the Finnish Red Cross Blood Service for skilful technical help.

References

- Petersdorf EW: **HLA matching in allogeneic stem cell transplantation.** *Current Opinion in Hematology* 2004, **11**:386-391.
- Velardi A, Ruggeri L, Capanni M, Mancusi A, Perruccio K, Aversa F, Martelli MF: **Immunotherapy with alloreactive natural killer cells in haploidentical haematopoietic transplantation.** *Hematol J* 2004, **5**(Suppl 3):S87-S90.
- Dickinson AM, Middleton PG, Rocha V, Gluckman E, Holler E: **Genetic polymorphisms predicting the outcome of bone marrow transplants.** *Br J Haematol* 2004, **127**:479-490.
- Bonnet D, Warren EH, Greenberg PD, Dick JE, Riddell SR: **CD8(+) minor histocompatibility antigen-specific cytotoxic T lymphocyte clones eliminate human acute myeloid leukemia stem cells.** *Proc Natl Acad Sci USA* 1999, **96**:8639-8644.
- Moore KW, de Waal MR, Coffman RL, O'Garra A: **Interleukin-10 and the interleukin-10 receptor.** *Annu Rev Immunol* 2001, **19**:683-765.
- Kim DH, Lee NY, Sohn SK, Baek JH, Kim JG, Suh JS, Lee KB, Shin IH: **IL-10 promoter gene polymorphism associated with the occurrence of chronic GVHD and its clinical course during systemic immunosuppressive treatment for chronic GVHD after allogeneic peripheral blood stem cell transplantation.** *Transplantation* 2005, **79**:1615-1622.
- Lin MT, Storer B, Martin PJ, Tseng LH, Grogan B, Chen PJ, Zhao LP, Hansen JA: **Genetic variation in the IL-10 pathway modulates severity of acute graft-versus-host disease following hematopoietic cell transplantation: synergism between IL-10 genotype of patient and IL-10 receptor beta genotype of donor.** *Blood* 2005, **106**:3995-4001.
- Karabon L, Wysoczanska B, Bogunia-Kubik K, Suchnicki K, Lange A: **IL-6 and IL-10 promoter gene polymorphisms of patients and donors of allogeneic sibling hematopoietic stem cell transplants associate with the risk of acute graft-versus-host disease.** *Human Immunology* 2005, **66**:700-710.
- Cavet J, Middleton PG, Segall M, Noreen H, Davies SM, Dickinson AM: **Recipient tumor necrosis factor-alpha and interleukin-10 gene polymorphisms associate with early mortality and acute graft-versus-host disease severity in HLA-matched sibling bone marrow transplants.** *Blood* 1999, **94**:3941-3946.
- Akdis CA, Blaser K: **Mechanisms of interleukin-10-mediated immune suppression.** *Immunology* 2001, **103**:131-136.
- Nolan KF, Strong V, Soler D, Fairchild PJ, Cobbold SP, Croxton R, Gonzalo JA, Rubio A, Wells M, Waldmann H: **IL-10-conditioned dendritic cells, decommissioned for recruitment of adaptive immunity, elicit innate inflammatory gene products in response to danger signals.** *J Immunol* 2004, **172**:2201-2209.
- Moore KW, O'Garra A, de Waal MR, Vieira P, Mosmann TR: **Interleukin-10.** *Annu Rev Immunol* 1993, **11**:165-190.
- Reitamo S, Remitz A, Tamai K, Uitto J: **Interleukin-10 modulates type I collagen and matrix metalloprotease gene expression in cultured human skin fibroblasts.** *J Clin Invest* 1994, **94**:2489-2492.
- Donnelly RP, Sheikh F, Kotenko SV, Dickensheets H: **The expanded family of class II cytokines that share the IL-10 receptor-2 (IL-10R2) chain.** *J Leukoc Biol* 2004, **76**:314-321.
- Socie G, Loiseau P, Tamouza R, Janin A, Busson M, Gluckman E, Charon D: **Both genetic and clinical factors predict the development of graft-versus-host disease after allogeneic hematopoietic stem cell transplantation.** *Transplantation* 2001, **72**:699-706.
- Lin MT, Storer B, Martin PJ, Tseng LH, Gooley T, Chen PJ, Hansen JA: **Relation of an interleukin-10 promoter polymorphism to graft-versus-host disease and survival after hematopoietic cell transplantation.** *New England Journal of Medicine* 2003, **349**:2201-2210.
- Cavet J, Dickinson AM, Norden J, Taylor PRA, Jackson GH, Middleton PG: **Interferon-gamma and interleukin-6 gene polymorphisms associate with graft-versus-host disease in HLA-matched sibling bone marrow transplantation.** *Blood* 2001, **98**:1594-1600.

18. Bertinetto FE, Dall'Omo AM, Mazzola GA, Rendine S, Berrino M, Bertola L, Magistroni P, Caropreso P, Falda M, Locatelli F, et al.: **Role of non-HLA genetic polymorphisms in graft-versus-host disease after haematopoietic stem cell transplantation.** *International Journal of Immunogenetics* 2006, **33**:375-384.
19. Storek J, Dawson MA, Storer B, Stevens-Ayers T, Maloney DG, Marr KA, Witherspoon RP, Bensinger W, Flowers ME, Martin P, et al.: **Immune reconstitution after allogeneic marrow transplantation compared with blood stem cell transplantation.** *Blood* 2001, **97**:3380-3389.
20. Olkinuora H, Talvensaaari K, Kaartinen T, Siitonen S, Saarinen-Pihkala U, Partanen J, Vettentranta K: **T cell regeneration in pediatric allogeneic stem cell transplantation.** *Bone Marrow Transplant* 2007, **39**:149-156.
21. Przepiorka D, Weisdorf D, Martin P, Klingemann HG, Beatty P, Hows J, Thomas ED: **1994 Consensus Conference on Acute GVHD Grading.** *Bone Marrow Transplant* 1995, **15**:825-828.
22. Tseng LH, Chen PJ, Lin MT, Singleton K, Martin EG, Yen AH, Chuang SM, Martin PJ, Hansen JA: **Simultaneous genotyping of single nucleotide polymorphisms in the IL-6, IL-10, TNFalpha and TNFbeta genes.** *Tissue Antigens* 2002, **59**:280-286.
23. Woolley N, Mustalahti K, Maki M, Partanen J: **Cytokine gene polymorphisms and genetic association with coeliac disease in the Finnish population.** *Scandinavian Journal of Immunology* 2005, **61**:51-56.
24. Suarez A, Castro P, Alonso R, Mozo L, Gutierrez C: **Interindividual variations in constitutive interleukin-10 messenger RNA and protein levels and their association with genetic polymorphisms.** *Transplantation* 2003, **75**:711-717.
25. Crawley E, Kay R, Sillibourne J, Patel P, Hutchinson I, Woo P: **Poly-morphic haplotypes of the interleukin-10 5' flanking region determine variable interleukin-10 transcription and are associated with particular phenotypes of juvenile rheumatoid arthritis.** *Arthritis Rheum* 1999, **42**:1101-1108.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

