

## Collagen Binding, Elastase Production, and Slime Production Associated with Coagulase-Negative Staphylococci Isolated from Bovine Intramammary Infections†

J. L. WATTS,<sup>1\*</sup> A. S. NAIDU,<sup>2</sup> AND T. WADSTRÖM<sup>3</sup>

*Mastitis Research Laboratory, Hill Farm Research Station, Louisiana Agricultural Experiment Station, Louisiana State University Agricultural Center, Route 1, Box 10, Homer, Louisiana 71040,<sup>1</sup> and Department of Medical Microbiology, Malmö General Hospital, University of Lund, S-214 01 Malmö,<sup>2</sup> and Department of Medical Microbiology, University of Lund, S-22362 Lund,<sup>3</sup> Sweden*

Received 6 September 1989/Accepted 15 November 1989

Collagen binding, elastase production, slime production, and associated somatic cell counts were determined with 160 strains of coagulase-negative staphylococci isolated from bovine intramammary infections. Mean binding values for type I and II collagen with *Staphylococcus epidermidis*, *S. chromogenes*, and *S. hyicus* strains were 5.8, 6.6, and 7.4 and 4.3, 4.2, and 4.9%, respectively. Eleven of 28 (39.3%) *S. epidermidis*, 1 of 38 (2.6%) *S. chromogenes*, and 1 of 94 (1.1%) *S. hyicus* strains were elastase positive. Slime production was noted with 12 (42.9%) *S. epidermidis*, 1 (2.6%) *S. chromogenes*, and 11 (11.7%) *S. hyicus* strains. No differences in somatic cell counts were observed with type I or type II collagen binding, elastase production, or slime production with *S. epidermidis* or *S. chromogenes*. However, somatic cell counts associated with *S. hyicus* strains with collagen type I binding affinities of >5 and type II binding affinities of >3 were  $320.7 \times 10^3$  compared with  $163.9 \times 10^3$  for strains with lower binding affinities.

Historically, virulence of mastitis pathogens has been determined by the magnitude of specific cellular immune responses (14, 19). Polymorphonuclear leukocytes preferentially migrate into the udder in response to bacterial infection (18), increasing the somatic cell counts (SCC), which are highly predictive of infection and inversely correlated with milk production (14, 19). Past recommendations (20) considered SCC of normal milk to be  $500 \times 10^3$  or less. A recent study (14) indicated that normal SCC may be as low as  $50 \times 10^3$  and that significant milk losses occur at SCC above this level.

Coagulase-negative staphylococci are the organisms most frequently isolated from bovine intramammary infections (IMI) (1, 5, 6). Associated SCC levels have been placed at  $462 \times 10^3$  per ml (1, 3, 5). Since these SCC levels were within the  $500 \times 10^3$  level previously considered normal, coagulase-negative staphylococci were classified as nonpathogenic or minor pathogens (3). The realization that normal milk SCC were much lower than realized (14) has resulted in reclassification of the coagulase-negative staphylococci as mammary gland pathogens.

Delineation of virulent coagulase-negative staphylococci from nonvirulent resident flora has been difficult because virulence factors have not been well defined (8, 9). Gemmell and Schumacher-Perdreau (9) determined that coagulase-negative staphylococci produce as many as eight toxins and enzymes that could contribute to virulence. Of these, an enzyme resembling *Staphylococcus aureus* delta-toxin demonstrated the greatest biological activity (9). Hébert and Hancock (11) determined that this toxin could be readily detected by using a synergistic hemolysin assay. Watts and Owens (23) determined that elevated SCC levels were asso-

ciated with synergistic hemolysin assay-positive strains of *S. epidermidis* but not with strains of animal-associated, coagulase-negative staphylococci such as *S. hyicus* or *S. chromogenes*. Other workers (4, 13, 16, 20) have determined that coagulase-negative staphylococci produce a variety of substances capable of degrading or adhering to normal mammalian tissue components. However, the role of these substances in bovine mastitis has not been elucidated.

The purpose of this study was to determine the ability of coagulase-negative staphylococci isolated from bovine IMI to bind collagen, degrade elastin, and produce slime. Furthermore, SCC associated with collagen binding, elastase production, and slime production were determined.

(This paper was presented in part at the International Conference on Mastitis, 29 May-2 June 1989, St. Georgen, Austria.)

### MATERIALS AND METHODS

**Bacteria.** A total of 160 coagulase-negative staphylococci, including 28 *S. epidermidis*, 38 *S. chromogenes*, and 94 *S. hyicus* strains, were used. All isolates were obtained from bovine IMI. The Staph-Trac system (Analytab Products, Plainview, N.Y.) was used to identify all isolates. Accuracy of the system was established previously (22). Isolates demonstrating atypical test results were identified by a previously described conventional method (22).

All isolates were stored in Trypticase soy broth (BBL Microbiology Systems, Cockeysville, Md.) supplemented with 20% glycerin at  $-70^\circ\text{C}$ . Prior to testing, all isolates were subcultured twice on Trypticase soy agar (BBL) supplemented with 5% bovine blood and 0.1% esculin (Sigma Chemical Co., St. Louis, Mo.) for 24 h at  $35^\circ\text{C}$ .

**Collagen binding.** Determination of type I collagen (CN-I) and type II collagen (CN-II) (Collagen Corp., Palo Alto, Calif.) binding was performed as described previously (18). Briefly, CN-I and CN-II were labeled with  $^{125}\text{I}$  by the

\* Corresponding author.

† Approved for publication by the Director of the Louisiana Agricultural Experiment Station as manuscript 89-80-3312.




FIG. 1. Binding of  $^{125}\text{I}$ -labeled type I (●) and II (○) collagens to three species of coagulase-negative staphylococci isolated from bovine mastitis. Radioisotope protein binding assay was performed with bacterial cells grown in Trypticase soy broth, as described in Materials and Methods. A binding value of <5% of the total added radioactivity was considered negative.

chloramine-T method (10). Binding to staphylococcal cells was quantified by the method of Fröman et al. (7). Bacteria were grown in Trypticase soy broth at 37°C for 18 h with constant shaking. Cells were washed once in 0.15 M phosphate-buffered saline, pH 7.2, and the density was adjusted photometrically to  $10^{10}$  cells per ml.  $^{125}\text{I}$ -labeled CN-I and CN-II were diluted to a radioactivity measurement between 15,000 and 25,000 cpm in phosphate-buffered saline, pH 7.2, containing 1% bovine serum albumin. Labeled CN-I or CN-II (50  $\mu\text{l}$ ) was mixed with 100  $\mu\text{l}$  of bacterial suspension in a polystyrene centrifuge tube and kept at room temperature for 1 h. After addition of 2 ml of ice-cold phosphate-buffered saline containing 0.05% sodium azide and 0.1% Tween 20, tubes were centrifuged at  $4,500 \times g$  for 10 min and supernatants were aspirated. Pellet radioactivity was measured in a 1260-Multigamma counter (LKB-Wallac, Turku, Finland). *S. aureus* strains Cowan 1, Newman, and Wood 46 and two *Micrococcus luteus* strains were used as positive and negative controls, respectively, during the binding assay. Binding assays were performed in triplicate and repeated four times.

**Elastase production.** Elastase production was determined by the method of Janda (13). Briefly, each isolate was streaked radially onto a brain heart infusion agar (BBL) plate supplemented with 0.3% elastin purified from bovine neck ligament (Sigma). A total of eight isolates were streaked on each plate, and plates were incubated for 48 h at 35°C and then transferred to room temperature for an additional 8 days. Plates were observed for elastolytic activity after 24, 48, 72, 120, 168, 216, and 240 h. Clearing of elastin particles adjacent to bacterial colonies was considered positive.

**Slime production.** Slime production was determined by the method of Christensen et al. (4). Briefly, a single well-isolated colony from a 24-h bovine blood agar culture was

inoculated into 10 ml of Trypticase soy broth and incubated overnight at 37°C. The test tubes were emptied and stained with safranin. An isolate was considered positive for slime production when a visible film lined the tube walls. Ring formation at the liquid-air interface was considered negative.

**SCC.** SCC were performed as described previously (23). After microbiological analysis, milk samples containing sufficient volume were prepared for SCC determination by adding potassium dichromate (Nasco, Fort Atkinson, Wis.) to a final concentration of 0.2%. Samples were refrigerated at 2 to 8°C for at least 24 h to permit proper fixation of the somatic cells. All SCC were performed within 96 h of collection with a Fossomatic milk analyzer (A. S. Foss, Hillerød, Denmark) and reported as  $\text{SCC} \times 10^3$  per ml.

**Statistical analysis.** Least-squares analysis of variance was utilized, using the General Linear Models procedure of SAS (SAS Institute, Cary, N.C.). SCC were analyzed for effects of collagen binding, elastase production, and slime production.

## RESULTS

Mean CN-I binding values for *S. epidermidis*, *S. chromogenes*, and *S. hyicus* were 5.8, 6.6, and 7.4%, respectively. CN-II binding was lower, with values of 4.3, 4.2, and 4.9% for *S. epidermidis*, *S. chromogenes*, and *S. hyicus*, respectively (Fig. 1). Differences in associated SCC and collagen were observed with *S. hyicus* strains, which yielded CN-I binding of >5 and CN-II binding of >3, but not with *S. epidermidis* or *S. chromogenes* strains (Table 1). Only 7 of 28 (25.0%) *S. epidermidis* strains yielded CN-I binding values of >5 and CN-II binding values of >3. Mean SCC associated with high collagen binding strains was 59.7 compared with 84.0 for low collagen binding strains. Of 38 *S.*

TABLE 1. SCC associated with collagen-binding, coagulase-negative staphylococci isolated from bovine IMI

Organism	CN-I > 5, CN-II > 3		CN-I < 5, CN-II < 3	
	No. of isolates	SCC/ml (mean ± SE) (10 <sup>3</sup> )	No. of isolates	SCC/ml (mean ± SE) (10 <sup>3</sup> )
<i>S. epidermidis</i>	7	59.7 ± 32.6	21	84.0 ± 18.25
<i>S. chromogenes</i>	20	182.4 ± 71.5	18	152.5 ± 57.49
<i>S. hyicus</i>	43	320.7 ± 57.77	51	163.9 ± 40.45

*chromogenes* strains tested, 20 (52.8%) with high collagen binding values had SCC values of 182.4 compared with 152.5 for strains with lower binding affinities. SCC values for 43 (45.7%) *S. hyicus* strains with high collagen binding values were 320.7 compared with a mean SCC of 163.9 for strains with low collagen binding values. Differences between high and low collagen binding strains for each species were not statistically significant.

Elastase production and associated SCC of coagulase-negative staphylococci isolated from bovine IMI are presented in Table 2. Eleven of 28 (39.3%) *S. epidermidis* strains were elastase positive compared with only 1 of 38 (2.6%) *S. chromogenes* strains and 1 of 94 (1.1%) *S. hyicus* strains. SCC associated with elastase-positive *S. epidermidis* strains were higher (93.7) than SCC for elastase-negative strains (67.7). A single elastase-positive *S. chromogenes* strain was associated with an SCC of 89.0 compared with 170.4 for the elastase-negative strains. The elastase-positive *S. hyicus* strain was associated with an SCC of 656.0 compared with 231.1 for elastase-negative strains. Differences in SCC between elastase-positive and elastase-negative strains were not significant.

Slime production and associated SCC for the coagulase-negative staphylococcal species tested are presented in Table 3. Of the three species tested, 12 (42.9%) *S. epidermidis*, 1 (2.6%) *S. chromogenes*, and 11 (11.7%) *S. hyicus* strains were positive for slime production. However, SCC for the slime production-positive *S. epidermidis*, *S. chromogenes*, and *S. hyicus* strains were 85.1, 89.0, and 257.9 compared with 72.6, 170.4, and 232.7, respectively, for slime-negative strains. These differences were not statistically significant.

## DISCUSSION

*S. epidermidis*, *S. chromogenes*, and *S. hyicus* are the most frequently isolated coagulase-negative staphylococcal species from bovine IMI (1, 5, 6; reviewed in reference 21). *S. epidermidis* is considered a human-associated staphylococcal species and is the coagulase-negative staphylococcal species most often associated with abnormal milk or elevated SCC (1). *S. chromogenes* and *S. hyicus* are considered normal flora for the skin of dairy cattle and are frequently

TABLE 2. SCC associated with elastase-positive and -negative, coagulase-negative staphylococci isolated from bovine IMI

Organism	Elastase positive		Elastase negative	
	No. of isolates	SCC/ml (mean ± SE) (10 <sup>3</sup> )	No. of isolates	SCC/ml (mean ± SE) (10 <sup>3</sup> )
<i>S. epidermidis</i>	11	93.7 ± 30.56	17	67.7 ± 17.15
<i>S. chromogenes</i>	1	89.0	37	170.4 ± 47.11
<i>S. hyicus</i>	1	656.0	93	231.1 ± 35.18

TABLE 3. SCC associated with slime-positive and -negative, coagulase-negative staphylococci isolated from bovine IMI

Organism	Slime positive		Slime negative	
	No. of isolates	SCC/ml (mean ± SE) (10 <sup>3</sup> )	No. of isolates	SCC/ml (mean ± SE) (10 <sup>3</sup> )
<i>S. epidermidis</i>	12	85.1 ± 25.51	16	72.6 ± 20.48
<i>S. chromogenes</i>	1	89.0	37	170.4 ± 47.11
<i>S. hyicus</i>	11	257.9 ± 96.06	83	232.7 ± 37.82

isolated from bovine teat skin and teat canals as well as bovine IMI (2, 5, 6).

Results of the present study revealed differences among the three species in ability to bind CN-I and CN-II. Strains of *S. hyicus* and *S. chromogenes* demonstrated higher CN-I binding values than *S. epidermidis* strains. Miedzobrodzki et al. (16) reported that more *S. epidermidis* strains bound CN-I (76.7%) than did either *S. hyicus* (74.3%) or *S. chromogenes* (68.1%) strains. These results may be due to assay differences, as the particle agglutination assay used in the previous study (16) detected CN-I binding whereas the collagen binding assay used in the present study determined the degree of binding.

Mamö et al. (15) determined that the mean binding values for CN-II with three *S. hyicus* and three *S. chromogenes* strains were 5.3 and 13.0%, respectively. These findings are similar to results for the 94 *S. hyicus* strains (4.96%) in the present study. However, these findings are much higher than the 4.19% binding obtained with 38 *S. chromogenes* strains in the present study. Mamö et al. (15) did not include *S. epidermidis* strains: our study revealed a 4.29% mean binding value for this species.

No relationship between CN-I and CN-II binding and SCC for *S. epidermidis* and *S. chromogenes* was observed. However, SCC for *S. hyicus* strains with CN-I binding values of >5 and CN-II binding values of >3 were twice as high as for strains with CN-I and CN-II binding values below these thresholds. It appears that collagen binding may play a role in the virulence of *S. hyicus* but not *S. chromogenes* or *S. epidermidis*.

The ability of coagulase-negative staphylococci to degrade elastin may play an important role in tissue invasion. Janda (13) examined 161 staphylococci of human origin for elastolytic activity and determined that, of 12 coagulase-negative staphylococcal species tested, only strains of *S. epidermidis* were capable of producing elastase. Results of the present study revealed that 39.3% of *S. epidermidis* strains of bovine origin were elastase positive compared with 100% of strains in the previous study (13). Also, one strain each of *S. chromogenes* and *S. hyicus* was elastase positive. Comparison of associated SCC between elastase-positive and -negative *S. epidermidis* strains revealed only minor differences. These findings indicate that, while elastase may play a role in tissue invasion, this enzyme probably has a minor role in the pathogenicity of *S. epidermidis* in bovine IMI.

Slime production has been determined to be associated with *S. epidermidis* in disease and is used for recognition of pathogenic coagulase-negative staphylococci isolated from human sources (4, 11, 12). Christensen et al. (4) determined that slime production was associated with gentamicin resistance and symptomatic infections. Ishak et al. (12) determined that 13 of 14 clinically significant coagulase-negative staphylococci isolates produced slime, whereas only 3 of 13 contaminants and 4 of 27 skin isolates were slime producers.

Furthermore, these workers (12) found that *S. epidermidis* produced slime more frequently than other species. Hébert and Hancock (11) determined that 83% of 672 *S. epidermidis* strains of human origin were slime producers and that the slime production test may be useful in defining clinical significance. Results of the present study showed that a larger percentage of *S. epidermidis* strains (42.9%) than *S. chromogenes* (2.6%) or *S. hyicus* (11.7%) produced slime. Comparison of SCC associated with slime-positive and slime-negative strains revealed only slight differences. Thus, it appeared that slime production was not a reliable indicator of pathogenicity for coagulase-negative staphylococci isolated from bovine IMI.

Virulence factors associated with coagulase-negative staphylococci isolated from bovine IMI remain undefined. Results of the present study indicate that elastase, slime production, and collagen binding were not associated with increased SCC. A previous study (23) determined that production of a delta-like toxin by *S. epidermidis* strains was associated with a threefold increase in SCC. Thus, it appears that toxin production by *S. epidermidis* is the most reliable indicator of pathogenicity. Elastase and slime production occurred with only one *S. chromogenes* strain, and SCC with all strains were low. Furthermore, a previous study (23) indicated that relatively few *S. chromogenes* strains produced a delta-like toxin. Thus, *S. chromogenes* may be relatively nonpathogenic and, since this organism readily colonizes teat canals (2), IMI produced by this organism may be a consequence of opportunity rather than virulence. Increased SCC were observed with *S. hyicus* strains with high collagen binding affinities but not with elastase or slime production in the present study or toxin production in a previous study (23). Whether the increased collagen binding or another substance common to these strains was responsible for virulence has not been elucidated.

#### LITERATURE CITED

- Baba, E., T. Fukata, and H. Matsumoto. 1980. Ecological studies on coagulase-negative staphylococci in and around bovine udder. *Bull. Osaka Pref.* 32:69-75.
- Boddie, R. L., S. C. Nickerson, W. E. Owens, and J. L. Watts. 1987. Udder microflora in non-lactating heifers. *AGRI-Practice* 8:22-25.
- Bramley, A. J., and C. F. H. Dodd. 1984. Reviews of the progress of dairy science: mastitis control-progress and prospects. *J. Dairy Res.* 51:481-512.
- Christensen, G. D., J. T. Parisi, A. L. Bisno, W. A. Simpson, and E. H. Beachey. 1983. Characterization of clinically significant strains of coagulase-negative staphylococci. *J. Clin. Microbiol.* 18:258-269.
- Devriese, L. A., and H. De Keyser. 1980. Prevalence of different species of coagulase-negative staphylococci on teats and in milk samples from dairy cows. *J. Dairy Res.* 47:155-158.
- Devriese, L. A., and J. Deryeke. 1979. *Staphylococcus hyicus* in cattle. *Res. Vet. Sci.* 26:356-358.
- Fröman, G., L. M. Switalski, A. Faris, T. Wadström, and M. Höök. 1984. Binding of *Escherichia coli* to fibronectin. A mechanism of adherence. *J. Biol. Chem.* 23:14899-14905.
- Gemmell, C. G. 1983. Extracellular toxins and enzymes of coagulase-negative staphylococci, p. 809-827. In C. S. F. Easmon and C. Adlam (ed.), *Staphylococci and staphylococcal infections*. Academic Press, Inc., New York.
- Gemmell, C. G., and F. Schumacher-Perdreau. 1986. Extracellular toxins and enzymes elaborated by coagulase-negative staphylococci, p. 109-121. In P. A. Mardh and K. H. Schleifer (ed.), *Coagulase-negative staphylococci*. Almquist and Wiksell, Stockholm.
- Greenwood, F. C., W. M. Hunter, and J. S. Glover. 1963. The preparation of <sup>125</sup>I-labeled human growth hormone of high specific radioactivity. *Biochem. J.* 89:114-123.
- Hébert, G. A., and G. A. Hancock. 1985. Synergistic hemolysis exhibited by species of staphylococci. *J. Clin. Microbiol.* 22:409-415.
- Ishak, M. A., D. H. M. Gröschel, G. L. Mandell, and R. P. Wenzel. 1985. Association of slime with pathogenicity of coagulase-negative staphylococci causing nosocomial septicemia. *J. Clin. Microbiol.* 22:1025-1029.
- Janda, J. M. 1986. Elastolytic activity among staphylococci. *J. Clin. Microbiol.* 24:945-946.
- Jones, G. M., R. E. Pearson, G. A. Clabaugh, and C. W. Heald. 1984. Relationship between somatic cell counts and milk production. *J. Dairy Sci.* 67:1823-1831.
- Mamö, W., G. Fröman, and T. Wadström. 1988. Interaction of sub-epithelial connective tissue components with *Staphylococcus aureus* and coagulase-negative staphylococci from bovine mastitis. *Vet. Microbiol.* 18:163-176.
- Miedzobrodzki, J., A. S. Naidu, J. L. Watts, P. Ciborowski, K. Palm, and T. Wadström. 1988. Effect of milk on fibronectin and collagen type I binding to *Staphylococcus aureus* and coagulase-negative staphylococci isolated from bovine mastitis. *J. Clin. Microbiol.* 27:540-544.
- Naidu, A. S., J. Ekstrand, and T. Wadström. 1989. Binding of type-I and type-II collagens in *Staphylococcus aureus* strains isolated from patients with toxic shock syndrome compared to other staphylococcal infections. *FEMS Microbiol. Immunol.* 47:219-228.
- Nickerson, S. C., M. J. Paape, R. J. Harmon, and G. Ziv. 1986. Mammary leukocyte response to drug therapy. *J. Dairy Sci.* 69:1733-1742.
- Poutrel, B., and P. Rainard. 1982. Predicting the probability of quarter infection (by major pathogens) from somatic cell concentrations. *Am. J. Vet. Res.* 43:1296-1299.
- Wadström, T. 1987. Molecular aspects on pathogenesis of wound and foreign body infections due to staphylococci. *Zentralbl. Bakteriol. Parasitenkd. Infektionskr. Hyg. Abt. 1 Orig. Reihe A* 266:191-211.
- Watts, J. L. 1988. Etiological agents of bovine mastitis. *Vet. Microbiol.* 16:41-66.
- Watts, J. L., and S. C. Nickerson. 1986. A comparison of STAPH-Ident and STAPH-Trac systems to conventional methods in the identification of staphylococci isolated from bovine udders. *Vet. Microbiol.* 12:179-187.
- Watts, J. L., and W. E. Owens. 1987. Synergistic hemolysis associated with coagulase-negative staphylococci isolated from bovine mammary glands. *J. Clin. Microbiol.* 25:2037-2039.