

# Costs to Medicare of the Informatics for Diabetes Education and Telemedicine (IDEATel) Home Telemedicine Demonstration

Findings from an independent evaluation

LORENZO MORENO, PHD<sup>1</sup>  
STACY B. DALE, MPA<sup>1</sup>

ARNOLD Y. CHEN, MD<sup>1</sup>  
CAROL A. MAGEE, PHD<sup>2</sup>

**OBJECTIVE** — To estimate the impacts on Medicare costs of providing a particular type of home telemedicine to eligible Medicare beneficiaries with type 2 diabetes.

**RESEARCH DESIGN AND METHODS** — Two cohorts of beneficiaries ( $n = 1,665$  and  $504$ , respectively) living in two medically underserved areas of New York between 2000 and 2007 were randomized to intensive nurse case management via televisits or usual care. Medicare service use and costs covering a 6-year follow-up period were drawn from claims data. Impacts were estimated using regression analyses.

**RESULTS** — Informatics for Diabetes Education and Telemedicine (IDEATel) did not reduce Medicare costs in either site. Total costs were between 71 and 116% higher for the treatment group than for the control group.

**CONCLUSIONS** — Although IDEATel had modest effects on clinical outcomes (reported elsewhere), it did not reduce Medicare use or costs for health services. The intervention's costs were excessive (over \$8,000 per person per year) compared with programs with similar-sized clinical impacts.

*Diabetes Care* 32:1202–1204, 2009

Home telemedicine delivers monitoring, educational, and therapeutic services to people through telecommunications technology. It may be a promising way to deliver such services to individuals with poor access to high-quality care due to language, culture, low educational attainment, disempowerment, and lack of social reinforcement for healthy behaviors.

The congressionally mandated Informatics for Diabetes Education and Telemedicine (IDEATel) tested the clinical outcomes of providing a particular type of home telemedicine to Medicare beneficiaries with type 2 diabetes. A consortium

led by Columbia University implemented IDEATel in two 4-year phases (February 2000 to February 2008) (1).

## RESEARCH DESIGN AND METHODS

### Goals

For phase I, participants aimed to control blood glucose, high blood pressure, and abnormal lipid levels and reduce or eliminate obesity and physical inactivity. Physicians aimed to increase guideline-based diabetes care. For phase II, the consortium addressed phase I lessons learned and pursued the original goals.

### Recruitment

Between December 2000 and October 2002, the Consortium recruited and randomized 1,665 cohort 1 Medicare beneficiaries (775 in New York City; 890 in upstate NY). Subsequently, between December 2004 and October 2005, the Consortium recruited and randomized 504 cohort 2 beneficiaries (174 in New York City; 330 in upstate NY). For both cohorts, eligibility was limited to beneficiaries aged  $\geq 55$  years who were being treated for diabetes by diet, oral medications, or insulin; living in a medically underserved or health professional shortage area in New York state; and English or Spanish speaking. Poor-health exclusion criteria also applied. After consenting, beneficiaries underwent an in-person baseline assessment by Consortium staff. The Consortium randomly assigned beneficiaries in both cohorts, in equal proportions, to a treatment or control group.

### The intervention

After randomization, treatment and control group members continued receiving diabetes care from their primary care physicians, but treatment group members' physicians received recommendations from the IDEATel diabetologists concerning the care of participants. Treatment group members were offered installation of a home telemedicine unit (HTU) and training in its use.

For phase I, the HTU was a desktop-model PC, connected to a regular telephone line, with a monitor, keyboard, and mouse; video camera; speakers; microphone; and glucose and blood pressure meters. Participants could use the HTU components to 1) measure and monitor blood glucose and blood pressure readings; 2) interact with an IDEATel nurse case manager, in English or Spanish, through scheduled two-way video conferences; and 3) access web-based educational materials. For phase II, the Con-

From <sup>1</sup>Mathematica Policy Research, Inc., Princeton, New Jersey; and the <sup>2</sup>Centers for Medicare & Medicaid Services, Baltimore, Maryland.

Corresponding author: Lorenzo Moreno, lmoreno@mathematica-mpr.com.

Received 18 January 2009 and accepted 30 March 2009.

Published ahead of print at <http://care.diabetesjournals.org> on 14 April 2009. DOI: 10.2337/dc09-0094.

© 2009 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

**Table 1—Estimated annual per-person expenditures for Medicare-covered services, intervention-related costs, and costs for total services, by site, evaluation group, and cohort**

	New York City			Upstate New York		
	Treatment group	Control group	Difference (P)	Treatment group	Control group	Difference (P)
Cohort 1 (both phases)						
Total expenditures for						
Medicare-covered services	13,845	12,961	884 (0.476)	9,566	8,450	1,116 (0.094)
Medicare Part A	8,446	7,502	945 (0.344)	5,136	4,539	597 (0.247)
Medicare Part B	5,399	5,459	−59 (0.870)	4,430	3,911	519 (0.025)
Total intervention-related costs*	8,662	0	NA	8,662	0	NA
Total costs	22,507	12,961	9,546 (0.001)	18,228	8,450	9,778 (0.000)
Cohort 2 (only phase II)						
Total expenditures for						
Medicare-covered services	11,906	11,661	245 (0.931)	6,450	8,694	−2,244 (0.132)
Medicare Part A	7,296	6,886	410 (0.867)	2,991	4,957	−1,966 (0.118)
Medicare Part B	4,610	4,775	−165 (0.799)	3,458	3,736	−278 (0.443)
Total intervention-related costs	8,437	0	NA	8,437	0	NA
Total costs	20,343	11,661	8,682 (0.000)	14,877	8,694	6,183 (0.000)
Sample sizes						
Cohort 1	379	358	—	446	442	—
Cohort 2	82	84	—	161	164	—

Data are means, in dollars. Source: IDEATel tracking status file and 1999–2006 Medicare claims and enrollment records. Means were predicted with ordinary least-squares regression models, which controlled for enrollees' baseline characteristics and the pre-enrollment value of the outcome measure. Estimates reflect annualized expenditures for the period from each sample member's randomization through the end of the study follow-up period (31 December 2006). Observations are weighted by the fraction of the follow-up period that the enrollee was alive and not in an HMO; expenditures are excluded during months the enrollee was in an HMO or not alive. Three control group members were dropped from the analysis in New York City because they are missing control variables used in the regression analysis. See the online appendix for details on the data sources for and construction of the intervention-related cost estimates. \*Total intervention-related costs for cohort 1 are based on the arithmetic average of demonstration costs for phase I and phase II, weighted by the average length of time that phase I participants were enrolled during each phase (see online appendix). NA, not applicable.

sortium redesigned the HTU to address its large size and difficulty of use.

### Hypotheses

Nurse education and coaching through televisits and self-tracking of progress through other HTU functions could have improved participants' self-care behaviors, adhering to diet, exercise, foot care, and medication regimes. IDEATel's guideline-based recommendations to physicians could have promoted better prescribing patterns. These improvements could help participants avoid long-term health complications that could reduce use of acute care services, primarily hospitalizations, and Medicare costs.

### Outcomes

Use of Medicare-covered services and Medicare costs by type of service, total Medicare costs for health care services, and total Medicare costs for both health care services and the intervention.

### Data

The Consortium extracted Medicare claims without identifying information. Follow-up data were available for up to 6

years, from randomization through December 2006. An intention-to-treat analysis included 1,625 cohort 1 and 491 cohort 2 enrollees with complete data.

### Impact estimation

Site-specific impacts were estimated with linear regression (ordinary least-squares) models that controlled for baseline socio-demographic characteristics, experience with computers, diabetes control, and a measure of the outcome in question. Outcomes were annualized and weighted by their months of enrollment in fee-for-service Medicare because no claims data exist for HMO enrollees. The reported treatment and control group means were predicted from the coefficients of the estimated models.

### Demonstration costs

The budget for the demonstration's first and second phases was \$28,159,066 and \$28,812,419, respectively (2). Estimates of the intervention's costs are summarized in Table A4 (available in an online-only appendix at <http://care.diabetesjournals.org/cgi/content/full/dc09-0094/DC1>).

**RESULTS** — In both sites, and for both cohorts, treatment and control group members were similar, on average, on all baseline characteristics, as expected under random assignment (online appendix Tables A1 and A2). However, enrollees' characteristics varied by site and cohort.

Only for cohort 1 were mean annual total Medicare Part B expenditures significantly higher (13% of the control group mean;  $P = 0.025$ ) for treatment group members than for control group members in upstate New York (Table 1).

Total intervention costs were \$8,924 and \$8,437 per person per year for phases I and II, respectively. The costs during phase II were lower than during phase I because the costs were spread over a longer period.

The savings in total Medicare expenditures in any site or cohort were either nonexistent or too small to offset the high costs of the intervention. Total per-person costs were between \$9,500 and \$9,800 higher for treatment group than for control group members for cohort 1 and \$6,200 to \$8,700 for cohort 2 ( $P < 0.001$ ).

The study is limited because it cannot

definitively attribute impacts to telemedicine, case management, or both. Several sensitivity tests were conducted, with no change in findings.

**CONCLUSIONS**— IDEATel's intervention-related costs were excessive because of the size of the budget allocated to its operations and the costly HTUs. The demonstration's costs were higher than the costs of comparable telemedicine programs (\$415 to \$1,830 per participant per year) that served people with diabetes, used televisits with nurse case managers and in-home visits, and had the "potential to effect costs savings" (3,4).

Given the absence of effects on service use (2), finding no effects on Medicare costs was not surprising. The higher Medicare expenditures for the treatment group may have been due to chance or because IDEATel identified the need for some health services among medically underserved beneficiaries.

For IDEATel to be cost-effective, the intervention-related costs would have to be drastically reduced, while maintaining clinical impacts. Less expensive telephonic interventions (5,6) and diabetes case-management programs have yielded comparable improvements in beneficiaries' clinical outcomes to IDEATel's impacts (7). Even if intervention costs were

halved and the program reduced hospitalizations by 50%, both unlikely scenarios, the program would still increase costs to the government.

**Acknowledgments**— This research was funded under contract between the Centers for Medicare & Medicaid Services and Mathematica Policy Research (MPR) (contract no. HHSM-500-2004-00022C). There were no industry sponsors of this study. The authors are either salaried employees of Mathematica Policy Research (L.M., S.B.D., A.Y.C.) or of the Centers for Medicare & Medicaid Services (C.A.M.) and received no compensation from any other source. They do not own stock in the program being evaluated or stand to profit in any way, directly or indirectly, from particular findings in this article.

No potential conflicts of interest relevant to this article were reported.

Randall Brown from MPR provided useful suggestions and Jeffrey Holt, Matthew Jacobus, and Amy Zambrowski, also from MPR, provided excellent computer support. Two anonymous reviewers gave insightful observations.

#### References

1. Shea S, Weinstock RS, Starren J, Teresi J, Palmas W, Field L, Morin P, Goland R, Izquierdo RE, Wolff LT, Ashraf M, Hilliman C, Silver S, Meyer S, Holmes D, Petkova E, Capps L, Lantigua RA. A randomized trial comparing telemedicine case management with usual care in older, ethnically diverse medically underserved patients with diabetes mellitus. *J Am Med Informat Assoc* 2006;13:40–51
2. Moreno L, Shapiro R, Dale SB, Foster L, Chen A. *Final Report to Congress on the Informatics for Diabetes Education and Telemedicine (IDEATel) Demonstration, Phases I and II*. Princeton, NJ, Mathematica Policy Research, Inc., 2008
3. Dansky K, Palmer L, Shea D, Bowles KH. Cost analysis of Telehomecare. *Telem J E-Health* 2001;7:225–232
4. Johnston B, Wheeler L, Dueser J, Sousa K. Outcomes of the Kaiser Permanente Tele-Home Health Research Project. *Arch Fam Med* 2000;9:40–45
5. Piette JD, Weinberger M, McPhee SJ, Mah CA, Kraemer FB, Crapo LM. Can automated calls with nurse follow-up improve self-care and glycemic control among vulnerable patients with diabetes? A randomized controlled trial. *Am J Med* 2000;108:20–17
6. Piette JD, Weinberger M, Kraemer FB, McFee SJ. Impact of automated calls with nurse follow-up on diabetes treatment outcomes in a Department of Veterans Affairs health care system: a randomized controlled trial. *Diabetes Care* 2001;24:202–208
7. Gilmer TP, Roze S, Valentine WJ, Emy-Albrecht K, Ray JA, Cobden D, Nicklasson L, Philis-Tsimikas A, Palmer AJ. Cost-effectiveness of diabetes case management for low-income populations. *Health Serv Res* 2007;42:1943–1959