



Published in final edited form as:

Physiol Behav. 2009 May 25; 97(2): 143–145. doi:10.1016/j.physbeh.2008.12.002.

Introduction: The End of Sex As We Once Knew It

Bruce S. McEwen, Ph.D.

Laboratory of Neuroendocrinology, The Rockefeller University, 1230 York Avenue, New York, NY 10065

Abstract

This special issue of *Physiology and Behavior* is devoted to papers summarizing sex differences that are now recognized to include not only reproductive behaviors, but also non-reproductive processes and phenomena such as feeding, thirst, pain, sensory processes, mood, cognitive function, the effects of stress, and the propensity for drug abuse. The purpose of this brief introduction is to trace some of the main themes and historical highlights in the discovery that the entire nervous system appears to be a target of reproductive hormones, as well as being subject to developmentally programmed sex differences. A semi-popular article with a similar title provides a complementary perspective** to the following thoughts [1].

Keywords

sex differences; gonadal hormones; brain; sexual differentiation; genomic; non-genomic

The study of sex hormone action and sex differences in the brain began with investigating the effects of testicular secretions on reproductive behavior and function [2]. Yet, until the 1960's, there was doubt that sex hormones even entered the brain [3]. However, the use of discrete hormone implants into the hypothalamus demonstrated that estrogens and androgens activate reproductive behaviors [4,5]. Concurrently, the use of tritiated steroid hormones and the technique of steroid autoradiography revealed the presence of binding sites for these steroids that resemble the cell nuclear receptors that were demonstrated by tritiated steroid binding in uterus and other reproductive organs (see [6]).

Sex differences in reproductive function were first uncovered by manipulations of gonadal hormones in rodents immediately after birth [7,8]. Subsequent studies revealed morphological sex differences in hypothalamus [9,10] and also in spinal cord [11]. Subsequent work extended these findings to the human hypothalamus [12-14]. All of this important knowledge has now

**Quote from End of Sex As We Know It: by B.S.McEwen and E.N.Lasley (2005)

Contact information: Bruce S. McEwen, Ph.D., Alfred E. Mirsky Professor, The Rockefeller University, Box 165, 1230 York Avenue, New York, NY 10065, Tel: 212 327 8624, FAX: 212 327 8634, Email: E-mail: mcewen@rockefeller.edu.

“Right now, the research is raising as many questions as it is answering, at least when it comes to definite conclusions about human behavior and many brain functions. It goes without saying, or should, that the existence of important sex-based brain differences does not mean in any way, shape, or form, that a stronger or more intelligent sex exists. Several animal studies, however, are clarifying the specific differences in males and females and how chemistry, genetics, and the environment play their parts in creating divergent results depending on sex. Scientists suspect that the observable differences in performance of men and women in various areas are produced by some combination of genetic and environmental influences that bring about the distinct, sometimes dramatic, divergence in pathways of brain and behavior that is emerging from new research.”

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

been supplemented and enriched by findings outside of the hypothalamus. Indeed, the notion that the hypothalamus and sexually dimorphic nucleus of the spinal cord are not the only areas of the CNS where sex hormones act and where sex differences exist was initially suggested by findings in Parkinson's Disease, the neuroanatomy of the corpus callosum and estrous cyclicity of seizure sensitivity in the hippocampus.

The Parkinson's connection arose with the observation that high dose estrogen treatment used in the initial contraceptive preparations exacerbated symptoms of Parkinson's Disease in women [15]. This was very unexpected for those who believed in the nuclear estrogen receptor story because there are no cell nuclear estrogen receptors in the rodent striatum, and yet tiny unilateral implants of estradiol in the rodent striatum elicited unilateral rotation associated with imbalanced dopaminergic function [16]. Furthermore, estradiol regulates dopamine release from striatum in a sexually-dimorphic manner [17].

The corpus callosum and anterior commissure emerged as brain regions that differ in shape between men and women [18,19] and which, therefore, may be influenced by hormone in early development. More recent studies also showed morphological sex differences in the cellular organization of the human temporal lobe [20,21].

Studies on the hippocampus revealed large differences across the rodent estrous cycle in the seizure threshold - with proestrus females showing greater sensitivity (lower thresholds) than diestrus females [22]. This was followed by the finding of estrogen-induced spine synapse induction in hippocampus [23,24], which also was surprising because, like the striatum, there were very few nuclear estrogen receptors in the hippocampus [25,26]. The hippocampus has also turned out to be a target of sexual differentiation, with sex differences in spatial memory acquisition [27] and in sex hormone activation of synaptogenesis [28] that are reversed by the presence or absence of testosterone in neonates [29]. But the hippocampus has very few cell nuclear estrogen receptors except in inhibitory interneurons [30] and this finding led to further studies of estrogen's mechanism of action.

Indeed, a critical advance in the puzzle of gonadal hormone actions where there are no detectable cell nuclear receptors was the finding, using electron microscopic immunocytochemistry that estrogen receptors occur outside the nucleus in dendrites, presynaptic terminals and astroglial processes [31]. These receptors, which have recently been shown to bind radioactive estradiol [32], appear to signal rapid non-genomic actions via phosphorylation of second messenger systems such as PI3 kinase and Akt, LIM kinase1, MAP kinases, trk B neurotrophin receptors and CREB [31,33-39]. Non-genomic estrogen receptors were also detected in brain regions, such as corpus striatum and cerebellum [31,40]. Moreover, a growing body of evidence indicates the presence of receptors for testosterone, progesterone and glucocorticoids in non-genomic sites [41-43]. Thus, non-genomic signalling pathways are likely to be involved in the actions of these steroid hormones on the nervous system.

A related aspect of steroid hormone actions on structural plasticity in the brain is their dependence on concurrent excitatory amino acid neurotransmission, as revealed by studies showing effects of estrogens on synapse formation [44] and of glucocorticoids on dendritic remodeling [45,46]. Excitatory amino acid neurotransmission is vital to the ability of the brain to encode memories of experiences, and experience is also capable of changing the brain both structurally and functionally [47]. Because of their actions on the brain, as described above, hormones, through their synergistic actions with neurotransmitters, are able to bias the nervous system towards adopting certain strategies for learning or other behavioral responses (e.g., see [48]). Likewise, the process of sexual differentiation interacts with experience to produce sexually dimorphic patterns of behavior [for a masterful discussion of this topic, see [49].

Thus, recent advances in the study of sex hormone action and sex differences are challenging long-held dogma that only the reproductive aspects of brain function are affected by sexual differentiation and by sex hormones. Rather, we now are beginning to understand that the entire brain is, to varying degrees, a target of gonadal hormones and a potential site of developmentally-programmed sex differences. This is made even more interesting and complex by the realization that there are also sex differences that are chromosomally linked [50,52]. The future prospects for this line of research are both challenging and highly relevant to many aspects of neuroscience, behavioral biology and medicine [53,54].

References

1. McEwen, BS.; Lasley, EN. Cerebrum. The Dana Forum on Brain Science. Vol. 7. Dana Press; 2005. The End of Sex as We Know It.
2. Berthold AA. Transplantation der Hoden. Arch Anat Physiol Wiss Med 1849;16:42–60.
3. Young, W. The Hormones and Mating Behavior. In: Young, W., editor. Sex and Internal Secretions. Vol. 7. Baltimore: Williams and Wilkins; 1961. p. 1173-239.
4. Lisk R. Diencephalic placement of estradiol and sexual receptivity in female rat. Am J Physiol 1962;203:493–6. [PubMed: 14465798]
5. Barfield R. Activation of copulatory behavior by androgen implanted into the preoptic area of the male fowl. Horm Behav 1969;1:37–52.
6. Pfaff, DW. Estrogens and Brain Function. N.Y.: Springer-Verlag; 1980.
7. Harris GW, Levine S. Sexual differentiation of the brain and its experimental control. J Physiol 1965;181:379–400. [PubMed: 5893725]
8. Goy, R.; McEwen, BS. Sexual Differentiation of the Brain. Cambridge: MIT Press; 1980. Notes: Pages: -223
9. Raisman G, Field P. Sexual dimorphism in the preoptic area of the rat. Science 1971;173:731–3. [PubMed: 5568506]
10. Gorski RA, Gordon JH, Shryne JE, Southam AM. Evidence for a morphological sex differences within the medial preoptic area of the rat brain. Brain Res 1978;148:333–46. [PubMed: 656937]
11. Breedlove SM, Arnold AP. Sexually dimorphic motor nucleus in the rat lumbar spinal cord: Response to adult hormone manipulation, absence in androgen-insensitive rats. Brain Res 1981;225:297–307. [PubMed: 7306791]
12. Allen L, Hines M, Shryne J, Gorski R. Two sexually dimorphic cell groups in the human brain. J Neurosci 1989;9:497–506. [PubMed: 2918374]
13. Swaab DF, Gooren LJJ, Hofman MA. The human hypothalamus in relation to gender and sexual orientation. Progress in Brain Research 1992;93:205–19. [PubMed: 1480750]
14. LeVay S. A difference in hypothalamic structure between heterosexual and homosexual men. Science 1991;253:1036–8.
15. Bedard PJ, Langelier P, Villeneuve A. Estrogens and the extrapyramidal system. The Lancet 1977;2:1367–8.
16. Van Hartesveldt C, Joyce J. Effects of estrogen on the basal ganglia. Neurosci Biobehav Rev 1986;10:1–14. Notes: User Def 1: social environment User Def 2: X5X
17. Becker JB. Direct effect of 17 β -estradiol on striatum: Sex differences in dopamine release. Synapse 1990;5:157–64. [PubMed: 2309159]
18. Witelson S. Hand and sex differences in the isthmus and genu of the human corpus callosum. Brain 1989;112:799–835. [PubMed: 2731030]
19. Allen L, Gorski R. Sexual dimorphism of the anterior commissure and massa intermedia of the human brain. Journal Comp Neurol 1991;312:97–104.
20. Witelson SF, Glezer II, Kigar DL. Women have greater density of neurons in posterior temporal cortex. J Neurosci 1995;15(5 Pt 1):3418–28. [PubMed: 7751921]
21. Alonso-Nanclares L, Gonzalez-Soriano JG, Rodriguez JR, DeFelipe J. Gender differences in human cortical synaptic density. Proc Natl Acad Sci USA 2008;105:14615–9. [PubMed: 18779570]

22. Terasawa E, Timiras P. Electrical activity during the estrous cycle of the rat: cyclic changes in limbic structures. *Endocrinology* 1968;83:207–16. [PubMed: 4874282]
23. Gould E, Woolley C, Frankfurt M, McEwen BS. Gonadal steroids regulate dendritic spine density in hippocampal pyramidal cells in adulthood. *J Neurosci* 1990;10:1286–91. [PubMed: 2329377]
24. Woolley C, McEwen BS. Estradiol mediates fluctuation in hippocampal synapse density during the estrous cycle in the adult rat. *Journal Neuroscience* 1992;12:2549–54.
25. Pfaff DW, Keiner M. Atlas of estradiol-concentrating cells in the central nervous system of the female rat. *J Comp Neurol* 1973;151:121–58. [PubMed: 4744471]
26. Loy R, Gerlach J, McEwen BS. Autoradiographic localization of estradiol-binding neurons in rat hippocampal formation and entorhinal cortex. *Dev Brain Res* 1988;39:245–51.
27. Williams CL, Meck WH. The organizational effects of gonadal steroids on sexually dimorphic spatial ability. *Psychoneuroendocrinology* 1991;16(13):155–76. [PubMed: 1961837]
28. Leranath C, Hajszan T, MacLusky NJ. Androgens increase spine synapse density in the CA1 hippocampal subfield of ovariectomized female rats. *J Neurosci* 2004;24:495–9. [PubMed: 14724248]
29. Lewis C, McEwen BS, Frankfurt M. Estrogen-induction of dendritic spines in ventromedial hypothalamus and hippocampus: effects of neonatal aromatase blockade and adult castration. *Devel Brain Res* 1995;87:91–5. [PubMed: 7554237]
30. McEwen BS, Alves SH. Estrogen Actions in the Central Nervous System. *Endocrine Rev* 1999;20:279–307. [PubMed: 10368772]
31. McEwen BS, Milner TA. Hippocampal formation: Shedding light on the influence of sex and stress on the brain. *Brain Res Rev* 2007;55:343–55. [PubMed: 17395265]
32. Milner TA, Lubbers LS, Alves SE, McEwen BS. Nuclear and extranuclear estrogen binding sites in the rat forebrain and autonomic medullary areas. *Endocrinology* 2008;149:3306–12. [PubMed: 18356271]
33. Akama KT, McEwen BS. Estrogen stimulates postsynaptic density-95 rapid protein synthesis via the Akt/protein kinase B pathway. *J Neurosci* 2003;23:2333–9. [PubMed: 12657692]
34. Dominguez R, Liu R, Baudry M. 17- β -Estradiol-mediated activation of extracellular-signal regulated kinase, phosphatidylinositol 3-kinase/protein kinase B-Akt and N-methyl-D-aspartate receptor phosphorylation in cortical synaptoneurosome. *J Neurochem* 2007;101:232–40. [PubMed: 17250656]
35. Kelly MJ, Levin ER. Rapid actions of plasma membrane estrogen receptors. *Trends Endo & Metab* 2001;12:152–6.
36. Znamensky V, Akama KT, McEwen BS, Milner TA. Estrogen levels regulate the subcellular distribution of phosphorylated Akt in hippocampal CA1 dendrites. *J Neurosci* 2003;23:2340–7. [PubMed: 12657693]
37. Lee SJ, Campomanes CR, Sikat PT, Greenfield AT, Allen PB, McEwen BS. Estrogen induces phosphorylation of cyclic AMP response element binding (pCREB) in primary hippocampal cells in a time-dependent manner. *Neuroscience* 2004;124:549–60. [PubMed: 14980726]
38. Spencer JL, Waters EM, Milner TA, McEwen BS. Estrous cycle regulates activation of hippocampal Akt, LIM kinase, and neurotrophin receptors in C57BL/6 mice. *Neuroscience* 2008;155:1106–19. [PubMed: 18601981]
39. Yildirim M, Janssen WGM, Tabori NE, et al. Estrogen and aging affect synaptic distribution of phosphorylated LIM kinase (pLIMK) in CA1 region of female rat hippocampus. *Neuroscience* 2008;152:360–70. [PubMed: 18294775]
40. Milner, TA.; Alves, SE.; Hayashi, S.; McEwen, BS. The Identities of Membrane Steroid Receptors. Vol. Chapter 3. Boston, MA: Kluwer Academic Publishers; 2003. An expanded view of estrogen receptor localization in neurons; p. 21-5.
41. Tabori NE, Stewart LS, Znamensky V, et al. Ultrastructural evidence that androgen receptors are located at extranuclear sites in the rat hippocampal formation. *Neuroscience* 2005;130:151–63. [PubMed: 15561432]
42. Johnson LR, Farb C, Morrison JH, McEwen BS, LeDoux JE. Localization of glucocorticoid receptors at postsynaptic membranes in the lateral amygdala. *Neuroscience* 2005;136:289–99. [PubMed: 16181741]

43. Waters EM, Torres-Revon AT, McEwen BS, Milner TA. Extranuclear progesterone receptor immunoreactivity in the rat hippocampal formation. *J Comp Neurol* 2008;511:34–46. [PubMed: 18720413]
44. Woolley C, McEwen BS. Estradiol regulates hippocampal dendritic spine density via an N-methyl-D-aspartate receptor dependent mechanism. *Journal Neuroscience* 1994;14:7680–7.
45. Magarinos AM, McEwen BS. Stress-Induced Atrophy of Apical Dendrites of Hippocampal CA3c Neurons: Comparison of Stressors. *Neuroscience* 1995;69(1):83–8. [PubMed: 8637635]
46. Magarinos AM, McEwen BS. Stress-induced atrophy of apical dendrites of hippocampal CA3c neurons: involvement of glucocorticoid secretion and excitatory amino acid receptors. *Neuroscience* 1995;69(1):89–98. [PubMed: 8637636]
47. Bennett E, Diamond M, Krech D, Rosenzweig M. Chemical and anatomical plasticity of brain. *Science* 1964;146:610–9. [PubMed: 14191699]
48. Levy LJ, Astur RS, Frick KM. Men and women differ in object memory but not performance of a virtual radial maze. *Behav Neurosci* 2005;119:853–62. [PubMed: 16187814]
49. Goy, RW. Early hormonal influences on the development of sexual and sex-related behavior. In: Schmitt, FO., editor. *The Neurosciences: Second Study Program*. New York: Rockefeller University Press; 1970. p. 196-206.
50. Reisert I, Pilgrim C. Sexual differentiation of monoaminergic neurons -- genetic or epigenetic? *TINS* 1996;14(10):468–73. [PubMed: 1722367]
51. Arnold AP. Genetically triggered sexual differentiation of brain and behavior. *Horm Behav* 1996;3:495–505. [PubMed: 9047274]
52. Carruth LL, Reisert I, Arnold AP. Sex chromosome genes directly affect brain sexual differentiation. *Nature Neurosci* 2002;5:933–4. [PubMed: 12244322]
53. NRC Report. *Exploring the Biological Contributions to Human Health: Does Sex Matter*. Natl Acad Sci 2001:1–8.
54. Becker, JB.; Berkley, KJ.; Geary, N.; Hampson, E.; Herman, JP.; Young, E. *Sex Differences in the Brain: From Genes to Behavior*. Oxford University Press; 2007.