



Published in final edited form as:

*Bioorg Med Chem Lett.* 2009 June 1; 19(11): 3023–3026. doi:10.1016/j.bmcl.2009.04.058.

## Isoxazole Moiety in the Linker Region of HDAC Inhibitors Adjacent to the Zn-chelating Group: Effects on HDAC Biology and Antiproliferative Activity

Subhasish Tapadar<sup>a</sup>, Rong He<sup>a</sup>, Doris N. Savoy<sup>b</sup>, Daniel D. Billadeau<sup>b</sup>, and Alan P. Kozikowski<sup>a</sup>

<sup>a</sup> Department of Medicinal Chemistry & Pharmacognosy, University of Illinois at Chicago, 833 S. Wood St. Chicago, IL 60612, USA

<sup>b</sup> Division of Oncology Research, College of Medicine, Mayo Clinic, Rochester, MN 55905, USA

### Abstract

A series of hydroxamic acid based histone deacetylase inhibitors **6–15**, containing an isoxazole moiety adjacent to the Zn-chelating hydroxamic acid, is reported herein. Some of these compounds showed nanomolar activity in the HDAC isoform inhibitory assay and exhibited micro molar inhibitory activity against five pancreatic cancer cell lines.

Epigenetic regulation of gene expression is partly controlled by chromatin remodeling which is initiated either by post-translational modification of histone proteins<sup>1,2</sup> or DNA methylation.<sup>3</sup> Several different modifications that occur in histone amino termini are defined as acetylation, methylation, ubiquitinylation, and glycosylation. Histone acetylation-deacetylation is a reversible phenomenon and is tightly regulated by two competing enzymes known as histone acetyltransferases (HATs) and histone deacetylases (HDACs).<sup>4</sup> In general, HATs catalyze the acetylation of *N*-terminal lysine  $\epsilon$ -amino groups in nuclear histones, resulting in neutralization of the positive charges on the histones and a more open, transcriptionally active chromatin structure while the HDACs catalyze the reverse reaction and suppress transcription. Altered HDAC activity is seen to be associated with many cancers, validating HDACs as promising targets for cancer therapy. Several small molecule HDAC inhibitors<sup>5–7</sup> (Figure 1) such as suberoylanilide hydroxamic acid (SAHA), depsipeptide FK228, MS-275, and valproic acid have been found to have promising antitumor activities in clinical studies and in 2006, the US FDA approved SAHA for the treatment of a rare cancer, cutaneous T-cell lymphoma (CTCL).<sup>8</sup> These compounds owe their antiproliferative action to their ability to allow the transcription and expression of repressed genes including tumor suppressor genes. Beside their antitumor activity, HDAC inhibitors are also found to have potential applications in the treatment of neurodegenerative disorders such as Parkinson's and Huntington's diseases, as well as malaria.<sup>9</sup> Human HDACs are divided into four classes based on their function and structural homology. Class I (HADC1, 2, 3, and 8), class II (HDAC4, 5, 6, 7, 9 and 10), and class IV (HDAC11) HDACs are zinc-dependent enzymes while class III HDACs are sirtuin related proteins and require the cofactor NAD<sup>+</sup> for their deacetylase function.<sup>7</sup> To learn more about the role the individual HDACs play in cell growth and differentiation, neuronal protection, and apoptosis, it is important to develop agents that show selectivity for individual isoforms or a small subset

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

of these isoforms. To date, a large number of structurally diverse, natural and synthetic, hydroxamic acid based HDAC inhibitors (Figure 1) have been reported which include trichostatin A (TSA), SAHA, depsipeptide, and tubacin etc., but the number of isoform selective HDAC inhibitors are very limited.<sup>5,6</sup> Recently, we have reported hydroxamate-based HDAC inhibitors bearing substituted aryl-isoxazoles (Figure 2) as the cap groups. Some of these compounds were found to be highly potent and selective at HDAC6 as well as HDAC3 (compound **1** and **2**; Table 1).<sup>10</sup>

We anticipated that the exceptional potency and selectivity of those compounds were due to the presence of the isoxazole moiety whose mode of interaction with surface residues is still under investigation. In this paper, we report a series of hydroxamic acid-based HDAC inhibitors with an isoxazole moiety in the linker region adjacent to the hydroxamic acid group (Figure 3). These compounds were designed with the idea that the more rigid and bulkier zinc binding group might allow for a different selectivity. The choice of the cap groups was based on the HDAC inhibitors previously explored by us<sup>11,12</sup> (Figure 2; Table 1) and Miyata *et al.*<sup>13</sup> In this context, the hydroxamic acid was suitably replaced by the other functionalities like alcohol, carboxylic acid, and amide to explore their Zn-binding capabilities. All the synthesized compounds were subjected to *in vitro* HDAC inhibition study by using catalytically active recombinant HDACs, and their antiproliferative activities were tested on various pancreatic cancer cell lines as well.

Different heteroaromatic, aromatic and alicyclic amino cap groups, namely 2-amino-4-phenylthiazole (**16**), 4-(3-nitrophenyl)thiazol-2-ylamine (**17**), 3-aminoquinoline (**23**), 4-aminobiphenyl (**24**), cyclopentylamine (**29**), and cycloheptylamine (**30**) were coupled with 6-heptynoic acid under peptide coupling conditions to yield terminal acetylenic compounds **18**, **19**, **25**, **26**, **31**, and **32**, respectively. Subsequently the terminal acetylenic group of these compounds was converted by the use of nitrile oxide cycloaddition chemistry<sup>14</sup> to the substituted-isoxazole carboxylic acid ethyl ester (**20**, **21**, **27**, **28**, **33**, and **34**). Lastly, these ester compounds were reacted with hydroxylamine hydrochloride in the presence of potassium hydroxide in accord with a known literature procedure (Schemes 1 and 2)<sup>15</sup> to provide hydroxamic acids **6**, **7**, and **11** – **14**. In case of the synthesis of compound **7**, the intermediate **21** bearing a nitro cap group was reduced to the aniline derivative **22** by catalytic hydrogenation (Scheme 1), followed by hydroxamic acid formation. Sodium borohydride reduction of compound **20** led to alcohol **8**. The carboxylic acid **9** was synthesized by saponification (Scheme 1) of the corresponding ester **20**. The amide **10** was obtained from **9** by an active ester method (Scheme 1). A different strategy was applied to synthesize compound **15** as shown in Scheme 3. Sequential saponification and amide coupling<sup>10</sup> of **35** gave the intermediate **38** via the ester **36**. The ester **38** was converted to hydroxamic acid **15** following the same general method used to prepare previous hydroxamic acid compounds.

All the compounds **6**–**15** were tested against both the class I (1, 2, and 3) and class II (6 and 10) HDACs and their IC<sub>50</sub> values are shown in Table 2. Compound **6**, **7**, and **11**, with a phenylthiazole, an *m*-aminosubstituted phenylthiazole, and a quinoline moiety as cap groups, were the most active HDAC inhibitors in this series. These compounds **6**, **7**, and **11** were comparatively more active at HDAC3 and HDAC6 than HDAC1, HDAC2, and HDAC10. The hydroxamic acids **12**–**15** are very poor HDAC inhibitors except for compound **12**, which was weakly active at HDAC1 and HDAC6. The inhibitors **6**, **7** and **11** were less potent than those compounds with an isoxazole moiety in the cap region<sup>10</sup> (Table 1) and also SAHA. Placement of the isoxazole group in the linker region decreased the inhibitory activity of compounds **6** and **7**, when compared to those hydroxamic acids **3** and **4** bearing the same cap groups and a linear linker (Figure 2). At the same time, these structural alterations made **6** and **7** relatively more selective as HDAC3 and HDAC6 inhibitors, respectively (**6** vs **3**, and **7** vs **4**; Table 1 and Table 2). Attempted replacement of the hydroxamate group by the other functional groups such

as alcohol, carboxylic acid, and amide, failed to provide any significant HDAC inhibitory activity (Compound **8**, **9**, and **10**). This is not surprising in light of other publications.<sup>16,17</sup>

We also investigated the ability of this set of compounds **6–15** to inhibit cancer cell growth using different pancreatic cancer cell lines. Pancreatic cancer is the fourth leading cause of cancer death in the United States, and essentially remains an incurable disease with a five-year survival rate of < 5%.<sup>18</sup> This cancer fails to respond to widely-used chemotherapeutic drugs like 5-fluorouracil and gemcitabine,<sup>19</sup> and thus there is an urgent need for more efficient drugs. All HDAC inhibitors **6–15**, along with the currently marketed drug SAHA were tested on five pancreatic cancer cell lines using the MTT assay. These data are provided in the Table 3. As is apparent from these data, compound **6** nicely inhibited all the five transformed cell lines and its IC<sub>50</sub> values were similar to those of SAHA. The relatively weak HDAC inhibitor **12** showed only moderate activity against three of the five transformed cell lines, namely, BxPC-3, HuPT3 and Su.86.86. In contrast, the most potent HDAC inhibitor in this series, compound **7**, only moderately inhibited the five pancreatic cancer cell lines. Compound **11** failed to show similar antiproliferative activity compared to compound **6**, although its IC<sub>50</sub> values in the *in vitro* HDAC inhibitory assay are very similar to those of **6**. The weak inhibitory activities of compound **7** and **11** against different pancreatic cancer cell lines might be due to their poor cell permeability. Compounds **8–10** and **13–15** exhibited poor antiproliferative activities against all five pancreatic cancer cell lines, which are consistent with their poor HDAC inhibitory activities.

In summary, we synthesized a series of hydroxamic acid based HDAC inhibitors wherein the isoxazole moiety was embedded in the linker region and directly attached to the Zn-chelating hydroxamic acid group. The present findings suggest that no significant isozyme selectivity is gained using the more rigid isoxazole hydroxamate as zinc binding group.


## Acknowledgments

This work was supported in part by gift funds from an anonymous donor, by a grant (no. 271210) from the ADDF/Elan, and by the Mayo Foundation and the Pancreatic Cancer SPORE P50 CA102701.


## References and notes

1. Grunstein M. *Nature* 1997;389:349. [PubMed: 9311776]
2. Wolffe AP, Guschin D. *J Struct Biol* 2000;129:102. [PubMed: 10806063]
3. Abreu PA, Dellamora-Ortiz G, Leao-Ferreira LR, Gouveia M, Braggio E, Zalcborg I, Santos DO, Bourguinhon S, Cabral LM, Rodrigues CR, Castro HC. *Expert Opin Ther Targets* 2008;12:1035. [PubMed: 18620524]
4. Marks PA, Richon VM, Rifkind RA. *J Natl Cancer Inst* 2000;92:1210. [PubMed: 10922406]
5. Butler KV, Kozikowski AP. *Curr Pharm Des* 2008;14:505. [PubMed: 18336297]
6. Paris M, Porcelloni M, Binaschi M, Fattori D. *J Med Chem* 2008;51:1505. [PubMed: 18247554]
7. Bolden JE, Peart MJ, Johnstone RW. *Nat Rev Drug Discov* 2006;5:769. [PubMed: 16955068]
8. Mann BS, Johnson JR, Cohen MH, Justice R, Pazdur R. *Oncologist* 2007;12:1247. [PubMed: 17962618]
9. Morrison BE, Majdzadeh N, Zhang X, Lyles A, Bassel-Duby R, Olson EN, D'Mello SR. *Mol Cell Biol* 2006;26:3550. [PubMed: 16611996]
10. Kozikowski AP, Tapadar S, Luchini DN, Kim KH, Billadeau DD. *J Med Chem* 2008;51:4370. [PubMed: 18642892]
11. Kozikowski AP, Chen Y, Gaysin AM, Savoy DN, Billadeau DD, Kim KH. *ChemMedChem* 2008;3:487. [PubMed: 18181121]
12. Chen B, Petukhov PA, Jung M, Velena A, Eliseeva E, Dritschilo A, Kozikowski AP. *Bioorg Med Chem Lett* 2005;15:1389. [PubMed: 15713393]

13. Itoh Y, Suzuki T, Kouketsu A, Suzuki N, Maeda S, Yoshida M, Nakagawa H, Miyata N. *J Med Chem* 2007;50:5425. [PubMed: 17929798]
14. Simoni D, Invidiata FP, Rondanin R, Grimaudo S, Cannizzo G, Barbusca E, Porretto F, D' Alessandro N, Tolomeo M. *J Med Chem* 1999;42:4961. [PubMed: 10585206]
15. Gediya LK, Chopra P, Purushottamachar P, Maheshwari N, Njar VC. *J Med Chem* 2005;48:5047. [PubMed: 16033284]
16. Wahhab A, Smil D, Ajamian A, Allan M, Chantigny Y, Therrien E, Nguyen N, Manku S, Leit S, Rahil J, Petschner AJ, Lu AH, Nicolescu A, Lefebvre S, Montcalm S, Fournel M, Yan TP, Li Z, Besterman JM, Deziel R. *Bioorg Med Chem Lett* 2009;19:336. [PubMed: 19084395]
17. Hanessian S, Vinci V, Auzzas L, Marzi M, Giannini G. *Bioorg Med Chem Lett* 2006;16:4784. [PubMed: 16870438]
18. Li D, Xie K, Wolff R, Abbruzzese JL. *Lancet* 2004;363:1049. [PubMed: 15051286]
19. Haller DG. *Semin Oncol* 2003;30:3. [PubMed: 14506598]




**Figure 1.**  
Structures of some known HDAC inhibitors




**Figure 2.** HDAC inhibitors containing a substituted isoxazole, a substituted thiazole, or a biphenyl group as the cap

CAP: different aromatic,  
heteroaromatic,  
and alicyclic groups

Zn<sup>++</sup>-binding domain




**Figure 3.**  
General structural of HDAC inhibitors 6-15




**Scheme 1.**

Synthesis of compounds **6–10**. Reagents and conditions: (a) 7-Heptynoic acid,  $\text{POCl}_3$ , pyridine,  $-13\text{ }^\circ\text{C}$  to rt, 30 min. (b) Ethyl chlorooxamidoacetate, triethylamine, THF, rt, 16 h. (c)  $\text{H}_2$ , 10 % Pd/C, EtOAc, EtOH,  $60\text{ }^\circ\text{C}$ , 30 min. (d)  $\text{NH}_2\text{OH}\cdot\text{HCl}$ , KOH, MeOH, rt, 15 min. (e)  $\text{NaBH}_4$ , MeOH,  $0\text{ }^\circ\text{C}$  to rt, 2 h. (f) LiOH, THF:MeOH:H<sub>2</sub>O (3:1:1),  $0\text{ }^\circ\text{C}$ , 30 min. (g) Ethyl chloroformate, triethylamine, THF,  $0\text{ }^\circ\text{C}$ , 30 min, then aqueous  $\text{NH}_3$ , rt, 12 h.



**Scheme 2.**

Synthesis of compounds **11–14**. Reagents and conditions: (a) 7-Heptynoic acid,  $\text{POCl}_3$ , pyridine,  $-13\text{ }^\circ\text{C}$  to rt, 30 min. (b) Ethyl chlorooxamidoacetate,  $\text{Et}_3\text{N}$ , THF, rt, 16 h. (c)  $\text{NH}_2\text{OH}\cdot\text{HCl}$ , KOH, MeOH, rt, 15 min. (d) EDCI, HOBT, DIEA, amine, DMF,  $0\text{ }^\circ\text{C}$  to rt, 12 h.

**Scheme 3.**

Synthesis of compound **15**. Reagents and conditions: (a) LiOH, THF:MeOH:H<sub>2</sub>O (3:1:1), 0 °C, 30 min. (b) EDCl, HOBT, DIEA, glycine ethyl ester hydrochloride, DMF, 0 °C to rt, 12 h. (c) LiOH, THF:MeOH:H<sub>2</sub>O (3:1:1), 0 °C, 30 min. (d) TFA, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C to rt, 2 h. (e) **37**, POCl<sub>3</sub>, pyridine, -13 °C to rt, 30 min. (f) NH<sub>2</sub>OH·HCl, KOH, MeOH, rt, 15 min.

**Table 1**  
*In vitro* HDAC Inhibitory Activities of Compounds **1–5**.

Compds	IC <sub>50</sub> (nM)		
	HDAC1	HDAC3	HDAC6
<b>1</b> <sup>a</sup>	100 ± 16	< 0.002	< 0.002
<b>2</b> <sup>a</sup>	271 ± 23	0.42 ± 0.08	0.002
<b>3</b> <sup>b</sup>	3	ND <sup>b</sup>	3
<b>4</b> <sup>b</sup>	4	ND <sup>b</sup>	3
<b>5</b> <sup>b</sup>	33	7	5

<sup>a</sup>Ref. 10.

<sup>b</sup>Ref. 11

**Table 2**  
*In vitro* HDAC inhibitory assay results for compounds 6–15.

Comps	IC <sub>50</sub> (nM)						
	HDAC1	HDAC2	HDAC3	HDAC6	HDAC10	HDAC10	HDAC10
<b>6</b>	303 ± 22	430 ± 37	30 ± 1	68 ± 5	254 ± 25		
<b>7</b>	139 ± 4	164 ± 4	25 ± 1	82 ± 5	250 ± 6		
<b>8</b>	>30,000	>30,000	>30,000	>30,000	>30,000		
<b>9</b>	>30,000	>30,000	10,600 ± 829	>30,000	>30,000		
<b>10</b>	>30,000	>30,000	>30,000	>30,000	>30,000		
<b>11</b>	328 ± 10	469 ± 18	102 ± 3	51 ± 0.9	471 ± 20		
<b>12</b>	885	3,750	7,410	885	ND <sup>a</sup>		
<b>13</b>	17,100	> 30,000	nd <sup>a</sup>	17,100	ND <sup>a</sup>		
<b>14</b>	12,300	> 30,000	> 30,000	12,300	ND <sup>a</sup>		
<b>15</b>	9,200 ± 670	>30,000	1,920 ± 73	1710 ± 82	13,300 ± 560		
<b>SAHA<sup>b</sup></b>	96	282	17	14	72		

<sup>a</sup>ND = Not Detected.

<sup>b</sup>SAHA data provided by Amphora Inc.

**Table 3**  
*In vitro* growth inhibition assay results against pancreatic cancer cell lines for compounds **6–15**.

Comps	IC <sub>50</sub> (μM)					Su.86.86
	BxPC-3	HupT3	Mia Paca-2	Panc 04.03		
<b>6</b>	3	1	4	4		1
<b>7</b>	10.5	28.4	9	14.4		15
<b>8</b>	> 50	> 50	> 50	> 50		> 50
<b>9</b>	> 50	> 50	> 50	> 50		> 50
<b>10</b>	> 50	> 50	> 50	> 50		> 50
<b>11</b>	36	49	24	44		25
<b>12</b>	9	7	30	> 50		9
<b>13</b>	> 50	> 50	24	30		43
<b>14</b>	> 50	> 50	> 50	> 50		> 50
<b>15</b>	> 50	> 50	> 50	> 50		> 50
SAHA	5	0.8	1	1		1