

Research

Open Access

Psychiatric morbidity among adult patients in a semi-urban primary care setting in Malaysia

Ruzanna ZamZam*¹, Maniam Thambu^{†1}, Marhani Midin^{†1},
Khairani Omar^{†2} and Pervesh Kaur^{†2}

Address: ¹Department of Psychiatry, Faculty of Medicine, Universiti Kebangsaan Malaysia, Kuala Lumpur 56000, Malaysia and ²Department of Family Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia, Kuala Lumpur 56000, Malaysia

Email: Ruzanna ZamZam* - ruzanna@ppukm.ukm.my; Maniam Thambu - tmaniam@yahoo.com; Marhani Midin - marhani@ppukm.ukm.my; Khairani Omar - khairani@ppukm.ukm.my; Pervesh Kaur - Pervesh.Kaur@shell.com

* Corresponding author †Equal contributors

Published: 18 June 2009

Received: 17 February 2009

International Journal of Mental Health Systems 2009, **3**:13 doi:10.1186/1752-4458-3-13

Accepted: 18 June 2009

This article is available from: <http://www.ijmhs.com/content/3/1/13>

© 2009 ZamZam et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Screening for psychiatric disorders in primary care can improve the detection rate and helps in preventing grave consequences of unrecognised and untreated psychiatric morbidity. This is relevant to the Malaysian setting where mental health care is now also being provided at primary care level. The aim of this paper is to report the prevalence of psychiatric illness in a semi-urban primary care setting in Malaysia using the screening tool Patient Health Questionnaire (PHQ).

Methods: This is a cross-sectional study carried out in a semi-urban primary healthcare centre located south of Kuala Lumpur. Systematic random sampling was carried out and a total of 267 subjects completed the PHQ during the study period.

Results: The proportion of respondents who had at least one PHQ positive diagnosis was 24.7% and some respondents had more than one diagnosis. Diagnoses included depressive illness (n = 38, 14.4%), somatoform disorder (n = 32, 12.2%), panic and anxiety disorders (n = 17, 6.5%), binge eating disorder (n = 9, 3.4%) and alcohol abuse (n = 6, 2.3%). Younger age (18 to 29 years) and having a history of stressors in the previous four weeks were found to be significantly associated (p = 0.036 and p = 0.044 respectively) with PHQ positive scores.

Conclusion: These findings are broadly similar to the findings of studies done in other countries and are a useful guide to the probable prevalence of psychiatric morbidity in primary care in other similar settings in Malaysia.

Introduction

Most of the psychiatric morbidity in the community is seen at the primary care level [1-5]. Studies using screening instruments have reported prevalence rates ranging from 16 to 43% of general practice attenders [1,6-8]. A study of psychiatric morbidity using the General Health

Questionnaire (GHQ) reported rates up to 30% [1]. In a Malaysian primary health setting the prevalence of psychiatric morbidity using GHQ-30 item version was 26.7% [9]. In the general population, as measured by GHQ-12 in the Malaysian National Health and Morbidity Survey, the prevalence of psychiatric disorder was 11% [10].

It has been reported that primary care practitioners miss about one third of psychiatrically ill people [4,11,12]. A number of reasons have been adduced for this. Patients seeing their primary care doctors tend to somatize their emotional distress, presenting with physical symptoms rather than overt psychological symptoms [13]. Medical history is often taken in conditions of little privacy thereby discouraging patients from sharing sensitive aspects of their distress [6]. Primary care practitioners (PCP) may also not be confident in diagnosing and treating psychiatric disorders [14,15] Moreover PCP tend to have limited time in which to obtain a psychiatric history [15].

The consequences of psychiatric morbidity such as depression when it is not identified and treated can be severe. These include suicide, loss of jobs and relationships, and deterioration in physical health including higher risk of myocardial infarction [16-18]. Early detection is important because it would reduce not only the above mentioned consequences but also unnecessary suffering for the patients [19,20]. In view of the serious consequences of psychiatric morbidity it would be useful for us to explore the prevalence of psychiatric morbidity in our own local setting.

The use of self-report screening tools in primary care settings is potentially helpful in view of the impediments to recognition and treatment of mental disorder in primary care. A number of screening tools exist, including GHQ, Self-rating Questionnaire (SRQ) and the Patient Health Questionnaire (PHQ) [21-23]. The aim of this paper is to report the prevalence of psychiatric disorder in a semi-urban primary care setting using the PHQ.

Methods

This is a cross-sectional study carried out in a semi-urban primary healthcare center located south of Kuala Lumpur. The center is a government run healthcare facility which is easily accessible by public transport. The center caters for the health needs of a population of 603,800 people of whom a majority are from the lower middle socio-economic group. The center provides general medical outpatient care, emergency services and special clinics for antenatal, diabetes, hypertension, psychiatric illnesses and HIV-related diseases. The average attendance in a month is about 5,000 cases (100 to 150 patients per day), a majority of whom are antenatal cases seen by nurses and midwives. Other facilities include laboratory investigations, ECG, radiography and ultrasound. The clinics are run by a primary healthcare physician, one medical officer, eight nurses, two midwives, two health attendants, two assistant pharmacists and one laboratory technician.

Patients aged from 18–70 years were selected based on their registration number and subjected to a systematic random sampling. Exclusion criteria were patients who did not consent and those who were not literate.

Instruments

The PHQ was translated into Bahasa Malaysia and back translated. It was pre-tested to ensure that the original meanings were retained. The questionnaire consists of items that enable identification of eight psychiatric diagnoses including somatoform disorder, major depressive disorder, panic disorder, binge eating disorder, bulimia nervosa, alcohol abuse, other anxiety disorders and other depressive disorders.

The PHQ was validated against the GHQ-30 which has previously been validated for a Malaysian population [24]. The results of the validation will be reported in a subsequent paper. The respondents were also requested to complete a data form consisting of sociodemographic data, psychiatric history and recent life events.

Procedure

The patients completed the forms while waiting to see the doctor. Once they were in the consultation room, the doctor verified positive responses and applied the diagnostic algorithm. The time taken by respondents to complete the questionnaire was 5 to 15 minutes. The ethics approval for the study was obtained from the Research Committee of the Department of Family Medicine, National University of Malaysia.

Results

The sociodemographic characteristics of the sample are shown in Table 1. A total of 267 subjects returned questionnaires during the study period. Four questionnaires were incomplete and were excluded from analysis. The sample consisted of 97 (36.9%) young adults aged 18 to 29 years, 91 (34.6%) aged 30 to 39 years, and 75 (28.5%) aged 40 to 70 years.

Of the 263 respondents who returned completed questionnaires, 233 (88.6%) were married. The majority of the respondents were female ($n = 188$, 71.7%). Malays formed the majority ethnic group ($n = 199$, 75.7%), followed by Indians ($n = 32$, 12%), Chinese ($n = 29$, 11%) and other races ($n = 3$, 2%).

Most of the respondents had at least secondary school education (64.6%) and 69% were employed. Almost half (46%) were from the lower income group.

About twenty percent had a previously known medical illness (including diabetes mellitus, hypertension, coronary heart disease or bronchial asthma). One respondent had

Table 1: Demographic data of respondents

Demographic Data		Respondents (n = 263)	
		No.	Percentage (%) against total
Age group	18 ~ 29 years	97	36.9
	30 ~ 39 years	91	34.6
	40 ~ 49 years	51	19.4
	50 ~ 59 years	15	5.7
	60 ~ 70 years	9	3.4
Sex	Female	188	71.5
	Male	75	28.5
Race/Ethnic	Malay	199	75.7
	Chinese	29	11.0
	Indian	31	11.8
	Others	4	1.5
Education Level	None/Primary	27	10.3
	Secondary	170	64.6
	College/Tertiary	66	25.1
Marital Status	Single	26	9.9
	Married	233	88.6
	Widow/Divorced	4	1.5
Occupation	Professional, tech. & rel. work	78	29.7
	Clerical & Sales	69	26.2
	Production workers & laborers	35	13.3
	Economically inactive	81	30.8
Income Group	≤ RM1, 000	39	14.8
	RM1, 001 ~ RM1, 999	121	46.0
	RM2, 000 ~ RM2, 999	57	21.7
	≥ RM3, 000	46	17.5
Clinical Data		Respondents (n = 263)	
		No.	Percentage (%) of total respondents
PHQ	Positive	65	24.7
GHQ	Positive	71	27.0
Previously known medical illness	Yes	52	19.8
Previous history of psychiatric illness	Yes	1	0.4
Family history of psychiatric illness	Yes	1	0.4
History of stresses in the last 4 weeks	Yes	18	6.8
History of menstrual problems	Yes	25	9.5

previous psychiatric illness while another reported a family history of psychiatric illness (0.4%). Only 6.8% reported having a history of stress or losses in the previous four weeks.

It was found that 24.7% of respondents had at least one PHQ positive diagnosis. Some had more than one diagnosis. The diagnoses included depressive illness (n = 38,14.4%), somatoform disorder (n = 32, 12.2%), panic and anxiety disorders (n = 17, 6.5%), binge eating disorder (n = 9,3.4%), alcohol abuse (n = 6, 2.3%) (Table 2).

To analyze the association between PHQ positive status and sociodemographic and clinical characteristics, chi-square test was done with $p < 0.05$ taken as statistically significant.

Table 3 shows that age group was the only sociodemographic factor significantly associated with positive PHQ. It was significantly associated with young age group between 18 to 29 years. ($p = 0.036$)

Having a history of stressors in the last four weeks was found to be significantly associated with PHQ positive ($p = 0.044$)

Discussion

Screening for psychiatric disorders in primary care is an important step to improve services. It should prompt physicians to consider further full diagnostic interview and referral to specialized psychiatric services whenever necessary [5]

An earlier study [25] used the GHQ-30 in detecting psychiatric morbidity in an urban Malaysian primary care population. In this study the PHQ is used because it is designed to provide diagnoses of mental disorders in primary care [15,26]

In this study it was found that the prevalence of psychiatric disorder among adult patients attending a primary care center was 24.7%. This is similar to studies done in other countries which reported prevalence between 25% and 35% [2,27-29]. A study done in Taiwan showed a higher prevalence of 38.2% [30]

A previous Malaysian study of probable mental disorder in primary care, using GHQ, and carried out in an urban population attending private outpatient clinics in Kuala Lumpur, reported a prevalence of 29.9% [25].

The overall prevalence of psychiatric disorder (24.7%) and the prevalence of depressive illness (14.4%) are within the expected range in a primary care setting [2,27-29]. The prevalence of somatoform disorder (12.2%) and the lower than expected prevalence of anxiety disorders (6.5%) perhaps reflect the tendency of Asian patients to present their psychological distress in somatic symptoms [31-33]. An interesting finding from this study is a finding of the higher than expected prevalence of binge eating disorder (3.4%) which is, to the best of our knowledge, the first estimate of prevalence of this disorder in a primary care setting in Malaysia.

Other studies found that being female, unemployed, separated or divorced is associated with a higher probability of psychiatric disorder [1,2,4,10,29,34-36]. However in this study, the only two factors found to be associated with psychiatric morbidity were age between 18 to 29 and a history of recent stressors.

Inferences from this study should be drawn with caution because of several limitations. The questionnaire was validated against the GHQ and not against a structured clinical interview which would have given better data on psychometric properties of the PHQ. There was a preponderance of Malays and women in this sample rendering the findings not applicable to primary care settings with

Table 2: Clinical data of respondents

PHQ Diagnosis	Respondents (n = 263)		
		No.	Percentage (%) of total respondents
Somatoform disorder	Positive	32	12.2
Major depression & other depressive Syndrome	Positive	38	14.4
Panic Syndrome & other anxiety syndrome	Positive	17	6.5
Binge eating disorder	Positive	9	3.4
Alcohol abuse	Positive	6	2.3

Table 3: Association of demographic data with PHQ score

Demographic Data		Respondents (n = 263)		P
		PHQ positive (% of total respondents)	PHQ negative (% of total respondents)	
Age	18 ~ 29 years	25 (9.5)	72 (27.5)	0.036
	30 ~ 39 years	17 (6.5)	74 (28.1)	
	40 ~ 49 years	20 (7.6)	31 (11.8)	
	50 ~ 59 years	1 (0.4)	14 (5.3)	
	60 ~ 70 years	2 (0.8)	7 (2.7)	
Sex	Female	47 (17.9)	141 (53.6)	0.865
	Male	18 (6.8)	57 (21.7)	
Race/Ethnic	Malay	47 (17.9)	152 (53.6)	0.230
	Chinese	5 (1.9)	24 (9.1)	
	Indian	11 (4.2)	20 (7.6)	
	Others	2 (0.8)	2 (0.8)	
Education Level	None/Primary	8 (3.0)	19 (7.2)	0.822
	Secondary	41 (15.6)	129 (49.0)	
	College/Tertiary	16 (6.1)	50 (19.0)	
Marital Status	Single	7 (2.7)	18 (7.2)	0.472
	Married	56 (21.3)	188 (67.3)	
	Widow/Divorced	2 (0.8)	2 (0.8)	
Occupation	Professional, tech. & rel. work	18 (6.8)	60 (22.8)	0.574
	Clerical & Sales	16 (6.1)	53 (20.2)	
	Production workers & labourers	12 (4.6)	23 (8.7)	
	Economically inactive	19 (7.2)	62 (23.6)	
Income Group	≤ RM1, 000	8 (3.0)	31 (11.8)	0.740
	RM1, 001 ~ RM1, 999	33 (12.5)	88 (33.5)	
	RM2, 000 ~ RM2, 999	12 (4.6)	45 (17.1)	
	≥ RM3, 000	12 (4.6)	34 (12.9)	

Note: total % exceeds 24.7% because some patients have more than one diagnosis.

different ethnic composition. It is more representative of semi-urban primary care centers with a preponderance of Malay attenders from the lower and lower middle income groups.

Systematic random sampling was carried out thereby reducing sampling bias. The sample size was adequate.

Conclusion

The healthcare centre in which the study was carried out shares common features with many other centres in rural Malaysia. Hence the study is potentially useful in estimating the prevalence of psychiatric disorder in similar settings.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

RZ, MM and MT were involved in writing of this article. KO and MT were also involved in planning and supervision of the study. PK was involved in data collection.

Acknowledgements

We would like to thank Prof. Robert Spitzer for permission to use the PHQ.

References

1. Goldberg DP, Kay C, Thompson L: **Psychiatric Morbidity in General Practice and The Community.** *Psychological Medicine* 1976, **6**:565-569.
2. Barrett JE, Barrett JA, Oxman TE, Gerber PD: **The Prevalence of Psychiatric Disorders in a Primary Care Practice.** *Archives of General Psychiatry* 1988, **45**:1100-1106.
3. Schulberg HC: **Mental Disorders in the Primary Care Setting.** *General Hospital Psychiatry* 1991, **13**:156-164.
4. Badamgarav E, Badamgarav E, Weingarten SR, Henning JM, Knight K, Hasselblad V, Gano A Jr, Ofman JJ: **Effectiveness of disease management programs in depression: a systematic review.** *American Journal of Psychiatry* 2003, **160**(12):2080-2090.

5. Halverson J, Chan C: **Screening for Psychiatric Disorders in Primary Care.** *Wisconsin Medical Journal* 2004, **103(6)**:46-51.
6. Goldberg D: **Identifying Psychiatric Illness among General Medical Patients.** *Br Med J (Clin Res Ed)* 1995, **291(6489)**:161-162.
7. Amin SM, Hushaimi B, Syed Hassan A: **Recognizing people with psychiatry problems in community.** *Malaysian Journal of Psychiatry* 1997, **5**:50-54.
8. Eisenberg L: **Treating Depression and Anxiety in Primary Care.** *The New England Journal of Medicine* 1992, **326**:1080-1083.
9. Varma SL, Azhar MZ: **Psychiatry Symptomatology in a Primary Health Setting in Malaysia.** *Med J Malaysia* 1995, **50**:11-16.
10. Weich S, Lewis G: **Poverty: Unemployment, and Common Mental Disorders: Population Based Cohort Study.** *British Medical Journal* 1998, **317**:115-119.
11. Indran SK: **Current Perspective on Community Psychiatry.** *Malaysian Journal of Psychiatry* 1994, **2**:90-94.
12. Deva MP: **Psychiatry for General Practitioner.** *Med J Malaysia* 1997, **52**:99-101.
13. Scicchitano J, Lovell P, Pearce P, Marley J, Pilowsky I: **Illness Behavior and Somatization in General Practice.** *J Psychosom Res* 1996, **41(3)**:247-254.
14. Pini S, Berardi D, Rucci P, Piccinelli M, Neri C, Tansella M, Ferrari G: **Identification of Psychiatric Distress by Primary Care Physicians.** *General Hospital Psychiatry* 1997, **19**:411-418.
15. Spitzer RL, Willaims JBW, Kroenke K, Linzer M, Verloin F, Hahn SR, Brody D, Johnson JG: **Utility of a New Procedure for Diagnosing Mental Disorders in Primary Care.** *JAMA* 1994, **272**:1749-1756.
16. World Health Organization, Reducing Risks, Promoting Healthy Life: **The World Health Report 2002.** Geneva 2002.
17. Ustun TB, Ayuso-Mateos JL, Chatterji S, et al.: **Global burden of depressive disorders: methods and data sources.** *British Journal of Psychiatry* 2004, **184**:386-392.
18. Chisholm D, Sanderson K, Ayuso-Mateos JL, Saxena S: **Reducing the global burden of depression. Population-level analysis of intervention cost-effectiveness in 14 world regions.** *Br J Psychiatry* 2004, **184**:393-403.
19. Pignone MP, Gaynes BN, Rushton JL, Burchell CM, Orleans CT, Mulrow CD, Lohr KN: **Screening for depression in adults: a summary of the evidence for the U.S. Preventive Services Task Force.** *Ann Intern Med* 2002, **136**:765-776.
20. Wittchen HU, Holsboer F, Jacobi F: **Met and unmet needs in the management of depressive disorder in the community and primary care: the size and breadth of the problem.** *J Clin Psychiatry* 2001; **62 Suppl 26**:23-8 2001, **62 Suppl 26**:23-28.
21. Goldberg DP, Gater R, Sartorius N, Ustun TB, Piccinelli M, Gureje O, Rutter C: **The Validity of two Versions of the GHQ in the WHO Study of Mental Illness in General Health Care.** *Psychological Medicine* 1997, **27**:191-197.
22. Salleh MR: **Psychiatric Morbidity in Schizophrenic Relatives – Use of Self Reporting Questionnaire (SRQ).** *Singapore Med J* 1990, **31(5)**:457-462.
23. Ramli H, Ohn K, Krishnaswamy S, Kasmini K, Hassan S: **A Prevalence Survey of Psychiatric Morbidity in a Rural Malaysian Village.** *Singapore Med J* 1987, **28(6)**:530-533.
24. Abdul Hamid AR, Mohamed Hatta SA: **Validation Study of the Malay language Translation of General Health Questionnaire-30 (GHQ-30).** *Malaysian Journal of Psychiatry* 1996, **2(4)**:118-122.
25. Maniam T: **Psychiatric Morbidity in an Urban General Practice.** *Med J Malaysia* 1994, **49**:242-246.
26. Carlat DJ: **The Psychiatric review of Symptoms: A Screening Tool for Family Physicians** 1998 [<http://www.aafp.org/afp/981101ap/carlat.html>]. The American Academy of Family Physician
27. Kessler LG, Cleary PD, Burke JD: **Psychiatric Disorders in Primary Care: Results of Follow-up Study.** *Archives of General Psychiatry* 1985, **42**:583-587.
28. Bellantuono C, Fiorio R, Williams P, Cortina P: **Psychiatry Morbidity in an Italian Practice.** *Psychological Medicine* 1987, **17**:243-247.
29. Vazquez-Barquero JL, Garcia J, Simon JA, Iglesias C, Montejo J, Herran A, Dunn G: **Mental Health in Primary Care: An Epidemiological Study of Morbidity and Use of Health Resources.** *British Journal of Psychiatry* 1997, **170**:529-535.
30. Liu SII: **The prevalence of psychiatric morbidity and its associated factors in general health care in Taiwan.** *Psychol Med* 2002, **32(4)**:629-637.
31. Chou JCY: **Treating Chinese Psychiatric Patients.** *7th International Conference on Health Problems Related to the Chinese* 1994.
32. Markus H, Kitayama S: **Cultural variation in the self concept.** In *Multidisciplinary perspectives on the self* Edited by: Goethals GR, Strauss J. New York, Springer-Verlag; 1991:224-253.
33. Kamil MA, Beng KS: **Cultural health beliefs in a rural family practice: A Malaysian perspective.** *Aust J Rural Health* 2006, **14**:2-8.
34. Finlay-Jones RA, Burvill PW: **The Prevalence of Minor Psychiatric morbidity in the Community.** *Psychol Med* 1977, **7(3)**:475-489.
35. Pini S, Berardi D, Rucci P, Piccinelli M, Neri C, Tansella M, Ferrari G: **Identification of Psychiatric Distress by Primary Care Physicians.** *General Hospital Psychiatry* 1997, **19**:411-418.
36. Salleh MR: **Somatization of Depression in Psychiatric Out-Patients.** *Med J Malaysia* 1989, **44**:275-282.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

