

Functional Analysis of Transcription Factors in Arabidopsis

Nobutaka Mitsuda and Masaru Ohme-Takagi*

Research Institute of Genome-Based Biofactory, National Institute of Advanced Industrial Science and Technology (AIST), Central 4, Higashi 1-1-1, Tsukuba, 305-8562 Japan

Transcription factors (TFs) regulate the expression of genes at the transcriptional level. Modification of TF activity dynamically alters the transcriptome, which leads to metabolic and phenotypic changes. Thus, functional analysis of TFs using 'omics-based' methodologies is one of the most important areas of the post-genome era. In this mini-review, we present an overview of Arabidopsis TFs and introduce strategies for the functional analysis of plant TFs, which include both traditional and recently developed technologies. These strategies can be assigned to five categories: bioinformatic analysis; analysis of molecular function; expression analysis; phenotype analysis; and network analysis for the description of entire transcriptional regulatory networks.

Keywords: Arabidopsis • Bioinformatics • Functional analysis • Methodology • Transcription factor.

Abbreviations: AD, activation domain; CaMV, cauliflower mosaic virus; CD, conserved domain; ChIP, chromatin immunoprecipitation; CHX, cycloheximide; CRES-T, chimeric repressor silencing technology; DBD, DNA-binding domain; DEG, differentially expressed gene; DEX, dexamethasone; EAR, ERF-associated amphiphilic repression; ER, estrogen receptor; FDR, false discovery rate; FOX, full-length cDNA overexpressor; GFP, green fluorescent protein; GO, gene ontology; GR, glucocorticoid receptor; GUS, β -glucuronidase; LUC, luciferase; miRNA, microRNA; PPI, protein–protein interaction; RD, repression domain; RT-PCR, reverse transcription–PCR; SRDX, modified EAR motif plant-specific repression domain showing strong repression activity; ta-siRNA, trans-acting small interfering RNA; TF, transcription factor; Y1H, yeast one-hybrid screening.

Introduction

It is evident that transcriptional regulation plays a pivotal role in the control of gene expression in plants. Intensive

studies of plant mutants have revealed that informative phenotypes are often caused by mutations in genes for transcription factors (TFs), and a number of TFs have been identified that act as key regulators of various plant functions. TFs, which regulate the first step of gene expression, are usually defined as proteins containing a DNA-binding domain (DBD) that recognize a specific DNA sequence. In addition, proteins without a DBD, which interact with a DNA-binding protein to form a transcriptional complex, are often categorized as TFs. Although some metabolic enzymes have been suggested to regulate gene expression directly in yeast (Hall et al. 2004), we do not focus on such multifunctional proteins in this review. In 2000, the entire genome sequence of *Arabidopsis thaliana* was determined and the genome was predicted to contain 25,498 protein-coding genes (Arabidopsis Genome Initiative 2000). Based on sequence conservation with known DBDs, Riechmann et al. (2000) reported that around 1,500 of these genes encode TFs, and more recent analyses have recognized >2,000 TF genes in the Arabidopsis genome (Davuluri et al. 2003, Guo et al. 2005, Iida et al. 2005, Riano-Pachon et al. 2007).

In contrast to Arabidopsis, the number of TF genes found in *Drosophila melanogaster* and *Caenorhabditis elegans*, which have similar sized genomes to that of Arabidopsis, is around 600, which is significantly less than that in Arabidopsis (Riechmann et al. 2000). The ratio of TF genes to the total number of genes in Arabidopsis is 5–10% depending on databases, which is higher than that of *D. melanogaster* (4.7%) and of *C. elegans* (3.6%) (Riechmann et al. 2000), although it is comparable with that of human (6.0%) (Venter et al. 2001). In addition to the larger number of TF genes in Arabidopsis, there is a greater variety of TFs, with a greater diversity of DNA binding specificities, compared with *D. melanogaster* or *C. elegans* (see later for more details). These characteristic features of Arabidopsis TFs suggest that transcriptional regulation plays more important roles in plants than in animals. Because transcriptional regulation is

*Corresponding author: E-mail, m-takagi@aist.go.jp; Fax, +81-29-861-3026.

Plant Cell Physiol. 50(7): 1232–1248 (2009) doi:10.1093/pcp/pcp075, available online at www.pcp.oxfordjournals.org

© The Author 2009. Published by Oxford University Press on behalf of Japanese Society of Plant Physiologists. All rights reserved.

The online version of this article has been published under an open access model. Users are entitled to use, reproduce, disseminate, or display the open access version of this article for non-commercial purposes provided that: the original authorship is properly and fully attributed; the Journal and the Japanese Society of Plant Physiologists are attributed as the original place of publication with the correct citation details given; if an article is subsequently reproduced or disseminated not in its entirety but only in part or as a derivative work this must be clearly indicated. For commercial re-use, please contact journals.permissions@oxfordjournals.org

the first step of gene expression and could affect various 'omes', namely the proteome, metabolome and phenome, the functional analysis of TFs is important and necessary for omics studies and for the elucidation of whole functional networks in plants. Although much effort has been made to identify the function of TFs, most of their functions remain to be clarified. In this mini-review, we present an overview of Arabidopsis TFs and describe strategies for the functional analysis of plant TFs, which include both traditional and recently developed technologies.

Overview of Arabidopsis transcription factors

According to The Arabidopsis Information Resources (TAIR, <http://www.arabidopsis.org>), there are 27,235 protein-coding genes in the Arabidopsis genome (ftp://ftp.arabidopsis.org/home/tair/Genes/TAIR8_genome_release/README). Four independent reports have recently shown that approximately 2,000 genes encode TFs (**Table 1**). These four representative databases of Arabidopsis TFs are: RARTF (<http://rarge.gsc.riken.jp/rartf/>) (Iida et al. 2005), AGRIS (<http://arabidopsis.med.ohio-state.edu/AtTFDB/>) (Davuluri et al. 2003), DATF (<http://datf.cbi.pku.edu.cn/>) (Guo et al. 2005) and PlnTFDB (http://plntfdb.bio.uni-potsdam.de/v2.0/index.php?sp_id=ATH) (Riano-Pachon et al. 2007). Each database classified TFs into families based on their own classification criteria, and the number of loci in each family is different among the four databases. A total of 51, 51, 64 and 67 families (72 families in total) and 1,965, 1,837, 1,914 and 1,949 loci (2,620 loci in total), respectively, were identified. A total of 1,318 loci are recognized by all four databases (**Table 1** and **Supplementary Table S1**). These differences are mainly due to the different definition of a TF in each database. For example, the AGRIS database does not include AUX/IAA proteins as TFs as they do not directly bind to DNA but repress auxin-mediated gene transcription by interacting with ARF transcription factors (Ouellet et al. 2001), whereas the other three databases classify them as being TFs (**Table 1**).

Arabidopsis TFs are characterized by a large number of genes and by the variety of gene families when compared with those of *D. melanogaster* or *C. elegans*. For example, zinc-finger TFs represent more than half of all TFs in *D. melanogaster* or *C. elegans*, whereas those in Arabidopsis represent around 20% (Riechmann et al. 2000). Around half of Arabidopsis TFs are plant specific and possess DBDs found only in plants (Riechmann et al. 2000 and **Table 1**). AP2-ERF, NAC, Dof, YABBY, WRKY, GARP, TCP, SBP, ABI3-VP1 (B3), EIL and LFY are plant-specific TFs. The three-dimensional structures of several plant-specific DBDs, i.e. NAC, WRKY, SBP, EIL, B3 and AP2-ERF, have been determined (Allen et al. 1998, Ernst et al. 2004, Yamasaki et al. 2004a, Yamasaki et al. 2004b, Yamasaki et al. 2005a, Yamasaki et al. 2005b). Most Arabidopsis TFs form large families, which share similar DBD

structures. For example, AP2-ERF and NAC domain families contain >100 loci each (**Table 1**). MYB, MADS box, bHLH (basic helix-loop helix), bZIP and HB, which are not plant-specific families, also form large families. These families, such as the MADS box family, which includes a number of ABC floral homeotic genes (Riechmann et al. 1996), play important roles in the control of plant growth and development.

TFs act as transcriptional activators or repressors. In common with other eukaryotes, TFs containing domains rich in the acidic amino acids glutamine or proline, such as TOC1, DREBs, ARFs and GBF1, are transcriptional activators (Schindler et al. 1992, Ulmasov et al. 1999, Strayer et al. 2000, Sakuma et al. 2002). In addition, the AHA motif, which has a characteristic pattern of aromatic and large hydrophobic amino acid residues embedded in an acidic context, was shown to act as an activation domain (AD) in plant heat shock factors (Döring et al. 2000).

On the other hand, transcriptional repressors in plants were not elucidated until the ERF-associated amphiphilic repression (EAR) motif was identified in tobacco ETHYLENE RESPONSIVE ELEMENT BINDING FACTOR 3 (EREBP3) (Ohta et al. 2000). Transcriptional repressors are roughly categorized into passive or active repressors. Passive repressors have neither an AD nor a repression domain (RD). Some repress transcription by binding to the promoter of the target gene, thereby competing with an activator that interacts with the same *cis*-element. Maize Dof2 is known to be a passive repressor (Yanagisawa and Sheen 1998). Arabidopsis CAPRICE (CPC), TRIPTYCHON (TRY), ENHANCER OF TRY AND CPC1 (ETC1), ETC2 and ETC3, which are all small MYB proteins with a single R3-MYB domain, are negative regulators involved in the development of epidermal cells (reviewed in Simon et al. 2007) and are likely to act as passive repressors. They compete with other R2-R3 MYB proteins such as GLABRA1 (GL1) and WEREWOLF (WER) that positively regulate epidermal cell development for interaction with bHLH proteins (Esch et al. 2004, Simon et al. 2007, Tominaga et al. 2007). The active repressors possess distinct RDs that confer repressive activity to the TF. The EAR motif is a plant-specific repression domain. The minimum unit of the EAR-motif RD is only six amino acids, which comprise an amphiphilic feature composed of leucine and acidic amino acids (Hiratsu et al. 2004). Because fusion of the EAR motif RD can convert a transcriptional activator into a strong repressor (Hiratsu et al. 2003), TFs that contain this motif are assumed to be transcriptional repressors, although experimental validation is required. Database analysis revealed that the EAR motif RD is found in 404 loci among 2,620 putative TFs (**Supplementary Table S1**). Interestingly, RDs are over-represented in the C2H2 zinc finger (68/136), AUX-IAA (28/29) and HB (32/93) families (**Supplementary Table S1**). Most RDs are conserved in various plants, including dicots and monocots, but are not obviously over-represented in TFs of other

Table 1 Comparison of plant TF databases

| | RARTF | | AGRIS | | DATF | | PInTFDB | |
|-----|----------------------|---------|---|---------------------|---|---------------------|------------------------------|----------|
| | Family | Loci | Family | Loci | Family | Loci | Family | Loci |
| 1. | ABI3/VP1 | 51 | ABI3VP1 REM | 11 21 | ABI3-VP1 | 60 | ABI3VP1 | 56 |
| 2. | Alfin-like | 47 | Alfin-like | 7 | Alfin | 7 | Alfin-like | 7 |
| 3. | AP2/EREBP | 93 | AP2-EREBP | 136 | AP2-EREBP | 146 | AP2-EREBP | 146 |
| | ERF | 19 | | | | | | |
| | Pti4 | 5 | | | | | | |
| | Pti5 | 5 | | | | | | |
| | Pti6 | 18 | | | | | | |
| 4. | ARF | 71 | ARF RAV | 22 11 | ARF | 23 | ARF | 23 |
| 5. | ARID | 6 | ARID | 7 | ARID | 10 | ARID | 10 |
| 6. | AT-hook | 31 | – | – | – | – | – | – |
| 7. | – | – | – | – | AS2 | 42 | – | – |
| 8. | Aux/IAA | 21 | – | – | AUX-IAA | 28 | AUX/IAA | 27 |
| 9. | – | – | BBR/BPC | 7 | BBR-BPC | 7 | BBR/BPC | 7 |
| 10. | – | – | BZR | 6 | BES1 | 8 | BES1 | 8 |
| 11. | bHLH | 157 | bHLH | 162 | bHLH | 127 | bHLH | 134 |
| 12. | – | – | – | – | – | – | bHSH | 1 |
| 13. | bZIP | 56 | bZIP | 73 | bZIP | 72 | bZIP | 71 |
| | TGA3 | 27 | | | | | | |
| 14. | C2C2(Zn)-CO-like | 51 | C2C2-CO-like | 30 | C2C2-CO-like | 37 | C2C2-CO-like Pseudo ARR-B | 17 5 |
| 15. | C2C2(Zn)-Dof | 33 | C2C2-Dof | 36 | C2C2-Dof | 36 | C2C2-Dof | 36 |
| 16. | C2C2(Zn)-GATA | 37 | C2C2-Gata | 30 | C2C2-GATA | 26 | C2C2-GATA | 29 |
| 17. | C2C2(Zn)-YABBY | 5 | C2C2-YABBY | 6 | C2C2-YABBY | 5 | C2C2-YABBY | 6 |
| 18. | C2H2(Zn) | 177 | C2H2 | 211 | C2H2 | 134 | C2H2 | 96 |
| 19. | C3H-type 1(Zn) | 37 | C3H | 165 | C3H | 59 | C3H | 67 |
| 20. | – | – | CAMTA | 6 | CAMTA | 6 | CAMTA | 6 |
| 21. | CBF5 | 2 | – | – | – | – | – | – |
| 22. | CCAAT | 37 | CCAAT-DR1 CCAAT-HAP2 CCAAT-HAP3 CCAAT-HAP5 | 2 10 10 13 | CCAAT-Dr1 CCAAT-HAP2 CCAAT-HAP3 CCAAT-HAP5 | 2 10 11 13 | CCAAT | 43 |
| 23. | CPP(ZN) | 8 | CPP | 8 | CPP | 8 | CPP | 8 |
| 24. | – | – | – | – | – | – | CSD | 4 |
| 25. | – | – | – | – | – | – | DBP | 4 |
| 26. | – | – | – | – | – | – | DDT | 4 |
| 27. | E2F/DP | 8 | E2F-DP | 8 | E2F-DP | 8 | E2F-DP | 7 |
| 28. | EIL | 6 | EIL | 6 | EIL | 6 | EIL | 6 |
| 29. | – | – | – | – | FHA | 16 | FHA | 17 |
| 30. | GARP | 51 | G2-like ARR-B | 40 15 | GARP-G2-like GARP-ARR-B | 42 10 | G2-like ARR-B | 39 13 |
| 31. | – | – | GeBP | 16 | GeBP | 21 | GeBP | 20 |
| 32. | – | – | – | – | GIF | 3 | – | – |
| 33. | GRAS | 32 | GRAS | 31 | GRAS | 33 | GRAS | 33 |
| 34. | – | – | GRF | 9 | GRF | 9 | GRF | 9 |
| 35. | HB PAIRED(w/o HB) | 97 2 | Homeobox | 91 | HB | 87 | HB | 91 |
| 36. | HMG-box | 11 | – | – | HMG | 11 | HMG | 11 |

continued

Table 1 Continued

| RARTF | | AGRIS | | DATF | | PlnTFDB | |
|---------------------|------|-------------|------|-------------|------|--------------|------|
| Family | Loci | Family | Loci | Family | Loci | Family | Loci |
| 37. – | – | HRT | 3 | HRT-like | 2 | HRT | 2 |
| 38. HSF | 27 | HSF | 21 | HSF | 23 | HSF | 23 |
| 39. C3H-type 2(Zn) | 10 | JUMONJI | 5 | JUMONJI | 17 | Jumonji | 17 |
| | 13 | | | | | | |
| 40. LFY | 3 | LFY | 1 | LFY | 1 | LFY | 1 |
| 41. LIM-domain | 6 | – | – | LIM | 13 | LIM | 6 |
| 42. – | – | – | – | LUG | 2 | LUG | 2 |
| 43. MADS | 106 | MADS | 109 | MADS | 102 | MADS | 102 |
| 44. – | – | – | – | MBF1 | 3 | MBF1 | 3 |
| 45. MYB superfamily | 189 | MYB | 130 | MYB | 149 | MYB | 145 |
| | | MYB-related | 67 | MYB-related | 49 | MYB-related | 64 |
| 46. NAC | 106 | NAC | 94 | NAC | 105 | NAC | 101 |
| 47. Nin-like | 14 | NLP | 9 | Nin-like | 14 | RWP-RK | 14 |
| | | AtRKD | 5 | | | | |
| 48. – | – | – | – | NZZ | 1 | NOZZLE | 1 |
| 49. PcG; E(z) class | 32 | | | PcG | 34 | SET | 33 |
| PcG; Esc class | 3 | | | | | | |
| 50. PHD-finger | 9 | PHD | 11 | PHD | 55 | PHD | 43 |
| 51. – | – | – | – | PLATZ | 10 | PLATZ | 11 |
| 52. – | – | – | – | – | – | RB | 1 |
| 53. – | – | – | – | S1Fa-like | 3 | S1Fa-like | 3 |
| 54. – | – | – | – | SAP | 1 | SAP | 1 |
| 55. SBP | 17 | SBP | 16 | SBP | 16 | SBP | 16 |
| 56. Sir2 | 2 | – | – | – | – | – | – |
| 57. – | – | – | – | – | – | Sigma70-like | 6 |
| 58. – | – | – | – | SRS | 10 | SRS | 10 |
| 59. – | – | – | – | – | – | SNF2 | 38 |
| 60. SW13 | 6 | – | – | – | – | – | – |
| 61. Swi4/Swi6 | 1 | – | – | – | – | – | – |
| 62. – | – | – | – | TAZ | 9 | TAZ | 8 |
| 63. TCP | 24 | TCP | 26 | TCP | 23 | TCP | 24 |
| 64. Trihelix | 31 | Trihelix | 29 | Trihelix | 26 | Trihelix | 23 |
| 65. TUB | 11 | TUB | 10 | TLP | 11 | TUB | 10 |
| 66. – | – | – | – | ULT | 2 | ULT | 2 |
| 67. – | – | VOZ | 2 | VOZ | 2 | VOZ | 2 |
| 68. VIP3 | 1 | – | – | – | – | – | – |
| 69. – | – | Whirly | 3 | Whirly | 2 | PBF-2-like | 3 |
| 70. WRKY(Zn) | 72 | WRKY | 72 | WRKY | 72 | WRKY | 72 |
| 71. – | – | ZF-HD | 15 | ZF-HD | 16 | zf-HD | 17 |
| 72. – | – | ZIM | 2 | ZIM | 18 | ZIM | 15 |
| Other | 81 | Other | 1 | | | Other | 69 |
| Total | 1965 | Total | 1837 | Total | 1914 | Total | 1949 |

The number of loci in each database is shown. '–' indicates that no corresponding TF family is defined in the database.

organisms, such as yeast (N. Mitsuda et al. unpublished results). These suggest that the EAR motif RD and its mechanism of action is plant specific. Recently, novel RDs that could not be categorized according to the EAR motif (Hiratsu et al. 2004) were identified in AtMYBL2 and B3 DBD TFs (Matsui et al. 2008, Ikeda and Ohme-Takagi 2009). This suggests that unidentified transcriptional repressors with novel RDs may be encoded in plant genomes. Activators and repressors act antagonistically to control the fine-tuning of gene expression. The molecular mechanism of transcriptional repression via the EAR motif RD remains to be clarified. Chromatin remodeling may be involved because the EAR motif interacts with TOPLESS (TPL), and mutations in *HISTONE ACETYLTRANSFERASE GNAT SUPERFAMILY 1* suppress the *tpl-1* phenotype (Long et al. 2006, Szemenyei et al. 2008). In animals, bifunctional TFs have been reported, which can act as transcriptional activators or repressors, depending on the environment or target genes (Adkins et al. 2006). In plants, WRKY53 has been shown to act as either a transcriptional activator or repressor depending on the sequence surrounding the W-box (Miao et al. 2004).

The activities of most TFs are controlled at the transcriptional level by other TFs, while several TFs are regulated post-transcriptionally, such as EIN3 (Yanagisawa et al. 2003). Small RNAs that target TF genes are also important regulators of gene expression (see **Table 3** and Bioinformatic analysis). TFs that are regulated at the post-transcriptional level may be regulators that act at early stages of transcriptional cascades.

Bioinformatic analysis

The functional analysis of TFs using bioinformatic techniques has become an important and effective strategy. Databases concerned with the functional analysis of TFs are listed in **Table 2**. Initially, amino acid sequence analysis should be performed to find evolutionarily conserved domains (CDs), including DBDs. Some TFs possess two DBDs. For example, RAV1 (At1g13260) group members have both AP2-ERF and B3 DBDs (Kagaya et al. 1999). Normally, a TF has a DBD and a transcriptional AD or RD. TFs having only a DBD are likely to be passive repressors, as reported for CPC and TRY, which interfere with the activity of the transcriptional activator (complex) (Simon et al. 2007). CD searches against known motifs can be performed using many web-based programs. One of the most useful services is InterProScan provided by the European Bioinformatics Institute (EBI) (<http://www.ebi.ac.uk/Tools/InterProScan/>) (Quevillon et al. 2005). This search is comprehensively performed against various CD databases and provides sophisticated graphical output. Finding known or unknown CDs among a set of proteins can be performed by MEME (<http://meme.sdsc.edu/meme/intro.html>) (Bailey et al. 2006). The SALAD database, which was developed specifically for plant proteins, also provides

MEME-based CD searching with various other useful tools (<http://salad.dna.affrc.go.jp/salad/en/>).

Homology searches, for example performed by BLAST, are also important for the bioinformatic study of TFs. Proteins which share high homology not only in their DBDs but also in other regions are likely to be functionally redundant, at least in tissues where they are co-expressed. However, it is frequently observed that proteins with high homology only in the DBD also function redundantly. For example, although CUP-SHAPED COTYLEDON1 (CUC1) and CUC3 are known to function redundantly, there is no significant sequence similarity outside the DBD (Vroemen et al. 2003, Hibara et al. 2006). BLAST searches against Arabidopsis can be performed through TAIR (<http://www.arabidopsis.org/Blast/index.jsp>). BLAST searches against multiple species, such as Arabidopsis and rice, can also be informative to assess functional redundancy. If three Arabidopsis proteins correspond to one rice protein, these three proteins might function redundantly. BLAST searches against a favorite combination of species can be performed through the NCBI website (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

Analysis of the subcellular localization of putative TFs is important because TFs cannot function outside the nucleus. Some NAC domain TFs possess a transmembrane motif at the C-terminus and are liberated by proteolytic cleavage to move into the nucleus (Kim et al. 2006, Kim et al. 2008). Subcellular localization of proteins can be predicted by computer programs such as SubLoc (Hua and Sun 2001), TargetP (Emanuelsson et al. 2000) and WoLF PSORT (Horton et al. 2007). Predictions of subcellular localization using 10 different computer programs and also from experimental evidence can be retrieved from the SUBAII database (<http://www.plantenergy.uwa.edu.au/suba2/>) (Heazlewood et al. 2007). This database also provides hydropathy plots of all Arabidopsis proteins.

The investigation of proteins that interact with TFs is also of great importance. Many TFs are known to form functional complexes. For example, some NAC TFs and MADS TFs form homo- or hetero-dimeric or tetrameric complexes (Honma and Goto 2001, Ernst et al. 2004, Heazlewood et al. 2007). MYB TFs and bHLH TFs often form complexes (Zimmermann et al. 2004a). A number of TFs are known to interact with kinases, resulting in TF phosphorylation (He et al. 2002, Furihata et al. 2006, Robertson et al. 2008). Predicted or experimentally validated protein–protein interactions (PPIs) among Arabidopsis proteins can be retrieved from the TAIR, EBI and AtPID databases. The 'Arabidopsis predicted interactome', stored at TAIR, provides a set of >20,000 PPIs based on ortholog matching (<http://www.arabidopsis.org/portals/proteome/proteinInteract.jsp>) (Geisler-Lee et al. 2007). EBI provides the IntAct database, which stores continuously updated PPI information of all organisms based on literature curation (<http://www.ebi.ac.uk/intact/site/index.jsf>).

Table 2 List of useful databases for the functional analysis of TFs

| Category/database name | URL | Comment |
|---|---|---|
| Plant (<i>Arabidopsis</i>) transcription factors | | |
| RARTF | http://rarge.gsc.riken.jp/rartf/ | |
| AGRIS | http://arabidopsis.med.ohio-state.edu/AtTFDB/ | |
| DATF | http://datf.cbi.pku.edu.cn/ | A part of a plant transcription factor database |
| PlnTFDB | http://plntfdb.bio.uni-potsdam.de/v2.0/index.php?sp_id=ATH | Data of other plants are also stored |
| Conserved domain search | | |
| InterProScan | http://www.ebi.ac.uk/Tools/InterProScan/ | For known motifs |
| MEME | http://meme.sdsc.edu/meme/intro.html | For discovering unknown motifs |
| SALAD database | http://salad.dna.affrc.go.jp/salad/en/ | For known and unknown motifs |
| Homology search | | |
| TAIR BLAST | http://www.arabidopsis.org/Blast/index.jsp | For <i>Arabidopsis</i> only |
| NCBI BLAST | http://blast.ncbi.nlm.nih.gov/Blast.cgi | For multispecies search |
| Prediction of subcellular localization | | |
| SUBAII | http://www.plantenergy.uwa.edu.au/suba2/ | Experimental data are also stored |
| Protein–protein interaction | | |
| <i>Arabidopsis</i> predicted interactome | http://www.arabidopsis.org/portals/proteome/proteinInteract.jsp | |
| EBI IntAct | http://www.ebi.ac.uk/intact/site/index.jsf | For all organisms |
| AtPID | http://atpid.biosino.org/index.php | |
| Small RNAs | | |
| ASRP | http://asrp.cgrb.oregonstate.edu/db/ | Includes data of miRNA, siRNA and ta-siRNA |
| Repository of microarray data | | |
| NCBI GEO | http://www.ncbi.nlm.nih.gov/geo/ | |
| EBI ArrayExpress | http://www.ebi.ac.uk/microarray-as/ae/ | |
| NASCArrays | http://affymetrix.arabidopsis.info/narrays/experimentbrowse.pl | |
| Browsing microarray data and co-expression analysis | | |
| ATTED-II | http://atted.jp/ | |
| Genevestigator | https://www.genevestigator.com/gv/index.jsp | |
| BAR eFP browser | http://bbc.botany.utoronto.ca/efp/cgi-bin/efpWeb.cgi | |
| Finding novel <i>cis</i> -elements | | |
| TAIR motif analysis | http://www.arabidopsis.org/tools/bulk/motiffinder/index.jsp | |
| Database of known <i>cis</i> -elements | | |
| PLACE | http://www.dna.affrc.go.jp/PLACE/ | No longer updated after 2007 |
| AGRIS ATCISDB | http://arabidopsis.med.ohio-state.edu/AtcisDB/ | |
| GO categorization | | |
| TAIR GO annotation search | http://www.arabidopsis.org/tools/bulk/go/index.jsp | |

(Kerrien et al. 2007). The *Arabidopsis thaliana* Protein Interactome Database (AtPID) provides a search facility with graphical output against a predicted and literature-curated *Arabidopsis* PPI data set (<http://atpid.biosino.org/index.php>) (Cui et al. 2008).

microRNA (miRNA) is also an important regulator of TF activity. According to the *Arabidopsis* small RNA Project (ASRP) (<http://asrp.cgrb.oregonstate.edu/db/>) (Gustafson et al. 2005, Backman et al. 2008), 200 genes are predicted to be targets of known miRNAs. Interestingly, 69 of these genes

Table 3 List of TF genes targeted by miRNA

| miRNA family | miRNA locus | Target family | Target locus |
|---------------------|---------------------|----------------------|----------------------|
| miR156/miR157 | AT2G25095 (miR156A) | SBP | AT5G43270 (SPL2) |
| | AT4G30972 (miR156B) | | AT2G33810 (SPL3) |
| | AT4G31877 (miR156C) | | AT1G53160 (SPL4) |
| | AT5G10945 (miR156D) | | AT3G15270 (SPL5) |
| | AT5G11977 (miR156E) | | AT1G69170 (SPL6) |
| | AT5G26147 (miR156F) | | AT2G42200 (SPL9) |
| | AT2G19425 (miR156G) | | AT1G27370 (SPL10) |
| | AT5G55835 (miR156H) | | AT1G27360 (SPL11) |
| | AT1G66783 (miR157A) | | AT5G50570 (SPL13) |
| | AT1G66795 (miR157B) | | AT3G57920 (SPL15) |
| | AT3G18217 (miR157C) | | |
| AT1G48742 (miR157D) | | | |
| miR159/miR319 | AT1G73687 (miR159A) | MYB | AT5G06100 (ATMYB33) |
| | AT1G18075 (miR159B) | | AT3G11440 (ATMYB65) |
| | AT2G46255 (miR159C) | | AT4G26930 (ATMYB97) |
| | AT4G23713 (miR319A) | | AT2G32460 (ATMYB101) |
| | AT5G41663 (miR319B) | | AT2G26950 (ATMYB104) |
| | AT2G40805 (miR319C) | AT5G55020 (ATMYB120) | |
| | | TCP | AT3G60460 (DUO1) |
| | | | AT4G18390 (TCP2) |
| | | | AT1G53230 (TCP3) |
| | | | AT3G15030 (TCP4) |
| | | | AT2G31070 (TCP10) |
| | AT1G30210 (TCP24) | | |
| miR160 | AT2G39175 (miR160A) | ARF | AT2G28350 (ARF10) |
| | AT4G17788 (miR160B) | | AT4G30080 (ARF16) |
| | AT5G46845 (miR160C) | | AT1G77850 (ARF17) |
| miR164 | AT2G47585 (miR164A) | NAC | AT3G15170 (CUC1) |
| | AT5G01747 (miR164B) | | AT5G53950 (CUC2) |
| | AT5G27807 (miR164C) | | AT5G07680 (ATNAC4) |
| | | | AT5G61430 (ATNAC5) |
| | | | AT1G56010 (NAC1) |
| | AT5G39610 (ORE1) | | |
| | AT3G12977 | | |
| miR165/miR166 | AT1G01183 (miR165A) | HB | AT2G34710 (PHB) |
| | AT4G00885 (miR165B) | | AT1G30490 (PHV) |
| | AT2G46685 (miR166A) | | AT1G52150 (CAN) |
| | AT3G61897 (miR166B) | | AT5G60690 (REV) |
| | AT5G08712 (miR166C) | | AT4G32880 (ATHB8) |
| | AT5G08717 (miR166D) | | |
| | AT5G41905 (miR166E) | ARF | AT1G30330 (ARF6) |
| | AT5G43603 (miR166F) | | AT1G37020 (ARF8) |
| | AT5G63715 (miR166G) | | |
| | AT3G22886 (miR167A) | | |
| | AT3G63375 (miR167B) | | |
| AT3G04765 (miR167C) | | | |
| AT1G31173 (miR167D) | | | |
| miR169 | AT3G13405 (miR169A) | CCAAT | AT5G06510 (NF-YA10) |
| | AT5G24825 (miR169B) | | AT1G72830 (HAP2C) |
| | AT5G39635 (miR169C) | | AT1G17590 (NF-YA8) |
| | AT1G53683 (miR169D) | | AT1G54160 (NF-YA5) |
| | AT1G53687 (miR169E) | | AT5G12840 (HAP2A) |
| | AT3G14385 (miR169F) | | AT3G20910 (NF-YA9) |
| | AT4G21595 (miR169G) | | AT3G05690 (HAP2B) |

continued

Table 3 Continued

| miRNA family | miRNA locus | Target family | Target locus |
|---------------|---------------------|---------------|--|
| miR170/miR171 | AT1G19371 (miR169H) | GRAS | AT2G45160 AT3G60630 AT4G00150 (SCL6) |
| | AT3G26812 (miR169I) | | |
| | AT3G26813 (miR169J) | | |
| | AT3G26815 (miR169K) | | |
| | AT3G26816 (miR169L) | | |
| | AT3G26818 (miR169M) | | |
| | AT3G26819 (miR169N) | | |
| miR172 | AT5G66045 (miR170) | AP2-EREBP | AT2G28550 (TOE1) AT5G60120 (TOE2) AT5G67180 (TOE3) AT4G36920 (AP2) AT2G39250 (SNZ) AT3G54990 (SMZ) |
| | AT3G51375 (miR171A) | | |
| | AT1G11735 (miR171B) | | |
| | AT1G62035 (miR171C) | | |
| | AT2G28056 (miR172A) | | |
| miR393 | AT2G39885 (miR393A) | bHLH | AT3G23690 (bHLH077) |
| | AT3G55734 (miR393B) | | |
| miR396 | AT2G10606 (miR396A) | GRF | AT2G22840 (ATGRF1) AT4G37740 (ATGRF2) AT2G36400 (ATGRF3) AT3G52910 (ATGRF4) AT5G53660 (ATGRF7) AT4G24150 (ATGRF8) AT2G45480 (ATGRF9) |
| | AT5G35407 (miR396B) | | |
| miR778 | AT2G41616 (miR778A) | SET | AT2G35160 (SGD9) AT2G22740 (SDG23) |
| miR824 | AT4G24415 (miR824A) | MADS | AT3G57230 (AGL16) |
| miR828 | AT4G27765 (miR828A) | MYB | AT1G66370 (ATMYB113) |
| miR858 | AT1G71002 (miR858A) | MYB | AT2G47460 (ATMYB12) AT3G08500 (ATMYB83) |

(35%) encode putative TFs (**Table 3** and **Supplementary Table S1**), despite TFs representing only 5–10% of all genes. For example, miRNAs that target TCP, NAC and SBP TFs are known to play very important roles in the control of plant growth and development (Palatnik et al. 2003, Wu and Poethig 2006, Ori et al. 2007, Schommer et al. 2008, Kim et al. 2009, Larue et al. 2009). ASRP also provides non-coding small RNA information (reviewed in Ramachandran and Chen 2008), such as for small interfering RNAs (siRNAs) and trans-acting siRNAs (ta-siRNAs), in addition to that for miRNAs. Six TF genes are listed as targets of ta-siRNAs in ASRP.

The spatial and temporal expression profile of a gene and the expression in response to varying conditions is fundamental to its biological function. Development of microarray technologies and public data repositories, such as NCBI

Gene Expression Omnibus (GEO) (<http://www.ncbi.nlm.nih.gov/geo/>) (Barrett et al. 2007), EBI ArrayExpress (<http://www.ebi.ac.uk/microarray-as/ae/>) (Parkinson et al. 2009) and NASCArrays (<http://affymetrix.arabidopsis.info/narrays/experimentbrowse.pl>) (Craigon et al. 2004), enables us to access many kinds of microarray data easily. The expression profile of a gene of interest can also be easily accessed on many web sites, such as ATTED-II (<http://atted.jp/>) (Obayashi et al. 2009), Genevestigator (<https://www.genevestigator.com/gv/index.jsp>) (Zimmermann et al. 2004b, Grennan 2006) and BAR eFP browser (<http://bbc.botany.utoronto.ca/efp/cgi-bin/efpWeb.cgi>) (Winter et al. 2007). These web sites also provide information regarding co-expression analysis. They provide lists of genes whose expression profiles are positively or negatively correlated with the

query gene (Zimmermann et al. 2004b, Toufighi et al. 2005, Grennan 2006, Obayashi et al. 2009). Co-expression analysis is particularly important for the functional analysis of TFs because co-expressed genes might encode proteins that are functionally related and/or are putative interacting proteins. They might also be downstream and/or upstream genes in the context of a transcriptional cascade. Hirai et al. (2007) identified MYB TFs as the key regulators of aliphatic glucosinolate biosynthesis by co-expression analysis. From promoter regions of co-expressed genes, short sequences that are statistically over-represented and may represent TF-binding sites can be identified using TAIR Motif Analysis (<http://www.arabidopsis.org/tools/bulk/motiffinder/index.jsp>). It is valuable to compare these sequences with known *cis*-elements stored in *cis*-element databases, such as PLACE (<http://www.dna.affrc.go.jp/PLACE/>) (Higo et al. 1999) and AGRIS ATCISDB (<http://arabidopsis.med.ohio-state.edu/AtcisDB/>). Furthermore, functional characteristics of these genes can be analyzed using the TAIR Gene Ontology (GO) annotation search (<http://www.arabidopsis.org/tools/bulk/go/index.jsp>). A set of favorite genes can be categorized based on a limited GO term (Ashburner et al. 2000) and shown as a graphical pie chart. In addition, we can compare these data with the results of functional categorization of all Arabidopsis proteins. These analyses help us to speculate on the biological processes in which the set of genes and the query TF are involved.

Molecular analysis

The molecular analysis of TFs involves the characterization of their activation or repression activities. To this end, analysis using reporter and effector genes is often employed (Ohta et al. 2001; Fig. 1). A commonly used effector gene consists of a chimeric construct, in which a TF-coding sequence is fused to that of a heterogeneous DBD, such as the GAL4 DNA-binding domain (GAL4DB) from yeast, and which is driven by a strong promoter, such as cauliflower mosaic virus (CaMV) 35S. The reporter gene is usually the firefly luciferase (LUC) or *Escherichia coli* β -glucuronidase (GUS) gene, which is driven by a minimal promoter with upstream repeated *cis*-elements, such as the binding sequence of the GAL4DB. Another reporter construct, containing a constitutively expressed reporter, such as sea pansy *LUC*, is used as an internal control (reference). These reporter, effector and reference constructs are transiently co-expressed by particle bombardment of leaf tissues or by polyethylene glycol-mediated transformation of leaf protoplasts or cultured cells. By assaying the activity of the reporter gene following co-expression of the effector gene, the activation activity of a TF can be examined. A transient expression assay using particle bombardment into leaf tissue is simple and reproducible and has several advantages for analyzing the molecular function of TFs (Ueki et al. 2009). Once a TF is identified




Fig. 1 Schematic drawing summarizing the molecular analysis of TFs. The effector and reporter plasmids are co-introduced into Arabidopsis leaf by particle bombardment. Reporter activity is measured to examine whether a TF is an activator or a repressor.

as a transcriptional activator, the AD can be determined by investigating the activities of truncated TF proteins. Activation activity of TFs can also be assessed in a yeast system. However, ADs identified in a plant system can sometimes differ from those identified in a yeast system (Ohta et al. 2000).

Some Arabidopsis TFs are known to act as transcriptional repressors. To analyze whether the TF of interest is a repressor, the repressive activity of the effector construct, using a reporter gene containing a transcriptional enhancer in the promoter, such as that of the CaMV 35S promoter, is utilized. As in the case of an activator, by analyzing the repressive activities of truncated proteins, it is possible to identify the RD of the TF. This strategy for the molecular analysis of TFs is summarized in Fig. 1.

Expression analysis

Expression analysis aims to identify the cells and tissues that express TFs and to define the levels and temporal patterns of expression. As described above, the expression profile of each TF gene has been analyzed for various tissues, under certain conditions and chemical treatments by large-scale microarray analyses (Schmid et al. 2005, Kilian et al. 2007, Goda et al. 2008). Although it has long been believed that the expression levels of TFs are lower relative to those of non-TFs, these large-scale microarray data revealed that this is not always the case. The distribution of signal values was not significantly different between TFs and non-TFs. Although the frequency of non-TF genes that are expressed at extremely high levels is higher than that of TFs, the frequency of genes expressed at low levels is almost the same between TFs and non-TFs (Fig. 2).

Analysis of gene expression by quantitative reverse transcription-PCR (RT-PCR) has become routine, especially in the last decade. However, many concerns about quantitative RT-PCR have recently been pointed out (Czechowski et al. 2005, Gutierrez et al. 2008, Udvardi et al. 2008). In particular, selection of a reference gene(s) is a critical issue. Although traditional housekeeping genes, such as *ACT2*, *TUB6*, *EF-1 α* , *UBQ10* and cytosolic *GAPDH*, are often used as a reference, they are not always uniformly expressed in various tissues (Czechowski et al. 2005). Thus, for quantitative RT-PCR analysis, it is important to select a suitable reference gene.

Promoter-reporter experiments using GUS or green fluorescent protein (GFP) are often employed as a powerful tool to obtain detailed expression profiles. However, it is well known that exogenous promoter activities do not always reflect the genuine expression pattern of a gene of interest due to post-transcriptional regulation and/or lack of necessary gene regulatory elements in the promoter. Overestimation would occur in the case of miRNA target genes. In contrast, underestimation would occur if the promoter region used in the experiment does not contain the *cis*-element(s) required for sufficient expression of the gene. Although promoters of 3 kb or longer are empirically considered to be sufficient to reflect accurate expression in *Arabidopsis*, some genes contain regulatory elements in their introns (Deyholos and Sieburth 2000) or in downstream or upstream regions, at significant distances from the coding region. For the accurate profiling of expression patterns, it is recommended to use the largest genomic fragment possible, containing upstream and downstream regions, in which the GUS or GFP gene should be inserted into the coding region to generate a fusion protein. Such a construct would be expected to reflect an accurate expression pattern, even if the gene is a target of miRNA (Wu and Poethig 2006, Schwarz et al. 2008).

In situ RNA hybridization is a reliable method to examine the gene expression profile at the cellular level.




Fig. 2 Histogram from a large-scale microarray experiment showing distributions of averaged, normalized signals of 21,099 non-TF genes and 2,155 TF genes among a 'developmental set' of genes (Schmid et al. 2005). The distribution is not drastically different between non-TFs and TFs. The frequency of highly expressed genes is higher in non-TFs than in TFs (indicated by a green bracket).

Immunohistochemistry may be more suitable to analyze cells where the protein product actually works, because some TFs, such as SHORT-ROOT (SHR), CPC and KNAT1, are known to move from cell to cell (Nakajima et al. 2001, Kim et al. 2003, Kurata et al. 2005). However, these methods are not suitable for obtaining expression profiles at the organ level or for high-throughput analysis.

Phenotypic analysis

Phenotypic analysis of TFs involves the identification of a phenotype that is regulated by a TF. For this analysis, it is necessary to obtain mutants or transgenic plants with informative phenotypes that lead to the identification of biological function of TFs. As mentioned above, a number of plant TFs act as key regulators and directly regulate various plant functions. Therefore, manipulation of TFs provides the possibility to induce phenotype changes more efficiently than the manipulation of other factors. Phenotypes include not only visible morphological and metabolic changes, but also 'hidden' changes that are only visible under certain conditions. To induce phenotypic changes by manipulating TFs, two strategies, 'gain of function' and 'loss of function', are usually applied. The 'gain-of-function' method induces an abnormal phenotype by the ectopic expression of a TF gene using a constitutive promoter, such as the CaMV 35S promoter, or by enhancing an activating activity of a TF by fusion of an exogenous AD, such as the VP16 AD of herpes simplex virus. Conditional expression of a gene using an inducible promoter or a hormone receptor system (Severin and Schoffl 1990, Takahashi et al. 1992, Aoyama and Chua 1997, Caddick et al. 1998, Zuo et al. 2000) is also a useful

strategy when overexpression of a TF induces lethality. By observing the phenotype induced by ectopic expression, the protein function of a TF can be deduced. For example, ectopic expression of *PAP1*, *AtMYB23* and *NST1* induces accumulation of anthocyanin, excessive trichomes and ectopic secondary wall thickenings, respectively, suggesting that the proteins encoded by these genes have the ability to induce these respective phenotypes (Kirik et al. 2001, Esch et al. 2004, Mitsuda et al. 2005). For systematic gain-of-function analysis in Arabidopsis and rice, the full-length cDNA overexpressor gene hunting system (FOX-hunting) has been applied at a large scale (Ichikawa et al. 2006, Kondou et al. 2009). This gain-of-function assay is easy and simple to apply and is effective for many genes; however, this system does not always reflect the native function of TFs due to ectopic induction. In addition, ectopic expression of a TF often fails to induce an informative phenotype, suggesting that ectopic expression of a single TF might be inadequate to activate the expression of target genes. Other factors that cooperate with the TF may regulate the expression of target genes.

In comparison with gain-of-function analysis, phenotypes induced by loss-of-function analysis should more directly reflect native gene function. For loss-of-function analysis, inactivation of genes or of a gene's activity is necessary. T-DNA-tagged lines of most Arabidopsis genes, in which the large T-DNA fragment is inserted into them, are now available (Alonso et al. 2003). T-DNA-tagged lines for genes of interest can be searched for in the T-DNA express database (<http://signal.salk.edu/cgi-bin/tdnaexpress>) and can be retrieved from the Arabidopsis Biological Resource Center (ABRC) and the European Arabidopsis Stock Centre (NASC). However, a significant number of genes, especially genes with short coding regions, have no tag in the coding region. Furthermore, tagged lines in which a tag is inserted into the 3' side of a coding region or into an intron are frequently not null alleles. Expression of complementary RNA, namely antisense RNA and RNA interference (RNAi) strategies, is frequently adopted for the inactivation of target genes. Furthermore, the use of artificial miRNA (amiRNA) has also been recently proposed (Niu et al. 2006). However, only a small number of transgenic plants show aberrant phenotypes and the phenotype is unstable in many cases.

A major obstacle for loss-of-function approaches is functional gene redundancy. The plant genome has frequently experienced segmental duplication, and Arabidopsis TF genes form large families that share similar DBDs. These domains can have similar DNA-binding specificities, which could act redundantly with other members. Such functional redundancy often interferes with efforts to identify TF functions. To overcome this difficulty, a novel gene silencing system, called Chimeric REpressor Gene-Silencing Technology (CRES-T), was developed (Hiratsu et al. 2003). In the CRES-T system, a chimeric repressor, produced by fusion of a

TF to a modified EAR motif plant-specific repression domain (SRDX), dominantly suppresses the expression of target genes over the activity of endogenous and functionally redundant TFs. As a result, the transgenic plant that expresses the chimeric repressor exhibits a phenotype similar to the loss-of-function phenotype of the TF and of its functionally redundant paralog, even if there are endogenous functionally redundant TFs. It should be noted, however, that an abnormal phenotype induced by the chimeric repressor driven by a constitutive promoter does not always reflect its naive function. For example, the chimeric repressor of a TF driven by a constitutive promoter sometimes induces an aberrant phenotype where the native TF is not expressed. Therefore, the expression of a chimeric repressor, driven by its own promoter, is desirable to investigate the precise biological function of a TF. The CRES-T system functions successfully for various TFs and has succeeded in inducing distinct phenotypes that have not been identified with single gene knockout lines or by the expression of complementary RNA (Kubo et al. 2005, Mitsuda et al. 2005, Ishida et al. 2007, Ito et al. 2007, Koyama et al. 2007, Groszmann et al. 2008, Kawamura et al. 2008, Soyano et al. 2008). In addition, this system is available not only for dicots, such as Arabidopsis, but also for monocots, such as rice (Mitsuda et al. 2006). The advantages of the CRES-T system are that (i) it is simple; (ii) it works in a dominant fashion; and (iii) cloning of the gene of interest is not necessarily required when applied to species for which the genome sequence is unavailable. The CRES-T construct, which is based on Arabidopsis TFs, can work effectively in other species, sometimes without any modification (unpublished result). This is a great benefit for the application of this technology to agronomically important crops. The CRES-T system is also applicable to traditional forward genetic strategies. Screening interesting phenotypes from the seed pool of CRES-T lines would find novel factors because a chimeric repressor works dominantly, even in the presence of functionally redundant TFs.

The CRES-T system is less effective for transcriptional repressors because it is expected that the CRES-T line would have a similar or enhanced phenotype compared with that of ectopic overexpression lines. For loss-of-function analysis of the repressor, the expression of a fusion protein between the repressor and the VP16 AD might be effective. Because a strong RD may overcome the effect of VP16 and the fusion molecule behaves as a repressor (Ohta et al. 2001), the RD should, if possible, be removed from the construct.

Network analysis

Unraveling the entire transcriptional regulatory network is one of the final goals of TF research. Experimental approaches toward this end can be classified into genetic and non-genetic approaches. Traditional genetic approaches, namely analysis of mutant lines, provide high quality information

if mutants or transgenic plants provide informative phenotypes. To find suppressors that reduce the phenotypic abnormality of mutant or transgenic plants is one of the most reliable approaches to identify gene(s) acting downstream of the gene of interest. For example, transgenic plants expressing TCP3SRDX are very small and have a disordered appearance (Koyama et al. 2007). This aberrant phenotype was reduced drastically when crossed to a mutant of *CUC* genes, suggesting that TCP TFs negatively regulate *CUC* genes (Koyama et al. 2007). The *fruitfull* (*ful*) mutant bears a very short silique full of seeds and *FUL* encodes a MADS-box TF (Gu et al. 1998). The phenotype of *ful* can be mostly restored by introducing a mutation into *INDEHISCENT* (*IND*), *SHATTERPROOF1* (*SHP1*) and *SHP2*, suggesting that *FUL* negatively regulates the expression of *IND*, *SHP1* and *SHP2* (Liljegren et al. 2004). The use of a promoter-reporter also helps to reveal genetic relationships. For example, an enhancer trap line of *CUC1* (the expression of GFP is driven by the *CUC1* promoter) showed ectopic fluorescence after the introduction of *35S::TCP3SRDX*, supporting the genetic relationship that TCPs negatively regulate CUCs (Koyama et al. 2007). Promoter activity of *IND* is ectopically expanded in plants overexpressing *SHP1* and *SHP2*, suggesting that *SHP1* and *SHP2* positively regulate the expression of *IND* (Liljegren et al. 2000, Liljegren et al. 2004).

Non-genetic methods, especially large-scale high-throughput experiments, have developed rapidly in recent years. The most successfully applied method to find TFs that act upstream of the TF of interest is yeast one-hybrid screening (Y1H) (Luo et al. 1996). This system uses a tandem repeat of a putative short *cis*-element as bait to screen a cDNA library prepared for Y1H. However, it is laborious to identify a putative short *cis*-element from a promoter region. Another weakness of this system is the cDNA library. Normally, cDNA libraries are not normalized, and include unnecessary non-TF genes, and only one-sixth of the library is successfully fused, in-frame, to the transcriptional AD encoded in the vector. To overcome these problems, a novel approach was introduced, in which a promoter region of <500 bp was directly employed as bait and the cDNA library consisted only of TFs (Deplancke et al. 2006, Pruneda-Paz et al. 2009). A novel component of the circadian system, *CCA1 HIKING EXPEDITION* (*CHE*)/*TCP21* was identified as a direct regulator of *CIRCADIAN CLOCK ASSOCIATED 1* (*CCA1*) by this approach (Pruneda-Paz et al. 2009). These strategies have great potential to contribute to the analysis of gene regulatory networks.

To uncover genes acting downstream of a TF, microarray approaches are the most suitable. The correct statistical analysis of microarray data is critical to exploit the power of this technology. To compare microarray data among two or more samples, cross-chip normalization is required. This normalization procedure normally contains two steps.

Median scaling is a method in which each signal value is divided by the median of all signals within each array. The resulting median signal is scaled to 1 for each array. It should be noted, however, that a recent study reported that 75th percentile scaling is more robust than median scaling (Shippy et al. 2006). Quantile normalization is an optional non-linear conversion of signal values, making signal distribution of all arrays the same (Bolstad et al. 2003). Some data processing software, such as RMA (Irizarry et al. 2003) and GCRMA (Seo et al. 2006) for Affymetrix GeneChip arrays, contain quantile normalization as a default. To select differentially expressed genes (DEGs) between two samples, statistical significance (*P*-value), calculated by the *t*-test, is often used as a cut-off criterion, in combination with a fold change threshold. It should be noted that experiments need to be performed at least in triplicate for effective statistical analysis. Furthermore, multiple testing correction was proposed to be needed to control the false discovery rate (FDR) when selecting DEGs (Dudoit et al. 2002). A stepwise correction of the *P*-value based on *P*-value rank is often used to overcome this problem (Benjamini and Hochberg 1995). Another method has been proposed to transform a *P*-value into a *Q*-value to represent an FDR when the value is used as a threshold (Storey and Tibshirani 2003). One of the basic analyses of microarray data, when comparing two samples, is to find statistically over-represented and under-represented gene groups with specific characteristics among DEGs by statistical tests such as Fisher's exact test. For example, heat-responsive genes were shown to be apparently over-represented among up-regulated genes in a plant overexpressing a constitutively active form of DREB2A, leading to the discovery of cross-talk between the drought stress response, salt stress response and the heat stress response (Sakuma et al. 2006). Microarray experiments are also effective for dissecting distinct roles of related TFs. Rashotte et al. (2006) succeeded in characterizing the precise roles of newly identified CYTOKININ RESPONSE FACTORS (CRFs) by comparing transcriptomes of multiple *crf* mutants with the double mutant of *ARABIDOPSIS RESPONSE REGULATOR 1* (*ARR1*) and *ARR12*. Morohashi and Grotewold (2009) proposed distinct roles for *GLABRA 1* (*GL1*) and *GL3*, which form a heterodimer to regulate trichome initiation.

The identification of direct TF targets plays a central role in unraveling transcriptional regulatory networks. There are several major experimental approaches to achieve this. The TF-binding sequence can be determined using purified protein and random oligonucleotide selection (Wright et al. 1991). It is easy to find genes with determined sequence in their promoters in genome-sequenced species. However, this method is only usable if the TF protein can be purified and the possibility cannot be excluded that the determined sequence does not reflect the actual *in vivo* binding sequence. Furthermore, this method is not suitable for high-throughput

analysis due to the difficulties of protein expression and purification.

The inducible expression of transgenes by a system using a chimeric TF, consisting of a yeast GAL4DB, the transcriptional VP16 AD and the glucocorticoid receptor (GR) or estrogen receptor (ER) has been developed in the last decade (Aoyama and Chua 1997, Zuo et al. 2000). Direct fusion of a TF protein to a GR or ER is sometimes sufficient for TF functional analysis and exploration of TF target genes (Sablowski and Meyerowitz 1998). Expression of downstream genes is rapidly induced after dexamethasone (DEX) or estrogen treatment of the transgenic plant constitutively expressing TF–GR or TF–ER fusion proteins. When the plant is treated simultaneously with DEX and cycloheximide (CHX), which inhibits protein synthesis, expression of the direct TF target gene only is induced. *SUPPRESSOR OF OVER-EXPRESSION OF CO 1 (SOC1)* and *FLOWERING LOCUS T (FT)* were identified as direct targets of CONSTANS (CO) by this approach (Samach et al. 2000). If this system is combined with microarray approaches or next-generation sequencing technology, direct target genes of TFs can be comprehensively identified. However, it should be noted that the expression of some genes involved in the stress response are induced by DEX and/or CHX treatment.

Chromatin immunoprecipitation (ChIP) is a rapidly emerging technology to identify genomic fragments that bind to a DNA-binding protein (O'Neill and Turner 1996). This method is suitable for confirming that a TF binds directly to the promoter region of a putative target gene. It is considered strong evidence if the promoter region of the putative target gene is enriched in the collected pool of genomic fragments. A combination of this technique with whole genome tiling arrays or next-generation sequencing technology (ChIP-chip or ChIP-seq) has great potential to identify comprehensively the region bound by a TF (reviewed in Buck and Lieb 2004, Jiang and Pugh 2009). Some pioneering studies applying ChIP-chip to plant TFs have already been accomplished (Morohashi and Grotewold 2009, Lee et al. 2007). Another approach, utilizing yeast screening, has been performed to identify comprehensively the genomic regions bound by TFs (Zeng et al. 2008).

Recent development of these technologies allows us to identify putative direct and indirect downstream genes of TFs more easily; nevertheless, the contribution of these genes to a plant's phenotype should be carefully examined. The challenge of trying to incorporate this vast amount of information into one huge gene regulatory network has been launched under the umbrella of systems biology. AtRegNet (<http://arabidopsis.med.ohio-state.edu/RGNet/>), hosted in AGRIS, is one of most sophisticated frameworks addressing this issue. It stores information of TF to gene regulations, and users can add original data. In this internet era, 'collective

intelligence' is expected to facilitate the drawing of the entire picture.

Contribution of TF functional analysis to omics studies

Omics studies include the study of various 'omes', such as the transcriptome, proteome, metabolome and phenome. To perform omics studies effectively, it is necessary to prepare useful lines, such as mutant lines, which provide informative phenotypes. In plants, transcriptional regulation plays a major role in the control of gene expression, and a number of plant TFs are known to act as key regulators of various functions. Manipulation of a TF often induces drastic phenotype changes and alters the proteome, metabolome and phenome, in addition to the transcriptome. By clarifying the complete functional network of TFs, it may be possible to predict events in the transcriptome, metabolome and phenome induced by manipulation of a gene of interest. Development of novel innovative tools for functional analysis, in addition to CRES-T, ChIP, microarray analysis and next-generation sequencing, will provide new avenues for the functional analysis of TFs.

Supplementary data

Supplementary data are available at PCP online.

Acknowledgments

We express appreciation to Ms. Yuko Takiguchi for her skillful assistance with computational searches.

References

- Adkins, N.L., Hall, J.A. and Georgel, P.T. (2006) GAGA protein: a multifaceted transcription factor. *Biochem. Cell Biol.* 84: 559–567.
- Allen, M.D., Yamasaki, K., Ohme-Takagi, M., Tateno, M. and Suzuki, M. (1998) A novel mode of DNA recognition by a β -sheet revealed by the solution structure of the GCC-box binding domain in complex with DNA. *EMBO J.* 17: 5484–5496.
- Alonso, J.M., Stepanova, A.N., Leisse, T.J., Kim, C.J., Chen, H., Shinn, P., et al. (2003) Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science* 301: 653–657.
- Aoyama, T. and Chua, N.H. (1997) A glucocorticoid-mediated transcriptional induction system in transgenic plants. *Plant J.* 11: 605–612.
- Arabidopsis Genome Initiative (2000) Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408: 796–815.
- Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., et al. (2000) Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat. Genet.* 25: 25–29.
- Backman, T.W., Sullivan, C.M., Cumbie, J.S., Miller, Z.A., Chapman, E.J., Fahlgren, N., et al. (2008) Update of ASRP: the *Arabidopsis* small RNA project database. *Nucleic Acids Res.* 36: D982–D985.

- Bailey, T.L., Williams, N., Mischak, H. and Li, W.W. (2006) MEME: discovering and analyzing DNA and protein sequence motifs. *Nucleic Acids Res.* 34: W369–W373.
- Barrett, T., Troup, D.B., Wilhite, S.E., Ledoux, P., Rudnev, D., Evangelista, C., et al. (2007) NCBI GEO: mining tens of millions of expression profiles—database and tools update. *Nucleic Acids Res.* 35: D760–D765.
- Benjamini, Y. and Hochberg, Y. (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. R. Stat. Soc. Ser. B* 57: 289–300.
- Bolstad, B.M., Irizarry, R.A., Astrand, M. and Speed, T.P. (2003) A comparison of normalization methods for high density oligonucleotide array data based on variance and bias. *Bioinformatics* 19: 185–193.
- Buck, M.J. and Lieb, J.D. (2004) ChIP-chip: considerations for the design, analysis and application of genome-wide chromatin immunoprecipitation experiments. *Genomics* 83: 349–360.
- Caddick, M.X., Greenland, A.J., Jepson, I., Krause, K.P., Qu, N., Riddell, K.V., et al. (1998) An ethanol inducible gene switch for plants used to manipulate carbon metabolism. *Nat. Biotechnol.* 16: 177–180.
- Craigon, D.J., James, N., Okyere, J., Higgins, J., Jotham, J. and May, S. (2004) NASCArrays: a repository for microarray data generated by NASC's transcriptomics service. *Nucleic Acids Res.* 32: D575–D577.
- Cui, J., Li P., Li G., Xu F., Zhao, C., Li Y., et al. (2008) AtPID: *Arabidopsis thaliana* protein interactome database—an integrative platform for plant systems biology. *Nucleic Acids Res.* 36: D999–D1008.
- Czechowski, T., Stitt, M., Altmann, T., Udvardi, M.K. and Scheible, W.R. (2005) Genome-wide identification and testing of superior reference genes for transcript normalization in *Arabidopsis*. *Plant Physiol.* 139: 5–17.
- Davuluri, R.V., Sun, H., Palaniswamy, S.K., Matthews, N., Molina, C., Kurtz, M., et al. (2003) AGRIS: *Arabidopsis* gene regulatory information server, an information resource of *Arabidopsis* cis-regulatory elements and transcription factors. *BMC Bioinformatics* 4: 25.
- Deplancke, B., Mukhopadhyay, A., Ao W., Elewa, A.M., Grove, C.A., Martinez, N.J., et al. (2006) A gene-centered *C. elegans* protein–DNA interaction network. *Cell* 125: 1193–1205.
- Deyholos, M.K. and Sieburth, L.E. (2000) Separable whorl-specific expression and negative regulation by enhancer elements within the AGAMOUS second intron. *Plant Cell* 12: 1799–1810.
- Döring, P., Treuter, E., Kistner, C., Lyck, R., Chen, A. and Nover, L. (2000) The role of AHA motifs in the activator function of tomato heat stress transcription factors HsfA1 and HsfA2. *Plant Cell* 12: 265–278.
- Dudoit, S., Yang, Y.H., Speed, T.P. and Callow, M.J. (2002) Statistical methods for identifying differentially expressed genes in replicated cDNA microarray experiments. *Stat. Sin.* 12: 111–139.
- Emanuelsson, O., Nielsen, H., Brunak, S. and von Heijne, G. (2000) Predicting subcellular localization of proteins based on their N-terminal amino acid sequence. *J. Mol. Biol.* 300: 1005–1016.
- Ernst, H.A., Olsen, A.N., Larsen, S. and Lo Leggio, L. (2004) Structure of the conserved domain of ANAC, a member of the NAC family of transcription factors. *EMBO Rep.* 5: 297–303.
- Esch, J.J., Chen, M.A., Hillestad, M. and Marks, M.D. (2004) Comparison of TRY and the closely related at1g01380 gene in controlling *Arabidopsis* trichome patterning. *Plant J.* 40: 860–869.
- Furihata, T., Maruyama, K., Fujita, Y., Umezawa, T., Yoshida, R., Shinozaki, K., et al. (2006) Abscisic acid-dependent multisite phosphorylation regulates the activity of a transcription activator AREB1. *Proc. Natl Acad. Sci. USA* 103: 1988–1993.
- Geisler-Lee, J., O'Toole, N., Ammar, R., Provart, N.J., Millar, A.H. and Geisler, M. (2007) A predicted interactome for *Arabidopsis*. *Plant Physiol.* 145: 317–329.
- Goda, H., Sasaki, E., Akiyama, K., Maruyama-Nakashita, A., Nakabayashi, K., Li, W., et al. (2008) The atGenExpress hormone and chemical treatment data set: experimental design, data evaluation, model data analysis and data access. *Plant J.* 55: 526–542.
- Grennan, A.K. (2006) Genevestigator. Facilitating web-based gene-expression analysis. *Plant Physiol.* 141: 1164–1166.
- Groszmann, M., Paicu, T. and Smyth, D.R. (2008) Functional domains of SPATULA, a bHLH transcription factor involved in carpel and fruit development in *Arabidopsis*. *Plant J.* 55: 40–52.
- Gu, Q., Ferrándiz, C., Yanofsky, M. F. and Martienssen, R. (1998) The FRUITFULL MADS-box gene mediates cell differentiation during *Arabidopsis* fruit development. *Development* 125: 1509–1517.
- Guo, A., He K., Liu, D., Bai, S., Gu X., Wei, L., et al. (2005) DATF: a database of *Arabidopsis* transcription factors. *Bioinformatics* 21: 2568–2569.
- Gustafson, A.M., Allen, E., Givan, S., Smith, D., Carrington, J.C. and Kasschau, K.D. (2005) ASRP: the *Arabidopsis* small RNA project database. *Nucleic Acids Res.* 33: D637–D640.
- Gutierrez, L., Mauriat, M., Pelloux, J., Bellini, C. and Van Wuytswinkel, O. (2008) Towards a systematic validation of references in real-time RT-PCR. *Plant Cell* 20: 1734–1735.
- Hall, D.A., Zhu, H., Zhu, X., Royce, T., Gerstein, M. and Snyder, M. (2004) Regulation of gene expression by a metabolic enzyme. *Science* 306: 482–484.
- He J.X., Gendron, J.M., Yang, Y., Li J. and Wang, Z.Y. (2002) The GSK3-like kinase BIN2 phosphorylates and destabilizes BZR1, a positive regulator of the brassinosteroid signaling pathway in *Arabidopsis*. *Proc. Natl Acad. Sci. USA* 99: 10185–10190.
- Heazlewood, J.L., Verboom, R.E., Tonti-Filippini, J., Small, I. and Millar, A.H. (2007) SUBA: the *Arabidopsis* subcellular database. *Nucleic Acids Res.* 35: D213–D218.
- Hibara, K., Karim, M.R., Takada, S., Taoka, K., Furutani, M., Aida, M., et al. (2006) *Arabidopsis* CUP-SHAPED COTYLEDON3 regulates postembryonic shoot meristem and organ boundary formation. *Plant Cell* 18: 2946–2957.
- Higo, K., Ugawa, Y., Iwamoto, M. and Korenaga, T. (1999) Plant cis-acting regulatory DNA elements (PLACE) database: 1999. *Nucleic Acids Res.* 27: 297–300.
- Hirai, M.Y., Sugiyama, K., Sawada, Y., Tohge, T., Obayashi, T., Suzuki, A., et al. (2007) Omics-based identification of *Arabidopsis* Myb transcription factors regulating aliphatic glucosinolate biosynthesis. *Proc. Natl Acad. Sci. USA* 104: 6478–6483.
- Hiratsu, K., Matsui, K., Koyama, T. and Ohme-Takagi, M. (2003) Dominant repression of target genes by chimeric repressors that include the EAR motif, a repression domain, in *Arabidopsis*. *Plant J.* 34: 733–739.
- Hiratsu, K., Mitsuda, N., Matsui, K. and Ohme-Takagi, M. (2004) Identification of the minimal repression domain of SUPERMAN shows that the DLELRL hexapeptide is both necessary and sufficient for repression of transcription in *Arabidopsis*. *Biochem. Biophys. Res. Commun.* 321: 172–178.

- Honma, T. and Goto, K. (2001) Complexes of MADS-box proteins are sufficient to convert leaves into floral organs. *Nature* 409: 525–529.
- Horton, P., Park, K.J., Obayashi, T., Fujita, N., Harada, H., Adams-Collier, C.J., et al. (2007) WoLF PSORT: protein localization predictor. *Nucleic Acids Res.* 35: W585–W587.
- Hua, S. and Sun, Z. (2001) Support vector machine approach for protein subcellular localization prediction. *Bioinformatics* 17: 721–728.
- Ichikawa, T., Nakazawa, M., Kawashima, M., Iizumi, H., Kuroda, H., Kondou, Y., et al. (2006) The FOX hunting system: an alternative gain-of-function gene hunting technique. *Plant J.* 48: 974–985.
- Iida, K., Seki, M., Sakurai, T., Satou, M., Akiyama, K., Toyoda, T., et al. (2005) RARTF: database and tools for complete sets of *Arabidopsis* transcription factors. *DNA Res.* 12: 247–256.
- Ikeda, M. and Ohme-Takagi, M. (2009) A novel group of transcriptional repressors in *Arabidopsis*. *Plant Cell Physiol.* 50: 970–975.
- Irizarry, R.A., Hobbs, B., Collin, F., Beazer-Barclay, Y.D., Antonellis, K.J., Scherf, U., et al. (2003) Exploration, normalization and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4: 249–264.
- Ishida, T., Hattori, S., Sano, R., Inoue, K., Shirano, Y., Hayashi, H., et al. (2007) *Arabidopsis* TRANSPARENT TESTA GLABRA2 is directly regulated by R2R3 MYB transcription factors and is involved in regulation of GLABRA2 transcription in epidermal differentiation. *Plant Cell* 19: 2531–2543.
- Ito, T., Nagata, N., Yoshida, Y., Ohme-Takagi, M., Ma, H. and Shinozaki, K. (2007) *Arabidopsis* MALE STERILITY1 encodes a PHD-type transcription factor and regulates pollen and tapetum development. *Plant Cell* 19: 3549–3562.
- Jiang, C. and Pugh, B.F. (2009) Nucleosome positioning and gene regulation: advances through genomics. *Nat. Rev. Genet.* 10: 161–72.
- Kagaya, Y., Ohmiya, K. and Hattori, T. (1999) RAV1, a novel DNA-binding protein, binds to bipartite recognition sequence through two distinct DNA-binding domains uniquely found in higher plants. *Nucleic Acids Res.* 27: 470–478.
- Kawamura, M., Ito, S., Nakamichi, N., Yamashino, T. and Mizuno, T. (2008) The function of the clock-associated transcriptional regulator CCA1 (CIRCADIAN CLOCK-ASSOCIATED 1) in *Arabidopsis thaliana*. *Biosci. Biotechnol. Biochem.* 72: 1307–1316.
- Kerrien, S., Alam-Faruque, Y., Aranda, B., Bancarz, I., Bridge, A., Derow, C., et al. (2007) IntAct—open source resource for molecular interaction data. *Nucleic Acids Res.* 35: D561–D565.
- Kilian, J., Whitehead, D., Horak, J., Wanke, D., Weinl, S., Batistic, O., et al. (2007) The atGenExpress global stress expression data set: protocols, evaluation and model data analysis of UV-B light, drought and cold stress responses. *Plant J.* 50: 347–363.
- Kim, J.H., Woo, H.R., Kim, J., Lim, P.O., Lee, I.C., Choi, S.H., et al. (2009) Trifurcate feed-forward regulation of age-dependent cell death involving miR164 in *Arabidopsis*. *Science* 323: 1053–1057.
- Kim, J.Y., Yuan, Z. and Jackson, D. (2003) Developmental regulation and significance of KNOX protein trafficking in *Arabidopsis*. *Development* 130: 4351–4362.
- Kim, S.G., Lee, A.K., Yoon, H.K. and Park, C.M. (2008) A membrane-bound NAC transcription factor NTL8 regulates gibberellic acid-mediated salt signaling in *Arabidopsis* seed germination. *Plant J.* 55: 77–88.
- Kim, Y.S., Kim, S.G., Park, J.E., Park, H.Y., Lim, M.H., Chua, N.H., et al. (2006) A membrane-bound NAC transcription factor regulates cell division in *Arabidopsis*. *Plant Cell* 18: 3132–3144.
- Kirik, V., Schnittger, A., Radchuk, V., Adler, K., Hulskamp, M. and Baumlein, H. (2001) Ectopic expression of the *Arabidopsis* AtMYB23 gene induces differentiation of trichome cells. *Dev. Biol.* 235: 366–377.
- Kondou, Y., Higuchi, M., Takahashi, S., Sakurai, T., Ichikawa, T., Kuroda, H., et al. (2009) Systematic approaches to using the FOX hunting system to identify useful rice genes. *Plant J.* 57: 883–894.
- Koyama, T., Furutani, M., Tasaka, M. and Ohme-Takagi, M. (2007) TCP transcription factors control the morphology of shoot lateral organs via negative regulation of the expression of boundary-specific genes in *Arabidopsis*. *Plant Cell* 19: 473–484.
- Kubo, M., Udagawa, M., Nishikubo, N., Horiguchi, G., Yamaguchi, M., Ito, J., et al. (2005) Transcription switches for protoxylem and metaxylem vessel formation. *Genes Dev.* 19: 1855–1860.
- Kurata, T., Ishida, T., Kawabata-Awai, C., Noguchi, M., Hattori, S., Sano, R., et al. (2005) Cell-to-cell movement of the CAPRICE protein in *Arabidopsis* root epidermal cell differentiation. *Development* 132: 5387–5398.
- Larue, C.T., Wen, J. and Walker, J.C. (2009) A microRNA–transcription factor module regulates lateral organ size and patterning in *Arabidopsis*. *Plant J.* 58: 450–463.
- Lee, J., He K., Stolc, V., Lee, H., Figueroa, P., Gao, Y., et al. (2007) Analysis of transcription factor HY5 genomic binding sites revealed its hierarchical role in light regulation of development. *Plant Cell* 19: 731–749.
- Liljegen, S.J., Ditta, G.S., Eshed, Y., Savidge, B., Bowman, J.L. and Yanofsky, M.F. (2000) SHATTERPROOF MADS-box genes control seed dispersal in *Arabidopsis*. *Nature* 404: 766–770.
- Liljegen, S.J., Roeder, A.H.K., Kempin, S.A., Gremski, K., Ostergaard, L., Guimil, S., et al. (2004) Control of fruit patterning in *Arabidopsis* by INDEHISCENT. *Cell* 116: 843–853.
- Long, J.A., Ohno, C., Smith, Z.R. and Meyerowitz, E.M. (2006) TOPLESS regulates apical embryonic fate in *Arabidopsis*. *Science*. 312: 1520–1523.
- Luo, Y., Vijaychander, S., Stile, J. and Zhu, L. (1996) Cloning and analysis of DNA-binding proteins by yeast one-hybrid and one-two-hybrid systems. *Biotechniques* 20: 564–568.
- Matsui, K., Umemura, Y. and Ohme-Takagi, M. (2008) AtMYB2, a protein with a single MYB domain, acts as a negative regulator of anthocyanin biosynthesis in *Arabidopsis*. *Plant J.* 55: 954–967.
- Miao, Y., Laun, T., Zimmermann, P. and Zentgraf, U. (2004) Targets of the WRKY53 transcription factor and its role during leaf senescence in *Arabidopsis*. *Plant Mol. Biol.* 55: 853–857.
- Mitsuda, N., Hiratsu, K., Todaka, D., Nakashima, K., Yamaguchi-Shinozaki, K. and Ohme-Takagi, M. (2006) Efficient production of male and female sterile plants by expression of a chimeric repressor in *Arabidopsis* and rice. *Plant Biotechnol. J.* 4: 325–332.
- Mitsuda, N., Seki, M., Shinozaki, K. and Ohme-Takagi, M. (2005) The NAC transcription factors NST1 and NST2 of *Arabidopsis* regulate secondary wall thickenings and are required for anther dehiscence. *Plant Cell* 17: 2993–3006.
- Morohashi, K. and Grotewold, E. (2009) A systems approach reveals regulatory circuitry for *Arabidopsis* trichome initiation by the GL3 and GL1 selectors. *PLoS Genet.* 5: e1000396.

- Nakajima, K., Sena, G., Nawy, T. and Benfy, P.N. (2001) Intracellular movement of the putative transcription factor SHR in root patterning. *Nature* 413: 307–311.
- Niu, Q.W., Lin, S.S., Reyes, J.L., Chen, K.C., Wu, H.W., Yeh, S.D., et al. (2006) Expression of artificial microRNAs in transgenic *Arabidopsis thaliana* confers virus resistance. *Nat. Biotechnol.* 24: 1420–1428.
- O'Neill, L.P. and Turner, B.M. (1996) Immunoprecipitation of chromatin. *Methods Enzymol.* 274: 189–197.
- Obayashi, T., Hayashi, S., Saeki, M., Ohta, H. and Kinoshita, K. (2009) ATTED-II provides coexpressed gene networks for *Arabidopsis*. *Nucleic Acids Res.* 37: D987–D991.
- Ohta, M., Matsui, K., Hiratsu, K., Shinshi, H. and Ohme-Takagi, M. (2001) Repression domains of class II ERF transcriptional repressors share an essential motif for active repression. *Plant Cell* 13: 1959–1968.
- Ohta, M., Ohme-Takagi, M. and Shinshi, H. (2000) Three ethylene-responsive transcription factors in tobacco with distinct transactivation functions. *Plant J.* 22: 29–38.
- Ori, N., Cohen, A.R., Etzioni, A., Brand, A., Yanai, O., Shleizer, S., et al. (2007) Regulation of LANCEOLATE by miR319 is required for compound-leaf development in tomato. *Nat. Genet.* 39: 787–791.
- Ouellet, F., Overvoorde, P.J. and Theologis, A. (2001) IAA17/AXR3. Biochemical insight into an auxin mutant phenotype. *Plant Cell* 13: 829–842.
- Palatnik, J.F., Allen, E., Wu X., Schommer, C., Schwab, R., Carrington, J.C., et al. (2003) Control of leaf morphogenesis by microRNAs. *Nature* 425: 257–263.
- Parkinson, H., Kapushesky, M., Kolesnikov, N., Rustici, G., Shojatalab, M., Abeygunawardena, N., et al. (2009) ArrayExpress update—from an archive of functional genomics experiments to the atlas of gene expression. *Nucleic Acids Res.* 37: D868–D872.
- Pruneda-Paz, J.L., Breton, G., Para, A. and Kay, S.A. (2009) A functional genomics approach reveals CHE as a component of the *Arabidopsis* circadian clock. *Science* 323: 1481–1485.
- Quevillon, E., Silventoinen, V., Pillai, S., Harte, N., Mulder, N., Apweiler, R., et al. (2005) InterProScan: protein domains identifier. *Nucleic Acids Res.* 33: W116–W120.
- Rashotte, A.M., Mason, M.G., Hutchison, C.E., Ferreira, F.J., Schaller, G.E. and Kieber, J.J. (2006) A subset of *Arabidopsis* AP2 transcription factors mediates cytokinin responses in concert with a two-component pathway. *Proc. Natl Acad. Sci. USA* 103: 11081–11085.
- Ramachandran, V. and Chen, X. (2008) Small RNA metabolism in *Arabidopsis*. *Trends Plant Sci.* 13: 368–374.
- Riano-Pachon, D.M., Ruzicic, S., Dreyer, I. and Mueller-Roeber, B. (2007) PlnTFDB: an integrative plant transcription factor database. *BMC Bioinformatics* 8: 42.
- Riechmann, J.L., Krizek, B.A. and Meyerowitz, E.M. (1996) Dimerization specificity of *Arabidopsis* MADS domain homeotic proteins APETALA1, APETALA3, PISTILLATA and AGAMOUS. *Proc. Natl Acad. Sci. USA* 93: 4793–4798.
- Riechmann, J.L., Heard, J., Martin, G., Reuber, L., Jiang, C., Keddie, J., et al. (2000) *Arabidopsis* transcription factors: genome-wide comparative analysis among eukaryotes. *Science* 290: 2105–2110.
- Robertson, M., Helliwell, C.A. and Dennis, E.S. (2008) Post-translational modifications of the endogenous and transgenic FLC protein in *Arabidopsis thaliana*. *Plant Cell Physiol.* 49: 1859–1866.
- Sablowski, R.W. and Meyerowitz, E.M. (1998) A homolog of *NO APICAL MERISTEM* is an immediate target of the floral homeotic genes *APETALA3/PISTILLATA*. *Cell* 92: 93–103.
- Sakuma, Y., Liu, Q., Dubouzet, J.G., Abe, H., Shinozaki, K. and Yamaguchi-Shinozaki, K. (2002) DNA-binding specificity of the ERF/AP2 domain of *Arabidopsis* DREBs, transcription factors involved in dehydration- and cold-inducible gene expression. *Biochem. Biophys. Res. Commun.* 290: 998–1009.
- Sakuma, Y., Maruyama, K., Qin, F., Osakabe, Y., Shinozaki, K. and Yamaguchi-Shinozaki, K. (2006) Dual function of an *Arabidopsis* transcription factor DREB2A in water-stress-responsive and heat-stress-responsive gene expression. *Proc. Natl Acad. Sci. USA* 103: 18822–18827.
- Samach, A., Onouchi, H., Gold, S.E., Ditta, G.S., Schwarz-Sommer, Z., Yanofsky, M.F., et al. (2000) Distinct roles of CONSTANS target genes in reproductive development of *Arabidopsis*. *Science* 288: 1613–1616.
- Schindler, U., Terzaghi, W., Beckmann, H., Kadesch, T. and Cashmore, A.R. (1992) DNA binding site preferences and transcriptional activation properties of the *Arabidopsis* transcription factor GBF1. *EMBO J.* 11: 1275–1289.
- Schmid, M., Davison, T.S., Henz, S.R., Pape, U.J., Demar, M., Vingron, M., et al. (2005) A gene expression map of *Arabidopsis thaliana* development. *Nat. Genet.* 37: 501–506.
- Schommer, C., Palatnik, J.F., Aggarwal, P., Chetelat, A., Cubas, P., Farmer, E.E., et al. (2008) Control of jasmonate biosynthesis and senescence by miR319 targets. *PLoS Biol.* 6: e230.
- Schwarz, S., Grande, A.V., Bujdosó, N., Saedler, H. and Huijser, P. (2008) The microRNA regulated SBP-box genes *SPL9* and *SPL15* control shoot maturation in *Arabidopsis*. *Plant Mol. Biol.* 67: 183–195.
- Severin, K. and Schoffl, F. (1990) Heat-inducible hygromycin resistance in transgenic tobacco. *Plant Mol. Biol.* 15: 827–833.
- Seo, J., Gordish-Dressman, H. and Hoffman, E.P. (2006) An interactive power analysis tool for microarray hypothesis testing and generation. *Bioinformatics* 22: 808–814.
- Shippy, R., Fulmer-Smentek, S., Jensen, R.V., Jones, W.D., Wolber, P.K., Johnson, C.D., et al. (2006) Using RNA sample titrations to assess microarray platform performance and normalization techniques. *Nat. Biotechnol.* 24: 1123–1131.
- Simon, M., Lee, M.M., Lin, Y., Gish, L. and Schiefelbein, J. (2007) Distinct and overlapping roles of single-repeat MYB genes in root epidermal patterning. *Dev. Biol.* 311: 566–578.
- Soyano, T., Thitamadee, S., Machida, Y. and Chua, N.H. (2008) ASYMMETRIC LEAVES2-LIKE19/LATERAL ORGAN BOUNDARIES DOMAIN30 and ASL20/LBD18 regulate tracheary element differentiation in *Arabidopsis*. *Plant Cell* 20: 3359–3373.
- Storey, J.D. and Tibshirani, R. (2003) Statistical significance for genomewide studies. *Proc. Natl Acad. Sci. USA* 100: 9440–9445.
- Strayer, C., Oyama, T., Schultz, T.F., Raman, R., Somers, D.E., Más, P., et al. (2000) Cloning of the *Arabidopsis* clock gene *TOC1*, an autoregulatory response regulator homolog. *Science* 289: 768–771.
- Szemenyei, H., Hannon, M. and Long, J.A. (2008) TOPLESS mediates auxin-dependent transcriptional repression during *Arabidopsis* embryogenesis. *Science* 319: 1384–1386.
- Takahashi, T., Naito, S. and Komeda, Y. (1992) The *Arabidopsis* *HSP18.2* promoter/*GUS* gene fusion in transgenic *Arabidopsis* plants: a powerful tool for the isolation of regulatory mutants of the heat-shock response. *Plant J.* 2: 751–761.

- Tominaga, R., Iwata, M., Okada, K. and Wada, T. (2007) Functional analysis of the epidermal-specific MYB genes CAPRICE and WEREWOLF in *Arabidopsis*. *Plant Cell* 19: 2264–2277.
- Toufighi, K., Brady, S.M., Austin, R., Ly E. and Provart, N.J. (2005) The botany array resource: e-northern, expression angling and promoter analyses. *Plant J.* 43: 153–163.
- Udvardi, M.K., Czechowski, T. and Scheible, W.R. (2008) Eleven golden rules of quantitative RT–PCR. *Plant Cell* 20: 1736–1737.
- Ueki, S., Lacroix, B., Krichevsky, A., Lazarowitz, S.G. and Citovsky, V. (2009) Functional transient genetic transformation of *Arabidopsis* leaves by biolistic bombardment. *Nat. Protoc.* 4: 71–77.
- Ulmasov, T., Hagen, G. and Guilfoyle, T.J. (1999) Activation and repression of transcription by auxin-response factors. *Proc. Natl Acad. Sci. USA.* 96: 5844–5849.
- Venter, J.C., Adams, M.D., Myers, E.W., Li, P.W., Mural, R.J., Sutton, G.G., et al. (2001) The sequence of the human genome. *Science* 291: 1304–1351.
- Vroemen, C.W., Mordhorst, A.P., Albrecht, C., Kwaaitaal, M.A. and de Vries, S.C. (2003) The CUP-SHAPED COTYLEDON3 gene is required for boundary and shoot meristem formation in *Arabidopsis*. *Plant Cell* 15: 1563–1577.
- Winter, D., Vinegar, B., Nahal, H., Ammar, R., Wilson, G.V. and Provart, N.J. (2007) An ‘electronic fluorescent pictograph’ browser for exploring and analyzing large-scale biological data sets. *PLoS ONE* 2: e718.
- Wright, W.E., Binder, M. and Funk, W. (1991) Cyclic amplification and selection of targets (CASTing) for the myogenin consensus binding site. *Mol. Cell Biol.* 11: 4104–4110.
- Wu G. and Poethig, R.S. (2006) Temporal regulation of shoot development in *Arabidopsis thaliana* by miR156 and its target SPL3. *Development* 133: 3539–3547.
- Yamasaki, K., Kigawa, T., Inoue, M., Tateno, M., Yamasaki, T., Yabuki, T., et al. (2004a) A novel zinc-binding motif revealed by solution structures of DNA-binding domains of *Arabidopsis* SBP-family transcription factors. *J. Mol. Biol.* 337: 49–63.
- Yamasaki, K., Kigawa, T., Inoue, M., Tateno, M., Yamasaki, T., Yabuki, T., et al. (2004b) Solution structure of the B3 DNA binding domain of the *Arabidopsis* cold-responsive transcription factor RAV1. *Plant Cell* 16: 3448–3459.
- Yamasaki, K., Kigawa, T., Inoue, M., Tateno, M., Yamasaki, T., Yabuki, T., et al. (2005a) Solution structure of an *Arabidopsis* WRKY DNA binding domain. *Plant Cell* 17: 944–956.
- Yamasaki, K., Kigawa, T., Inoue, M., Yamasaki, T., Yabuki, T., Aoki, M., et al. (2005b) Solution structure of the major DNA-binding domain of *Arabidopsis thaliana* ethylene-insensitive3-like3. *J. Mol. Biol.* 348: 253–264.
- Yanagisawa, S. and Sheen, J. (1998) Involvement of maize dof zinc finger proteins in tissue-specific and light-regulated gene expression. *Plant Cell* 10: 75–89.
- Yanagisawa, S., Yoo, S.D. and Sheen, J. (2003) Differential regulation of EIN3 stability by glucose and ethylene signaling in plants. *Nature* 425: 521–525.
- Zeng, J., Yan, J., Wang, T., Mosbrook-Davis, D., Dolan, K.T., Christensen, R., et al. (2008) Genome wide screens in yeast to identify potential binding sites and target genes of DNA-binding proteins. *Nucleic Acids Res.* 36: e8.
- Zimmermann, I.M., Heim, M.A., Weisshaar, B. and Uhrig, J.F. (2004a) Comprehensive identification of *Arabidopsis thaliana* MYB transcription factors interacting with R/B-like BHLH proteins. *Plant J.* 40: 22–34.
- Zimmermann, P., Hirsch-Hoffmann, M., Hennig, L. and Gruissem, W. (2004b) GENEVESTIGATOR. *Arabidopsis* microarray database and analysis toolbox. *Plant Physiol.* 136: 2621–2632.
- Zuo, J., Niu, Q.W. and Chua, N.H. (2000) Technical advance: an estrogen receptor-based transactivator XVE mediates highly inducible gene expression in transgenic plants. *Plant J.* 24: 265–273.

(Received April 8, 2009; Accepted May 26, 2009)