

The StkP/PhpP Signaling Couple in *Streptococcus pneumoniae*: Cellular Organization and Physiological Characterization^{∇†}

Makoto Osaki,^{1,2} Tania Arcondéguy,¹ Amandine Bastide,¹ Christian Touriol,¹
Hervé Prats,¹ and Marie-Claude Trombe^{1*}

INSERM, U858, Toulouse, F-31432 France, and IFR31 Université Toulouse III Paul-Sabatier, Toulouse, France,¹ and
National Institute of Animal Health, National Agriculture and Food Research Organization,
Tsukuba, Ibaraki 305-0856, Japan²

Received 13 February 2009/Accepted 28 May 2009

In *Streptococcus pneumoniae*, *stkP* and *phpP*, encoding the eukaryotic-type serine-threonine kinase and PP2C phosphatase, respectively, form an operon. PhpP has the features of a so-called “soluble” protein, whereas StkP protein is membrane associated. Here we provide the first genetic and physiological evidence that PhpP and StkP, with antagonist enzymatic activities, constitute a signaling couple. The StkP-PhpP couple signals competence upstream of the competence-specific histidine kinase ComD, receptor for the oligopeptide pheromone “competence stimulating peptide.” We show that PhpP activity is essential in a *stkP*⁺ genetic background, suggesting tight control of StkP activity by PhpP. Proteins PhpP and StkP colocalized to the cell membrane subcellular fraction and likely belong to the same complex, as revealed by coimmunoprecipitation in cellular extracts. Specific coimmunoprecipitation of the N-kinase domain of StkP and PhpP recombinant proteins by PhpP-specific antibodies demonstrates direct interaction between these proteins. Consistently, flow cytometry analysis allowed the determination of the cytoplasmic localization of PhpP and of the N-terminal kinase domain of StkP, in contrast to the periplasmic localization of the StkP C-terminal PASTA (penicillin-binding protein and serine-threonine kinase associated) domain. A signaling route involving interplay between serine, threonine, and histidine phosphorylation is thus described for the first time in this human pathogen.

Signal transduction via the transfer of phosphoryl groups and transient protein phosphorylation involves the concerted activities of kinases and phosphatases and controls various cellular functions (for a review, see reference 18). In eubacteria, the histidine kinases of two-component systems (TCSs) and the phosphatases with which they interact are involved in cellular adaptation to environmental conditions (for a review, see reference 44). Serine-threonine kinases and PP2C phosphatases also contribute to regulatory and developmental processes, as described in *Bacillus subtilis* (1, 2, 13, 23, 29, 33, 43). In mycobacteria, a membrane-associated PP2C phosphatase controls the activity of several serine-threonine kinases (4, 6, 52), and the transcriptional activator EmbR has been identified as a phosphorylation target for PknH kinase (3, 4, 42). In *Streptococcus agalactiae*, the signaling pair Stk1-Stp1 plays a role in virulence, by modifying cytotoxin production and purine metabolism (34, 35, 36). In *Streptococcus pyogenes*, a histone-like protein is phosphorylated by the serine-threonine kinase, generating a substrate for the kinase-related phosphatase (21). The human pathogen *Streptococcus pneumoniae* has only one gene encoding a PP2C-type phosphatase, PhpP, located upstream from the *stkP* gene, which encodes the only membrane-associated serine-threonine kinase, StkP (17). The *phpP* and *stkP* genes overlap by 4 bp and form an operon (31). In a mouse model of infection, null mutations

affecting StkP greatly attenuate tissue and bloodstream invasion (17). In cultures, these mutations are highly pleiotropic, presenting a notably important impact on competence development for genetic transformation (17, 40). In vitro studies have shown that autophosphorylated recombinant StkP (StkP-P) is a substrate for recombinant PhpP, suggesting that StkP and PhpP may function in a coordinated manner (31). However, the regulators controlling the cellular level of StkP-P and the network involving StkP in growing cultures remain ill defined. Recent studies of strain TIGR4 revealed that the transcriptional regulator RitR (45) is phosphorylated by StkP and dephosphorylated by PhpP (50). In order to get better insight into StkP signaling, we investigated the specific role of PhpP phosphatase on bacterial growth and on the development of competence for genetic transformation in cultures, by using genetic analysis. Specific null mutation in PhpP was not obtained, indicating the essentiality of the phosphatase in the control of StkP activity. It is thus likely that PhpP and StkP constitute a functional couple. Data on cellular organization and coimmunoprecipitation suggest that PhpP and StkP proteins can interact within a submembrane complex facing the cytoplasm. Furthermore, results from physiological and genetic studies demonstrate that the StkP-PhpP couple belongs to the competence-signaling network, upstream of the ComDE TCS target for the competence stimulating peptide pheromone, CSP.

(Part of this work was presented at the Europneumo meeting, Oeiras, Portugal, April 2007; at the Gram+ meeting, Pisa, Italy, June 2007; and at the Experimental Workshop on Signal Transduction in Host-Bacterial Interactions, Jerusalem, Israel, October 2007.)

* Corresponding author. Mailing address: INSERM, U858, Toulouse, F-31432 France. Phone: 33(0)561322974. Fax: 33(0)561322620. E-mail: Marie-Claude.Trombe@inserm.fr.

† Supplemental material for this article may be found at <http://jbb.asm.org/>.

∇ Published ahead of print on 5 June 2009.

TABLE 1. Strains and plasmids

Strain or plasmid	Relevant genotype	Phenotype	Source or reference
Strain			
<i>E. coli</i>			
XL1-Blue	F ['] ::Tn10 <i>proA</i> ⁺ <i>B</i> ⁺ <i>lacI</i> ^q Δ(<i>lacZ</i>) <i>M15/recA1 endA1 gyrA96</i> (Nal ^r) <i>thi</i> <i>hsdR17</i> (r _K ⁻ m _K ⁺) <i>supE44 relA1 lac</i>		Stratagene
BL21(DE3)	F ⁻ <i>ompT gal [dcm] [lon] hsdS_B</i> (r _B ⁻ m _B ⁻) (DE3)		Novagen
Top10	F ⁻ <i>mcrA</i> Δ(<i>mrr-hsdRMS-mcrBC</i>) φ80 <i>lacZ</i> Δ <i>M15</i> Δ <i>lacX74 recA1</i> <i>araD139</i> Δ(<i>ara leu</i>)7697 <i>galU galK rpsL</i> (StrR) <i>endA1 nupG</i>		Invitrogen
<i>S. pneumoniae</i>			
Cp1015	RX derivative; <i>str1 hexA</i>	Sm ^r	30
Cp1016	Cp1015, but <i>rif23</i> and <i>stgF</i>	Rif ^r , Stg ^r	5
R119	R6 <i>hex</i> , but <i>rif23 stgF str1 fusA nov1 opt12 gln tdr</i>	Sm ^r , Fus ^r , Rif ^r , Stg ^r , Nov ^r , Opt ^r	46
Cp1008	Cp1015, but <i>comA::spc</i>	Sp ^r	30
Cp1095	Cp1015, but <i>lytA::ery</i>	Ery ^r	47, 49
Cp7000	Cp1015, but Δ <i>stkP::cat</i>	Cam ^r	31
Cp9000	Cp1015, but <i>stkP::aphA-3</i>	Km ^r	17
Cp9090	Cp1095, but <i>stkP::aphA-3</i>	Km ^r , Ery ^r	17
Cp7004	Cp7000, but <i>phpPD1</i>	Cam ^r	This work
Cp9091	Cp9090, but <i>stkP</i> ⁺	Kan ^s	This work
Cp6600	Cp1015, but <i>comD</i> (D299N) <i>rif23</i>	Rif ^r	14
Cp6607	Cp6600, but Δ <i>stkP::cat</i>	Rif ^r , Cam ^r	This work
Plasmid			
pCR2.1TOPO	Cloning vector	Amp ^r , Km ^r	Invitrogen
pWSK29	Cloning vector derived from pBluescript	Amp ^r	51
pET28b	His tag expression vector; Kan ^r , f1 <i>ori</i>	Km ^r	Novagen
pBluescript ISK ⁺ /KS ⁺	Amp ^r , <i>lacZ</i> , f1 ± <i>ori</i> , ColE1 <i>ori</i>	Amp ^r	Stratagene
pPJ1	pUC derivative containing a HincII fragment carrying the kanamycin resistance gene	Amp ^r , Km ^r	32
pPHK29	2.96-kb EcoRI/SalI amplicon (primers PHL and PCR) from <i>S. pneumoniae</i> Cp1015 DNA containing <i>phpP</i> and <i>stkP</i> genes cloned into pWSK29	Amp ^r	17
pPK29	2.3-kb EcoRI/SalI amplicon (primers PCL and PCR) from <i>S. pneumoniae</i> Cp1015 DNA containing <i>stkP</i> gene in pWSK29	Amp ^r	This work
pEXstkP-N	0.825-kb NdeI/EcoRI amplicon (primers STKP-F and STKP-RT) from pPHK29 containing fragment (kinase domain) of <i>stkP</i> gene inserted into pET28b	Km ^r	This work
pEX-PhpP	0.74-kb amplicon (<i>phpP</i>) inserted into pET28b	Km ^r	31
pEX-PASTA	0.9-kb NcoI/EcoRI amplicon (primers STKP-PF and STKP-R2) from pPHK29 containing <i>stkP</i> gene inserted into pET28b	Km ^r	This work
pMUT1	2.6-kb amplicon of <i>phpPD1 stkP</i> ⁺	Km ^r	This work
pMUT2	2.6-kb amplicon of <i>phpPD2 stkP</i> ⁺	Km ^r	This work
pMUT3	2.6-kb amplicon of <i>phpPD1 ΔstkP::cat</i>	Km ^r	This work
pMUT4	2.6-kb amplicon of <i>phpPD2 ΔstkP::cat</i>	Km ^r	This work

MATERIALS AND METHODS

Bacterial strains and growth conditions. The *Streptococcus pneumoniae* strains described in Table 1 were grown in casein tryptone medium (CAT), and the cultures were stored at -80°C in 12% (vol/vol) glycerol. For competence studies, bacteria from frozen cultures were grown in CAT transformation medium (CTM), made from CAT medium supplemented with 0.2% (wt/vol) bovine serum albumin (BSA) and 0.1 mM CaCl₂. The pH was then adjusted to 6.5 or 8 (5, 9, 49). *S. pneumoniae* transformants were selected on appropriate antibiotics: kanamycin, 50 mg liter⁻¹; rifampin (rifampicin), 2 mg liter⁻¹; and chloramphenicol, 10 mg liter⁻¹. The *Escherichia coli* strains described in Table 1 were propagated in Luria broth in the presence of appropriate antibiotics: ampicillin, 100 mg liter⁻¹; kanamycin, 50 mg liter⁻¹; and rifampin, 400 mg liter⁻¹.

Construction of mutagenic plasmids and mutant strains. *Escherichia coli* strains XL1-Blue and Top10 were used as hosts for recombinant vectors (Table 1). Isogenic mutant strains of *S. pneumoniae* were generated by allelic exchange. Briefly, the *S. pneumoniae* recipient strain was transformed with nonreplicative chimeric plasmids carrying the relevant pneumococcal allele(s). In the absence of direct selection with antibiotics, 40 clones were selected and the chromosomal DNA of each clone was extracted with Instagene Matrix (Bio-Rad) for diagnostic

PCR and restriction analysis of the corresponding amplicon (5). One clone carrying the relevant mutation was verified by DNA sequencing with the primers listed in Table 2 and was maintained under standard growth conditions. The insertion/deletion mutation of *stkP* has been described elsewhere (31) and was used to construct strain Cp7000, carrying the Δ*stkP::cat* allele. Plasmids pMUT1 to -4 were used as DNA donors in transformation of the wild-type strain Cp1015, in order to obtain the single mutants *phpPD1* and *phpPD2* and the double mutants *phpPD1 stkP::cat* and *phpPD2 stkP::cat*. The *phpPD1* allele containing the point mutation D241A and a StuI restriction site was associated with *stkP*⁺ or Δ*stkP::cat* DNA fragments, respectively, and the *phpPD2* allele carrying only the StuI restriction site was associated with either *stkP*⁺ or Δ*stkP::cat* DNA fragments to create plasmids pMUT1 to -4 (Table 1). Plasmids pMUT1 to -4 were constructed using standard protocols (39). The 5' region of *phpP* was amplified from the genomic DNA of Cp1015 with the UFKFP and MUT4 primers, and the 3' region of *phpP* was amplified with the MUT2 and DSKR primers. The amplified fragments were mixed and fused with UFKFP and DSKR by a second PCR. The fused fragment was inserted into pCR2.1TOPO to yield pMUT1. Similarly, a DNA fragment carrying only the StuI restriction site was amplified by PCR from the genomic DNA of Cp1015, using the UFKFP-plus-

TABLE 2. Primers

Name	Sequence	Unique restriction site
DSKR	5'-CGGATAAAATTTTCCACATAGAGG	
UFKFP	5'-CGCAAGATATCGGATTAGGAA	
MUT2	5'-GCAGGAGGCCTAGCCAACATTACG	StuI
MUT4	5'-AATGTTGGCTAGGCCTCCTGC	StuI
MUT3	5'-GCAGGAGGCCTAGACAACATTACG	StuI
MUT5	5'-AATGTTGTCTAGGCCTCCTGC	StuI
DFKRP	5'-CGCGGATCCTCATAATATCACGGA CCGCAT	
stkP-PF	5'-AGGATGCCATATGAGATCTCCTGC AACCATTGCCATT	NdeI
stkP-R2	5'-TTGATTATGAATTCGCTTTTAAGGA GTAGCTGAAGTTG	EcoRI
STKP-RT	5'-GTAGGACAGAATTCAAGACAAGTC TACATAA	EcoRI
STKP-F	5'-AGGATGCCATATGATCCAAATCG GCAA	NdeI
PCL	5'-CCAGCTGAATCCAGGCAGTGAGAT TCGCTG	EcoRI
PCR	5'-CCATACGTCGACCGGCATTTAATT CTTTTG	Sall

MUT3 and MUT5-plus-DSKR primer combinations, to yield pMUT2. Similarly, using Cp7000 genomic DNA as a template and primers MUT2 to -5, UFKFP, and DFKRP, we constructed pMUT3 and pMUT4. The sequences of all mutated DNA fragments were verified using universal primers and specific primers of the generated sequences (Table 2).

Production of specific rabbit antibodies. Specific antibodies against PhpP phosphatase, the kinase domain (StkP-N), and the PASTA (penicillin-binding protein and serine-threonine kinase associated) domain of StkP were produced by Eurogentec (France), using the corresponding recombinant proteins produced in *E. coli*. Recombinant proteins were obtained as oligohistidine fusions in the *E. coli* BL21(DE3) strain (Table 1) and purified on TALON metal affinity resin (Clontech) according to the manufacturer's protocol. Antibodies that rose against PhpP or the kinase or PASTA domain and the corresponding preimmune sera were purified on Tgel (Pierce) according to the manufacturer's instructions.

Cellular fractionation and localization of StkP and PhpP. Cultures of the wild-type strain and the StkP null mutant were collected by centrifugation at 4°C, washed with phosphate-buffered saline, resuspended in phosphate-buffered saline, and sonicated. Subcellular fractions were obtained by centrifugation (100,000 × *g* for 1 h), resulting in a pellet fraction containing cellular envelopes and in a supernatant fraction containing the cytoplasm. Proteins (10 µg) from each fraction were subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis, blotted onto nitrocellulose, probed with specific antibodies, and detected with an ECL system (Amersham). Specific membrane localization of StkP has already been demonstrated (31) and has been considered the fractionation control. Nox immunoblots have been used as the loading control, because this protein is found in both the membrane and the cytoplasmic fractions (5; also our unpublished data).

Immunoprecipitation. For cellular extract analysis, clarified lysates (500 µg of proteins in lysis buffer: 50 mM Tris-HCl, pH 8, 1% Igepal, and antiprotease cocktail from Roche) were incubated in the presence or absence of PhpP-specific antibodies (2 µg) for 18 h at 4°C. Recombinant proteins (2 µg in 150 µl of lysis buffer containing 0.5% [wt/vol] BSA) were incubated during 90 min at 4°C before the addition of 20 µg of PhpP antibodies or the corresponding preimmune serum and incubation at 4°C under gentle agitation was allowed during 18 h. Protein A/G-coated Bio-Ademabeads were subsequently added and used as recommended by the supplier (Ademtech), by performing washing steps with the lysis or alternatively the RipA buffer from Ademtech. Elution was obtained in loading buffer in denaturing and reducing conditions. Rabbit preimmune serum was used as a specificity control of immunoprecipitation, and nonspecific binding to the Bio-Ademabeads was evaluated in antibody-free assays when indicated. Total samples and the fraction eluted from the Bio-Ademabeads were loaded onto a 4 to 20% gradient gel and analyzed by Western blotting with antibodies raised against the recombinant PASTA and N-kinase domain proteins.

Flow cytometry analysis. Bacteria and protoplasts were prepared from fresh cultures in CAT medium at an optical density at 400 nm (OD₄₀₀) of 0.2. Proto-

plasts were obtained by treating exponentially growing cultures for 30 min at room temperature with 500 mg liter⁻¹ of lysozyme, 20% (wt/vol) sucrose, 0.1 M Tris, and 10 mM MgSO₄ at pH 7, as described elsewhere (48). Where required, permeabilization of the bacteria was established by treatment of a suspension of heat-killed bacteria (60°C for 30 min) with a Cytofix/Cytoperm kit from BD Biosciences, according to the manufacturer's instructions. Cell permeability was evaluated by assessing the chromosome labeling induced by incubation with fluorescent propidium iodide. For flow cytometry analysis, suspensions of 1 × 10⁷ bacteria and protoplasts in 100 µl of 0.1 M Tris, 20% (wt/vol) sucrose, and 10 mM MgSO₄ at pH 7 were incubated for 30 min at 4°C with antibodies (5 µg/ml) raised against the kinase domain, the PASTA domain, and PhpP phosphatase. Cells and protoplast suspensions were washed three times with 1 ml of Tris-sucrose-MgSO₄ buffer and subsequently incubated for 30 min in the dark at 4°C with goat anti-rabbit-specific immunoglobulin G labeled with phycoerythrin. Two washes were performed with a 1-ml solution of Tris, sucrose, and MgSO₄, and fluorescence was assessed in a Becton Dickinson FACSCalibur spectrofluorimeter (excitation at 488 nm and analysis at 585 nm).

Competence measurements. The level of competence development in cultures was evaluated using the "transformation test," as described elsewhere (14, 30). In brief, cultures in CTM rendered either acidic or alkaline (9) were incubated at 37°C. At an OD of 0.1 to 0.2, 100-µl aliquots were withdrawn and mixed with the DNA from strain Cp1016 or R119 (1 µg ml⁻¹) used for transformation (Table 1). Cells were incubated for 20 min at 20°C, mixed with blood agar medium, and incubated for 2 h at 37°C. A layer of agar with or without selective antibiotics was then poured over the surface of the plate. Colonies were counted after overnight incubation at 37°C, and transformation efficiencies were calculated as the ratio of resistant transformed colonies to total viable counts. For CSP-induced competence, cultures in the early exponential growth phase (OD₄₀₀ of 0.05) in CTM were activated by incubation with 0.1 to 100 ng/ml of CSP for 10 min at 37°C. Competence was assessed with the "transformation test," as described above.

Mutation rate determination. The mutation rates of strains Cp1015 and Cp7000 were evaluated with the P0 test of Luria and Delbruck (27). For each strain, 50 independent 1-ml cultures in CAT medium at pH 7 with a starting inoculum of 1 × 10² CFU (*N*₀) and containing zero mutants resistant to 2 µg/ml of rifampin were cultured for 8 h at 37°C. At the end of the growth period, the total CFU content of the culture (*N*) was determined by plating serial dilutions from one culture of each strain and by plating the other cultures on blood agar supplemented with rifampin. The frequency of cultures containing 0 CFU on rifampin plates was determined (*P*₀) and the mutation frequency (μ) was calculated using the following equation: $\mu = -\ln [P_0/(N - N_0)]$.

Genes *stkP/PhpP* were given NCBI registration number AF285441 on 7 July 2000. They are named *pkn2/spr1577* (and *phpP* is named *pppL/spr1578*) in the R6 genome databases.

RESULTS

The essential nature of PhpP. Biochemical studies of recombinant proteins have shown that phosphorylated StkP is a substrate for PhpP (31), although the cellular function of PhpP has not been elucidated. We carried out a mutational analysis to specifically assess the physiological role of PhpP phosphatase in *S. pneumoniae*. Attempts to construct an in-frame deletion of the *phpP* gene were unsuccessful, a result reminiscent of previous observations in *S. pyogenes* (21). In contrast, *phpP* mutants carrying cassette insertion and stop codon mutations in *phpP* leading to a downstream polar effect on *stkP* could be obtained (data not shown). These results suggested that functional PhpP was probably essential in the wild-type but not in the StkP⁻ genetic background. In order to test this hypothesis, the wild-type strain Cp1015 was transformed with pMUT1 to -4 plasmids encoding either D231A PhpP (*phpPD1*), which has been shown to have no phosphatase activity in vitro (31), or a functional D231D PhpP (*phpPD2*) protein, associated with both the wild-type *stkP* and Δ *stkP::cat* allele, in the vicinity of a newly created StuI restriction site (see Materials and Methods). Transformed clones resulting in allelic exchange ("transformants") harboring the StuI restriction site were detected and quantified. Transformants carrying the *phpPD2* allele in

TABLE 3. Allele *phpPD1* encoding inactive D231A PhpP phosphatase is lethal in the RX *stkP*⁺ genetic background^a

Donor DNA	Transformed colonies for the relevant allele (% CFU)		
	<i>phpP</i>	<i>cat</i>	<i>rif23</i>
pMUT1 <i>phpPD1</i> <i>stkP</i> ⁺	0 (40)		
pMUT2 <i>phpPD2</i> <i>stkP</i> ⁺	75 (40)		
pMUT3 <i>phpPD1</i> Δ <i>stkP::cat</i>	91 (35)	100 (32)	
pMUT4 <i>phpPD2</i> Δ <i>stkP::cat</i>	97 (40)		
R119 chromosomal DNA <i>rif23</i>			3

^a The recipient bacteria was Cp1015 in all cases. Parentheses indicate the number of colonies submitted to diagnostic PCR and *StuI* restriction (CFU). Cam, chloramphenicol; Rif, rifampin (both antibiotics were used at 2 μ g/ml).

addition to *StuI* were obtained with high frequency, regardless of the wild-type or mutant nature of the associated *stkP* allele on plasmids pMUT2 and pMUT4 (Table 3). In contrast, *phpPD1* transformants were exclusively recovered from the transformations involving the pMUT3 plasmid carrying the Δ *stkP::cat* allele. Moreover, all these transformants were also resistant to chloramphenicol, demonstrating cotransformation of *StuI* and Δ *stkP::cat* mutations (Table 3). These results provide the genetic demonstration that *phpPD1* encoding the D231A PhpP protein is lethal, specifically in the *stkP*⁺ genetic background, suggesting that the phosphatase activity of PhpP is essential in wild-type but not in *StkP* mutant bacteria. Our data imply that uncontrolled cellular levels of *StkP*-P are toxic in *S. pneumoniae* RX derivatives (Table 1). During bacterial growth, PhpP phosphatase may therefore play a key role in controlling phosphorylation levels of *StkP* and of its targets. This hypothesis incited us to assess the cellular organization of the membrane-associated serine-threonine kinase and the "so-called" soluble PhpP phosphatase in order to get further insight into the cellular role of the PhpP phosphatase with regard to *StkP* kinase.

The kinase domains of *StkP* and *PhpP* proteins interact within a submembrane complex. The membrane localization of *StkP* protein has been demonstrated elsewhere. In contrast, *PhpP* protein showed the features of a soluble protein (31). Western blot analysis of cellular envelopes and cytoplasmic subcellular fractions from the wild-type strain Cp1015, using specific antibodies directed against *StkP* or *PhpP* proteins, revealed that *PhpP* was mainly present in the envelopes in a manner similar to *StkP*. In contrast, in extracts from the *stkP* deletion strain Cp7000, *PhpP* was mainly present in the cytoplasmic fraction (Fig. 1). Cellular distribution of the Nox protein present both in the envelopes and in the cytoplasm (5; also our unpublished data) was not altered due to *StkP* mutation (Fig. 1), indicating that *StkP* protein affects the distribution of *PhpP* protein selectively. These results raised the hypothesis that *StkP* and *PhpP* may form a protein complex in living bacteria. This hypothesis was tested by coimmunoprecipitation experiments. Cleared bacterial lysates from exponential growing cultures of the wild-type and *stkP* mutant strains were incubated with antibodies directed against *PhpP* or the corresponding preimmune serum as indicated in Materials and Methods. Analysis of the immunocomplexes by Western blotting using *StkP*-specific antibodies clearly revealed a band at 72 kDa, corresponding to the *StkP* protein (calculated molecular

mass of 72.4 kDa), specifically in the *PhpP* antibody assay. This result indicates coimmunoprecipitation of *StkP* protein by *PhpP* antibodies in cellular extracts (Fig. 2A). In a parallel experiment, we observed coimmunoprecipitation of the recombinant *PhpP* and N-kinase domain but not the C-terminal PASTA domain proteins by *PhpP*-specific antibodies, suggesting direct and specific interaction between the kinase module of *StkP* and *PhpP* proteins (Fig. 2B; also data not shown). Cellular localization of the proteins by flow cytometry analysis using suspensions of whole bacteria, protoplasts, and permeabilized cells (see Materials and Methods) provides credence to this hypothesis. The results presented in Fig. 3 show, on the one hand, that in the wild-type strain, the C-terminal PASTA domain of *StkP* can be efficiently labeled in protoplasts as well as in permeabilized cells, whereas the N-terminal kinase domain of *StkP* and *PhpP* proteins were labeled specifically in permeabilized cells. On the other hand, labeling efficiencies are very low in nonpermeabilized cells, suggesting that antibodies were hampered by the cell wall. These results suggest that the C-terminal PASTA domain of *StkP* sticks out in the periplasmic space, beneath the peptidoglycan cell wall, and the kinase domain is located in the cytoplasm, like *PhpP* is. Such a colocalization of the *StkP* kinase domain and *PhpP* proteins is consistent with the proposal of *PhpP* and its kinase module target being organized close together and interacting within a membrane-associated protein complex facing the cytoplasm. However, localization of *PhpP* protein in the cytoplasmic fraction of *StkP* mutant strains (Fig. 1) raises the question of the *StkP*-independent activity of *PhpP* accounting for their phenotype (17, 40).

Physiological characterization of the *StkP*-*PhpP* signaling system. The loss-of-function mutation in *StkP* kinase is pleiotropic, and we have demonstrated a strong reduction of competence expression in cultures from *stkP* mutant strains (17). In brief, competence regulation involves signalization in response to the peptide pheromone CSP targeting the histidine kinase receptor of the TCS ComDE (19). The specific role of the




FIG. 1. Colocalization of *StkP* and *PhpP* in wild-type *S. pneumoniae*. (A) Cytoplasm (Cyto) and envelopes (Env) from sonicated bacteria were obtained as described in Materials and Methods. Proteins (10 μ g) of each fraction from the wild-type strain Cp1015 (lanes 1 and 3) and the Δ *stkP::cat* mutant strain Cp7000 (lanes 2 and 4) were analyzed by Western blotting with kinase- and phosphatase-specific antibodies. Nox immunoblots were taken as controls. (B) Quantification of the immunoreactive material corresponding to *StkP* and *PhpP* in extracts from the wild-type strain Cp1015 has been obtained by reference to Nox-immunoreactive material. The data are representative of two independent experiments.




FIG. 2. Coimmunoprecipitation of StkP and PhpP proteins. (A) Cellular lysates from the wild-type strain Cp1015 and the Δ stkP::cat mutant strain Cp7000 (Table 1), containing 500- μ g proteins, were incubated with PhpP-specific antibodies (2 μ g) or their corresponding preimmune serum in 150 μ l of lysis buffer. The immunocomplexes were analyzed by Western blotting with N-kinase-specific antibodies as described in Materials and Methods. The “Beads” control was obtained in the absence of PhpP antibodies. The total load represents 0.4% of the assay. The data are representative of five independent experiments. (B) Recombinant PhpP and N-kinase domain proteins (2 μ g) were mixed in 150 μ l of lysis buffer supplemented with BSA [0.5% (wt/vol)] and incubated with PhpP antibodies (20 μ g) or their corresponding preimmune serum. Controls were obtained with single protein assays containing either recombinant N-kinase protein or recombinant PhpP protein. Twenty percent of each sample was used for Western blot analysis with N-kinase domain-specific antibodies. The data are representative of two independent experiments. Ig, immunoglobulin; WT, wild type; IP, immunoprecipitation; r, recombinant; WB, Western blot.

phosphatase in this regulation has been evaluated by genetic dissection, using the isogenic strains Cp7000, Cp9090, Cp1095, and Cp7004 carrying the mutated alleles Δ stkP::cat, stkP::Apha3, stkP⁺, and Δ stkP::cat plus phpPD1, respectively (Table 1). Strain Cp9091 carrying a genetic reversion of stkP::Apha3 into stkP⁺ (Table 1) was added in the study because loss-of-function mutation in StkP has been related to a fivefold-increased mutation rate of the mutated—compared to the wild-type—strain and might culminate in genetic drift. Indeed, the calculated rates of mutation (27) conferring resistance to rifampin, in two independent experiments (see Materials and Methods), were 9.9×10^{-7} and 8.1×10^{-7} in the stkP mutant strain and 2.1×10^{-7} and 1.4×10^{-7} in the wild-type strain Cp1015.

The results presented in Table 4 show that a competence defect resulting from a loss-of-function mutation in StkP is not suppressed by the additional mutation PhpP(D231A) in strain




FIG. 3. Cellular organization of StkP kinase and PhpP phosphatase. Flow cytometry was performed with bacterial suspensions from exponentially growing cultures of the wild-type strain Cp1015. Nonpermeabilized and permeabilized bacteria as well as protoplast suspensions were analyzed as described in Materials and Methods, using specific antibodies directed against PhpP and the N-kinase (“Kinase”) and C-PASTA (“PASTA”) domains of StkP protein. Bacteria labeling was obtained as described in Materials and Methods. Nonspecific labeling was evaluated by measurements on the Δ stkP::cat mutant strain Cp7000 and represents $3.5\% \pm 5\%$ of total binding on wild-type bacteria. Specific labeling of the wild-type bacteria was the difference between total labeling and nonspecific labeling. In each run, 10,000 cells were analyzed. Rates of permeability to propidium iodide were 5% for bacterial suspensions, 40% for protoplast suspensions, and 95% for permeabilized bacteria. The data are representative of two independent experiments.

Cp7004 but is reversed due to allelic exchange at the *stkP*::Apha3 locus in the *stkP*⁺ strain Cp9091 (Table 1). Taken together, our results for strains Cp7004 and Cp9091 demonstrate that the transformation-negative phenotype of strain Cp9090 (17) is a direct result of the StkP loss-of-function mutation rather than a consequence of the activity of PhpP in the cytoplasm of *stkP* mutant cells. Therefore in this model, the phenotype of StkP mutant strains is not related to StkP-independent PhpP activity but to StkP loss-of-function mutation. It is thus likely that the dedicated PhpP target, StkP-P, is required for the trigger of competence.

It has already been established that the expression of competence in growing cultures represents a global response to several signals transduced by TCSs ComDE, CiaRH, and MicAB (10, 11, 14, 15, 16, 17, 25, 41). We further investigated the role of the StkP-PhpP couple in the competence-signaling network by introducing Δ stkP::cat mutations into strain Cp6600 expressing the *comD*(D299N) allele to obtain strain Cp6607

TABLE 4. Impact of *stkP* and *phpP* mutations on culture transformability^a

Strain	Relevant genotype	n	Transformed colonies (% Rif ^r /total CFU)
Cp1015	WT	6	0.86
Cp9090	<i>stkP</i> ::Apha-3	5	<0.01
Cp7000	Δ stkP::cat	6	<0.01
Cp9091	<i>stkP</i> ⁺	2	1.05
Cp7004	Δ stkP::cat <i>phpD1</i>	2	<0.01

^a Donor DNA (1 μ g/ml) was from strain Cp1016 carrying the *rif23* allele, conferring resistance to rifampin (Rif^r). Values are representative of “n” independent determinations. WT, wild type.

TABLE 5. Impact of gain-of-function mutation in the CSP receptor ComD on the transformability of *stkP* mutant strains^a

Strain	Relevant genotype	Transformed colonies (% Nov ^r /total CFU)	
		pH 6.5	pH 8
Cp1015	WT	<0.01	0.31
Cp6600	<i>comD(D299N)</i>	0.30	0.52
Cp6607	<i>comD(D299N) ΔstkP::cat</i>	0.25	0.62
Cp7000	<i>ΔstkP::cat</i>	<0.01	<0.01

^a Donor DNA (1 μg/ml) was from strain R119 carrying a *novI* mutation, conferring resistance to 2 μg/ml of novobiocin. Results are representative of three independent experiments. WT, wild type.

(Table 1). Mutation D299N in ComD is a gain-of-function mutation conferring CSP-independent competence development in various conditions that are nonpermissive for the wild-type strain Cp1015, including growth in CTM at pH 6.5 (9, 25). Transformation tests on strain Cp6607 indicated pH-independent competence development similar to that of strain Cp6600 (Table 5), and similar results were obtained for the Cp6600 derivative carrying the *stkP::Apha-3* allele (our unpublished data). Thus, the ComD(D299N) mutation has an epistatic effect on StkP loss-of-function mutations, suggesting that StkP intervenes upstream from the CSP receptor ComD. This strongly suggests that ComD activation is sufficient to bypass the effects of the StkP mutation. We tested this hypothesis by subjecting StkP mutant strains Cp7000 (*ΔstkP::cat*) and Cp7004 [*ΔstkP::cat phpP(D241A)*] to “transformation tests” in CSP-supplemented CTM (see Materials and Methods). Strain Cp1008, carrying the *comA::spc* mutation abolishing ComA-dependent CSP export and maturation (7, 9, 11, 20, 30), was used as a control, in addition to the wild-type strain Cp1015. Dose-response tests clearly show that the threshold CSP concentration activating strains Cp7000 and Cp7004 is similar to that for strain Cp1008 and 20- to 100-fold higher than that for the wild type (see the supplemental material). Taken together, the results in Table 5 and the data presented in the supplemental material indicate that activation of ComD is sufficient to trigger competence development in StkP mutant strains. Complementation of StkP deficiency by synthetic CSP targeting the ComD receptor suggests that during bacterial growth, StkP, under the control of PhpP, transduces signals culminating directly or indirectly in ComD activation.

DISCUSSION

Adaptation to environmental conditions represents a major challenge for living organisms. Transient histidine phosphorylation plays a key role in TCS-mediated signaling in eubacteria. We provide biological and genetic evidences in favor of a role for transient serine-threonine phosphorylation in the regulation of growth and competence in *Streptococcus pneumoniae*. The proteins encoded by the *stkP-phpP* operon (31) constitute a signaling couple acting upstream of the competence-specific histidine kinase receptor ComD. ComD is a target for the oligopeptide pheromone CSP and belongs to the TCS ComDE. It is thus likely that transient serine-threonine phosphorylation is involved in the activation of a TCS signaling pathway in *Streptococcus pneumoniae*.

PhpP contributes to the pattern of competence development through its essential control of StkP activity. We previously established, using recombinant proteins, that StkP-P is a substrate for PhpP (31). The essentiality of PhpP phosphatase in the RX *stkP*⁺ genetic background (Table 3) suggests strong control on the level of StkP-P during bacterial growth. In growing bacteria, cellular localization of PhpP protein has been found to be dependent on the presence or absence of the membrane-associated protein kinase StkP (Fig. 1). Consistently, in StkP⁺ cells, proteins PhpP and StkP coimmunoprecipitate, suggesting that they belong to the same complex. Furthermore, utilization of recombinant proteins allows us to propose direct interaction between recombinant StkP kinase domain and recombinant PhpP proteins (Fig. 2). The transmembrane organization of StkP and intracellular localization of PhpP (Fig. 3) suggest a submembrane complex facing the cytoplasm, involving the N-terminal kinase domain of StkP and PhpP phosphatase. Mutations in StkP are highly pleiotropic. Cytoplasmic relocation of PhpP as a soluble protein in StkP-deficient bacteria (Fig. 1) has not been related to the competence-negative phenotype of the strain and did not reveal StkP-independent activity of PhpP. Rather, physiological and genetic analyses in the RX strain Cp1015 and its isogenic derivatives highlight the role of StkP and PhpP as a functional couple. This couple is involved in competence signaling. CSP supplementation experiments on *stkP* mutant strains (see the supplemental material) and the epistasis of the ComD(D299N) gain-of-function mutation on the StkP loss-of-function mutation in strain Cp6607 (Table 5) demonstrate that StkP kinase activates competence upstream of the histidine kinase receptor, ComD. This finding allows the exclusion of the hypothesis of competence inhibition, due to PhpP phosphatase activity in the absence of its specific substrate(s) StkP-P- and StkP-dependent phosphorylated targets. In the TIGR4 strain of *S. pneumoniae*, in vitro experiments on recombinant proteins suggest that PhpP and StkP interact competitively with RitR, modulating RitR-dependent Piu heme transporter expression (50). It is possible that some phenotypes of the StkP mutant strains involve alteration of RitR regulation; however, so far the role of RitR in competence remains uncertain (50). We present the hypothesis that in exponentially growing cultures, permissive for the development of a competence peak(s), the time window of cellular StkP-P, resulting in the antagonistic activities of StkP kinase and PhpP phosphatase, might determine the competence burst. During the course of this work, a report was published showing that StkP mutation leads to CSP-independent competence development, under conditions nonpermissive for the wild-type strain (40). Considering the mutator phenotype of StkP mutant strains, genetic drift might account for the results in reference 40. Indeed, alleles *ciaR::spc* (17) and *comD(D299N)* (this work) were shown to mask the phenotype of *stkP* mutations; other unidentified mutations might interfere with the phenotypic expression of StkP loss-of-function mutations and account for the results presented in reference 40.

StkP-PhpP signaling. The StkP-PhpP signaling couple transduces signals culminating in the activation of ComD, as shown by the epistatic relationship of *comD(D299N)* mutation on StkP0 mutations (Table 5) and by the CSP complementation test. Flow cytometry analysis using antibodies specific to

the C-terminal PASTA domain or the N-terminal kinase domain of the StkP protein indicated a transmembrane organization of the protein with its PASTA domain located beneath the cell wall, e.g., in the periplasmic space (Fig. 3). This cellular compartment contains the penicillin-binding proteins involved in the last steps of peptidoglycan biosynthesis in both gram-negative and gram-positive bacteria (28). Furthermore, PASTA domains are signatures for penicillin-binding proteins and serine-threonine kinases in gram-positive species, leading to speculations that the PASTA-containing kinases could constitute receptors for unlinked peptidoglycan stem peptides (12, 53). Consistently, the StkP mutations confer hypersensitivity to inhibitors of cell wall metabolism in *S. pneumoniae* (17, 40), potentially reflecting an involvement of StkP in cell wall homeostasis, as proposed elsewhere (22). This observation is reminiscent of results obtained in *Enterococcus faecalis*, suggesting a role for the serine-threonine kinase in the response to cell wall stress (24). In *S. pneumoniae*, downregulation of several genes, including competence genes, was observed in cultures containing sublethal concentrations of penicillin, indicating that periplasmic signals may control competence (37). We postulate that StkP activity might allow the entry of “periplasmic signals” into the competence regulatory network, culminating in a competence trigger. In addition to the ComDE TCS, ComAB, the CSP-autoregulated signaling network, TCSs CiaRH and MicAB (11, 15, 26, 41), the global regulator RegR (8), and the putative AI2 autoinducer system (38) have been shown to modulate competence expression in *S. pneumoniae*, implying that integration of different signals determines the pattern of competence development in growing cultures. Taking into account that PhpP is essential in the wild-type *stkP* genetic background, indicating strong control on StkP activity (Table 3), the StkP-PhpP signaling couple also represents an important actor determining the kinetics of competence development during bacterial growth in *Streptococcus pneumoniae*.

In conclusion, the results presented in this work provide physiological and molecular evidence for a signaling system based on transient serine-threonine phosphorylation in the regulation of competence for genetic transformation in *S. pneumoniae*. This system acts upstream from the competence-specific receptor ComD of the TCS ComDE. The biochemical nature of the gating signals, determining the time window of cellular StkP-P in growing cultures and culminating in the trigger of the TCS competence-signaling pathway, remains to be elucidated. At any rate, it is proposed as a working hypothesis that during bacterial growth the cellular level of StkP-P is finely tuned to the specific cellular physiology by PhpP phosphatase. Given the pleiotropic aspect of loss-of-function mutations in StkP, with effects on *in vitro* growth (17, 40) and virulence in mice (17), the transient serine-threonine phosphorylation, depending on both StkP and PhpP activities, plays a key role in the homeostasis of this pathogen in cultures and in animals.

ACKNOWLEDGMENTS

We thank Jean-Claude Lepert (Flow Cytometry Platform, IFR31) and Marc Trombe for helpful suggestions concerning the flow cytometry experiments. We also thank Lila Delbos and Camille Rouviere (master's students at Paul Sabatier University, Toulouse, France), and Nicolas Gay from INSA Toulouse for their contributions. Alexandre

Dubrac was helpful in Western blot analysis, and Ursula Liebl kindly helped in editing the manuscript.

This work was funded by INSERM Unité 858, Equipe 15 in France and by NARO Japan to M.O.

REFERENCES

- Absalon, C., M. Obuchowski, E. Madec, D. Delattre, B. Holland, and S. J. Seror. 2009. CpgA, EF-Tu and the stressosome protein YezB are substrates of the Ser/Thr kinase/phosphatase couple, PrkC/PrpC, in *Bacillus subtilis*. *Microbiology* **155**:932–963.
- Adler, E., A. Donella-Deana, F. Arigoni, L. A. Pinna, and P. Stragier. 1997. Structural relationship between a bacterial developmental protein and eukaryotic PP2C protein phosphatases. *Mol. Microbiol.* **23**:57–62.
- Alderwick, L. J., V. Molle, L. Kremer, A. J. Cozzone, T. R. Dafforn, G. S. Besra, and K. Fütterer. 2006. Molecular structure of EmbR, a response element of Ser/Thr kinase signaling in *Mycobacterium tuberculosis*. *Proc. Natl. Acad. Sci. USA* **103**:2558–2563.
- Av-Gay, Y., and M. Everett. 2000. The eukaryotic-like Ser/Thr protein kinases of *Mycobacterium tuberculosis*. *Trends Microbiol.* **8**:238–244.
- Auzat, I., S. Chapuy-Regaud, G. Le Bras, D. Dos Santos, A. D. Ogunniyi, I. Le Thomas, J.-R. Garel, J. C. Paton, and M.-C. Trombe. 1999. The NADH oxidase of *Streptococcus pneumoniae*, its role in competence and virulence. *Mol. Microbiol.* **34**:1018–1028.
- Boitel, B., M. Ortiz-Lombardía, R. Durán, F. Pompeo, S. T. Cole, C. Cerveñansky, and P. M. Alzari. 2003. PknB kinase activity is regulated by phosphorylation in two Thr residues and dephosphorylation by PstP, the cognate phospho-Ser/Thr phosphatase, in *Mycobacterium tuberculosis*. *Mol. Microbiol.* **49**:1493–1508.
- Chandler, M. S., and D. A. Morrison. 1988. Identification of two proteins encoded by *com*, a competence control locus of *Streptococcus pneumoniae*. *J. Bacteriol.* **170**:3136–3141.
- Chapuy-Regaud, S., A. D. Ogunniyi, N. Diallo, Y. Huet, J.-F. Desnottes, J. C. Paton, S. Escaich, and M.-C. Trombe. 2003. RegR, a global LacI/GalR, modulates virulence and competence in *Streptococcus pneumoniae*. *Infect. Immun.* **71**:2615–2625.
- Chen, J. D., and D. A. Morrison. 1987. Modulation of competence for genetic transformation in *Streptococcus pneumoniae*. *J. Gen. Microbiol.* **133**:1959–1967.
- Cheng, Q., E. A. Campbell, A. M. Naughton, S. Johnson, and H. R. Masure. 1997. The *com* locus controls genetic transformation in *Streptococcus pneumoniae*. *Mol. Microbiol.* **23**:683–692.
- Dagkessamakaia, A., M. Moscoso, V. Hénard, S. Guiral, K. Overweg, M. Reuter, B. Martin, J. Wells, and J. P. Claverys. 2004. Interconnection of competence, stress and CiaR regulons in *Streptococcus pneumoniae*: competence triggers stationary phase autolysis of *ciaR* mutant cells. *Mol. Microbiol.* **51**:1071–1086.
- Dessen, A., N. Mouz, E. Gordon, J. Hopkins, and O. Dideberg. 2001. Crystal structure of PBP2x from a highly penicillin-resistant *Streptococcus pneumoniae* clinical isolate. *J. Biol. Chem.* **276**:45106–45112.
- Deutscher, J., R. Herro, A. Bourand, I. Mijakovic, and S. Poncet. 2005. P-Ser-HPr—a link between carbon metabolism and the virulence of some pathogenic bacteria. *Biochim. Biophys. Acta* **1754**:118–125.
- Echenique, J. R., S. Chapuy-Regaud, and M.-C. Trombe. 2000. Competence regulation by oxygen in *Streptococcus pneumoniae*: involvement of *ciaRH* and *comCDE*. *Mol. Microbiol.* **36**:688–696.
- Echenique, J. R., and M.-C. Trombe. 2001. Competence repression under oxygen limitation through the two-component MicAB signal-transducing system in *Streptococcus pneumoniae* and involvement of the PAS domain of MicB. *J. Bacteriol.* **183**:4599–4608.
- Echenique, J. R., and M.-C. Trombe. 2001. Competence modulation by the NADH oxidase of *Streptococcus pneumoniae* involves signal transduction. *J. Bacteriol.* **183**:768–772.
- Echenique, J., A. Kadioglu, S. Romao, P. W. Andrew, and M.-C. Trombe. 2004. Protein serine/threonine kinase StkP positively controls virulence and competence in *Streptococcus pneumoniae*. *Infect. Immun.* **72**:2434–2437.
- Hunter, T. 1995. Protein-kinases and phosphatases: The yin and yang of protein p phosphorylation and signaling. *Cell* **80**:225–236.
- Iannelli, F., M. R. Oggioni, and G. Pozzi. 2005. Sensor domain of histidine kinase ComD confers competence pherotype specificity in *Streptococcus pneumoniae*. *FEMS Microbiol. Lett.* **252**:321–326.
- Ishii, S., T. Yano, and H. Hayashi. 2006. Expression and characterization of the peptidase domain of *Streptococcus pneumoniae* ComA, a bifunctional ATP-binding cassette transporter involved in quorum sensing pathway. *J. Biol. Chem.* **281**:4726–4731.
- Jin, H., and V. Pancholi. 2006. Identification and biochemical characterization of a eukaryotic-type serine/threonine kinase and its cognate phosphatase in *Streptococcus pyogenes*: their biological functions and substrate identification. *J. Mol. Biol.* **357**:1351–1372.
- Jones, G., and P. Dyson. 2006. Evolution of transmembrane protein kinases implicated in coordinating remodeling of gram-positive peptidoglycan: inside versus outside. *J. Bacteriol.* **188**:7470–7476.

23. Kim, T. J., T. A. Gaidenko, and C. W. Price. 2004. In vivo phosphorylation of partner switching regulators correlates with stress transmission in the environmental signaling pathway of *Bacillus subtilis*. *J. Bacteriol.* **186**:6124–6132.
24. Kristich, C. J., C. L. Wells, and G. M. Dunny. 2007. A eukaryotic-type Ser/Thr kinase in *Enterococcus faecalis* mediates antimicrobial resistance and intestinal persistence. *Proc. Natl. Acad. Sci. USA* **104**:3508–3513.
25. Lacks, S. A., and P. Greenberg. 2001. Constitutive competence for genetic transformation in *Streptococcus pneumoniae* caused by mutation of a transmembrane histidine kinase. *Mol. Microbiol.* **42**:1035–1045.
26. Lange, R., C. Wagner, A. de Saizieu, N. Flint, J. Molnos, M. Stieger, P. Caspers, M. Kamber, W. Keck, and K. E. Amrein. 1999. Domain organization and molecular characterization of 13 two-component systems identified by genome sequencing of *Streptococcus pneumoniae*. *Gene* **237**:223–234.
27. Luria, S. E., and M. Delbruck. 1943. Mutation of bacteria from virus sensitivity to virus resistance. *Genetics* **28**:491–511.
28. Macheboeuf, P., C. Contreras-Martel, V. Job, O. Dideberg, and A. Dessen. 2006. Penicillin binding proteins: key players in bacterial cell cycle and drug resistance processes. *FEMS Microbiol. Rev.* **30**:673–691.
29. Madec, E., A. Laszkiewicz, A. Iwanicki, M. Obuchowski, and S. Serror. 2002. Characterisation of a membrane-linked Ser/Thr protein kinase in *Bacillus subtilis* implicated in developmental processes. *Mol. Microbiol.* **46**:571–586.
30. Morrison, D. A., M.-C. Trombe, M. K. Hayden, G. A. Waszak, and J. D. Chen. 1984. Isolation of transformation-deficient *Streptococcus pneumoniae* mutants defective in control of competence, using insertion-duplication mutagenesis with the erythromycin resistance determinant of pAM β 1. *J. Bacteriol.* **159**:870–876.
31. Nováková, L., L. Sasková, P. Pallová, J. Janecek, J. Novotná, A. Ulrych, J. Echenique, M.-C. Trombe, and P. Branny. 2005. Characterization of a eukaryotic type serine/threonine protein kinase and protein phosphatase of *Streptococcus pneumoniae* and identification of kinase substrates. *FEBS J.* **272**:1243–1254.
32. Peeters, B. P., J. H. de Boer, S. Bron, and G. Venema. 1988. Structural plasmid instability in *Bacillus subtilis*: effect of direct and inverted repeats. *Mol. Gen. Genet.* **212**:450–458.
33. Perego, M. 1998. Kinase and phosphatase competition regulates *Bacillus subtilis* development. *Trends Microbiol.* **6**:366–370.
34. Rajagopal, L., A. Clancy, and C. E. Rubens. 2003. A eukaryotic type serine/threonine kinase and phosphatase in *Streptococcus agalactiae* reversibly phosphorylate an inorganic pyrophosphatase and affect growth, cell segregation, and virulence. *J. Biol. Chem.* **278**:14429–14441.
35. Rajagopal, L., A. Vo, A. Silvestroni, and C. E. Rubens. 2005. Regulation of purine biosynthesis by a eukaryotic-type kinase in *Streptococcus agalactiae*. *Mol. Microbiol.* **56**:1329–1346.
36. Rajagopal, L., A. Vo, A. Silvestroni, and C. E. Rubens. 2006. Regulation of cytotoxin expression by converging eukaryotic-type and two-component signalling mechanisms in *Streptococcus agalactiae*. *Mol. Microbiol.* **62**:941–957.
37. Rogers, P. D., T. T. Liu, K. S. Barker, G. M. Hilliard, B. K. English, J. Thornton, E. Swiatlo, and L. S. McDaniel. 2007. Gene expression profiling of the response of *Streptococcus pneumoniae* to penicillin. *J. Antimicrob. Chemother.* **59**:616–626.
38. Romao, S., G. Memmi, M. R. Oggioni, and M.-C. Trombe. 2006. LuxS impacts on LytA-dependent autolysis and on competence in *Streptococcus pneumoniae*. *Microbiology* **152**:333–341.
39. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
40. Sasková, L., L. Nováková, M. Basler, and P. Branny. 2007. Eukaryotic-type serine/threonine protein kinase StkP is a global regulator of gene expression in *Streptococcus pneumoniae*. *J. Bacteriol.* **189**:4168–4179.
41. Sebert, M. E., K. P. Patel, M. Plotnick, and J. N. Weiser. 2005. Pneumococcal HtrA protease mediates inhibition of competence by the CiaRH two-component signaling system. *J. Bacteriol.* **187**:3969–3979.
42. Sharma, K., M. Gupta, A. Krupa, N. Srinivasan, and Y. Singh. 2006. EmbR, a regulatory protein with ATPase activity, is a substrate of multiple serine/threonine kinases and phosphatase in *Mycobacterium tuberculosis*. *FEBS J.* **273**:2711–2721.
43. Singh, K. D., S. Halbedel, B. Görke, and J. Stülke. 2007. Control of the phosphorylation state of the HPr protein of the phosphotransferase system in *Bacillus subtilis*: implication of the protein phosphatase PrpC. *J. Mol. Microbiol. Biotechnol.* **13**:165–171.
44. Stock, A. M., V. L. Robinson, and P. N. Goudreau. 2000. Two-component signal transduction. *Annu. Rev. Biochem.* **69**:183–215.
45. Throup, J. P., K. K. Koretke, A. P. Bryant, K. A. Ingraham, A. F. Chalker, Y. Ge, A. Marra, N. G. Wallis, J. R. Brown, D. J. Holmes, M. Rosenberg, and M. K. Burnham. 2000. A genomic analysis of two-component signal transduction in *Streptococcus pneumoniae*. *Mol. Microbiol.* **35**:566–576.
46. Tiraby, G., M. S. Fox, and H. Bernheimer. 1975. Marker discrimination in deoxyribonucleic acid-mediated transformation of various *Pneumococcus* strains. *J. Bacteriol.* **121**:608–618.
47. Tomasz, A., P. Moreillon, and G. Pozzi. 1988. Insertional inactivation of the major autolysin gene of *Streptococcus pneumoniae*. *J. Bacteriol.* **170**:5931–5934.
48. Trombe, M.-C. 1981. Characterisation de mutants de résistance à l'améthoptérine chez *Streptococcus pneumoniae*, alteration du potentiel transmembranaire; mise en évidence d'une cible membranaire pour l'améthoptérine, Ph.D. thesis. Université Paul Sabatier, Toulouse, France.
49. Trombe, M.-C., C. Clavé, and J. M. Manias. 1992. Calcium regulation of growth and differentiation in *Streptococcus pneumoniae*. *J. Gen. Microbiol.* **138**:77–84.
50. Ulijasz, A. T., S. P. Falk, and B. Weisblum. 2009. Phosphorylation of the RitR DNA-binding domain by a Ser-Thr phosphokinase: implications for global gene regulation in the streptococci. *Mol. Microbiol.* **71**:382–390.
51. Wang, R. F., and S. R. Kushner. 1991. Construction of versatile low-copy-number vectors for cloning, sequencing and gene expression in *Escherichia coli*. *Gene* **100**:195–199.
52. Wehenkel, A., M. Bellinzoni, M. Graña, R. Duran, A. Villarino, P. Fernandez, G. Andre-Leroux, P. England, H. Takiff, C. Cerveñansky, S. T. Cole, and P. M. Alzari. 2008. Mycobacterial Ser/Thr protein kinases and phosphatases: physiological roles and therapeutic potential. *Biochim. Biophys. Acta* **1784**:193–202.
53. Yeats, C., R. D. Finn, and A. Bateman. 2002. The PASTA domain: a beta-lactam-binding domain. *Trends Biochem. Sci.* **27**:438–440.