

Status of intestinal parasite infections among children in Bat Dambang, Cambodia

Seung Kyu PARK¹⁾, Dong-Heui KIM²⁾, Young-Kun DEUNG²⁾, Hun-Joo KIM³⁾, Eun-Ju YANG⁴⁾,
Soo-Jung LIM⁴⁾, Yong-Suk RYANG⁴⁾, Dan JIN⁵⁾ and Kyu-Jae LEE^{1)*}

¹⁾Department of Parasitology and Institute of Basic Medical Science, ²⁾Department of Biology,

³⁾Department of Neurosurgery, Wonju College of Medicine, Yonsei University, Wonju 220-701,

⁴⁾Department of Biomedical Laboratory Science and Institute of Health Science,

College of Health Science, Yonsei University 220-710,

⁵⁾Department of Microbiology, Wonju College of Medicine, Yonsei University, Wonju 220-701, Korea

Abstract: A survey was conducted to determine the extent of intestinal parasite infection in Bat Dambang, Cambodia in March 2004. A total of 623 fecal specimens was collected from kindergarten and schoolchildren and examined using the formalin-ether sedimentation technique. The overall infection rate of intestinal parasites was 25.7% (boys, 26.2%; girls, 25.1%), and the infection rates of intestinal helminthes by species were as follows: *Echinostoma* sp. 4.8%, hookworm 3.4%, *Hymenolepis nana* 1.3%, and *Rhabditis* sp. 1.3%. The infection rates of intestinal protozoa were; *Entamoeba coli* 4.8%, *Giardia lamblia* 2.9%, *Iodamoeba butschlii* 1.4%, *Entamoeba polecki* 1.1%, and *Entamoeba histolytica* 0.8%. There were no egg positive cases of *Ascaris lumbricoides* or *Trichuris trichiura*. All children infected were treated with albendazole, praziquantel, or metronidazole according to parasite species. The results showed that intestinal parasites are highly endemic in Bat Dambang, Cambodia.

Key words: Cambodia, intestinal parasites, children

Cambodia is recognized as an endemic area for malaria and schistosomiasis (Meek, 1988; Singhasivanon, 1999; Urbani et al., 2002), but the infection status of intestinal parasites in Cambodia has not been thoroughly investigated. However, several reports show that Cambodian children and refugees are infected with intestinal parasites including nematodes, trematodes, and protozoa (Nwanyanwu et al., 1989; Lurio et al., 1991; Gyorkos et al., 1992; Lee et al., 2002).

With the aid of a local Korean medical service in

Cambodia, we collected 623 stool samples from children at Watkor Kindergarten and from schoolchildren in Bat Dambang, Cambodia in March 2004; Bat Dambang is located about 250 km southeast of Phnom Penh, the capital city of Cambodia.

Formalin-ether sedimentation revealed intestinal helminth eggs of Hookworm, *Rhabditis* sp., *Hymenolepis nana*, and *Echinostoma* sp., and protozoan cysts of *Entamoeba histolytica*, *Entamoeba coli*, *Giardia lamblia*, *Iodamoeba butschlii*, and *Entamoeba polecki*. Of the 623 stool specimens, 160 (25.7%) were positive. The positive rates for boys and girls were 26.2% (90/344) and 25.1% (70/279), respectively. Data revealed a highest positive rate (4.8%) for *Echinostoma* sp., followed by 3.4% for Hookworm (Table 1). Multiple parasite infec-

• Received 21 October 2004, accepted after revision 8 November 2004.

*Corresponding author (e-mail: kjee@wonju.yonsei.ac.kr)

Table 1. Prevalences of intestinal parasite infections in Watkor kindergarten children and schoolchildren in Bat Dambang, Cambodia

	Boy (%)	Girl (%)	Total (%)
Echinostomatidae	7(2.0)	23(8.2)	30(4.8)
Hookworm	13(3.8)	8(2.9)	21(3.4)
<i>Rhabditis</i> sp.	6(1.7)	2(0.7)	8(1.3)
<i>Hymenolepis nana</i>	4(1.2)	4(1.4)	8(1.3)
<i>Entamoeba coli</i>	20(5.8)	10(3.6)	30(4.8)
<i>Entamoeba histolytica</i>	4(1.2)	1(0.4)	5(0.8)
<i>Entamoeba polecki</i>	3(0.9)	4(1.4)	7(1.1)
<i>Endolimax nana</i>	23(6.7)	17(6.1)	40(6.4)
<i>Giardia lamblia</i>	15(4.4)	3(1.1)	18(2.9)
<i>Iodamoeba butschlii</i>	5(1.5)	4(1.4)	9(1.4)

Table 2. Prevalences of intestinal parasite infection by school grade among Watkor kindergarten children and schoolchildren in Bat Dambang, Cambodia

Grade	Kindergarden	Primary school					Middle school			
		1	2	3	4	5	6	7	8	9
No. examined/ positive (%)	20/2 (10.0)	52/14 (26.9)	61/13 (21.3)	89/34 (38.2)	84/20 (23.8)	82/16 (19.5)	67/12 (17.9)	94/24 (25.5)	41/14 (34.1)	33/9 (27.3)
Echinostomatidae	-	1(1.9)	4(6.5)	6(6.7)	3(3.6)	5(6.1)	3(4.5)	1(1.1)	4(9.8)	3(9.1)
Hookworm	-	2(3.8)	2(3.3)	5(5.6)	4(4.8)	2(2.4)	-	3(3.2)	3(7.3)	-
<i>Rhabditis</i> sp.	-	2(3.8)	-	1(1.1)	1(1.2)	1(1.2)	1(1.5)	1(1.1)	-	1(3.0)
<i>Hymenolepis nana</i>	-	1(1.9)	-	2(2.2)	1(1.2)	1(1.2)	-	2(2.1)	1(2.4)	-
<i>Entamoeba coli</i>	-	1(1.9)	5(8.2)	7(7.8)	5(5.9)	1(1.2)	4(6.0)	2(2.1)	2(4.9)	3(9.1)
<i>Entamoeba histolytica</i>	-	-	-	3(3.4)	-	-	-	1(1.1)	-	1(3.0)
<i>Entamoeba polecki</i>	-	1(1.9)	1(1.6)	2(2.2)	1(1.2)	1(1.2)	1(1.5)	-	-	-
<i>Endolimax nana</i>	1(5.0)	5(9.6)	2(3.3)	4(4.5)	2(2.4)	6(7.3)	1(1.5)	13(13.8)	4(9.8)	2(6.0)
<i>Giardia lamblia</i>	1(5.0)	3(5.8)	2(3.3)	2(2.2)	3(3.6)	1(1.2)	4(6.0)	1(1.1)	1(2.4)	-
<i>Iodamoeba butschlii</i>	-	-	1(1.6)	6(6.7)	1(1.2)	-	-	1(1.1)	-	-

tions were found in 9 cases. Infection rates ranged from 10% to 38.2% by school grade; the highest rate was found among third grade of primary schoolchildren. The results of a comparison of trematode infection and eating habits were interesting. In general, children did not consume raw fresh water fish (Story et al., 1989); however, they were believed to have been infected by inadequately cooked fish. *Echinostoma* species are important trematode parasites and require treatment; anti-helminthics were sent to the Watkor primary school after the stool examination.

The school is located along the Steung Sangkae River, which is connected to Tonle Sab Lake. Although many children swam in this river during the stool examination period, we failed to find

Schistosoma eggs. Accordingly, in view of their swimming habits, this area is not considered to be a schistosomiasis endemic area.

The present study showed different patterns of intestinal parasite infection compared to the previous survey at Kampongcham (Lee et al., 2002); this survey showed no cases of *Ascaris lumbricoides*, *Trichuris trichiura* or *Opisthorchis* sp., while egg positive rates of *A. lumbricoides*, *T. trichiura*, and *Opisthorchis* sp. in Kampongcham were 26.3%, 1%, and 10%, respectively. However, children in Bat Dambang were found infected with *E. polecki* (1.1%) and *H. nana* (1.3%), whereas no cases of *E. polecki* or *H. nana* were recorded in the Kampongcham survey in 2002 (Lee et al., 2002). This study identified regional differences in

intestinal parasitic infections between in Bat Dambang and in Kampongcham.

Periodic surveys on intestinal parasitic infections, the treatment of infected cases, and education with respect to eating habits and improved sanitation conditions are highly recommended for parasite control in Cambodia.

ACKNOWLEDGEMENTS

We thank the medical service team at Wonju Christian Hospital for collection of stool samples, and the Harvest Mission Center in Cambodia for its cooperation during the visit to the survey area.

REFERENCES

Gyorkos TW, MacLean JD, Viens P, Chheang C, Kokoskin-Nelson E (1992) Intestinal parasite infection in the Kampuchean refugee population 6 years after resettlement in Canada. *J Infect Dis* **166**: 413-417.

Lee KJ, Bae YT, Kim DH, et al. (2002) Status of intestinal par-

asites infection among primary school children in Kampongcham, Cambodia. *Korean J Parasitol* **40**: 153-155.

Lurio J, Verson H, Karp S (1991) Intestinal parasites in Cambodians: comparison of diagnostic methods used in screening refugees with implications for treatment of populations with high rates of infestation. *J Am Board Fam Pract* **4**: 71-78.

Meek SR (1988) Epidemiology of malaria in displaced Khmers on the Thai-Kampuchean border. *Southeast Asian J Trop Med Public Health* **19**: 243-252.

Nwanyanwu OC, Moore JS, Adams ED (1989) Parasitic infections in Asian refugees in Fort Worth. *Tex Med* **85**: 42-45.

Singhasivanon P (1999) Mekong malaria. Malaria, multi-drug resistance and economic development in the greater Mekong subregion of Southeast Asia. *Southeast Asian J Trop Med Public Health* **30** (Suppl 4): i-iv, 1-101.

Story M, Harris LJ (1989) Food habits and dietary change of Southeast Asian refugee families living in the United States. *J Am Diet Assoc* **89**: 800-803.

Urbani C, Sinoun M, Socheat D, et al. (2002) Epidemiology and control of mekongi schistosomiasis. *Acta Trop* **82**: 157-168.

**APPRECIATIONS TO PEER-REVIEWERS
FOR VOLUME 42, 2004**

The Editorial Board of the Korean Journal of Parasitology appreciates the contribution as reviewers of papers in the Volume 42, 2004. Following scientists are contributors.

Contributors from foreign countries:

Nawa, Yukifumi (Miyazaki Med. Coll., Japan)

Klein, Terry A. (US Army)

Contributors from Korea:

Bae, Young-An (Sungkyunkwan Univ.)

Cho, Moon-Jae (Cheju Nat. Univ.)

Chung, Dong-Il (Kyungpook Nat. Univ.)

Eom, Keeseon S. (Chungbuk Nat. Univ.)

Hong, Sung-Tae (Seoul Nat. Univ.)

Hwang, Ui-Wook (Kyungpook Nat. Univ.)

Kim, Hong-Bin (Seoul Nat. Univ.)

Kong, Yoon (Sungkyunkwan Univ.)

Min, Duk-Young (Hanyang Univ.)

Nam, Ho-Woo (Catholic Univ.)

Park, Ki-Ho (Seoul Nat. Univ.)

Shin, Myeong Heon (Yonsei Univ.)

Yang, Hyun-Jong (Ewha Womans Univ.)

Yu, Jae-Ran (Konkuk Univ.)

Chai, Jong-Yil (Seoul Nat. Univ.)

Choi, Min-Ho (Seoul Nat. Univ.)

Chung, Young-Bae (Cheju Nat. Univ.)

Hong, Sung-Jong (Chung-Ang Univ.)

Huh, Sun (Hallym Univ.)

Kho, Weon-Gyu (Inje Univ.)

Kong, Hyun-Hee (Kyungpook Nat. Univ.)

Lee, Young-Ha (Chungnam Nat. Univ.)

Mook, In-Hee (Seoul Nat. Univ.)

Park, Joong-Ki (Chungbuk Nat. Univ.)

Shin, Ho-Joon (Ajou Univ.)

Sohn, Woon-Mok (Gyeongsang Nat. Univ.)

Yong, Tai-Soon (Yonsei Univ.)