

SHORT COMMUNICATION

Maternal plasma 25-hydroxyvitamin D concentration and birthweight, growth and bone mineral accretion of Gambian infants

Ann Prentice (ann.prentice@mrc-hnr.cam.ac.uk)^{1,2}, Landing MA Jarjou², Gail R Goldberg^{1,2}, Janet Bennett¹, Tim J Cole³, Inez Schoenmakers¹

1.MRC Human Nutrition Research, Elsie Widdowson Laboratory, Cambridge, UK

2.MRC Keneba, The Gambia

3.MRC Centre of Epidemiology for Child Health, UCL Institute of Child Health, London, UK

Correspondence

Ann Prentice Ph.D., MRC Human Nutrition Research, Elsie Widdowson Laboratory, Cambridge, UK, CB1 9NL.
Tel: +44-1223-426356 |
Fax: +44-1223-437515 |
Email: ann.prentice@mrc-hnr.cam.ac.uk

Received

22 February 2009;
accepted 24 April 2009.

DOI:10.1111/j.1651-2227.2009.01352.x

Re-use of this article is permitted in accordance with the Creative Commons Deed, Attribution 2.5, which does not permit commercial exploitation.

Maternal vitamin D deficiency during pregnancy is a recognized risk factor for rickets and osteomalacia in infancy (1). The circulating plasma concentration of 25-hydroxyvitamin D (25OHD), a long-lived metabolite of vitamin D, is used to judge vitamin D status; values below 25 nmol/L are associated with an increased risk of rickets and osteomalacia (1). There is evidence that a low maternal plasma 25OHD in pregnancy may influence the growth and bone mineral accrual of the offspring during foetal life, infancy and childhood. Positive associations have been reported between maternal vitamin D status in pregnancy and birthweight, birth length, length at 1 year and bone mineral accretion at 9 years (2–6), although evidence is conflicting (7,8). These relationships have been observed at concentrations of 25OHD higher than those associated with rickets and osteomalacia, and there are calls to raise the accepted lower threshold of vitamin D sufficiency for pregnant women, most recently to 80 nmol/L (9). On a population basis, plasma 25OHD concentrations above 80 nmol/L are relatively uncommon in countries at temperate latitudes but are more common among people living in the tropics who have abundant skin sunshine exposure (10). To contribute to the debate on the definition of vitamin D sufficiency in pregnancy, we have investigated the influence of maternal plasma 25OHD concentration on foetal and infant growth in a rural area of The Gambia, West Africa (13°N). In this region, there is tropical sunshine all year, the women are farmers who work out-of-doors for much of each day, and local female dress does not restrict regular sunshine

exposure to the face, neck, shoulders, arms and feet, especially during farm work and gardening.

The study was a secondary analysis of biochemical, anthropometric and bone data from a subset of 125 women and infants collected during a calcium supplementation study of blood pressure in pregnant Gambian women (International Trial Registry: ISRCTN96502494). No significant benefits for foetal and infant growth of maternal calcium supplementation were identified despite the low customary calcium intake in The Gambia (11). The protocol, methods, maternal characteristics and infant data from the detailed study have been published (11). Briefly, women from the rural villages of Keneba and Manduar, West Kiang, The Gambia were recruited at 20 weeks of pregnancy (P20) and randomized to a daily calcium supplement or a matching placebo tablet until parturition (1500 mg Ca as calcium carbonate and microcellulose-lactose, respectively; Nycomed Pharma AS, Asker, Norway). Fasting, early morning blood was collected and anthropometry performed at P20 and 36 weeks of pregnancy (P36). The mean (\pm SD) age, weight, height and dietary calcium intake of the women at P20 were 27.4 ± 7.5 years, 56.3 ± 6.7 kg, 1.61 ± 0.05 m and 356 ± 190 mg/day, respectively. The median parity (range) was 3 (0–10). Infant birthweight was measured within 24 h of delivery. Weight, crown-heel length and head circumference were measured at 2, 13 and 52 weeks postpartum. In addition, infant bone mineral content (BMC), bone mineral density (BMD) and bone width (BW) or bone area (BA), were measured by single

photon absorptiometry of the midshaft radius (Lunar SP2, Lunar Corporation, Madison, WI, USA) and, for a subset ($n = 44, 47$ and 52 at $2, 13$ and 52 weeks, respectively), by whole-body dual-energy X-ray absorptiometry (Lunar DPX+, software version 4.7b, Lunar Corporation).

Plasma 25OHD was measured using a radioimmuno-metric assay (Diasorin Ltd, Wokingham, Berks, UK), with assay performance monitored through the Vitamin D External Quality Assessment Scheme (DEQAS; Endocrine/Oncology Laboratory, Charing Cross Hospital, London, UK). The intra- and inter-assay coefficients of variation were 4% and <6%, respectively. Relationships between infant outcomes and maternal 25OHD concentration were explored using multiple linear regressions (DataDesk 6.2.1, Data Description Inc., Ithaca, NY, USA). The following potential confounders were included in the full models: maternal weight, weight gain, height, parity, supplement group, sex of infant and season. Nonsignificant variables were removed by backwards elimination to produce parsimonious models. Infant BMC was corrected for bone and body size by including infant weight, length and BW or BA, in the regression models (12). An interaction term, supplement group \times 25OHD, was added to investigate potential synergistic effects of calcium supplementation and maternal vitamin D status on infant growth outcomes. Other analyses of maternal 25OHD included comparisons between P20 and P36, and also as outcomes the proportions above and below the cut-off of 80 nmol/L. A conservative level of significance of $p \leq 0.01$ was adopted in recognition of the large number of separate statistical tests performed (>100); possible trends in the data with $p = 0.01$ – 0.1 were noted. The analysis was conducted on 123 mother–infant pairs; blood samples from two subjects were not available.

Mean \pm SD 25OHD (range) was: P20 = 103 ± 25 (53–167) nmol/L; P36 = 111 ± 27 (51–189) nmol/L. No subject had a 25OHD value <50 nmol/L, 20% and 16% had 25OHD <80 nmol/L, at P20 and P36, respectively. There was a high degree of within-subject consistency in 25OHD at P20 and P36 ($25\text{OHD}_{\text{P36}} = 33.2 + [0.79 \pm 0.07] \times 25\text{OHD}_{\text{P20}}$, $p \leq 0.001$, R^2 adjusted 51.5%, $n = 121$); 11% of women had 25OHD <80 nmol/L at both P20 and P36. The mean birthweight of the infants was 2.99 ± 0.36 kg. The infant anthropometric and bone measures during the first year are given in Table 1. No significant relationships or trends in the data were observed between maternal 25OHD concentration using the values at P20, P36 or the mean of the two and any of the following infant measures: birthweight, infant weight, length, head circumference, BMC, BW (or BA), BMD and size-adjusted BMC of the midshaft radius and whole body at any time postpartum. This is illustrated in Figure 1 for birthweight as a function of maternal 25OHD concentration at P20. Comparing the results for mothers with 25OHD above and below 80 nmol/L did not alter this finding.

No significant interaction between supplement group and maternal 25OHD concentration was observed for any infant variable. Trends in the data were observed in a few instances for a supplement group \times 25OHD interaction among the

Table 1 Anthropometric and bone measures of Gambian infants

Characteristic	Age (weeks)		
	2 (n = 125)	13 (n = 123)	52 (n = 121)
Weight (kg)	3.32 ± 0.47	5.84 ± 0.72	8.08 ± 0.97
Length (cm)	50.5 ± 1.9	59.6 ± 2.0	71.9 ± 2.8
Head circumference (cm)	35.5 ± 1.6	40.0 ± 1.3	44.6 ± 1.4
Weight SDS	-1.08 ± 0.86	-0.37 ± 0.98	-1.96 ± 1.15
Length SDS	-0.80 ± 0.91	-0.41 ± 0.92	-1.21 ± 1.11
Radius shaft*			
Bone mineral content (g/cm)	0.056 ± 0.014	0.068 ± 0.018	0.087 ± 0.022
Bone width (cm)	0.374 ± 0.066	0.470 ± 0.067	0.594 ± 0.092
Whole body†			
Bone mineral content (g)	50.9 ± 11.6	89.5 ± 21.4	169.6 ± 35.1
Bone area (cm ²)	105 ± 20	177 ± 34	279 ± 50

Data are mean \pm SD.

SDS, standard deviation score relative to British reference data (14).

* $n = 116, 118, 96$ at $2, 13$ and 52 weeks, respectively.

† $n = 44, 47, 52$ at $2, 13$ and 52 weeks, respectively.




Figure 1 Lack of a significant relationship between infant birthweight and maternal vitamin D status at 20 weeks of pregnancy ($p = 0.8$). Multiple regression model included season, maternal height, weight, weight gain, supplement group and sex of the infant.

bone measures but no consistent picture emerged and they were considered to have arisen by chance.

We conclude that there is no evidence for an influence of vitamin D status during pregnancy on infant growth and bone mineral accrual in the conditions prevailing in The Gambia. The children in this study, as is common in this region of The Gambia (13), were born small, grew well for the first months of life but experienced growth faltering during later infancy compared to Western children (11,14), as demonstrated by their weight and length SDS. The 25OHD concentrations of the women were >50 nmol/L in the second half of pregnancy, and no distinction could be drawn in infant outcomes between mothers with concentrations above or below 80 nmol/L. Thus, our study suggests that, for women with regular, adventitious UVB sunshine exposure and in situations where foetal and infant growth may

be constrained by multiple factors, there would be no benefit for foetal and infant growth or bone mineral accrual in aiming to increase the vitamin D status of individual mothers during pregnancy above 50 or 80 nmol/L.

ACKNOWLEDGEMENTS

We are grateful for the following contributions at MRC Human Nutrition Research: Dr. Ann Laskey for expert oversight of SPA and DXA measurements, Dr. Shailja Nigdikar and Ann Laidlaw for laboratory management, Sheila Levitt for data entry. Our thanks also go to the midwives, fieldworkers and lab staff at MRC Keneba who were involved in the study: Frances Foord, Ndey Haddy Bah, Fatou Sosseh, Ebou Jarjou, Morikebba Sanyang, Lamin Sanneh, Mariama Jammeh, Abdoulie Jaiteh, Karamo Camara, Bakary Darboe and Musa Colley.

References

1. Scientific Advisory Committee on Nutrition. Update on Vitamin D. London: The Stationery Office, 2007.
2. Brooke OG, Brown IRF, Bone CDM, Carter ND, Cleeve HJW, Maxwell JD, et al. Vitamin D supplements in pregnant Asian women: effects on calcium status and fetal growth. *Br Med J* 1980; 280: 751–4.
3. Brooke OG, Butters F, Wood C. Intrauterine vitamin D nutrition and postnatal growth in Asian infants. *Br Med J* 1981; 283: 1024.
4. Javaid MK, Crozier SR, Harvey NC, Gale GR, Dennison EM, Boucher BJ, et al. Maternal vitamin D status during pregnancy and childhood bone mass at age 9 years: a longitudinal study. *Lancet* 2006; 367: 36–43.
5. Marya RK, Rathee S, Dua V, Sangwan K. Effect of vitamin D supplementation during pregnancy on foetal growth. *Ind J Med Res* 1988; 88: 488–92.
6. Mannion CA, Gray-Donald K, Koski KG. Association of low intake of milk and vitamin D during pregnancy with decreased birth weight. *Can Med Assoc J* 2006; 174: 1273–7.
7. Morley R, Carlin JB, Pasco JA, Wark JD. Maternal 25-hydroxyvitamin D and parathyroid hormone concentrations and offspring birth size. *J Clin Endocrinol Metab* 2006; 91: 906–12.
8. Mallet E, Gugi B, Brunelle P, Henocq A, Basuyau JP, Lemeur H. Vitamin D supplementation in pregnancy: a controlled trial of two methods. *Obstet Gynecol* 1986; 68: 300–4.
9. Hollis BW, Wagner CL. Vitamin D deficiency during pregnancy: an ongoing epidemic. *Am J Clin Nutr* 2006; 84: 273.
10. Prentice A. Vitamin D deficiency: a global perspective. *Nutr Rev* 2008; 66: S153–64.
11. Jarjou LMA, Prentice A, Sawo Y, Laskey MA, Bennett J, Goldberg GR, et al. Randomized, placebo-controlled, calcium supplementation study in pregnant Gambian women: effects on breastmilk calcium concentrations and infant birth weight, growth, and bone mineral accretion in the first year of life. *Am J Clin Nutr* 2006; 83: 657–66.
12. Prentice A, Parsons TJ, Cole TJ. Uncritical use of bone mineral density in absorptiometry may lead to size-related artifacts in the identification of bone mineral determinants. *Am J Clin Nutr* 1994; 60: 837–42.
13. Prentice A, Cole TJ, Whitehead RG. Impaired growth in infants born to mothers of very high parity. *Hum Nutr: Clin Nutr* 1987; 41C: 319–25.
14. Freeman JV, Cole TJ, Chinn S, Jones PRM, White EM, Preece MA. Cross-sectional stature and weight reference curves for the UK, 1990. *Arch Dis Child* 1995; 73: 17–24.