

## Spread of Epidemic Keratoconjunctivitis Due to a Novel Serotype of Human Adenovirus in Japan<sup>▽</sup>

We have reported a novel human adenovirus (HAdV) that has caused a nationwide epidemic of keratoconjunctivitis (EKC) in Japan (11). This virus has been characterized serologically and genetically as a novel serotype, and we propose naming it HAdV-54. The GenBank accession number of the nucleotide sequence of HAdV-54 is AB333801.

Adenovirus EKC is typically caused by three serotypes of HAdV-D, namely, HAdV-8, -19, and -37, and one serotype of HAdV-E, namely, HAdV-4. These strains frequently cause nosocomial outbreaks (3–5, 18) and were originally isolated from patients with EKC in 1951, 1955, and 1976 (8, 9, 12). Although several variants, including a novel hexon-chimeric intermediate type (HAdV-22,37/H8), have also been described (1, 2, 6, 7, 10, 14, 17), no other new serotype has appeared. HAdV-54 was first isolated from an EKC patient in Kobe, western Japan, in 2000 and then caused an EKC epidemic in 2000 to 2007. During this epidemic, 343 HAdV isolates were obtained from patients with EKC in 16 nosocomial and 15 sporadic outbreaks. HAdV-54 caused 5 (31.3%) of the 16 nosocomial outbreaks and 5 (33.3%) of the 15 sporadic outbreaks. Four other serotypes accounted for 51.6% and HAdV-22,37/H8 was responsible for 16.1% of both types of infections.

HAdV-54 continued to cause EKC in 2008 (Fig. 1). It has been reported only in Japan, but it is not yet known whether HAdV-54 is limited to this country. Also, the origin and route

of transmission of HAdV-54 are unknown. The phylogenetic analysis indicates that HAdV-54 had already appeared in 1995 as the causative agent of EKC. On the other hand, HAdV-54 did not originate from the specimens collected in Saudi Arabia, Austria, Nepal, Bangladesh, and Vietnam (13, 15, 16). It is also possible that this virus is circulating in humans with asymptomatic infections, although no asymptomatic infection was detectable in the present study. The virus might gain pathogenesis by some unknown mechanism. In cell culture, HAdV-54 grows slowly, and the Kobe strain was isolated after eight blind passages. Because this strain is close to HAdV-8 according to both neutralization assays and phylogenetic analyses, it has been diagnosed occasionally as HAdV-8. In Japan, approximately 1,000 HAdV strains were isolated from patients with EKC in 2003 to 2007 (Infectious Agents Surveillance Report URL: <http://idsc.nih.gov/iasr/29/338/tpc338.html>). The major causative agents were HAdV-4, -19a, and -37, and several infections have also been caused by HAdV-8. As EKC infections due to HAdV-8 have apparently been rare since 2000, we undertook phylogenetic analyses to confirm the causative HAdV strains by using partial hexon sequences derived from all prototypes of HAdV serotypes, including the novel serotype HAdV-54. We found that HAdV-8 strains reported during 2003 to 2005 were actually HAdV-54. The novel serotype




FIG. 1. Nosocomial and sporadic EKC infections analyzed in Japan during 2000 to 2008. A total of 343 HAdV isolates were obtained from patients with EKC in 16 nosocomial (red circles) and 15 sporadic (blue circles) outbreaks during 2000 to 2008. HAdV-54 was isolated from patients with nosocomial (red) and sporadic (blue) infections. (The map serving as the basis for this figure is reproduced from <http://www.freemap.jp/>).

HAdV-54 should therefore be monitored for worldwide as an emerging adenoviral infection.

## REFERENCES

1. Adrian, T., U. Wolf, H. J. Lauer, and R. Wigand. 1990. Restriction site mapping of adenovirus type 8 genome types. *Res. Virol.* **141**:611–624.
2. Aoki, K., H. Ishiko, T. Konno, Y. Shimada, A. Hayashi, H. Kaneko, T. Ohguchi, Y. Tagawa, S. Ohno, and S. Yamazaki. 2008. Epidemic keratoconjunctivitis due to the novel hexon-chimeric-intermediate 22,37/H8 human adenovirus. *J. Clin. Microbiol.* **46**:3259–3269.
3. Aoki, K., M. Kato, H. Ohtsuka, K. Ishii, Nakazono, and H. Sawada. 1982. Clinical and aetiological study of adenoviral conjunctivitis, with special reference to adenovirus type 4 and 19 infections. *Br. J. Ophthalmol.* **66**:776–780.
4. Aoki, K., R. Kawana, I. Matsumoto, G. Wadell, and J. C. de Jong. 1986. Viral conjunctivitis with special reference to adenovirus type 37 and enterovirus 70 infection. *Jpn. J. Ophthalmol.* **30**:158–164.
5. Aoki, K., and Y. Tagawa. 2002. A twenty-one year surveillance of adenoviral conjunctivitis in Sapporo, Japan. *Int. Ophthalmol. Clin.* **42**:49–54.
6. Ariga, T., Y. Shimada, K. Ohgami, Y. Tagawa, H. Ishiko, K. Aoki, and S. Ohno. 2004. New genome type of adenovirus serotype 4 caused nosocomial infections associated with epidemic conjunctivitis in Japan. *J. Clin. Microbiol.* **42**:3644–3648.
7. Ariga, T., Y. Shimada, K. Shiratori, K. Ohgami, S. Yamazaki, Y. Tagawa, M. Kikuchi, Y. Miyakita, K. Fujita, H. Ishiko, K. Aoki, and S. Ohno. 2005. Five new genome types of adenovirus type 37 caused epidemic keratoconjunctivitis in Sapporo, Japan, for more than 10 years. *J. Clin. Microbiol.* **43**:726–732.
8. de Jong, J. C., R. Wigand, G. Wadell, D. Keller, C. J. Muzerie, A. G. Wermenbol, and G. J. Schaap. 1981. Adenovirus 37: identification and characterization of a medically important new adenovirus type of subgroup D. *J. Med. Virol.* **7**:105–118.
9. Desmyter, J., J. C. De Jong, K. W. Slaterus, and H. Verlaeckaert. 1974. Keratoconjunctivitis caused by adenovirus type 19. *Br. Med. J.* **iv**:406. (Letter.)
10. Engelmann, I., I. Madisch, H. Pommer, and A. Heim. 2006. An outbreak of epidemic keratoconjunctivitis caused by a new intermediate adenovirus identified by molecular typing. *Clin. Infect. Dis.* **43**:e64–e66.
11. Ishiko, H., Y. Shimada, T. Konno, A. Hayashi, T. Ohguchi, Y. Tagawa, K. Aoki, S. Ohno, and S. Yamazaki. 2008. Novel human adenovirus causing nosocomial epidemic keratoconjunctivitis. *J. Clin. Microbiol.* **46**:2002–2008.
12. Jawetz, E., S. Kimura, A. N. Nicholas, P. Thygeson, and L. Hanna. 1955. New type of APC virus from epidemic keratoconjunctivitis. *Science* **122**:1190–1191.
13. Jin, X. H., H. Ishiko, T. H. Nguyen, T. Ohguchi, M. Akanuma, K. Aoki, and S. Ohno. 2006. Molecular epidemiology of adenoviral conjunctivitis in Hanoi, Vietnam. *Am. J. Ophthalmol.* **142**:1064–1066.
14. Noda, M., Y. Miyamoto, Y. Ikeda, T. Matsui, and T. Ogino. 1991. Intermediate human adenovirus type 22/H10,19,37 as a new etiological agent of conjunctivitis. *J. Clin. Microbiol.* **29**:1286–1289.
15. Ohno, S. 2005. Studies on exogenous and endogenous factors associated with various ocular diseases. *J. Jpn. Ophthalmol. Soc.* **109**:885–915. (In Japanese.)
16. Teuchner, B., M. Nagl, A. Schidlbauer, H. Ishiko, E. Dragosits, H. Ulmer, K. Aoki, S. Ohno, N. Mizuki, W. Gottardi, and C. Larcher. 2005. Tolerability and efficacy of *N*-chlorotaurine in epidemic keratoconjunctivitis—a double-blind, randomized, phase-2 clinical trial. *J. Ocul. Pharmacol. Ther.* **21**:157–165.
17. Wadell, G., and J. C. De Jong. 1980. Restriction endonucleases in identification of a genome type of adenovirus 19 associated with keratoconjunctivitis. *Infect. Immun.* **27**:292–296.
18. World, W. S. M., and M. S. Horowitz. 2007. Adenoviridae, p. 2395–2436. *In* D. M. Knipe, P. M. Howley, D. E. Griffin, R. A. Lamb, M. A. Martin, B. Roizman, and S. E. Straus (ed.), *Fields virology*, fifth ed., vol. II. Lippincott Williams & Wilkins, Philadelphia, PA.

**Hiroaki Ishiko\***

*Host Defense Laboratory  
Mitsubishi Chemical Medicine Corporation  
Tokyo 108-8559, Japan*

**Koki Aoki**

*Department of Ophthalmology and Visual Sciences  
Hokkaido University Graduate School of Medicine  
Sapporo 060-8638, Japan*

\*Phone: 81-3-6722-4240

Fax: 81-3-6722-4231

E-mail: Ishiko.Hiroaki@mp.medicence.co.jp

<sup>v</sup>Published ahead of print on 6 May 2009.