

Bell-like homeodomain selectively regulates the high-irradiance response of phytochrome A

Roberto J. Staneloni^a, María José Rodríguez-Batiller^a, Danilo Legisa^a, María R. Scarpin^a, Adamantia Agalou^b, Pablo D. Cerdán^a, Annemarie H. Meijer^b, Pieter B. F. Ouwkerk^b, and Jorge J. Casal^{c,1}

^aFundación Instituto Leloir, Instituto de Investigaciones Bioquímicas Buenos Aires, Consejo Nacional de Investigaciones Científicas y Técnicas and Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, C1405BWE Buenos Aires, Argentina; ^bDepartment of Molecular and Developmental Genetics, Institute of Biology, Leiden University, Wassenaarseweg 64, 2333 AL, Leiden, The Netherlands; and ^cInstituto de Investigaciones Fisiológicas y Ecológicas Vinculadas a la Agricultura, Facultad de Agronomía, Universidad de Buenos Aires and Consejo Nacional de Investigaciones Científicas y Técnicas, Avenue San Martín 4453, 1417 Buenos Aires, Argentina

Communicated by Anthony R. Cashmore, University of Pennsylvania, Philadelphia, PA, June 14, 2009 (received for review December 11, 2007)

Plant responses mediated by phytochrome A display a first phase saturated by transient light signals and a second phase requiring sustained excitation with far-red light (FR). These discrete outcomes, respectively so-called very-low-fluence response (VLFR) and high-irradiance response (HIR), are appropriate in different environmental and developmental contexts but the mechanisms that regulate the switch remain unexplored. Promoter analysis of a light-responsive target gene revealed a motif necessary for HIR but not for VLFR. This motif is required for binding of the Bell-like homeodomain 1 (BLH1) to the promoter in *in vitro* and in yeast 1-hybrid experiments. Promoter substitutions that increased BLH1 binding also enhanced HIR. *blh1* mutants showed reduced responses to continuous FR and to deep canopy shadelight, but they retained normal responses to pulsed FR or red light and unfiltered sunlight. BLH1 enhanced *BLH1* expression and its promotion by FR. We conclude that BLH1 specifically regulates HIR and not VLFR of phytochrome A.

Arabidopsis thaliana | promoter | TALE

When dark-grown seedlings are exposed to light, stem extension is arrested while foliage growth, pigmentation, and photosynthetic capacity are promoted. This developmental transition is largely mediated by phytochrome B (phyB) when the seedlings are exposed to red light (R) and by phytochrome A (phyA) when the seedlings are exposed to far-red light (FR) (1, 2). phyA initiates 2 photobiologically discrete response phases (3). One, often called very-low-fluence response (VLFR), saturates with infrequent excitation of phyA (e.g., 1 brief pulse every hour) with very low light fluence rates. The other phase, called high-irradiance response (HIR), operates above a threshold of excitation frequency and requires higher fluence rates of FR (3–5).

Processing of the incoming information by the cellular signaling circuits results in a wide range of input-output relationships that serve specific functional purposes (6, 7). The duality of phyA-mediated photobiology adds tools to achieve a versatile control of development by light. The VLFR to brief light exposures, such as those experienced by seeds during soil tillage, stimulates germination under favorable conditions (8). The HIR initiates the synthesis of photoprotective pigments only when the seedlings experience prolonged exposures to higher fluence rates (9). The expression of light-harvesting chlorophyll a-b binding protein (*Lhcb*) genes can exhibit both VLFR and HIR, providing a rapid reaction in response to the first photons when the seedling begins to emerge from the soil (VLFR) and a stronger promotion when excitation becomes persistent (HIR) (10). The transition between VLFR and HIR provides a suitable and functionally meaningful model to investigate mechanisms of regulation of input-output relationships in multicellular organisms.

The *phyA-302* allele, harboring a Glu-777 to Lys substitution in the PAS2 domain of the C terminus, retains near-normal

VLFR and lacks HIR (11). Under continuous FR, the formation of nuclear speckles containing phyA, and phyA degradation are reduced in *phyA-302* providing correlative evidence for a role of these processes in HIR. EID1 is an F-box protein that negatively regulates the HIR of phyA and has little effect on VLFR (12, 13). During the early steps of de-etiolation under continuous FR, phyA targets a relatively large number of transcription factors (14). However, no transcription factor is known to affect selectively the HIR of its target genes. Previously we observed that a target gene promoter contained *cis*-acting elements, required for both VLFR and HIR, and a 42-bp region between –176 and –134 specifically required for HIR (10). Here, we refine the definition of this HIR-specific motif and use this information to identify a transcription factor by using yeast 1-hybrid screens. We also show that this factor, the Bell-like homeodomain 1 (BLH1) protein, is specifically involved in the HIR but not in VLFR.

Results

Identification of a Sequence Specifically Required for HIR in the *Lhcb1*2* Promoter. Hourly pulses of FR (3 min, 60 $\mu\text{mol m}^{-2} \text{s}^{-1}$) saturate the VLFR phase. The difference between the effects of hourly and continuous FR defines the HIR, which operates above a threshold of excitation frequency. We first narrowed down the sequence containing a *cis*-acting motif specifically required for HIR to the –152 to –134 region of the *Lhcb1*2* promoter (Fig. S1A) (10). Subsequent substitution analysis carried out in the –152 to +67 *Lhcb1*2* promoter context shows that none of the mutations affected GUS activity in seedlings incubated in darkness or under hourly R or FR. Substitution of TGGA between –138 and –135 bp by CCCC abrogated the HIR because it reduced GUS activity under continuous FR to the level observed under hourly FR and severely impaired the fluence-rate dependency typical of the HIR (Fig. 1A and Fig. S1B and C). The TGGA sequence is found in *Lhcb* genes of different species and shows a conserved position relatively to other motifs, including AAAATCT and GATA (15) (Fig. S2A). Substitution of GATA motifs and of ATCT within the AAAATCT motif in the context of the –453 to +67 *Lhcb1*2* promoter caused reduced GUS activity in transgenic *Arabidopsis* grown under continuous FR, hourly pulses of R or FR, or darkness (Fig. S2B). Thus, while TGGA is specifically required for HIR, other motifs are required for multiple phytochrome-mediated responses.

Author contributions: R.J.S. and J.J.C. designed research; R.J.S., M.J.R.-B., D.L., M.R.S., A.A., P.D.C., A.H.M., P.B.F.O., and J.J.C. performed research; R.J.S. and J.J.C. analyzed data; and R.J.S. and J.J.C. wrote the paper.

The authors declare no conflict of interest.

¹To whom correspondence should be addressed. E-mail: casal@ifeva.edu.ar.

This article contains supporting information online at www.pnas.org/cgi/content/full/0906598106/DCSupplemental.




Fig. 1. BLH1 binds the HIR box of the *Lhcb1*2* promoter of *Nicotiana plumbaginifolia*, which is required for HIR but not for VLFR of phyA or for phyB-mediated responses. (A) Etiolated seedlings were exposed to hourly pulses of FR, continuous FR ($60 \mu\text{mol m}^{-2} \text{s}^{-1}$), or hourly pulses of R or remained as dark controls. Substitutions to CCCC affecting the indicated 4 bp in the context of the -152 to $+67$ promoter define the sequence required for HIR as TGGGA between -138 and -135 bp (i.e., -135 , see sequences in Fig. S1B). Data are means and SE of 5 independent transgenic lines. For each line, activity was normalized to the values under R pulses, which was unaffected by the substitutions (see SI Text). The interaction between light and promoter is significant at $P < 0.0001$ (2-way ANOVA), and the difference between hourly and continuous FR is significant for all promoters except -135 ($P < 0.01$, Bonferroni post-tests). (B) BLH1 protein binds the HIR box in a yeast 1-hybrid system in a sequence-specific manner. Yeast strains transformed with pACT-BLH1 fail to grow in the absence of histidine and the presence of 2 mM 3-AT if the bait reporter construct carries a TGGGA to CCCC substitution between -138 and -135 bp. (C) BLH1 protein binds the HIR box in vitro in a sequence-specific manner. Left: Electrophoretic mobility shift assay with recombinant BLH1 ($5 \mu\text{g}$) protein and labeled probes (10 femtomol) containing the TGGGA motif (HIRa or HIRb) or the CCCC substitution (HIRa m or HIRb m). Right:

The Homeodomain Transcription BLH1 Factor Binds the TGGGA Motif In Vivo. We conducted a yeast 1-hybrid screening of an *Arabidopsis thaliana* cDNA library from dark-grown seedlings to identify proteins binding the motif required for HIR. As bait we used a tandem of 4 repeats of a 23-bp sequence composed of the -141 to -129 sequence of the promoter followed by 10 bp complementary to the -141 to -132 fragment in inverted order. The yeast reporter strain transformed with an empty *HIS3* reporter vector was able to grow on medium without histidine, but not if it contained 2 mM 3-amino-1,2,4 triazole (3-AT) (Fig. S3). A 1.5-kb clone (pACT-BLH1) was identified on 2 independent occasions, its sequence corresponds to the BLH1 (At2g35940) protein lacking the 50 aa at the amino terminal end and 168 aa at the carboxy terminal end, but retaining the Bell domain, the homeodomain, and extended sequences of poor similarity to other BLH proteins (16).

To further characterize the binding of BLH1, we designed another set of *HIS3* reporters based on the -141 to -119 bp sequence of the *Lhcb1*2* promoter. This resulted in 6 additional plasmids carrying various 4-bp substitutions equivalent to those used in Fig. 1A. When transformed with pACT-BLH1, all strains were able to grow in the absence of histidine (Fig. S3), but those carrying the TGGGA to CCCC substitution between -138 and -135 bp failed to grow in the presence of 2 mM 3-AT (135 in Fig. 1B and Fig. S3).

BLH1 Binds the TGGGA Motif in Vitro. Recombinant BLH1 derived from the 1.5-kb clone identified in yeast screenings was purified from *Escherichia coli* as a GST-BLH1 protein fusion. Gel shift analysis of recombinant BLH1 with 2 radioactively labeled HIR probe variants resulted in complex formation (HIRa and HIRb in Fig. 1C, Left). The same TGGGA to CCCC substitution that abolished activation in a yeast 1-hybrid assay strongly reduced in vitro binding of BLH1 (HIRa m and HIRb m in Fig. 1C, Left). Lanes with control GST extracts did not show any complexes. The HIR probes containing a TGGGA to CCCC substitution did not compete with the HIRb fragment for BLH1 binding (compare HIRa versus HIRa m and HIRb versus HIRb m in Fig. 1C, Right).

Super HIR Promoter Shows Enhanced Binding by BLH1. Before using the promoter-GUS fusions carrying 4-bp substitutions, we had constructed fusions bearing 6-bp substitutions in the -453 bp promoter context. Disruption of the TGGGA core motif in the -135 substitution reduced the HIR, but the effect was not complete, very likely because it created a TGGT in a close location (-135 in Fig. 1D and SI Materials and Methods). Noteworthy, the -129 substitution showed a HIR that more than doubled the HIR of the control promoter (Fig. 1D). The -129 substitution incorporated GGTACC between -134 and -129 bp, and this created a TGGAGGT sequence between -137 and -132 , which could account for the enhanced HIR via stronger binding of a transcription factor. Consistently with this inter-

Competition experiment where BLH1 ($5 \mu\text{g}$, except first lane with $0 \mu\text{g}$ and third lane with $10 \mu\text{g}$) was incubated with labeled HIRb in the presence of 1000-, 500-, or 100-fold excess of the indicated unlabeled probe. cx, complex; fp, free probe. (D) Super HIR *Lhcb1*2* promoter shows enhanced HIR and binding by BLH1. At least 5 independent transgenic lines were used for each 6-bp substitution in the -453 *Lhcb1*2* promoter context. The inset shows enhanced growth of yeast strains transformed with pACT-BLH1 in the absence of HIS and the presence of 25 mM 3-AT if the bait reporter construct carries a TGGAGGT sequence (Super HIR) repeated 4 times compared to the WT HIR sequence repeated 4 times. Relative intensity of yeast colonies is the brightness intensity of yeast colonies photographed against a dark background and scanned with Photoshop software, expressed relative to the brightness with 0 mM 3-AT (means and SE of 3 experiments, $P < 0.001$ with 25 mM 3-AT).




Fig. 2. Impaired HIR and normal VLFR in *blh1* mutants. (A) *Lhcb1*2* promoter activity. Plants of the *blh1-1* mutant and of the *BLH1* siblings were transformed with the -453 *Lhcb1*2* promoter or the -453 *Lhcb1*2* promoter where TGGA was substituted by CCCC (-135). Etiolated seedlings were exposed to hourly pulses of FR, continuous FR ($60 \mu\text{mol m}^{-2} \text{s}^{-1}$), or hourly pulses of R or remained as dark controls. Data are means and SE of 5 independent transgenic lines for each transgene/background combination. For each line, activity was normalized to the values under R pulses, unaffected by the -135 substitution or *blh1* mutation (SI Text). (B–E) Hypocotyl growth inhibition relative to the length in dark controls (length in darkness, mm: WT = 13.4 ± 0.1 , *blh1* = 13.1 ± 0.2 , $P > 0.1$), cotyledon angle (photographs under continuous FR), anthocyanin levels, and hypocotyl angle. One-day-old seedlings were grown for 3 days under hourly pulses of FR, continuous FR, or hourly pulses of R, or remained as dark controls before measurements. The fluence rate of continuous FR was 0.3 (B), 0.5 (C), 10 (D), or 0.5 (E) $\mu\text{mol m}^{-2} \text{s}^{-1}$. Data are means and SE of at least 3 replicate boxes. ANOVA followed by Bonferroni post-tests indicate significant effects of *blh1* ($P < 0.001$) under continuous FR but not under any other light condition. The *blh1* mutation had no effects in the *phyA-211* mutant background under continuous FR (dark control data included in brackets). Inhibition of hypocotyl growth: *phyA* = 0.02 ± 0.02 (0.00 ± 0.02); *phyA blh1* = 0.02 ± 0.03 (0.00 ± 0.01); cotyledon angle (degrees): *phyA* = 1 ± 0 (1 ± 1); *phyA blh1* = 1 ± 0 (1 ± 0); anthocyanin (absorbance units): *phyA* = 0.01 ± 0.00 (0.01 ± 0.00); *phyA blh1* = 0.01 ± 0.00 (0.01 ± 0.00); hypocotyl angle (randomization of growth orientation) (degrees): *phyA* = 6 ± 1 (8 ± 1); *phyA blh1* = 6 ± 1 (8 ± 1). (F) Fluence rate-response curve of anthocyanin levels. Data are means and SE of 3 replicate boxes. ANOVA followed by Bonferroni post-tests indicate significant effects of *blh1* under 10 ($P < 0.001$) and 100 ($P < 0.05$) $\mu\text{mol m}^{-2} \text{s}^{-1}$. (G) When expressed in the *blh1-1* background under the control of the CaMV 35S promoter, *BLH1* inhibits hypocotyl growth even under pulsed FR. One-day-old seedlings were grown for 3 days in darkness or under hourly FR or continuous FR ($0.1 \mu\text{mol m}^{-2} \text{s}^{-1}$) before measurements of hypocotyl length. Data are means and SE of at least 3 replicate boxes. Each transgenic line is significantly different ($P < 0.05$) from *blh1* under hourly FR and under continuous FR (symbols as in A). *BLH1* expression levels in *blh1-1*, WT, and different transgenic lines are shown at the top. (H) Phenotype of the *blh1-1* mutant under a dense canopy. One-day-old seedlings of the WT Col, the *phyA-201* mutant, and the *blh1-1* mutant were grown for 3 days under a dense canopy of *Cynodon dactylon* before measurements. At a clear midday, the FR irradiance beneath the canopy was in average $1 \mu\text{mol m}^{-2} \text{s}^{-1}$, and the R/FR ratio was 0.01. The seedlings were grown in the same plastic boxes used for indoor experiments. Dark controls were placed beneath the canopy but wrapped in black plastic and aluminum foil, and unfiltered sunlight controls were grown outside the canopy. Under the canopy, the inhibition observed in *blh1-1* is significantly smaller ($P < 0.01$) than in the WT according to ANOVA and a Bonferroni post-test. Data are means and SE of 6 replicate boxes from 2 independent experiments.

pretation, a combination of pACT-BLH1 with a construct containing the TGGAGGT sequence (super HIR box) allowed enhanced growth of yeast under increasing 3-AT concentrations when compared to the construct containing the wild-type (WT) promoter (Fig. 1D, Inset, and Fig. S3F).

Normal HIR Require BLH1. To investigate the physiological significance of BLH1, we obtained homozygous *blh1-1*, *blh1-4*, and *blh1-5* mutant seedlings, which show reduced *BLH1* expression, and the corresponding WT *BLH1* siblings (Fig. S4A and B). In *blh1* hypocotyl length, hypocotyl angle, cotyledon angle, anthocyanin levels, and *Lhcb1*2* promoter activity were WT in darkness and under hourly pulses of R or FR. However, *blh1* showed significantly reduced responses to continuous FR (Fig. 2 and Fig. S4C and D). The *phyA* and *phyA blh1* mutants were indistinguishable and blind to FR (Fig. 2). The promotion of the endogenous *Lhcb* expression measured by real-time PCR was also reduced in *blh1* under continuous FR (Darkness: WT = 1 ± 0 ; *blh1-1* = 1 ± 0 ; hourly FR: WT = 138 ± 19 ; *blh1-1* = 153 ± 33 ; continuous FR: WT = 467 ± 98 ; *blh1-1* = 246 ± 39 ; $P < 0.05$). The induction of seed germination by a single pulse of long-wavelength FR, which is a typical VLFR (17, 18), was normal in *blh1* (Fig. S4E). This demonstrates that *blh1* mutations

have specific effects on the HIR without altering the VLFR or phyB-mediated responses.

We produced independent transgenic lines of the *blh1-1* background expressing the *BLH1* gene under the control of the CaMV 35S promoter (construct 35S-BLH1) at higher levels than the WT (lines 25 and 10) or at a lower overall level than the WT (but with a different tissue distribution predicted by the use of the CaMV 35S promoter) (Fig. 2G). These lines showed HIR (difference between hourly and continuous FR) even at a fluence rate ($0.1 \mu\text{mol m}^{-2} \text{s}^{-1}$), where HIR was weak in the WT and fully absent in *blh1-1*, indicating that the *BLH1* transgene rescued the mutant phenotype (Fig. 2G). More importantly, although the *blh1* mutation did not affect the response to hourly pulses of FR, the transgenic lines ectopically expressing *BLH1* showed enhanced responses to hourly pulses of FR (Fig. 2G and Fig. S5). This indicates that overexpression of *BLH1* bypasses the need of continuous FR to make BLH1 physiologically effective. This gain-of-function phenotype of the *BLH1* transgenics was absent in the *phyA* mutant background (Fig. S5B), indicating that the enhanced response to hourly FR induced by overexpressed *BLH1* is mediated by phyA. Overexpression of *BLH1* did not affect phyB-mediated signaling under hourly R (Fig. S5C).

The *blh1* mutant showed reduced hypocotyl growth inhibition




Fig. 3. BLH1 promotes its own expression. (A) BLH1 enhances the expression of *BLH1* under FR. Transgenic seedlings bearing *BLH1::GUS* were grown in darkness for 2 days and then exposed to hourly FR, continuous FR ($60 \mu\text{mol m}^{-2} \text{s}^{-1}$), or no light for 24 h before harvest. *BLH1* expression was also analyzed by real-time PCR in WT seedlings exposed for 6 h to the different light/dark treatments. Data are means and SE of 3 biological replicates (real-time PCR) or of 5 (WT) or 3 (*blh1*) independent transgenic lines bearing the *BLH1* promoter fused to GUS. The effect of continuous FR is significant ($P < 0.001$) in the WT but not ($P > 0.05$) in *blh1*. (B) BLH1 enhances *BLH1::GUS* activity in darkness and its response to continuous FR. Data are means and SE of 6 biological replicates from representative transgenic lines in the WT or *blh1-1* background. The slopes are significantly different ($P < 0.002$).

under a dense canopy, a condition where phyA is required for de-etiolation (Fig. 2H). The latter observation highlights the biological significance of BLH1 control of *Arabidopsis* photomorphogenesis.

Transcriptome of *blh1*. To investigate the effect of *blh1* on the transcriptome, WT and *blh1-1* seedlings were grown in darkness for 2 days and then transferred to continuous FR ($60 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 6 h before harvest. Dark controls were harvested simultaneously without exposure to FR. The effects of FR were consistent with previous reports (Fig. S6A). On average, the *blh1* mutation caused a 60% reduction of the response to FR (Fig. S6B). We used factorial ANOVA followed by clustering to select the genes showing little difference between the WT and *blh1* in darkness and reduced response to FR in the mutant (Table S1). Three clusters (125 genes) grouped genes with reduced promotion of expression by FR in *blh1*. Plastid-related genes were highly overrepresented [normed frequency \pm bootstrap SD (19): 9.7 ± 2.1] in these clusters. Two clusters (89 genes) grouped genes with reduced inhibition of expression by FR in *blh1*. Genes related to endoplasmic reticulum (8.3 ± 3.8) and receptor binding or activity (4.0 ± 2.3) were highly overrepresented in these clusters. Among the genes significantly affected in *blh1*, each 1 of the 3 possible 6-mer promoter sequences of the *Lhcb1*2* promoter containing the functional TGGGA motif was overrepresented (GATGGA: $P < 0.006$, TGGAAAT: $P < 0.03$, and ATGGAT: $P < 0.03$).

BLH1 Promotes BLH1 Promoter Activity. The *BLH1* gene showed expression in darkness, and this is consistent with the fact that it was found in a library from dark-grown *Arabidopsis*. *BLH1* promoter activity was reduced in dark-grown seedlings of the *blh1* mutant compared to the WT (Fig. S7A). In the WT, continuous FR increased *BLH1* expression (20) to a larger extent than hourly pulses of FR (Fig. 3A). The *blh1* mutant showed reduced response to continuous FR (Fig. 3A and Fig. S7B). GUS activity driven by the *BLH1* promoter showed FR fluence rate

dependency over a wide range of fluence rates, and this response was severely reduced in the *blh1* background (Fig. 3B).

Discussion

We have narrowed-down the HIR-specific motif of *Lhcb1*2*, which is not required for VLFR or phyB-mediated responses, to the 4-bp core sequence TGGGA (Fig. 1). “Phylogenetic footprinting” indicates that the TGGGA motif is present at relatively conserved distance from CCAAT and GATA motifs (15) in a number of *Lhcb* genes from different species (Fig. S2A), suggesting that it might be a binding site conserved by natural selection (21). Using a bait based on this sequence in combination with a library from dark-grown *A. thaliana* seedlings in the yeast 1-hybrid system we identified the homeodomain transcription factor BLH1. BLH1 belongs to the 3-amino acid loop extension (TALE) superclass of homeobox genes. In vivo and in vitro binding experiments confirmed the specific interaction between BLH1 and TGGGA (Fig. 1B and C). The TGGGA core is in the TGATGGATTA sequence of the tobacco *Lhcb1*2* promoter, which (i) differs only 2 bases from the central sequence of the ATGATAGATAAT segment from the nitrate reductase 2 gene of *Arabidopsis*, previously shown to bind in vitro BLH1 protein obtained from nuclear extracts of *Arabidopsis* roots (22); (ii) contains ATTA, the canonical binding site for most homeodomains, preceded by GG, which provides binding specificity for homeodomains containing a Lys residue at the position 50 in the recognition helix (23); and (iii) presents a full match to TGATXXATTA, the consensus binding site of heterodimers of PBC/Hox TALE proteins (24). Noteworthy, when we replaced the bases following the TGGGA motif to obtain TGGAGGT, the promoter showed both stronger binding of BLH1 in yeast and stronger HIR (Fig. 1D), providing gain-of-function evidence for the role of BLH1 binding to the promoter in HIR. The human homeodomain CCAAT-displacement protein (CDP)/Cux binds ATGGAT (25), the same core motif reported here for BLH1. In yeast and in vitro binding experiments, we used the originally identified *BLH1* clone, which lacks some amino acids at both ends. The full-length *BLH1* clone failed to form complexes in in vitro binding assays and yeast 1-hybrid experiments, and this resembles the behavior of CDP, a well-established case of auto-inhibition involving the N-terminal end of the protein (25, 26). The full-length maize BLH KIP does not bind DNA in the absence of maize Knotted-1 (27). Since BLH1 in nuclear extracts does form complexes with DNA sequences similar to those reported here, and these complexes contain other (unidentified) homeodomain proteins (22), these proteins could be required for cooperative binding of full-length BLH1 to DNA (28). The presence of the ATGGAT sequence in the 500-bp upstream region of genes with altered expression in the *blh1* mutant is higher than expected by chance.

In *blh1* seedlings, hypocotyl length, hypocotyl angle, cotyledon angle, anthocyanin levels, and *Lhcb1*2* promoter activity were normal in darkness and showed normal responses to hourly pulses of FR or R, but reduced responses to continuous FR (Fig. 2). Gene expression responses to 6 h of high-fluence continuous FR were reduced in average in a 60% (Fig. S6B). The VLFR of seed germination was also normal in *blh1* (Fig. S4E). Therefore, *blh1* mutations and mutations of a DNA motif required for BLH1 binding had specific effects on the HIR without altering the VLFR or phyB-mediated responses. The partial effect of *blh1* on HIR could reflect redundancy, which is not uncommon among TALE proteins (29, 30). *BLH6* shows strong promotion of expression in response to continuous FR (20) and *BLH5* shows the strongest similarity to *BLH1*. Both *blh6* and *blh5* mutations reduce HIR (Fig. S8). The expression of *BLH1* and its promotion by continuous FR depend strongly on BLH1 (Fig. 3A and B, and Fig. S7).

The TALE superclass of homeobox genes is conserved among animals, fungi, and plants, where they are required for meristem maintenance and proper regulation of shoot and leaf development in response to endogenous cues (31, 32). Both, TALE proteins and the HIR of *phyA* have been implicated in plant evolution, the former in the evolution of land plants from green algal ancestors and the diversification of the plant body (32), the latter in the adaptation to the dimly lit understory habitats experienced by the earliest angiosperms (33). Here, we relate BLH1 to HIR, placing a TALE protein as a regulator of plant responses to the specific environmental cue experienced when a seedling emerges from the soil under a deep canopy.

Materials and Methods

Transgenic Lines Carrying Fusions Containing Different Fragments of the *Lhcb1*2* Promoter. The transgenic lines of *A. thaliana* accession Columbia carrying the -453, -176, and -134 to +67 fragments of the *Lhcb1*2* promoter of *Nicotiana plumbaginifolia* fused to the coding region of *gusA* have been described previously (10). We continued to use this tobacco promoter to take advantage of previous studies and because the distribution of motifs important for promoter activity is very similar to that of *Lhcb1*1* and *Lhcb1*2* of *A. thaliana* (Fig. S2). The procedures to obtain transgenic lines were basically as described and are presented in further detail in *SI Text*.

***blh1* T-DNA Mutant and BLH1 OX Transgenic Lines.** The *blh1-1* allele in Columbia [Salk-089095 (34)] containing a T-DNA insertion in exon 6 of *BLH1* was obtained from *Arabidopsis* Biological Resource Center (ABRC) through The *Arabidopsis* Information Resource at www.arabidopsis.org. To rescue *blh1-1* seedling with the *BLH1* gene, the *A. thaliana* full-length cDNA (ABRC stock # U18537) (35), cloned into the the *KpnI-Sall* sites of the pPZP221 and pPZP212-based binary vectors CHF1 and CHF3 (36) under the control of the CaMV 35S promoter, were introduced in *Agrobacterium tumefaciens* strain GV3101, and used to transform *blh1-1* plants using the floral dip method (37). Homozygous lines were selected by kanamycin or gentamycin resistance. The *blh1-4* allele in Landsberg *erecta* (CSHL.GT9784) containing a Ds insertion in exon 3 was obtained from Gene Trap CSHL (<http://genetrapp.cshl.org>). The *blh1-5* allele in Columbia (SM.3.16405) containing a T-DNA insertion in exon 3 was obtained from ABRC. The primers used to identify homozygous mutants and WT siblings are provided in *SI Text*.

Yeast 1-Hybrid. Yeast 1-hybrid assays were conducted according to the method described (38) using yeast strain Y187 (Clontech), containing the bait sequence cloned as a *NotI-XbaI* fragment in the pINT1-HIS3NB integrative reporter gene system (GenBank accession no. AY061966) integrated at the nonessential *PDC6* locus (39). We cloned the bait sequence at the *NotI-XbaI* restriction sites

of the integrative vector pINT1-HIS3NB. Yeast cells Y187 were transformed by a lithium acetate procedure, and transformants were selected on YAPD plates with 150 μ g/mL G418 for the presence of the dominant antibiotic *APT1* marker gene present in vector pINT1-HIS3NB. The following sequence was used as baits in cDNA library screenings: 5'-GGCCGCTGATGGATTACAATAATCCATCATGATGGATTACAATAATCCATCATGATGGATTACAATAATCCATCAAGATCTT-3'. Details of yeast transformation and library description are provided in *SI Text*.

Electrophoretic Mobility Shift Assays. Electrophoretic mobility shift assays (EMSA) were carried out essentially as described by (40). Details are provided in *SI Text*.

Physiological Experiments. For the experiments to measure GUS activity driven by the *Lhcb1*2* promoter constructs, inhibition of hypocotyl growth, cotyledon angle, anthocyanin levels, and hypocotyl angle (randomization of growth orientation), the seedlings were grown in plastic boxes (10, 11). Details are provided in *SI Text*.

Analysis of BLH1 and Endogenous Lhcb Expression. We analyzed endogenous *BLH1* and *Lhcb* (combined *Lhcb1*1* and *Lhcb1*2*) expression by real-time PCR. We also constructed transgenic plants bearing the full-length *BLH1* promoter fused to the GUS reporter (*SI Text*).

Microarray Experiment. Seedlings grown in darkness for 2 days after the induction of seed germination were either exposed to 6 h continuous FR or left as dark controls. Two independent samples were harvested in liquid nitrogen for each genotype by FR/dark combination, and total RNA was extracted by using the RNEASY Plant mini kit (Qiagen). cDNA and cRNA synthesis and hybridization to 22 K (ATH1) Affymetrix Gene Chips were performed according to Affymetrix instructions. The scaling tab of the Affymetrix microarray suite in the mode "all probe sets" was used to standardize the trimmed mean signal of each array to the "target signal" according to the manufacturer's instructions. Details of the statistical analysis of the data, the functional classification of the genes, and the analysis of upstream promoter sequences are given in *SI Text*.

ACKNOWLEDGMENTS. We thank the *Arabidopsis* Biological Resource Center (ABRC) for their provision of seeds carrying the Salk-089095 insert, seeds of the line SM.3.16405, the *Arabidopsis* library CD4-22, and the clone U18537 and Cold Spring Harbor Lab for the provision of the CSHL.GT9784 line. This work was financially supported by Agencia Nacional de Promoción Científica y Tecnológica (Argentina) grant nos. PME54 and PICT 11631 and 32492 to J.J.C., University of Buenos Aires grant no. AG021 to J.J.C., Consejo Nacional de Investigaciones Científicas y Técnicas (Argentina) grants 5958 to J.J.C. and 06082 to R.J.S. A.H.M. was supported by the EU FP5 Project TF-STRESS (QLK3-2000-00328). A.A. and P.B.F.O. were supported by the EU FP6 INCO-MPC2 project CEDROME (INCO-CT-2005-015468).

- Quail PH (2002) Phytochrome photosensory signalling networks. *Nature Rev Mol Cell Biol* 3:85-93.
- Chen M, Chory J, Fankhauser C (2004) Light signal transduction in higher plants. *Annu Rev Genet* 38:87-117.
- Casal JJ, Yanovsky MJ, Luppi JP (2000) Two photobiological pathways of phytochrome A activity, only one of which shows dominant negative suppression by phytochrome B. *Photochem Photobiol* 71:481-486.
- Schäfer E, Beggs CJ, Fukshansky L, Holmes MG, Jabben M (1981) A comparative study of the responsivity of *Sinapis alba* L. seedlings to pulsed and continuous irradiation. *Planta* 153:258-261.
- Shinomura T, Uchida K, Furuya M (2000) Elementary processes of photoperception by phytochrome A for high-irradiance response of hypocotyl elongation in *Arabidopsis*. *Plant Physiol* 122:147-156.
- Kramer BP, Fussenegger M (2005) Hysteresis in a synthetic mammalian gene network. *Proc Natl Acad Sci USA* 102:9517-9522.
- Kobayashi H, et al. (2004) Programmable cells: Interfacing natural and engineered gene networks. *Proc Natl Acad Sci USA* 101:8414-8419.
- Scopel AL, Ballaré CL, Sánchez RA (1991) Induction of extreme light sensitivity in buried weed seeds and its role in the perception of soil cultivations. *Plant Cell Environ* 14:501-508.
- Shin J, Park E, Choi G (2007) PIF3 regulates anthocyanin biosynthesis in an HY5-dependent manner with both factors directly binding anthocyanin biosynthetic gene promoters in *Arabidopsis*. *Plant J* 49:981-994.
- Cerdán PD, et al. (2000) Sustained but not transient phytochrome A signaling targets a region of a *Lhcb1*2* promoter not necessary for phytochrome B action. *Plant Cell* 12:1203-1211.
- Yanovsky MJ, et al. (2002) Missense mutation in the PAS2 domain of phytochrome A impairs subnuclear localization and a subset of responses. *Plant Cell* 14:1591-1603.
- Dieterle M, Zhou Y-C, Schäfer E, Funk M, Kretsch T (2001) EID1, an F-box protein involved in phytochrome A-specific light signaling. *Genes Dev* 15:939-944.
- Zhou Y-C, Dieterle M, Buche C, Kretsch T (2002) The negatively acting factors EID1 and SPA1 have distinct functions in phytochrome A-specific light signaling. *Plant Physiol* 128:1098-1108.
- Tepperman JM, Zhu T, Chang H-S, Wang X, Quail PH (2001) Multiple transcription-factor genes are early targets of phytochrome A signaling. *Proc Natl Acad Sci USA* 98:9437-9442.
- Terzaghi WB, Cashmore AR (1995) Light-regulated transcription. *Annu Rev Plant Physiol Plant Mol Biol* 46:445-474.
- Becker A, Bey M, Bürglin TR, Saedler H, Theissen G (2002) Ancestry and diversity of BEL1-like homeobox genes revealed by gymnosperm (*Gnetum gnemon*) homologs. *Dev Genes Evol* 212:452-457.
- Botto JF, Sánchez RA, Whitelam GC, Casal JJ (1996) Phytochrome A mediates the promotion of seed germination by very low fluences of light and canopy shade light in *Arabidopsis*. *Plant Physiol* 110:439-444.
- Shinomura T, et al. (1996) Action spectra for phytochrome A- and phytochrome B-specific photoinduction of seed germination in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 93:8129-8133.
- Provart N, Zhu T (2003) A browser-based functional classification SuperViewer for *Arabidopsis* genomics. *Curr Comput Mol Biol* 2003:271-272.
- Zimmermann P, Hirsch-Hoffmann M, Hennig L, Gruissem W (2004) GENEVESTIGATOR. *Arabidopsis* microarray database and analysis toolbox. *Plant Physiol* 136:2621-2632.
- Wray GA, et al. (2003) The evolution of transcriptional regulation in eukaryotes. *Mol Biol Evol* 20:1377-1419.
- Sherameti I, et al. (2005) The endophytic fungus *Piriformospora indica* stimulates the expression of nitrate reductase and starch-degrading enzyme glukon-water dikinase in tobacco and *Arabidopsis* roots through a homeodomain transcription factor that binds to a conserved motif in their promoters. *J Biol Chem* 280:26241-26247.
- Schier AF, Gehring WJ (1992) Direct homeodomain-DNA interaction in the autoregulation of the fushi tarazu gene. *Nature* 356:804-807.

24. Wray GA, et al. (2003) The evolution of transcriptional regulation in eukaryotes. *Mol Biol Evol* 20:1377–1419.
25. Huang C-J, Chang J-G, Wu S-C, Choo K-B (2005) Negative transcriptional modulation and silencing of the Bi-exonic Rnf35 gene in the preimplantation embryo. Binding of the CCAAT-displacement protein/Cux to the untranslated exon 1 sequence. *J Biol Chem* 280:30681–30688.
26. Truscott M, et al. (2004) The N-terminal region of the CCAAT displacement protein (CDP)/Cux transcription factor functions as an autoinhibitory domain that modulates DNA binding. *J Biol Chem* 279:49787–49794.
27. Smith HM, Boschke I, Hake S (2002) Selective interaction of plant homeodomain protein mediates high DNA-binding affinity. *Proc Natl Acad Sci USA* 99:9579–9584.
28. Chen H, Banerjee AK, Hannapel DJ (2004) The tandem complex of BEL and KNOX partners is required for transcriptional repression of *ga20ox1*. *Plant J* 38:276–284.
29. Byrne ME, Simorowski J, Martienssen RA (2002) ASYMMETRIC LEAVES1 reveals *knox* gene redundancy in *Arabidopsis*. *Devel* 129:1957–1965.
30. Selleri L, et al. (2004) The TALE homeodomain protein Pbx2 is not essential for development and long-term survival. *Mol Cell Biol* 24:5324–5331.
31. Hake S, et al. (2004) The role of KNOX genes in plant development. *Annu Rev Cell Dev Biol* 20:125–151.
32. Dolan L (2008) Plant evolution: TALES of development. *Cell* 133:771–773.
33. Mathews S, Burleigh JG, Donoghue MJ (2003) Adaptive evolution in the photosensory domain of phytochrome A in early angiosperms. *Mol Biol Evol* 20:1087–1097.
34. Alonso JM, et al. (2003) Genome-wide insertional mutagenesis of *Arabidopsis thaliana*. *Science* 301:653–657.
35. Yamada K, et al. (2003) Empirical analysis of transcriptional activity in the *Arabidopsis* genome. *Science* 302:842–846.
36. Staiger D, et al. (2003) The *Arabidopsis* SRR1 gene mediates phyB signaling and is required for normal circadian clock function. *Genes Dev* 17:256–268.
37. Clough SJ, Bent AF (1998) Floral dip: A simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* 16:735–743.
38. Ouwerkerk PBF, Meijer AH (2001) in *Current Protocols in Molecular Biology* (John Wiley & Sons), Vol. Supplement 55.
39. Meijer AH, Ouwerkerk PBF, Hoge JH (1998) Vectors for transcription factor isolation and target gene identification by means of genetic selection in yeast. *Yeast* 14:1407–1416.
40. Ouwerkerk PBF, Trimborn TO, Hilliou F, Memelink J (1999) Nuclear factors GT-1 and 3AF1 interact with multiple sequences within the promoter of the *Tdc* gene from Madagascar periwinkle: GT-1 is involved in UV light-induced expression. *Mol Gen Genet* 261:610–622.