

Published in final edited form as:


*Org Lett.* 2008 November 6; 10(21): 4731–4734. doi:10.1021/ol8017038.

## Is Donor-Acceptor Hydrogen Bonding Necessary for 4,6-*O*-Benzylidene-directed $\beta$ -Mannopyranosylation? Stereoselective Synthesis of $\beta$ -*C*-Mannopyranosides and $\alpha$ -*C*-Glucopyranosides

David Crich\* and Indrajeet Sharma

Department of Chemistry, Wayne State University, 5101 Cass Avenue, Detroit, MI 48202, and Department of Chemistry, University of Illinois at Chicago, 845 West Taylor Street, Chicago, IL 60607

### Abstract


 Gluco- $\alpha$ -selective; Manno- $\beta$ -selective

2,3-Di-*O*-benzyl-4,6-*O*-benzylidene-thiohexopyranosides, on activation with 1-benzenesulfinyl piperidine and triflic anhydride, react with allyl silanes and stannanes, and with silyl enoethers to give *C*-glycosides. In mannose the  $\beta$ -isomers are formed selectively whereas the glucose series provides the  $\alpha$ -anomers. This selectivity pattern parallels that of *O*-glycoside formation and eliminates the need to consider donor-acceptor hydrogen bonding in the formation of the *O*-glycosides.

4,6-*O*-Benzylidene protected mannopyranosyl triflates carrying ether-type blocking groups on O2 and O3 are highly  $\beta$ -selective in their reactions with alcohols.<sup>1,2</sup> This selectivity is independent of the nature of the triflate precursor, thioglycoside or sulfoxide, and extends to other classes of glycosyl donor including the imidates,<sup>3</sup> the 2-hydroxycarbonylbenzyl glycosides,<sup>4</sup> and the phosphites.<sup>5</sup> The selectivity, however, is highly dependent on the presence of the benzylidene acetal or related group, as the corresponding tetra-*O*-benzyl or alkyl donors are unselective under comparable conditions.<sup>1</sup> In terms of reaction mechanism, the covalent triflate<sup>6</sup> is understood to serve as a reservoir for a transient contact ion pair (CIP) and, thereafter, for a solvent separated ion pair (SSIP). The CIP, in which the triflate shields the  $\alpha$ -face of the oxocarbenium ion, is  $\beta$ -selective whereas the SSIP is  $\alpha$ -selective in agreement with the dictates of the anomeric effect (Scheme 1).<sup>7</sup> As established by Bols,<sup>8</sup> the benzylidene effect is due to the locking of the C5–C6 bond in the most electron-withdrawing trans-gauche conformation, which destabilizes the oxocarbenium ion and limits the concentration of the undesired,  $\alpha$ -selective SSIP.<sup>9</sup>

dcrich@chem.wayne.edu.

**Supporting Information Available** Full experimental details and copies of NMR spectra for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

In stark contrast the corresponding 4,6-*O*-benzylidene glucosyl donors are  $\alpha$ -selective.<sup>10</sup> With the help of a series of 2- or 3-deoxy and the corresponding deoxy fluoro gluco and mannosyl donors, we have rationalized the glucose/mannose stereoselectivity shift on the basis of the differing O2-C2-C3-O3 torsional interactions as the covalent triflates collapse to the ion pairs.<sup>11</sup> In mannose this torsion angle is compressed as the oxacarbenium ion is formed, whereas in glucose it is relaxed. Effectively, the formation of the glucosyl oxacarbenium ion is less endothermic than that of its mannosyl counterpart leading to a higher concentration of ion pairs in glucose than in mannose.<sup>12</sup>

A potential alternative explanation for the change in selectivity between the glucose and mannose series invokes hydrogen bonding between the OH group of the incoming nucleophilic alcohol and O2 of the donor as the major influence on reaction stereoselectivity, at least for the case of the benzylidene acetal protected donors.<sup>13,14</sup> As we now describe, to address this issue we have investigated the reaction of glucosyl and mannosyl triflates with C-nucleophiles, for which the possibility of hydrogen bonding does not exist.

Thioglycosides **4–5** (Fig. 1) were synthesized as previously described<sup>15</sup> and converted in dichloromethane at  $-60\text{ }^{\circ}\text{C}$  with the combination of 1-benzenesulfinyl piperidine (BSP, Fig. 2)<sup>1b</sup> and trifluoromethanesulfonic anhydride in the presence of the hindered base 2,4,6-*tert*-butylpyrimidine (TTBP, Fig. 2)<sup>16</sup> to the corresponding glycosyl triflates. The nucleophile was then added and the reaction mixture was stirred for a further 2 h at  $-60\text{ }^{\circ}\text{C}$ , before it was quenched at that temperature, leading to the results presented in Table 1.

It is immediately apparent from the results presented in Table 1 that the carbon nucleophiles follow the well-established pattern of the alcohols, with the 4,6-*O*-benzylidene protected mannosyl donors being  $\beta$ -selective, their gluco counterparts  $\alpha$ -selective, and the tetrabenzyl mannosyl donor relatively unselective, ergo, there is no requirement for hydrogen bonding to direct these systems.

Comparison of Table 1, entries 1 and 2 reveals that, not unexpectedly, the more reactive allylstannane gives better  $\beta$ -selectivity than the allylsilane in reactions with the benzylidene protected mannosyl triflate. In the glucoserries (Table 1, entries 7 and 8) complete  $\alpha$ -selectivity is observed regardless of the nature of the allylmetal employed. With the trimethylsilyl enolethers as nucleophiles excellent  $\beta$ -selectivity was again observed for the formation of benzylidene protected mannosyl *C*-glycosides (Table 1, entries, 3,4, and 5), whereas the opposite selectivity was observed for glucose (Table 1, entry 9). Interestingly, with the pinacolone silyl enolether a minor amount of an  $\alpha$ -*O*-glycosyl enolether **8b** was observed as byproduct in coupling to the mannosyl donor **4** (Table 1, entry 3). With the corresponding glucosyl donor **5**, the  $\alpha$ -enolether was formed in greater yield than the corresponding *C*-glycoside (Table 1, entry 9).<sup>17</sup> With trimethylsilyloxycyclohexene as nucleophile the major product **10a** was accompanied by the formation of a double adduct **10b** arising from a subsequent condensation step.

Importantly, the reaction of the tetra-*O*-benzyl thiomannoside **6** with allyl trimethylsilane afforded *C*-glycoside **11** as a mixture of anomers favoring the  $\alpha$ -anomer, thereby revealing the importance of the benzylidene acetal in the control of stereochemistry (Table 1, entry 6). Lucero and Woerpel have discussed the preferential formation of  $\alpha$ -*C*-mannopyranosides from tetra-*O*-benzyl mannopyranosyl acetate and allyl trimethylsilane with activation by trimethylsilyl trifluoromethanesulfonate in terms of a Curtin-Hammett kinetic scheme in which reaction takes place preferentially on the less populated <sup>4</sup>H<sub>3</sub> conformer of the intermediate oxacarbenium ion that is in equilibrium with the more stable <sup>3</sup>H<sub>4</sub> conformer.<sup>18</sup> While the result obtained with the tetra-*O*-benzyl system **6** (Table 1, entry 6) could be interpreted as consistent with the Woerpel model, the  $\beta$ -selective reactions (Table 1, entries 1–5) from the 4,6-*O*-benzylidene mannosyl

donor, for which the intermediate cation is locked in the  ${}^4H_3$  or a closely related conformer clearly require another explanation. Consistent with our explanation for the selective formation of  $\beta$ -*O*-mannosides (Scheme 1) we suggest that the explanation lies in the role of the counterion, with glycosylation occurring via the CIP in the benzylidene protected mannose series.

In conclusion, the benzylidene acetal effect extends beyond the formation of *O*-glycosides to that of *C*-glycosides. The stereoselective formation of  $\beta$ -*C*-mannosides and  $\alpha$ -*C*-glucosides by this chemistry strongly suggests that hydrogen bonding between the incoming acceptor and O2 of the donor, or indeed any other donor oxygen, or the departing triflate anion, does not exert a strong influence on the formation of the equivalent *O*-glycosides. The benzylidene acetal-controlled one step stereoselective formation of  $\beta$ -*C*-mannosides is an attractive alternative to existing methods.<sup>19,20</sup>

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


## Acknowledgment

We thank the NIH (GM 62160) for support of this work, and Professor Glenn McGarvey, University of Virginia, for a helpful exchange of information.


## References

1. (a) Crich D, Sun S. *Tetrahedron* 1998;54:8321–8348. (b) Crich D, Smith M. J. *Am. Chem. Soc* 2001;123:9015–9020. [PubMed: 11552809] (c) Crich D, Lim LBL. *Org. React* 2004;64:115–251.
2. For a review on cyclic protecting groups in glycoside synthesis see: Litjens REJN, van den Bos JL, Codée JDC, Overkleeft HS, van der Marel G. *Carbohydr. Res* 2007;342:419–429.429 [PubMed: 17007825]
3. (a) Weingart R, Schmidt RR. *Tetrahedron Lett* 2000;41:8753–8758. (b) Tanaka, S-i; Takashima, M.; Tokimoto, H.; Fujimoto, Y.; Tanaka, K.; Fukase, K. *Synlett* 2005:2325–2328.
4. Kim KS, Kim JH, Lee YJ, Lee YJ, Park J. J. *Am. Chem. Soc* 2001;123:8477–8481. [PubMed: 11525654]
5. Tsuda T, Sato S, Nakamura S, Sato S. *Heterocycles* 2003;59:509–515.
6. Crich D, Sun S. *J. Am. Chem. Soc* 1997;119:11217–11223.
7. Crich D, Chandrasekera NS. *Angew. Chem. Int. Ed* 2004;43:5386–5387.
8. Jensen HH, Nordstrom M, Bols M. J. *Am. Chem. Soc* 2004;123:9205–9213. [PubMed: 15281809]
9. The possibility that the benzylidene effect also contains a torsional component exists: Andrews CW, Rodebaugh R, Fraser-Reid B. *J. Org. Chem* 1996;61:5280–5289.5289
10. (a) Crich D, Cai W. *J. Org. Chem* 1999;64:4926–4930. [PubMed: 11674572] (b) Bousquet E, Khitri M, Lay L, Nicotra F, Panza L, Russo G. *Carbohydr. Res* 1998;311:171–181. [PubMed: 9825520]
11. (a) Crich D, Vinogradova O. *J. Org. Chem* 2006;71:8473–8480. [PubMed: 17064022] (b) Crich D, Li L. *J. Org. Chem* 2007;72:1681–1690. [PubMed: 17266375]
12. As discussed previously, 11a,b this rationale is based on the computed conformations of the 4,6-*O*-benzylidene-protected gluco- and mannopyranosyl oxacarbenium ions as provided by Whitfield and coworkers: Nukada T, Bérces A, Whitfield DM. *Carbohydr. Res* 2002;337:765–774.774 [PubMed: 11950473]
13. Hydrogen bonding between the acceptor OH and O3, but not O2, of the benzylidene protected mannosyl donor has been invoked previously on the basis of computations as one of several factors possibly affecting stereoselectivity in these glycosylations. Likewise, computational work has also raised the possibility of hydrogen bonding between the incoming acceptor and the departing triflate, in an  $S_Ni$ -like manner as a possible rationale for the  $\alpha$ -selective glycosylations: (a) Ref12. (b) Nukada T, Bérces A, Wang L, Zgierski MZ, Whitfield DM. *Carbohydr. Res* 2005;340:841–852.852 [PubMed: 15780250] (c) Ionescu AR, Whitfield DM, Zgierski MZ, Nukada T. *Carbohydr. Res*


- 2006;341:2912–2920.2920 [PubMed: 17069777] (d)Whitfield DM, Nukada T. Carbohydr. Res 2007;342:1291–1304.1304 [PubMed: 17477909]
14. For previous discussions of the potential effects of donor-acceptor hydrogen bonding in glycosylation reactions see: Bércecs A, Whitfield DM, Nukada T, do Santos I, Obuchoskwa A, Krepinsky JJ. *Can. J. Chem* 2004;82:1157–1171.1171
15. (a) Crich D, Sun S. *Tetrahedron* 1998;54:8321–8348. (b) Dubois E, Beau JM. *Carbohydr. Res* 1992;228:103–120. [PubMed: 1516081]
16. Crich D, Smith M, Yao Q, Picione J. *Synthesis* 2001:323–326.
17. Previous syntheses and applications of glycosyl enolethers, or vinyl glycosides: (a)Chenault HK, Castro A, Chafin LF, Yang Y. *J. Org. Chem* 1996;61:5024–5031.5031 (b)Boons G-J, Heskamp B, Hout F. *Angew. Chem. Int. Ed* 1996;35:2845–2847.2847 (c)Boons G-J, Isles S. *J. Org. Chem* 1996;61:4262–4271.4271 [PubMed: 11667325] (d)Yuan J, Lindner K, Frauenrath H. *J. Org. Chem* 2006;71:5457–5467.5467 [PubMed: 16839123] (e)Wang P, Haldar P, Wang Y, Hu H. *J. Org. Chem* 2007;72:5870–5873.5873 [PubMed: 17580902]
18. Lucero CG, Woerpel KA. *J. Org. Chem* 2006;71:2641–2647. [PubMed: 16555815]
19. Levy, DE.; Tang, C. *The Chemistry of C-Glycosides*. Oxford: Pergamon; 1995. Postema, MHD., editor. *C-Glycoside Synthesis*. Boca Raton: CRC; 1995. (c) Beau J-M, Gallagher T. *Top. Curr. Chem* 1997;187:1–54. (d) Nicotra F. *Top. Curr. Chem* 1997;187:55–83.
20. For the analogous C-glycoside formation in the absence of the benzylidene acetal see the accompanying paper: McGarvey GJ, Leclair CA, Schmidtman BA. submitted.



**Figure 1.**  
Thioglycosides Employed




BSP



TTBP

**Figure 2.**  
Coupling Reagents



**scheme 1.**  
General Glycosylation Mechanism

coupled product	isolated yield <sup>a</sup>	$\alpha$ : $\beta$ ratio <sup>b,c</sup>
	58%	1:5



$\alpha$ :  $\beta$  ratio<sup>b,c</sup>isolated yield<sup>d</sup>

coupled product

1:8


61%



$\alpha$ :  $\beta$  ratio<sup>b,c</sup>isolated yield<sup>d</sup>8a  $\beta$  only  
8b,  $\alpha$  only

63% (7:1, 8a:8b)

coupled product



$\alpha$ :  $\beta$  ratio<sup>b,c</sup>isolated yield<sup>d</sup>

coupled product


 $\beta$  only

80%



$\alpha$ :  $\beta$  ratio<sup>b,c</sup>isolated yield<sup>d</sup>51% (2:1, **10a**:**10b**)**10a**,  $\beta$  only<sup>d</sup>  
**10b**,  $\beta$  only<sup>d</sup>

coupled product






$\alpha$ :  $\beta$  ratio<sup>b,c</sup>isolated yield<sup>d</sup>

coupled product

2:1

70%




$\alpha$ : $\beta$ ratio <sup>b,c</sup>	isolated yield <sup>d</sup>	coupled product
$\alpha$ only	54%	 <p style="text-align: center;"><b>12</b></p>
$\alpha$ only	52%	 <p style="text-align: center;"><b>12</b></p>

coupled product

 $\alpha$ :  $\beta$  ratio<sup>b,c</sup>isolated yield<sup>d</sup>

13a, 22% 13b, 33%

13a,  $\alpha$  only  
13b,  $\alpha$  only

$\alpha$ :  $\beta$  ratio<sup>b,c</sup>

isolated yield<sup>d</sup>

coupled product