

Published in final edited form as:

Nat Rev Mol Cell Biol. 2008 September ; 9(9): 673–678. doi:10.1038/nrm2479.

## Endogenous small interfering RNAs in animals

Katsutomo Okamura and Eric C. Lai

Sloan-Kettering Institute, Department of Developmental Biology, 521 Rockefeller Research Laboratories, 1275 York, Avenue, BOX 252, New York, New York 10065, USA.

### Abstract

Until recently, only nematodes among animals had a well-defined endogenous small interfering RNA (endo-siRNA) pathway. This has changed dramatically with the recent discovery of diverse intramolecular and intermolecular substrates that generate endo-siRNAs in *Drosophila melanogaster* and mice. These findings suggest broad and possibly conserved roles for endogenous RNA interference in regulating host-gene expression and transposable element transcripts. They also raise many questions regarding the biogenesis and function of small regulatory RNAs in animals.

---

RNA interference (RNAi), the process by which double-stranded RNA (dsRNA) is processed into small interfering RNAs (siRNAs) that silence homologous transcripts, is both a fascinating cellular machinery and a powerful experimental technique. Despite an avalanche of RNAi research over the past decade, however, a nagging question remained mostly unanswered: what good is RNAi to the organism itself?

Substantial roles for RNAi in regulating endogenous gene expression have been difficult to ascertain because *Drosophila melanogaster*<sup>1,2</sup> and *Caenorhabditis elegans*<sup>3,4</sup> mutants that selectively inactivate RNAi seem to be normal and fertile. These mutants are hypersensitive to viruses, which suggests that RNAi defends against selfish and invasive nucleic acids<sup>5</sup>. But if RNAi had an ancestral role in virus restriction it seems to have been subsumed in vertebrates by the interferon pathway. In fact, the nonspecific capacity of dsRNA to activate the interferon response, thereby leading to the general inhibition of cellular translation, was widely perceived to preclude substantial roles for endogenous RNAi in vertebrates.

Eight concurrent papers from the Zamore, Sasaki, Siomi, Lai and Hannon laboratories recently described a rich diversity of endogenous siRNAs (endo-siRNAs) in mice<sup>6,7</sup> and *D. melanogaster*<sup>8–13</sup>. These studies introduce unanticipated complexity in small-RNA sorting pathways and in the biological roles of siRNAs. We highlight these new classes of endo-siRNAs and the pressing questions that are raised by their discovery.

### Argonaute-bound small RNAs

Argonaute proteins lie at the heart of related small-RNA pathways that operate in organisms as diverse as Archaea, plants and animals<sup>14</sup>. They bind various small RNAs that are < 32 nucleotides (nt) in length which guide the Argonaute complexes to their regulatory targets (FIG. 1).

Among animals, the AGO and Piwi subclasses constitute two main classes of conserved Argonaute proteins. AGO proteins bind to microRNAs (miRNAs)<sup>14,15</sup> — RNAs of ~ 22 nt

that derive from host transcripts with short (usually < 100 nt) inverted repeats. These repeats are processed by the RNase III enzymes Droscha (in the nucleus) and Dicer (in the cytoplasm) (FIG. 1a). Specialized AGO proteins with efficient 'slicing' activity are the carriers of 21 nt siRNAs<sup>2,16,17</sup>. Exogenous dsRNAs are processed into siRNAs in the cytoplasm by Dicer, and therefore they do not require Droscha (FIG. 1 b). Piwi-interacting RNAs (piRNAs) are slightly longer RNAs (~24–32 nt) that are bound by Piwi-family proteins, which also have slicer activity<sup>18–20</sup>. Although their biogenesis is not completely understood, a major pathway for piRNA production involves reciprocal cleavages of sense and antisense substrates by antisense and sense piRNAs, respectively<sup>19,21</sup> (FIG. 2a).

## Primary and secondary nematode siRNAs

Until recently, *C. elegans* was the only animal for which endo-siRNAs had been well characterized. Primary siRNAs that are processed by dicing dsRNA are exceedingly rare in this organism<sup>22–24</sup>. Instead, the 3' ends of targets that have been cleaved by primary siRNAs are recognized by an RNA-dependent RNA polymerase (RdRP), which generates abundant, untemplated, secondary siRNAs with distinctive 5' triphosphates (FIG. 3). Secondary siRNAs are then loaded into specialized secondary Argonautes (SAGOs)<sup>25</sup>. Because other animals do not seem to encode RdRP or SAGOs, it is not evident that the mechanism for worm siRNA biogenesis is broadly conserved. Nematodes also lack conventional piRNAs, as the Piwi homologue PRG-1 contains '21U' RNAs. The biogenesis of these 21 nt RNAs does not seem to be related to that of fly or vertebrate piRNAs<sup>22,26–28</sup> (FIG. 2b). Therefore, fundamental aspects of conserved animal small-RNA pathways have clearly been altered in *C. elegans*.

## Endo-siRNAs in flies and mice

Recent work now reveals diverse sources of endo-siRNAs in *D. melanogaster* and in mouse. Most of these endo-siRNA classes seem to be analogous between species, and include those derived from transposable elements, from complementary annealed transcripts, and from long 'fold-back' transcripts called hairpin RNAs (hpRNAs).

## siRNAs from transposable elements

Because of the mutagenic consequences of transposable elements (TEs), powerful mechanisms are needed to restrict their activity. Such protection is indispensable in the germ line to maintain faithful transmission of the genome. In this context, piRNAs mediate a major defence against TEs<sup>29</sup>. However, scattered reports in the literature indicated that canonical RNAi also influences TEs. This was most clear in *C. elegans*, because many RNAi-defective mutants also deregulate transposons<sup>3,30</sup>. It was proposed that transcriptional read-through across Tc1 transposable elements might produce intramolecular dsRNA between the terminal inverted repeats, the processing of which by RNAi could generate siRNAs that silence Tc1 elements in *trans*<sup>31</sup>.

A conundrum for mammalian piRNA studies was that although multiple mouse *Piwi*-gene mutants exhibit testicular defects, transposon activation and sterility, corresponding mutant ovaries were normal and functional<sup>32–34</sup>. Instead, *Dicer*-mutant ovaries and oocytes exhibit higher levels of certain retrotransposon transcripts<sup>35,36</sup>. This is consistent with either an miRNA-based system for TE control or perhaps the usage of endo-siRNAs. In fact, earlier small-scale sequencing from mouse oocytes and testes revealed that some siRNAs derived from retrotransposons<sup>37</sup>, which could silence long interspersed nuclear elements (LINEs) in *trans*<sup>38</sup>. Newer large-scale cloning provided clearer evidence for TE-siRNAs in mouse oocytes<sup>6,7</sup>. Many of these mapped to the same genomic locations as piRNA clusters, which raised the possibility that these specialized 'master loci' are involved in both piRNA-mediated and siRNA-mediated TE control. However, some transposon classes were apparently targeted

by only one of these RNA classes, which suggested that piRNAs or siRNAs preferentially control certain TEs. For example, several long-terminal repeat (LTR) retrotransposons were nearly exclusively targeted by siRNAs<sup>6,7</sup>.

In *D. melanogaster*, deep sequencing of the small RNAs that directly associate with AGO2 (the Argonaute that mediates RNAi) revealed that TEs are a substantial source of RNAs of precisely 21 nt<sup>11,12</sup>. Similar conclusions were reached by sequencing RNAs that were  $\beta$ -eliminated — this prevents RNAs from being ligated on their 3' ends, unless they bear a 3' modification, and thus enriches AGO2-loaded RNAs<sup>8</sup> — or by analysing total head or cultured-cell RNAs<sup>13</sup>. Their accumulation is dependent on DCR2 (one of the two Dicers in *D. melanogaster*, and the one that generates exogenous siRNAs<sup>14</sup>; FIG. 1c), and the depletion or mutation of either DCR2 or AGO2 elevates TE transcript levels<sup>8,11–13</sup>. The TE-siRNA response is extremely active in various lines of cultured cells and correlates with the strong genomic amplification of specific LTR retrotransposons in these cells. Therefore, both TE-siRNAs and TE-piRNAs repress transposon transcripts in flies and mammals (FIG. 4).

### siRNAs from cis-natural antisense transcripts

*Cis*-natural antisense transcript (*cis*-NAT) arrangements are genomic regions that encode exons on both DNA strands, and can involve 5', 3' or internal exons (FIG. 4). Careful analysis of small-RNA sequences in mouse oocytes<sup>7</sup> and *D. melanogaster* tissues and cultured cells<sup>8,9,11,12</sup> revealed that *cis*-NAT overlaps are favourable for siRNA production. The extent of 21 nt RNA production was limited to annotated exons that are transcribed bidirectionally, excluding adjacent introns. *D. melanogaster cis*-NAT-siRNAs are dependent on DCR2, and mouse *cis*-NAT-siRNAs are similarly Dicer-dependent. However, although virtually all *cis*-NAT-siRNAs in flies derived from 3' untranslated region (UTR) overlaps, one of the abundant mouse *cis*-NAT-siRNA loci involved *Pdzd 11/Kif4*, whose transcripts overlap on their 5' UTRs (FIG. 4).

The levels of the 3'-overlapping transcripts *Pdzd 11* and *Kif4* increased modestly in mouse *Dicer* mutants<sup>7</sup>, consistent with an autoregulatory activity of the siRNAs generated by this *cis*-NAT. *D. melanogaster cis*-NAT-siRNAs specifically load AGO2, but evidence for changes in their progenitor transcripts on loss of DCR2 or AGO2 was equivocal. However, *D. melanogaster cis*-NAT-siRNA genes (but not *cis*-NAT genes in general) exhibited striking enrichment for several nucleic-acid-based functions, including transcription cofactors, deoxyribo-nucleases and ribonucleases<sup>9</sup>. In addition, most co-expressed *cis*-NATs in *D. melanogaster* S2 cells did not generate siRNAs. These data indicate that only a subset of co-expressed *cis*-NAT pairs are selected for siRNA production, presumably reflecting an endogenous functional use. Intriguingly, one of the most highly expressed *cis*-NAT-siRNA loci in the entire genome involves the *CG7739/AgO2* gene pair<sup>8,9,12</sup> — thus AGO2 carries its own siRNAs.

A special class of *cis*-NAT-siRNAs come from the *D. melanogaster klarsicht* (*klar*) gene, which is involved in lipid-droplet transport and nuclear migration, and from the *thickveins* (*tkv*) gene, which is involved in transforming growth factor- $\beta$  signalling<sup>9,12</sup>. Although these loci produce 3' modified, 21 nt, AGO2-bound RNAs from both DNA strands, they seem to involve a specialized mechanism for extremely efficient *cis*-NAT-siRNA production over extended genomic intervals that are 5–10 kb in length<sup>9,12</sup>. In addition, *klar* and *tkv* are not 3' *cis*-NATs, but instead involve overlaps with 5' exons, internal transcript exons and/or annotated intronic regions. Therefore, the strategy for *klar* and *tkv* siRNA production seems to differ from that of conventional *cis*-NAT-siRNAs.

### siRNAs from mammalian pseudogene-gene pairs

Mammalian genomes encode large numbers of pseudogenes, which are presumed to be non-functional entities that will eventually be lost. Small-RNA cloning from mouse oocytes revealed an unexpected class of 'functional' pseudogenes. Multiple genes with antisense-transcribed pseudogenes were inferred to anneal with their complementary progenitors (as *trans*-NATs) and be diced into siRNAs<sup>6,7</sup>. The existence of siRNAs that bridge exon-exon junctions suggested that mature mRNAs constitute the dsRNA substrate, as suggested for *cis*-NAT-siRNA pairs. Microarray profiling and quantitative PCR analysis of *Dicer*-mutant oocytes revealed substantial upregulation of multiple genes with complementary siRNAs (FIG. 4), indicating that this system regulates endogenous gene expression<sup>6,7</sup>. It is unclear whether the dicing of targets during *trans*-NAT-siRNA biogenesis accounts for target regulation, or whether pseudogene-derived antisense siRNAs actively slice sense-strand mRNAs (FIG. 1c). In at least one case — histone deacetylase-1 (*Hdac1*) — siRNAs derived exclusively from sense-antisense pseudogene duplexes, which were inferred to repress functional *Hdac1* transcripts<sup>6</sup>.

Earlier functional tests showed that long dsRNA does not activate protein kinase R or the interferon response in oocytes, as it does in most other mammalian cells<sup>39,40</sup>. Therefore, oocytes might provide a favourable setting for the exploitation of endogenous RNAi to regulate host transcripts. Genes with complementary pseudogene siRNAs are heavily enriched for microtubule-related functions<sup>6</sup>. This suggests a regulatory focus to the *trans*-NAT-siRNA pathway.

### siRNAs from hpRNA transcripts

Although animal miRNA hairpins are usually < 100 nt, plant miRNA hairpins can be significantly longer<sup>41</sup>. Because of this property, the hairpin precursors of some plant miRNAs were not initially recognized. Likewise, some 'long' miRNA hairpins that are double the length of typical miRNAs were only recently identified in *D. melanogaster*<sup>42</sup>. Therefore, animal RNAs that map to inverted repeats might have escaped conventional miRNA annotation.

Bioinformatics studies in *D. melanogaster* revealed a number of candidate loci that produce small RNAs from extended inverted repeats that are termed hairpin RNAs (hpRNAs), the stems of which were up to 400 base pairs in length<sup>10</sup>. At least seven distinct loci generate siRNAs, and the hp-CG4068 locus alone encodes 20 tandem hairpins<sup>10–12</sup>. Despite their structural similarity to miRNAs, hpRNAs are processed by DCR2 instead of DCR1, and generate 3' blocked siRNAs that load AGO2 (REFS<sup>10–12</sup>) (FIG. 1). As with siRNAs from artificial long-inverted repeats, the siRNA duplexes derived from hpRNAs are phased and direct AGO2 to cleave targets.

One of the hp-CG4068 siRNAs is highly complementary to the coding region of mutagen-sensitive-308 (*mus308*), a DNA polymerase that is involved in the DNA-damage response, and can cleave this target site<sup>10,12</sup>. In this case, *mus308* is the only obvious target of the many siRNAs that are generated by hp-CG4068. However, hp-CG18854 is a pseudogene with substantial homology to *CG8289*, which encodes a chromodomain protein, and elevated hp-CG 18854 could repress *CG8289* in *trans*<sup>10</sup>. Curiously, several candidate hpRNA loci were identified in mouse, including a long-inverted repeat pseudogene of the Ran GTPase-activating protein-1 (*Rangap1*) gene<sup>6,7</sup>. It is unclear whether these hpRNA pathways are conserved or convergent, but they at least suggest that analogous systems operate in *D. melanogaster* and mammals. However, it is clear that entry into an endo-siRNA pathway can endow pseudogenes in both species with regulatory activity.

## Fly endo-siRNAs require Loquacious

There are two Dicers in *D. melanogaster* — DCR1 cleaves pre-miRNA hairpins into miRNA duplexes, whereas DCR2 cleaves long dsRNA into siRNA duplexes<sup>14</sup> (FIG. 1). Each Dicer directly binds to a dsRNA-binding domain (dsRBD) partner that aids its function. DCR2 interacts with R2D2 (whose name derives from the fact that it contains two dsRNA-binding domains (R2) and is associated with DCR2 (D2)), which is essential for the loading of siRNA into AGO2 (REFS<sup>43,44</sup>). DCR1 interacts with Loquacious (LOQS), which promotes its ability to cleave pre-miRNA hairpins, the products of which are preferentially loaded into AGO1 (REFS<sup>45–47</sup>).

Although the attractive symmetry of RNase III, dsRBD and AGO partnerships in the RNAi and miRNA pathways lent support to the proposed division of these pathways, genetic observations suggested that there are much more complex interactions among these factors. For example, unlike *Dcr2* mutants, *r2d2* mutants reveal its requirement for early development and female fertility. Moreover, *r2d2* (but not *Dcr2*) phenotypes are strongly enhanced on reduction of *Dcr1* (REF<sup>48</sup>). Reciprocally, *Dcr1* proved to be an RNAi-defective mutant<sup>1</sup>. There is substantial functional overlap between AGO1 and AGO2, as detected by double-mutant analysis<sup>49</sup>, and some miRNAs sort to both AGO1 and AGO2 (REFS<sup>11,12,50–52</sup>). Finally, *loqs* functions in inverted-repeat RNA-mediated silencing<sup>46</sup>. These findings indicate that there is substantial crosstalk between the RNAi and miRNA pathways.

Despite its original classification as a core component of the miRNA pathway, *loqs*-null mutants have only modest defects in the maturation of many miRNAs<sup>53</sup>. It seems that DCR1 can cleave pre-miRNAs without LOQS, albeit with lowered efficiency that varies between miRNAs<sup>53</sup>. Surprisingly, LOQS is essential for the accumulation of many endo-siRNAs<sup>9,10,12,13</sup> (FIG. 1c). At least some of the members of all of the siRNA classes — TE-siRNAs, *cis*-NAT-siRNAs and hpRNA-siRNAs — are dependent on LOQS. Although previous tests did not reveal a physical interaction between LOQS and DCR2, proteomic analysis of DCR2 complexes revealed that there is comparable coverage of LOQS and R2D2 peptides<sup>12</sup>. Therefore, LOQS is a component of both miRNA and RNAi pathways.

## Endo-siRNA biogenesis: open questions

The recent papers on endo-siRNAs raise fundamental questions regarding the biogenesis of small RNAs. Some of the most important questions concern mechanistic aspects of small-RNA sorting pathways. For example, how does LOQS work with DCR2? And given that R2D2 is needed to load exosRNAs into AGO2 (REFS<sup>43,44</sup>), to what extent do endo-siRNAs require R2D2 for loading? How are miRNA and hpRNA precursors distinguished? Some ‘long’ miRNAs and ‘short’ hpRNAs in *D. melanogaster* are indistinguishable in size and structure<sup>10,42</sup>. They are effectively sorted, however, as long miRNAs make only a single small-RNA duplex (as is typical for DCR1 substrates), whereas short hpRNAs produce multiple duplexes (as is typical for DCR2 substrates). How can the cell distinguish these hairpins?

The regulation of dsRNA formation is another mystery. For example, the *cis*-NAT-siRNA pathway accepts many substrates — at least 17 in mouse oocytes<sup>6,7</sup> and at least 140 in *D. melanogaster*<sup>8,9,12</sup>. However, *cis*-NAT-siRNA loci constitute only 25% of co-expressed *cis*-NATs in *D. melanogaster*<sup>9</sup>. Is there active selection for entry into the RNAi pathway, which could be mediated at the step of dsRNA formation? Conversely, how do co-expressed mammalian *cis*-NATs, and co-expressed pseudogene-gene complementary pairs, avoid triggering an interferon response outside of oocytes? Finally, although it seems evident that *cis*-NAT and *trans*-NAT siRNAs are generated from processed transcripts, it is not known whether the dsRNA substrate forms in the nucleus or cytoplasm, nor is it clear where the dsRNA encounters Dicer.

Valuable lessons were taught by the length and structure of primary hpRNA transcripts. Their dsRNA character was recognized only after genomic fragments of sufficient length were examined, and consequently their siRNAs were prone to being misannotated as having derived from shorter, unstructured precursors<sup>8</sup>. The stems of some plant miRNA hairpins are separated by long, unstructured terminal loops and even introns<sup>54</sup>, and we now recognize the same to be true for several hpRNAs<sup>10,12</sup>. It is therefore conceivable that the stems of some hpRNA precursors might be separated by kilobases or tens of kilobases. Do the structured precursors of any anonymous cloned small RNAs that are currently deposited in public databases await discovery?

Endogenous sources of mammalian dsRNA remain to be recognized outside of oocytes. As is the case with oocytes, introduction of long dsRNA into embryonic stem cells (ESCs) does not activate an interferon response<sup>55,56</sup>. Might ESCs also harbour endo-siRNAs, the action of which is relevant for maintaining pluripotency? Although endo-siRNAs were not previously found in ESCs<sup>57</sup> this possibility might deserve further study.

Finally, although small-RNA sorting pathways have received little attention in mammalian systems, there is growing recognition of their importance to siRNA and miRNA function in plants<sup>58,59</sup>, worms<sup>60,61</sup> and flies<sup>51,62</sup>. As only one of the four mammalian AGO proteins (AGO2) has slicer activity<sup>16,17</sup>, the directed sorting of mammalian siRNAs is presumably important for their ability to slice complementary targets<sup>63</sup>. Consequently, the elucidation of mammalian siRNA sorting rules might have important implications for attempts to improve siRNA efficacy for experimental and therapeutic purposes.

## The biology of endo-siRNAs

To return to the question posed at the beginning of this Perspective, what good is endogenous RNAi to an organism? The necessity to preserve RNAi in mammals has been somewhat of an enigma as they seem to have mostly dispensed with siRNAs for antiviral defence, and some aspects of mammalian biology can be rescued by slicer-defective AGO2 (REF. <sup>64</sup>). However, in addition to a few endogenous cleavage targets of miRNAs<sup>65</sup>, and a role for AGO2 in the biogenesis of select miRNAs<sup>66</sup>, the new studies suggest widespread usage of endo-siRNAs as endogenous regulators of gene expression.

However, it is safe to say that we do not understand the specific biological functions of endo-siRNAs well. Indeed, the question of endo-siRNA function remains mostly unanswered in worms<sup>22,67,68</sup>, and the discovery of abundant endo-siRNAs in flies and mammals only makes the understanding of this topic more pressing. The recent papers do show deregulation of retrotransposon transcripts, pseudogene-complementary transcripts and some *cis*-NAT pairs in *Dicer* and/or *Ago* mutants, and thus their regulation by endo-siRNAs is plausible, although this remains to be shown directly. Evidence for direct siRNA-mediated target regulation was only explicitly shown for some hpRNAs in *D. melanogaster*<sup>10,12</sup>, and such evidence would be desirable for other classes of endo-siRNAs.

The established targets of *D. melanogaster* hpRNAs encode DNA-binding proteins<sup>10,12</sup>. This seems reminiscent of the fact that *D. melanogaster cis*-NAT-siRNA loci are significantly enriched for DNA and RNA-binding proteins<sup>9</sup>, raising this as a substantial molecular axis for endo-siRNA regulation. It is relevant to note, therefore, that *D. melanogaster Dcr2* mutants exhibit abnormal nucleolar morphology<sup>69</sup>, whereas *Ago2* mutants were reported to have chromosome segregation defects<sup>70</sup>. These phenotypes are plausibly connected to the types of gene functions that are highly enriched in *cis*-NAT-siRNAs. The mouse oocyte pseudogene-siRNA system seems to preferentially target genes that are involved in microtubule dynamics<sup>6</sup>, and this is plausibly connected to the observation that *Dicer* loss in growing oocytes disrupts spindle formation and chromosome segregation<sup>35,71</sup>. Nevertheless, the endogenous

requirement of these systems remains to be demonstrated by specific knockouts of hpRNAs or siRNA-generating pseudogenes.

Overall, the fact that core RNAi pathway mutants in worms and flies are mostly normal and fertile, whereas core miRNA pathway mutants are lethal, suggests that the role of endogenous RNAi is fundamentally different than that of miRNA regulation. This is further suggested by the fact that many miRNAs are deeply conserved but most *D. melanogaster* hpRNA loci<sup>10, 12</sup> and most mouse pseudogenes that generate siRNAs<sup>6,7</sup> are poorly conserved. We must therefore think more openly about their usage. Is the usage of these RNAs a matter of fine-tuning gene expression, or perhaps a matter of maintaining fitness in an ever-changing environment? Is endogenous RNAi used for robustness in gene regulation, perhaps to canalize traits? Or is it a regulatory mechanism that generates species-specific characters during evolution? These are questions that remain for the future, but given the pace with which the field of endo-siRNAs has recently advanced, we might expect some answers to soon be forthcoming.

#### DATABASES

**Entrez Gene:** <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=gene>

Hdacl | klar | mus308 | Rangap1 | tkv

**UniProtKB:** <http://www.uniprot.org>

AGO1 | AGQ2 | DCR1 | DCR2 | LOQS | PRG-1 | R2D2

#### FURTHER INFORMATION

**Eric C. Lai's homepage:**

<http://www.mskcc.org/mskcc/html/52949.cfm>

**ALL LINKS ARE ACTIVE IN THE ONLINE PDF**

## References

1. Lee YS, et al. Distinct roles for *Drosophila* Dicer-1 and Dicer-2 in the siRNA/miRNA silencing pathways. *Cell* 2004;117:69–81. [PubMed: 15066283]
2. Okamura K, Ishizuka A, Siomi H, Siomi MC. Distinct roles for Argonaute proteins in small RNA-directed RNA cleavage pathways. *Genes Dev* 2004;18:1655–1666. [PubMed: 15231716]
3. Tabara H, et al. The *rde-1* gene, RNA interference, and transposon silencing in *C. elegans*. *Cell* 1999;99:123–132. [PubMed: 10535731]
4. Tabara H, Yigit E, Siomi H, Mello CC. The dsRNA binding protein RDE-4 interacts with RDE-1, DCR-1, and a DExH-box helicase to direct RNAi in *C. elegans*. *Cell* 2002;109:861–871. [PubMed: 12110183]
5. Ding SW, Voinnet O. Antiviral immunity directed by small RNAs. *Cell* 2007;130:413–426. [PubMed: 17693253]
6. Tam OH, et al. Pseudogene-derived small interfering RNAs regulate gene expression in mouse oocytes. *Nature* 2008;453:534–538. [PubMed: 18404147]
7. Watanabe T, et al. Endogenous siRNAs from naturally formed dsRNAs regulate transcripts in mouse oocytes. *Nature* 2008;453:539–543. [PubMed: 18404146]
8. Ghildiyal M, et al. Endogenous siRNAs derived from transposons and mRNAs in *Drosophila* somatic cells. *Science* 2008;320:1077–1081. [PubMed: 18403677]
9. Okamura K, Balla S, Martin R, Liu N, Lai EC. Two distinct mechanisms generate endogenous siRNAs from bidirectional transcription in *Drosophila*. *Nature Struct. Mol. Biol* 2008;15:581–590. [PubMed: 18500351]

10. Okamura K, et al. The *Drosophila* hairpin RNA pathway generates endogenous short interfering RNAs. *Nature* 2008;453:803–806. [PubMed: 18463630]
11. Kawamura Y, et al. *Drosophila* endogenous small RNAs bind to Argonaute2 in somatic cells. *Nature* 2008;453:793–797. [PubMed: 18463636]
12. Czech B, et al. An endogenous siRNA pathway in *Drosophila*. *Nature* 2008;453:798–802. [PubMed: 18463631]
13. Chung WJ, Okamura K, Martin R, Lai EC. Endogenous RNA interference provides a somatic defense against *Drosophila* transposons. *Current Biology* 2008;18:795–802. [PubMed: 18501606]
14. Farazi TA, Juranek SA, Tuschl T. The growing catalog of small RNAs and their association with distinct Argonaute/Piwi family members. *Development* 2008;135:1201–1214. [PubMed: 18287206]
15. Mourelatos Z, et al. miRNPs: a novel class of ribonucleoproteins containing numerous microRNAs. *Genes Dev* 2002;16:720–728. [PubMed: 11914277]
16. Liu J, et al. Argonaute2 is the catalytic engine of mammalian RNAi. *Science* 2004;305:1437–1441. [PubMed: 15284456]
17. Meister G, et al. Human Argonaute2 mediates RNA cleavage targeted by mi RNAs and siRNAs. *Mol. Cell* 2004;15:185–197. [PubMed: 15260970]
18. Saito K, et al. Specific association of Piwi with rasiRNAs derived from retrotransposon and heterochromatic regions in the *Drosophila* genome. *Genes Dev* 2006;20:2214–2222. [PubMed: 16882972]
19. Gunawardane LS, et al. A slicer-mediated mechanism for repeat-associated siRNA 5' end formation in *Drosophila*. *Science* 2007;315:1587–1590. [PubMed: 17322028]
20. Lau NC, et al. Characterization of the piRNA complex from rat testes. *Science* 2006;313:363–367. [PubMed: 16778019]
21. Brennecke J, et al. Discrete small RNA-generating loci as master regulators of transposon activity in *Drosophila*. *Cell* 2007;128:1089–1103. [PubMed: 17346786]
22. Ruby JG, et al. Large-scale sequencing reveals 21 U-RNAs and additional microRNAs and endogenous siRNAs in *C. elegans*. *Cell* 2006;127:1193–207. [PubMed: 17174894]
23. Pak J, Fire A. Distinct populations of primary and secondary effectors during RNAi in *C. elegans*. *Science* 2007;315:241–244. [PubMed: 17124291]
24. Sijen T, Steiner FA, Thijssen KL, Plasterk RH. Secondary siRNAs result from unprimed RNA synthesis and form a distinct class. *Science* 2007;315:244–247. [PubMed: 17158288]
25. Yigit E, et al. Analysis of the *C. elegans* Argonaute family reveals that distinct Argonautes act sequentially during RNAi. *Cell* 2006;127:747–757. [PubMed: 17110334]
26. Wang G, Reinke VA. *C. elegans* Piwi, PRG-1, regulates 21 U-RNAs during spermatogenesis. *Curr. Biol* 2008;18:861–867. [PubMed: 18501605]
27. Batista PJ, et al. PRG-1 and 21 U-RNAs interact to form the piRNA complex required for fertility in *C. elegans*. *Mol. Cell* 2008;31:67–78. [PubMed: 18571452]
28. Das PP, et al. Piwi and piRNAs act upstream of an endogenous siRNA pathway to suppress Tc3 transposon mobility in the *Caenorhabditis elegans* germline. *Mol. Cell* 2008;31:79–90. [PubMed: 18571451]
29. Aravin AA, Hannon GJ, Brennecke J. The Piwi-piRNA pathway provides an adaptive defense in the transposon arms race. *Science* 2007;318:761–764. [PubMed: 17975059]
30. Ketting RF, Haverkamp TH, van Luenen HG, Plasterk RH. Mut-7 of *C. elegans*, required for transposon silencing and RNA interference, is a homolog of Werner syndrome helicase and RNaseD. *Cell* 1999;99:133–141. [PubMed: 10535732]
31. Sijen T, Plasterk RH. Transposon silencing in the *Caenorhabditis elegans* germ line by natural RNAi. *Nature* 2003;426:310–314. [PubMed: 14628056]
32. Kuramochi-Miyagawa S, et al. Mili, a mammalian member of *piwi* family gene, is essential for spermatogenesis. *Development* 2004;131:839–849. [PubMed: 14736746]
33. Carmell MA, et al. MIWI2 is essential for spermatogenesis and repression of transposons in the mouse male germline. *Dev. Cell* 2007;12:503–514. [PubMed: 17395546]
34. Deng W, Lin H. *miwi*, a murine homolog of *piwi*, encodes a cytoplasmic protein essential for spermatogenesis. *Dev. Cell* 2002;2:819–830. [PubMed: 12062093]




35. Murchison EP, et al. Critical roles for Dicer in the female germline. *Genes Dev* 2007;21:682–693. [PubMed: 17369401]
36. Svoboda P, et al. RNAi and expression of retrotransposons MuERV-L and IAP in preimplantation mouse embryos. *Dev. Biol* 2004;269:276–285. [PubMed: 15081373]
37. Watanabe T, et al. Identification and characterization of two novel classes of small RNAs in the mouse germline: retrotransposon-derived siRNAs in oocytes and germline small RNAs in testes. *Genes Dev* 2006;20:1732–1743. [PubMed: 16766679]
38. Yang N, Kazazian HH Jr. L1 retrotransposition is suppressed by endogenously encoded small interfering RNAs in human cultured cells. *Nature Struct. Mol. Biol* 2006;13:763–771. [PubMed: 16936727]
39. Svoboda P, Stein P, Hayashi H, Schultz RM. Selective reduction of dormant maternal mRNAs in mouse oocytes by RNA interference. *Development* 2000;127:4147–4156. [PubMed: 10976047]
40. Stein P, Zeng F, Pan H, Schultz RM. Absence of non-specific effects of RNA interference triggered by long double-stranded RNA in mouse oocytes. *Dev. Biol* 2005;286:464–471. [PubMed: 16154556]
41. Reinhart BJ, Weinstein EG, Rhoades MW, Bartel B, Bartel DP. MicroRNAs in plants. *Genes Dev* 2002;16:1616–1626. [PubMed: 12101121]
42. Ruby JG, et al. Evolution, biogenesis, expression, and target predictions of a substantially expanded set of *Drosophila* microRNAs. *Genome Res* 2007;17:1850–1864. [PubMed: 17989254]
43. Liu Q, et al. R2D2, a bridge between the initiation and effector steps of the *Drosophila* RNAi pathway. *Science* 2003;301:1921–1925. [PubMed: 14512631]
44. Tomari Y, Matranga C, Haley B, Martinez N, Zamore PD. A protein sensor for siRNA asymmetry. *Science* 2004;306:1377–1380. [PubMed: 15550672]
45. Jiang F, et al. Dicer-1 and R3D1 -L catalyze microRNA maturation in *Drosophila*. *Genes Dev* 2005;19:1674–1679. [PubMed: 15985611]
46. Forstemann K, et al. Normal microRNA maturation and germ-line stem cell maintenance requires Loquacious, a double-stranded RNA-binding domain protein. *PLoS Biol* 2005;3:e236. [PubMed: 15918770]
47. Saito K, Ishizuka A, Siomi H, Siomi MC. Processing of pre-microRNAs by the Dicer-1 -Loquacious complex in *Drosophila* cells. *PLoS Biol* 2005;3:e235. [PubMed: 15918769]
48. Kalidas S, et al. *Drosophila* R2D2 mediates follicle formation in somatic tissues through interactions with Dicer-1. *Mech. Dev* 2008;125:475–485. [PubMed: 18299191]
49. Meyer WJ, et al. Overlapping functions of argonaute proteins in patterning and morphogenesis of *Drosophila* embryos. *PLoS Genet* 2006;2:e134.
50. Seitz H, Ghildiyal M, Zamore PD. Argonaute loading improves the 5' precision of both microRNAs and their miRNA strands in flies. *Curr. Biol* 2008;18:147–151. [PubMed: 18207740]
51. Forstemann K, Horwich MD, Wee L, Tomari Y, Zamore PD. *Drosophila* microRNAs are sorted into functionally distinct argonaute complexes after production by dicer-1. *Cell* 2007;130:287–297. [PubMed: 17662943]
52. Horwich MD, et al. The *Drosophila* RNA methyltransferase, DmHen 1, modifies germline piRNAs and single-stranded siRNAs in RISC. *Curr. Biol* 2007;17:1265–1272. [PubMed: 17604629]
53. Liu X, et al. Dicer-1, but not Loquacious, is critical for assembly of miRNA-induced silencing complexes. *RNA* 2007;13:2324–2329. [PubMed: 17928574]
54. Sunkar R, Girke T, Jain PK, Zhu JK. Cloning and characterization of microRNAs from rice. *Plant Cell* 2005;17:1397–1411. [PubMed: 15805478]
55. Paddison PJ, Caudy AA, Hannon GJ. Stable suppression of gene expression by RNAi in mammalian cells. *Proc. Natl Acad. Sci. USA* 2002;99:1443–1448. [PubMed: 11818553]
56. Yang S, Tutton S, Pierce E, Yoon K. Specific double-stranded RNA interference in undifferentiated mouse embryonic stem cells. *Mol. Cell. Biol* 2001;21:7807–7816. [PubMed: 11604515]
57. Calabrese JM, Seila AC, Yeo GW, Sharp PA. RNA sequence analysis defines Dicer's role in mouse embryonic stem cells. *Proc. Natl Acad. Sci. USA* 2007;104:18097–18102. [PubMed: 17989215]
58. Mi S, et al. Sorting of small RNAs into *Arabidopsis* argonaute complexes is directed by the 5' terminal nucleotide. *Cell* 2008;133:116–127. [PubMed: 18342361]

59. Montgomery TA, et al. Specificity of ARGONAUTE7-miR390 interaction and dual functionality in TAS3 *trans*-acting siRNA formation. *Cell* 2008;133:128–141. [PubMed: 18342362]
60. Jannot G, Boisvert ME, Banville IH, Simard MJ. Two molecular features contribute to the Argonaute specificity for the microRNA and RNAi pathways in *C. elegans*. *RNA* 2008;14:829–835. [PubMed: 18367718]
61. Steiner FA, et al. Structural features of small RNA precursors determine Argonaute loading in *Caenorhabditis elegans*. *Nature Struct. Mol. Biol* 2007;14:927–933. [PubMed: 17891148]
62. Tomari Y, Du T, Zamore PD. Sorting of *Drosophila* small silencing RNAs. *Cell* 2007;130:299–308. [PubMed: 17662944]
63. Azuma-Mukai A, et al. Characterization of endogenous human Argonautes and their miRNA partners in RNA silencing. *Proc. Natl Acad. Sci. USA* 2008;105:7964–7969. [PubMed: 18524951]
64. O’Carroll D, et al. A slicer-independent role for Argonaute 2 in hematopoiesis and the microRNA pathway. *Genes Dev* 2007;21:1999–2004. [PubMed: 17626790]
65. Yekta S, Shih IH, Bartel DP. MicroRNA-directed cleavage of HOXB8 mRNA. *Science* 2004;304:594–596. [PubMed: 15105502]
66. Diederichs S, Haber DA. Dual role for argonautes in microRNA processing and posttranscriptional regulation of microRNA expression. *Cell* 2007;131:1097–1108. [PubMed: 18083100]
67. Spike CA, Bader J, Reinke V, Strome S. DEPS-1 promotes P-granule assembly and RNA interference in *C. elegans* germ cells. *Development* 2008;135:983–993. [PubMed: 18234720]
68. Grishok A, Sharp PA. Negative regulation of nuclear divisions in *Caenorhabditis elegans* by retinoblastoma and RNA interference-related genes. *Proc. Natl Acad. Sci. USA* 2005;102:17360–17365. [PubMed: 16287966]
69. Peng JC, Karpen GH. H3K9 methylation and RNA interference regulate nucleolar organization and repeated DNA stability. *Nature Cell Biol* 2007;9:25–35. [PubMed: 17159999]
70. Deshpande G, Calhoun G, Schedl P. *Drosophila* argonaute-2 is required early in embryogenesis for the assembly of centric/centromeric heterochromatin, nuclear division, nuclear migration, and germ-cell formation. *Genes Dev* 2005;19:1680–1685. [PubMed: 16024657]
71. Tang F, et al. Maternal microRNAs are essential for mouse zygotic development. *Genes Dev* 2007;21:644–648. [PubMed: 17369397]


## Acknowledgements

K.O. was supported by the Charles Revson Foundation. E.C.L. was supported by the V Foundation for Cancer Research, the Sidney Kimmel Foundation for Cancer Research, the Alfred Bressler Scholars Fund and the National Institutes of Health (GM083300).



**Figure 1. Small RNA pathways in *Drosophila melanogaster***


In *Drosophila melanogaster*, micro (mi)RNAs are ~22 nucleotides (nt) long, have free hydroxy groups at their 3' ends, and associate primarily with the Argonaute protein AGO1. Small interfering (si)RNAs are ~21 nt, are methylated at their 3' ends, and associate primarily with AGO2. Three main protein families are denoted with RNase III enzymes (Drosha, Dicer-1 (DCR1) and DCR2; shown as hexagons), their dsRNA-binding domain (dsRBD) partners (Pasha, Loquacious (LOQS) and R2D2; shown as squares) and Argonaute proteins (AGO1 and AGO2; shown as ovals), **a** | miRNA pathway Endogenous transcripts that contain short inverted repeats are processed into ~21–22 nt RNAs that mostly function to repress endogenous targets by translational repression and deadenylation by AGO1. miRNA\* is the species on the other side of the hairpin to the miRNA. **b** | In *D. melanogaster*, viral dsRNA or artificial dsRNA produce exogenous siRNAs (exo-siRNAs) that are mostly sorted to AGO2 and restrict viral replication or cleave designed targets, **c** | *D. melanogaster* cells and mammalian oocytes produce several sources of endogenous dsRNA — transposable elements (TEs), *cis*-natural antisense transcripts (*cis*-NATs), *trans*-NATs and hairpin RNA transcripts—that are processed into endo-siRNAs that load mostly AGO2. These repress transposon transcripts or endogenous mRNAs. Note that a minority of miRNAs programme AGO2 and a small fraction of exo- and endo-siRNAs associate with AGO1, but the functional significance of this is currently unknown. 2Ome, 2'-*O*-methyl group.



**Figure 2. Specialized small-RNA regulatory pathways in the animal germ line**












These are mediated by Piwi-class Argonaute proteins (ovals), **a** | The Piwi-interacting (pi)RNA pathway operates in the *Drosophila melanogaster* and vertebrate germ line. A ‘ping-pong’ strategy amplifies piRNAs from complementary transcripts, in which the slicer activity of Piwi proteins (Piwi, Aubergine (AUB) and AGO3 in *D. melanogaster*) reciprocally define piRNA 5' ends. The mechanism that defines the 3' ends of piRNAs is not known. A conserved role of the piRNA pathway is to restrict transposon activity in the germ line; however, there might be other roles for abundant non-transposon-derived piRNAs that are found in mammals, **b** | Nematode 21U RNAs might be a functional analogue of piRNAs. These 21-nucleotide RNAs begin with U and are produced from genomic loci with a characteristic upstream motif

(CTGTTTCA), and they are bound by the Piwi protein PRG-1. The details of 21U biogenesis and function are unclear, but 21Us are linked to spermatogenesis and control of Tc3 transposition. 2Ome, 2'-*O*-methyl group.



**Figure 3. Nematode small interfering RNA pathways**

Processing of double-stranded (ds) substrates by a complex that includes an RNase III enzyme (Dicer-1 (DCR-1), a dsRNA-binding partner (dsRBD; RDE-4) and Dicer-related helicases (DRH-1/2) generates primary (1°) small interfering (si)RNAs with 5' monophosphates (P). These load into the RDE-1 Argonaute protein and can slice complementary transcripts. Cleaved transcripts, and possibly uncleaved transcripts, are substrates for an RNA-dependent RNA polymerase (RdRP) complex that generates secondary (2°) siRNAs that have 5' triphosphates (PPP). These are loaded into various secondary Argonautes (SAGOs) that lack slicer activity, or into the Argonaute slicer CSR-1.

siRNA	Gene structure	dsRNA structure	Loci collected	
			Fly	Mouse
TE-siRNA			Many	Many
<i>cis</i> -NAT-siRNA			140+	17 (+28*)
			Unknown	
		Unknown	2	
<i>trans</i> -NAT-siRNA			Unknown	15**
hpRNA			7 (+19?)	4

#### Figure 4. Substrates for endo-siRNA production in flies and mouse

The precise structure of the double-stranded (ds)RNA substrates of small interfering (si)RNAs derived from transposable elements (TEs) is unknown, but hundreds or thousands of TEs are inferred to directly generate siRNAs. siRNAs derived from *cis*-natural antisense transcripts (*cis*-NATs) involve bidirectional transcription across the same genomic DNA, and can be convergent, divergent or involve annotated introns and/or internal exons. *Drosophila melanogaster cis*-NAT-siRNAs derive almost exclusively from 3'-overlapping mRNAs, but two highly-expressed siRNA loci include annotated introns. Watanabe and colleagues describe 17 *cis* NAT siRNA loci<sup>7</sup>, but their precise categorization is ambiguous as many of them lack an annotated overlapping transcript. Tarn and colleagues describe another 28 mRNAs (\*) with siRNAs whose complementary transcript was not specifically described<sup>6</sup>. These might be *cis*-NATs, but some might represent *trans* natural antisense transcript (*trans* NAT) pairs. *Trans* NAT dsRNAs form between transcripts that are produced from distinct genomic locations, and usually comprise an mRNA and an antisense-transcribed pseudogene. Based on the cumulative data of Tarn *et al.*<sup>6</sup> and Watanabe *et al.*<sup>7,15</sup> *trans*-NAT gene-pseudogene pairs generate siRNAs (\*\*; some of the 28 mRNAs listed in the *cis*-NAT-siRNA category might have antisense pseudogenes that were not reported). siRNAs that are derived from hairpin RNA (hpRNAs) are long, inverted repeat transcripts whose double-stranded segment is typically much longer than that of miRNA precursors. In *D. melanogaster*, one of the 7 identified hpRNA loci encodes 20 hairpin direct repeats, which can function autonomously or as components of higher-order hairpins. The figure shows the numbers of loci that were collected from the recently published studies, but these numbers will probably increase with future studies.