

# The extremely slow and variable activity of dihydrofolate reductase in human liver and its implications for high folic acid intake

Steven W. Bailey<sup>1</sup> and June E. Ayling<sup>1,2</sup>

Department of Pharmacology, University of South Alabama, 307 North University Boulevard, Mobile, AL 36688

Edited by Bruce N. Ames, University of California, Berkeley, CA, and approved July 22, 2009 (received for review February 25, 2009)

Numerous clinical trials using folic acid for prevention of cardiovascular disease, stroke, cognitive decline, and neural tube defects have been completed or are underway. Yet, all functions of folate are performed by tetrahydrofolate and its one-carbon derivatives. Folic acid is a synthetic oxidized form not significantly found in fresh natural foods; to be used it must be converted to tetrahydrofolate by dihydrofolate reductase (DHFR). Increasing evidence suggests that this process may be slow in humans. Here we show, using a sensitive assay we developed, that the reduction of folic acid by DHFR per gram of human liver ( $n = 6$ ) obtained from organ donors or directly from surgery is, on average, less than 2% of that in rat liver at physiological pH. Moreover, in contrast to rats, there was almost a 5-fold variation of DHFR activity among the human samples. This limited ability to activate the synthetic vitamin raises issues about clinical trials using high levels of folic acid. The extremely low rate of conversion of folic acid suggests that the benefit of its use in high doses will be limited by saturation of DHFR, especially in individuals possessing lower than average activity. These results are also consistent with the reports of unmetabolized folic acid in plasma and urine.

folate | provitamin utilization | unmetabolized folic acid | vitamin supplements | nutrition

The physiological function of dihydrofolate reductase (DHFR) is the reduction of 7,8-dihydrofolate (7,8-DHF) (1) produced in the reaction of thymidylate synthase and possibly by autooxidation of tetrahydrofolate. It also may aid in the regeneration of tetrahydrobiopterin in endothelial cells needed for coupled nitric oxide synthase activity (2). However, since the synthesis of folic acid (FA, pteroylglutamic acid) as a provitamin in 1945, DHFR has taken on another role: reduction of FA to 7,8-DHF (Fig. 1). Folic acid, in itself, has no known cofactor activity and is not found to a significant extent in nature. By the time one-carbon metabolism was recognized as actually being carried out by derivatives of tetrahydrofolic acid (THF), folic acid already had been found to alleviate the anemia of folate deficiency and was in wide use in vitamin supplements. Folic acid has only a single chiral center in its glutamate residue and is therefore easier to synthesize than diastereomeric 6S-THF. Thus, there appeared to be little immediate reason to look beyond folic acid.

The U.S. Reference Daily Intake (Daily Value) for FA is 0.4 mg for adults to aid in the prevention of birth defects and anemia, or double this amount for pregnant or lactating women. Consumers of fortified ready-to-eat cereals can have an intake, due to nutrient overages, of up to 0.8 mg of FA per standard serving size (U.S. Department of Agriculture National Nutrient Database for Standard Reference, Release 18, [www.ars.usda.gov](http://www.ars.usda.gov)). Higher doses, up to 5 mg/d, have been used in clinical trials for secondary prevention of heart disease and stroke. All of these potential exposures are on top of that from the Food and Drug Administration mandated grain fortification program in the U.S. which has been estimated to provide up to 0.2 mg/d of FA to the average person (3, 4). One factor that has not been fully considered is whether DHFR has sufficient capacity in humans

to efficiently metabolize these potentially high loads of FA, particularly in light of a frequent misconception that FA is reduced as it crosses the intestinal wall [see review (5)].

Concern has been expressed that the presence of unmetabolized FA (pteroylglutamic acid in plasma or tissues not yet converted to active forms of folate) may be detrimental (6–9). Although folate deficiency has been linked to cancer susceptibility via increased uracil misincorporation (10), recent evidence suggests that high intake ( $\geq 1$  mg/d) of FA may exacerbate some preexisting cancers or progression of precancerous lesions, such as in the colon (11) or prostate (12). The activity of natural killer cells has been reported to be decreased in older women having higher plasma levels of unmetabolized FA (13). Unmetabolized FA has also been hypothesized to cause masking of B12 deficiency and/or aggravate the neurological damage associated with prolonged B12 deficiency (14–16). Therefore, it is important to develop a better understanding of the mechanism of accumulation of unmetabolized FA in humans.

The activity of DHFR in human liver and other tissues has not been well characterized. The few previous measurements mostly used autopsy tissue, which can lead to artifacts due to autolysis. Moreover, earlier DHFR activities were determined at a non-physiological pH, and/or by consumption of NADPH cofactor, which necessitates a large correction for nonspecific oxidases (17–19). Valid comparison of rates among species using values measured in different laboratories is greatly hindered by the use of variable reaction conditions (e.g., salt concentration, impure substrate preparations, and temperature, in addition to pH).

In this study, we determined the activity of DHFR at physiological pH in *fresh* samples from human livers with FA as substrate and compared this to the rate with 7,8-DHF. To accurately measure the very low activity with FA as substrate at physiological pH, a new procedure has been developed for the determination of DHFR based on analysis of the THF product by HPLC. This work reveals that the activity of DHFR in fresh human liver is extremely low in comparison to that in rat liver and suggests that it becomes limiting when FA is consumed at levels higher than the Tolerable Upper Intake Level (1 mg/d for adults).

## Results

**HPLC Assay for Tetrahydrofolate.** A linear response was observed down to the limit of detection of 3 fmol ( $S/N = 3$ ) [supporting information (SI) Fig. S1]. A typical chromatogram of a reaction with  $100,000 \times g$  supernatant of human liver as the source of

Author contributions: S.W.B. and J.E.A. designed research, performed research, contributed new reagents/analytic tools, analyzed data, and wrote the paper.

Conflict of interest: The authors declare no conflict of interest.


This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

<sup>1</sup>S.W.B. and J.E.A. contributed equally to this work.

<sup>2</sup>To whom correspondence should be addressed. E-mail: [jayling@jaguar1.usouthal.edu](mailto:jayling@jaguar1.usouthal.edu).

This article contains supporting information online at [www.pnas.org/cgi/content/full/0902072106/DCSupplemental](http://www.pnas.org/cgi/content/full/0902072106/DCSupplemental).



**Fig. 1.** The fully aromatic pteridine ring of folic acid (FA) presents a considerable barrier to the action of DHFR. Compared to 7,8-DHF, FA requires the loss of more resonance stabilization energy during the initial unsaturation of its pyrazine moiety. Thus, the  $V_{\max}$  for DHFR with FA is typically several orders-of-magnitude slower than with 7,8-DHF, regardless of the source of the enzyme. Glu, glutamate.

DHFR shows a well-isolated peak for THF (Fig. S2). Due to the lability of THF, especially at low concentrations, it is essential that the HPLC mobile phase contain a reducing agent (e.g., DTT) to prevent on-column oxidation. Measurement of the THF product of DHFR circumvents interference by nonspecific oxidases that hinder assays which monitor disappearance of NADPH. The HPLC/fluorescence assay is optimized by use of highly purified 7,8-DHF substrate to minimize background THF and inhibitory folic acid.

**Levels of DHFR Activity in Rat Liver with 7,8-DHF as Substrate.** DHFR activity in all rat liver extracts with 7,8-DHF as substrate was measured both spectrophotometrically and by HPLC. Activity was determined as a function of time and of DHFR concentration to ascertain that rates were measured within a valid range. The same activities were found whether or not the livers were frozen before homogenization, indicating the absence of significant DHFR in subcellular organelles. The ratio of the 2 methods, HPLC/spectrophotometric, was  $1.02 \pm 0.09$  for 100,000  $\times$  g supernatants from 7 different rats, or pooled rat preparations. The values ranged from 483 to 767 (average 577) nmol THF produced/min/gm wet weight liver (Table 1). All rats were adult and were from 2 different strains and from both sexes. Even so, there was less than a 2-fold variation in the activity of DHFR in their livers.

**Levels of DHFR Activity in Human Liver with 7,8-DHF as Substrate.** DHFR activity in extracts of human liver was found to be very low compared to rat. Measurements of activity in crude extracts were not sufficiently accurate by the spectrophotometric method since baseline rates due to nonspecific NADPH oxidation were high compared to utilization by DHFR. Therefore, all quantitative analyses were obtained using the HPLC procedure. The coefficient of variation for replicate measurements of DHFR

activity was 7%. Livers from 6 different donors were analyzed. As with the assay of enzyme from rat liver, activity was linear with reaction time for at least 10 min. (Fig. S3). The rate was also linear with the volume of extract added to the reaction (Fig. S4), demonstrating a lack of inhibition by endogenous compounds within the range examined. With 7,8-DHF as substrate, the levels of activity varied from 5.4 to 26 nmol/min/g wet weight liver (average 16.3) and did not appear to depend on age or gender (Fig. 2A). Formation of THF due to potential hydrolysis of endogenous THF polyglutamates was found to be insignificant in control reactions performed without added substrate. Even within these 6 liver samples, there is almost a 5-fold variation in the activity of DHFR in human liver. With the assumption that the preponderance of human liver DHFR activity is in the soluble fraction (as is thought to be the case with other mammalian species), the activity in human liver is 35 times lower on average (range: 22 to 106) than the average for rat liver with 7,8-DHF as substrate (Fig. 2B).

**Folic Acid as Substrate for DHFR.** Due to the slow rate of DHFR with FA, activity with this substrate was measured only by the HPLC method. The rate of THF formation by DHFR from rat liver with FA as substrate was 850 times slower than with 7,8-DHF (Table 2). The rate followed Michaelis-Menton kinetics; no substrate inhibition was observed, and  $K_m$  for FA was 1.8  $\mu$ M (Table 3).

The level of DHFR activity in human liver with FA as substrate was also found to be extremely low. The liver with the highest activity with FA was only 0.02 nmol/min/g wet weight liver, i.e., 1300 times slower than with 7,8-DHF as substrate (Table 2). The  $K_m$  for FA was found to be 0.5  $\mu$ M (Table 3). No substrate inhibition was observed, and the rate followed Michaelis-Menton kinetics. The activity in human liver is 56 times lower

**Table 1.** Rat liver DHFR activity with 7,8-DHF as substrate

Rat strain*	No. of rats <sup>†</sup>	Sex	No. of expts <sup>‡</sup>	Activity, nmol of THF produced per min per g wet weight of liver		
				HPLC assay	spec. assay <sup>§</sup>	HPLC/spec. <sup>§</sup>
CD (1)	1	M	1	767	840	0.91
CD (2)	1	M	5	737	769	0.96
CD (3)	1	M	1	526	516	1.02
CD (4)	1	M	2	508	491	1.03
CD (5)	1	M	1	483	508	0.95
Wistar	3	F	1	502	451	1.11
Wistar	16	M	3	513	437	1.17

\*Rats were adults of 300 to 480 g body weight.

<sup>†</sup>Extracts from Wistar rats were pooled from the number of animals listed.

<sup>‡</sup>Activity was measured  $\geq 4$  times per experiment, each of which was a separate homogenization.

<sup>§</sup>spec. = spectrophotometrically at 340 nm for the consumption of NADPH and 7,8-DHF.







quently not reported (17, 50). Inhibition of DHFR from human KB cells was found to be competitive at high concentrations of 7,8-DHF (51), as in this report. However, we also observed that FA behaves as a noncompetitive inhibitor of human DHFR at low concentrations of 7,8-DHF. Mammalian DHFR is a monomer with a single binding site for folate. We, therefore, suggest that at low 7,8-DHF the apparent noncompetitive inhibition by FA may be due to the binding of these 2 folates to separate species (e.g., FA to a DHFR/NADP complex and 7,8-DHF to DHFR/NADPH). At high 7,8-DHF concentrations both folates would compete for the same enzyme species. The physiological product of thymidylate synthase would probably be a polyglutamate of 7,8-DHF rather than the monoglutamate studied here. But, unlike most other folate dependent enzymes, DHFR does not have a substantially greater preference for the polyglutamylated forms (50–52). Whether unmetabolized FA accumulates within cells to the extent of meaningfully decreasing the activity of DHFR is currently unknown, and in any event is unlikely to occur as a result of doses at or below the U.S. Daily Value.

The activity measurements reported here indicate that humans may have a very limited ability to efficiently metabolize FA, especially in high doses. Numerous clinical trials examining the prevention of heart disease, stroke, and Alzheimer's disease using FA have used doses up to 5 mg/day (or even higher). Although supplements containing FA have in some trials reduced the risk of stroke, no effect on risk for coronary heart disease has yet been reported. The low activity of DHFR in human liver suggests that increasing the dose of FA may at some point, with individual variation, no longer produce further increases in the active folate pool. It is clear, however, that the Daily Value of folic acid can improve folate status (53) and helps to reduce the risk of neural tube birth defects.

Our data in no way imply that FA taken in this amount is inappropriate. Indeed, they suggest that most of the FA given in the amount of the U.S. Daily Value (0.4 mg) is converted to active folate in most individuals. Our results, however, predict that intakes higher than the Tolerable Upper Intake Level (1.0 mg), whether from a combination of supplements and fortified foods, or from high dose folic acid administered for therapeutic purposes will considerably escalate exposure to circulating unmetabolized FA. This is consistent with a recent report showing increased levels of plasma unmetabolized FA as a result of the U.S. fortification program and in consumers of supplements compared to nonconsumers (54).

## Materials and Methods

**Reagents.** FA, phenylmethylsulfonyl fluoride, NADPH, Tris base, and sodium ascorbate were from Sigma, 2-mercaptoethanol from Fluka, tetrahydrofolate from Schircks, trichloroacetic acid (TCA) from EM Science (TX1045), phosphoric acid from Baker (0260), DTT from Gold Biotechnology and HPLC grade methanol from Fisher. THF stock solutions were stored under argon at  $-80^{\circ}\text{C}$ . NADPH solutions were made in 25 mM Tris-HCl, pH 8.0, and stored frozen. Folic acid solutions were kept dark and cold and stored frozen.

Commercial sources of 7,8-DHF were found to be contaminated with FA, which inhibits the reaction, and/or with THF, which decreases the sensitivity of the HPLC assay. To avoid these problems, 7,8-DHF was synthesized in our laboratory by adaptation of the method of Blakley (see *SI Text*). Analysis by HPLC showed that after calcium precipitation the product contained less than 0.1% THF or FA. Solutions of 7,8-DHF were prepared daily in argon sparged 0.1 M sodium ascorbate, and the final concentration of ascorbate in all DHFR reactions was 4 mM.

**Human Liver DHFR.** Fresh human liver was obtained from organ donors for whom no recipient was available or in one case at surgery. All procedures were approved by the University of South Alabama Institutional Review Board. Liver was frozen in liquid nitrogen within 20 min of removal of life support and stored at  $-80^{\circ}\text{C}$  until processed. The sample of liver obtained at surgery was homogenized immediately after removal and extracted without prior freezing. To accurately determine DHFR levels in tissues, liver was extracted with minimal manipulation. Liver was homogenized on ice in a Potter-Elvehjem homogenizer with 1 g liver:3 mL homogenizing buffer. The buffer was 0.2 M Tris-HCl, pH 7.6 at  $4^{\circ}\text{C}$ , a pH and ionic strength at which DHFR was found to be maximally stable, and contained 10 mM DTT, and 0.2 mM phenylmethylsulfonyl fluoride. The homogenate was centrifuged at  $100,000 \times g$  for 60 min at  $4^{\circ}\text{C}$ , and the supernatant removed and stored in aliquots at  $-80^{\circ}\text{C}$  until assayed for activity. For measurement of  $K_m$  and  $K_i$  values, and ratio of activity with FA vs. 7,8-DHF, a 50% to 90% saturating ammonium sulfate fraction was prepared. This contained 97% of the activity and was then subjected to ultrafiltration with 0.1 M Tris-HCl, pH 7.4, to remove ammonium sulfate and endogenous THF from the retentate.

**Rat Liver DHFR.** Livers from CD or Wistar adult male or female rats were removed immediately after euthanasia. Livers were either immediately homogenized after removal or frozen in liquid nitrogen before homogenization. DHFR activity was measured in  $100,000 \times g$  supernatants of rat liver prepared exactly as for human liver.

**DHFR Activity Assay.** Two methods were used to monitor the activity of DHFR.

(i) The standard spectrophotometric assay was used when activities were sufficiently high to make this feasible. Reactions were run in water-jacketed cuvettes with the temperature maintained at  $27^{\circ}\text{C}$ , and NADPH plus 7,8-DHF consumption monitored at 340 nm with a Cary 300 spectrophotometer. Formation of THF product was calculated using a millimolar extinction coefficient of  $12.3 \text{ mM}^{-1}\text{cm}^{-1}$  with 7,8-DHF as substrate (55).

(ii) To measure very low activities in crude extracts a column switching HPLC method was developed in which the product, THF, was detected by its native fluorescence. A Perisorb C18 guard column was attached to a Kromasil C18,  $5 \mu\text{m}$ ,  $50 \times 4.6 \text{ mm}$  column in series with a Kromasil C18,  $5 \mu\text{m}$ ,  $250 \times 4.6 \text{ mm}$  column. The columns (in water jackets maintained at  $30^{\circ}\text{C}$ ) were eluted at 1.1 mL/min with 15 mM sodium phosphate, pH 2.3/methanol (83:17), containing 5 mM DTT, made fresh daily. Samples ( $20 \mu\text{L}$ ) were injected onto the first column with a Varian 420 autosampler set at  $4^{\circ}\text{C}$ . Just after all of the THF had entered the second column, a Rheodyne LabPro column switching valve reversed the flow on the first column which was flushed with the same eluant at 0.7 mL/min by a second pump while continuing the forward elution of material on the longer column. THF was detected with a Jasco FP-1520 or Waters 2475 fluorometer at 290 nm ex, 360 nm em. Data were collected and analyzed with Waters Empower software.

**DHFR Reaction Conditions.** Reactions were run in 0.1 M Tris-HCl, pH 7.4 at  $27^{\circ}\text{C}$ , 0.1 mM NADPH, 50 mM 2-mercaptoethanol, and  $60 \mu\text{M}$  FA or 7,8-DHF (except where noted), and varying amounts of human or rat liver extract as the source of DHFR. All components of the reaction mixture except substrate were temperature equilibrated for 5 min before initiation of reaction with 7,8-DHF or FA. For the spectrophotometric assay the reaction was monitored during this time to observe any baseline rate due to nonspecific oxidation of NADPH. To measure the reaction rate by HPLC, aliquots were removed every 2 min for a total of 10 min and added to a solution of ascorbic acid and TCA, final concentrations 0.1 M and 0.55 M, respectively. The precipitated protein was removed by centrifugation at  $4^{\circ}\text{C}$  for 10 min at  $30,000 \times g$ , and the supernatant immediately adjusted to pH 4.0 to 4.5 to optimize the effectiveness of ascorbate in maintaining THF. Samples were stored at  $-80^{\circ}\text{C}$  until analyzed by HPLC. THF dilutions for HPLC standards were made fresh daily in the same solution composition and pH as in the DHFR reaction samples, and treated similarly to the samples. For  $K_m$  curves samples were taken every 30 s for 2 min. Further details of the assay are available in *SI Text*.

**ACKNOWLEDGMENTS.** We acknowledge the technical assistance of Mary C. Syslo, and valuable discussions with James Appleman regarding DHFR mechanisms. This research was supported by American Heart Association Grant 0150942B and National Institutes of Health Grant HL68165.

1. Blakley RL (1984) in *Folates and Pterins*, eds Blakley RL, Benkovic SJ (John Wiley & Sons, Hoboken, NJ) Vol 1, pp 197–201.
2. Chalupsky K, Cai H (2005) Endothelial dihydrofolate reductase: Critical for nitric oxide bioavailability and role in angiotensin II uncoupling of endothelial nitric oxide synthase. *Proc Natl Acad Sci USA* 102:9056–9061.

3. Choumenkovitch SF, Selhub J, Wilson PF, Rader JI, Rosenberg IH, Jacques PF (2002) Folic acid intake from fortification in the United States exceeds predictions. *J Nutr* 132:2792–2798.
4. Quinlivan EP, Gregory JF III. (2003) Effect of food fortification on folic acid intake in the United States. *Am J Clin Nutr* 77:221–225.

5. Wright AJ, Dainty JR, Finglas PM (2007) Folic acid metabolism in human subjects revisited: Potential implications for proposed mandatory folic acid fortification in the UK. *Br J Nutr* 98:667–675.
6. Mason JB, et al. (2007) A temporal association between folic acid fortification and an increase in colorectal cancer rates may be illuminating important biological principles: A hypothesis. *Cancer Epidemiol Biomarkers Prev* 16:1325–1329.
7. Young-In Kim (2007) Folate and colorectal cancer: An evidence-based critical review. *Mol Nutr Food Res* 51:267–292.
8. Solomons NW (2007) Food fortification with folic acid: Has the other shoe dropped? *Nutr Rev* 65:512–515.
9. Smith AD, Kim YI, Refsum H (2008) Is folic acid good for everyone? *Am J Clin Nutr* 87:517–533.
10. Blount BC, et al. (1997) Folate deficiency causes uracil misincorporation into human DNA and chromosome breakage: implications for cancer and neuronal damage. *Proc Natl Acad Sci USA* 94:3290–3295.
11. Cole BF, et al. (2007) Folic acid for the prevention of colorectal adenomas: A randomized clinical trial. *J Am Med Assoc* 297:2351–2359.
12. Figueiredo JC, et al. (2009) Folic acid and risk of prostate cancer: Results from a randomized clinical trial. *J Natl Cancer Inst* 101:432–435.
13. Troen AM, et al. (2006) Unmetabolized folic acid in plasma is associated with reduced natural killer cell cytotoxicity among postmenopausal women. *J Nutr* 136:189–194.
14. Morris MS, Jacques PF, Rosenberg IH, Selhub J (2007) Folate and vitamin B-12 status in relation to anemia, macrocytosis, and cognitive impairment in older Americans in the age of folic acid fortification. *Am J Clin Nutr* 85:193–200.
15. Reynolds EH (2002) Benefits and risks of folic acid to the nervous system. *J Neurol Neurosurg Psychiatry* 72:567–571.
16. Dickinson JC (1995) Does folic acid harm people with vitamin B12 deficiency? *Q J Med* 88:357–364.
17. Jarabak J, Bachur NR (1971) A soluble dihydrofolate reductase from human placenta: Purification and properties. *Arch Biochem Biophys* 142:417–425.
18. Roberts DW, Hall TC (1965) Studies on folic reductase. II. Enzyme activity of embryonic organs of the chicken, rat, and human. *Cancer Res* 25:1894–1898.
19. Rimet O, Chauvet M, Bourdeaux M, Briand C, Sastre B (1990) Activity measurements in human tissues of the methotrexate molecular target: A novel fluorometric assay. *Cancer Biochem Biophys* 1:239–245.
20. Chio LC, Queener SF (1993) Identification of highly potent and selective inhibitors of Toxoplasma gondii dihydrofolate reductase. *Antimicrob Agents Chemother* 37:1914–1923.
21. Appleman JR, et al. (1990) Unusual transient- and steady-state kinetic behavior is predicted by the kinetic scheme operational for recombinant human dihydrofolate reductase. *J Biol Chem* 265:2740–2748.
22. Nakano T, Spencer HT, Appleman JR, Blakley RL (1994) Critical role of phenylalanine 34 of human dihydrofolate reductase in substrate and inhibitor binding and in catalysis. *Biochemistry* 33:9945–9952.
23. Segel IH (1993) *Enzyme Kinetics: Behavior and Analysis of Rapid Equilibrium and Steady-State Enzyme Systems* (Wiley Classics Library, Hoboken, NJ) pp 166–170.
24. Whitehead VM, Kamen BA, Beaulieu D (1987) Levels of dihydrofolate reductase in livers of birds, animals, primates, and man. *Cancer Drug Deliv* 4:185–189.
25. Cooperman JM, Pesci-Bourel A, Luhby AL (1970) Urinary excretion of folic acid activity in man. *Clin Chem* 16:375–381.
26. Perry J, Chanarin I (1970) Intestinal absorption of reduced folate compounds in man. *Br J Haematol* 18:329–339.
27. Melikian V, Paton A, Leeming RJ, Portman-Graham H (1971) Site of reduction and methylation of folic acid in man. *Lancet* 2:955–957.
28. Whitehead VM, Cooper BA (1967) Absorption of unaltered folic acid from the gastrointestinal tract in man. *Br J Haematol* 13:679–686.
29. Rogers LM, Pfeiffer CM, Bailey LB, Gregory JF (1997) A dual-label stable-isotopic protocol is suitable for determination of folate bioavailability in humans: Evaluation of urinary excretion and plasma folate kinetics of intravenous and oral doses of [<sup>13</sup>C<sub>5</sub>] and [<sup>2</sup>H<sub>2</sub>]folic acid. *J Nutr* 127:2321–2327.
30. Wright AJ, et al. (2005) Differential kinetic behavior and distribution for pteroylglutamic acid and reduced folates: A revised hypothesis of the primary site of PteGlu metabolism in humans. *J Nutr* 135:619–623.
31. Kelly P, McPartlin J, Goggins M, Wier D, Scott J (1997) Unmetabolized folic acid in serum: Acute studies in subjects consuming fortified food and supplements. *Am J Clin Nutr* 65:1790–1795.
32. Appleman JR, et al. (1989) Atypical transient state kinetics of recombinant human dihydrofolate reductase produced by hysteretic behavior. Comparison with dihydrofolate reductases from other sources. *J Biol Chem* 264:2625–2633.
33. Webber S, Whiteley JM (1985) Comparative activity of rat liver dihydrofolate reductase with 7,8-dihydrofolate and other 7,8-dihydropteridines. *Arch Biochem Biophys* 236:681–690.
34. Skacel N, et al. (2005) Identification of amino acids required for the functional up-regulation of human dihydrofolate reductase protein in response to antifolate Treatment. *J Biol Chem* 280:22721–22721.
35. Tai N, Schmitz JC, Chen TM, O'Neill MB, Chu E (2008) Identification of a cis-acting element of human dihydrofolate reductase mRNA. *Biochem Biophys Res Commun* 369:795–800.
36. Sowers R, et al. (2003) mRNA expression levels of E2F transcription factors correlate with dihydrofolate reductase, reduced folate carrier, and thymidylate synthase mRNA expression in osteosarcoma. *Mol Cancer Ther* 2:535–541.
37. Hjortoe GM, Weilguny D, Willumsen BM (2005) Elk3 from hamster—a ternary complex factor with strong transcriptional repressor activity. *DNA Cell Biol* 24:35–42.
38. Schwabe RF, et al. (2003) c-Jun-N-terminal kinase drives cyclin D1 expression and proliferation during liver regeneration. *Hepatology* 37:824–832.
39. Schaffner F (1991) Structural and functional aspects of regeneration of human liver. *Dig Dis Sci* 36:1282–1286.
40. Maguire M, et al. (2008) MDM2 regulates dihydrofolate reductase activity through monoubiquitination. *Cancer Res* 68:3232–3242.
41. Uchida C, et al. (2005) Enhanced Mdm2 activity inhibits pRB function via ubiquitin-dependent degradation. *EMBO J* 24:160–169.
42. Gellekink H, Blom HJ, van der Linden IJ, den Heijer M (2007) Molecular genetic analysis of the human dihydrofolate reductase gene: Relation with plasma total homocysteine, serum and red blood cell folate levels. *Eur J Hum Genet* 15:103–109.
43. Johnson WG, et al. (2004) New 19 bp deletion polymorphism in intron-1 of dihydrofolate reductase (DHFR): A risk factor for spina bifida acting in mothers during pregnancy? *Am J Med Genet A* 124:339–345.
44. Parle-McDermott A, et al. (2007) The 19-bp deletion polymorphism in intron-1 of dihydrofolate reductase (DHFR) may decrease rather than increase risk for spina bifida in the Irish population. *Am J Med Genet A* 143A:1174–1180.
45. Xu X, et al. (2007) A functional 19-base pair deletion polymorphism of dihydrofolate reductase (DHFR) and risk of breast cancer in multivitamin users. *Am J Clin Nutr* 85:1098–1102.
46. Kalmbach RD, et al. (2008) A 19-base pair deletion polymorphism in dihydrofolate reductase is associated with increased unmetabolized folic acid in plasma and decreased red blood cell folate. *J Nutr* 138:2323–2327.
47. Dulucq S, et al. (2008) DNA variants in the dihydrofolate reductase gene and outcome in childhood ALL. *Blood* 111:3692–3700.
48. Mishra PJ, et al. (2007) A miR-24 microRNA binding-site polymorphism in dihydrofolate reductase gene leads to methotrexate resistance. *Proc Natl Acad Sci USA* 104:13513–13518.
49. Fierke CA, Johnson KA, Benkovic SJ (1987) Construction and evaluation of the kinetic scheme associated with dihydrofolate reductase from Escherichia coli. *Biochemistry* 26:4085–4092.
50. Morales DR, Greenberg DM (1964) Purification and properties of dihydrofolate reductase of sheep liver. *Biochim Biophys Acta* 85:360–376.
51. Domin BA, Cheng YC, Hakala MT (1982) Properties of dihydrofolate reductase from a methotrexate-resistant subline of human KB cells and comparison with enzyme from KB parent cells and mouse S180 AT/3000 cells. *Mol Pharmacol* 21:231–238.
52. Coward JK, Parameswaran KN, Cashmore AR, Bertino JR (1974) 7,8-Dihydropteroyl oligo-gamma-L-glutamates: synthesis and kinetic studies with purified dihydrofolate reductase from mammalian sources. *Biochemistry* 13:3899–3903.
53. Lamers Y, Prinz-Langenohl R, Brämwig S, Pietrzik K (2006) Red blood cell folate concentrations increase more after supplementation with [6S]-5-methyltetrahydrofolate than with folic acid in women of childbearing age. *Am J Clin Nutr* 84:156–161.
54. Kalmbach RD, et al. (2008) Circulating folic acid in plasma: Relation to folic acid fortification. *Am J Clin Nutr* 88:763–768.
55. Hillcoat BL, Nixon PF, Blakley RL (1967) Effect of substrate decomposition on the spectrophotometric assay of dihydrofolate reductase. *Anal Biochem* 121:178–189.