



Published in final edited form as:

J Am Chem Soc. 2007 June 20; 129(24): 7514–7516. doi:10.1021/ja071974v.

Regio- and Stereoselective Cross Coupling of Substituted Olefins and Imines. A Convergent Stereoselective Synthesis of Saturated 1,5-Aminoalcohols and Substituted Piperidines

Masayuki Takahashi and Glenn C. Micalizio*

Department of Chemistry, Yale University, New Haven, Connecticut 06520-8107

The development of reactions that facilitate the syntheses of nitrogen-containing small molecules continues to be an intense area of research in organic chemistry. Whereas many powerful methods exist for the preparation of such targets, recent accomplishments in reductive coupling chemistry have provided new strategies of untapped potential for the convergent synthesis of secondary carbinolamines. To date, methods have been described for bimolecular alkyne–imine coupling that provide a convenient route to substituted allylic amines.¹ The corresponding cross coupling reaction between internal olefins and imines represents a significantly more complex and potentially more powerful bond construction.² In addition to overcoming the lower reactivity of substituted olefins in bimolecular carbometalation reactions, a synthetically useful cross coupling process must address the additional complexity that results from the need to simultaneously control site selectivity and diastereoselection in the carbon–carbon bond forming event ($2+3\rightarrow 4-7$, Figure 1). The development of a synthetic method to accomplish the cross coupling of internal olefins and imines in a regio- and stereoselective manner would represent a significant advance, and provide a fundamentally new strategy for the synthesis of stereodefined secondary carbinolamines.

Here we describe a method for cross coupling of aromatic imines with terminal, 1,1-disubstituted, (*E*)- and (*Z*)-disubstituted olefins. Additionally, we describe the use of this coupling reaction in single- and double asymmetric synthesis, as well as define a novel route for convergent stereoselective synthesis of substituted piperidines.

Previously, we have described titanium alkoxide-mediated reactions for alkyne–alkyne,³ alkyne–alkene,⁴ and alkyne–imine⁵ cross coupling. These methods have provided a means to overcome the sluggish reactivity of highly substituted π -systems in reductive coupling chemistry, and defined a means to control regioselection in bimolecular carbometalation that is independent of the differential steric environment around an unsymmetrically substituted alkyne or alkene. As depicted in Figure 2, the present study explores alkoxide-directed, titanium-mediated, reductive cross coupling between homoallylic alkoxides **8** and imines **9** as a method for the synthesis of acyclic aminoalcohols (i.e. **10** or **12**) and substituted heterocycles (i.e. **11** or **13**). In contrast to established reductive cross coupling reactions between alkynes and heterosubstituted π -systems (alkyne–aldehyde, alkyne–imine), this alkene–imine cross coupling process is significantly more complex as the carbon–carbon bond formation proceeds with the simultaneous generation of two stereogenic sp^3 carbon centers.

To examine the potential of this proposed reaction pathway for regio- and diastereoselective C–C bond formation, a study was initiated to explore the reactivity of azametallacyclopropanes (generated in situ from aromatic imines) with homoallylic alkoxides of general structure **8**. As

E-mail: glenn.micalizio@yale.edu.

depicted in entry 1 of Table 1, deprotonation of homoallylic alcohol **14** (*n*-BuLi, Et₂O) and addition to a preformed azametallacyclopropane derived from imine **15** (Ti(O*i*-Pr)₄, *c*-C₅H₉MgCl, Et₂O, -70 to -40 °C), followed by protonation of the presumed bicyclic azametallacyclopentane resulted in formation of the 1,5-amino alcohol **16** in 76% yield (dr ≥ 95:5). The success of this coupling reaction not only defines a convergent route to aminoalcohols, but also provides a convenient means to access substituted piperidines. Treatment of 1,5-aminoalcohol **16** with PPh₃ and CCl₄ (reflux) resulted in the formation of the substituted piperidine **17** in 85% yield.⁶

More highly substituted olefins were well tolerated in this cross coupling reaction. As illustrated in entry 2, coupling of the (*E*)-disubstituted homoallylic alcohol **18** with imine **15** provided the 4,5-*anti*-1,5-aminoalcohol **19** in 73% yield (dr ≥ 95:5; rr ≥ 95:5), and on cyclization, the 2,3-*trans*-disubstituted piperidine **20** in 81% yield. Coupling of the related (*Z*)-disubstituted homoallylic alcohol **21** with imine **15** provided the 4,5-*anti*-1,5-aminoalcohol **22** in 63% yield (dr ≥ 95:5; rr ≥ 95:5), and on cyclization, the 2,3-*trans*-disubstituted piperidine **23** in 77% yield (entry 3).

As depicted in Figure 3, the stereoconvergence observed in these coupling reactions can be rationalized by assuming preferential reaction by way of transition state structures that avoid 1,2-steric interactions in a developing *cis*-fused bicyclo[3.3.0] ring system (**B** and **C**).^{7, 8}

The homoallylic alcohol **24**, bearing a 1,1-disubstituted olefin, was also an effective coupling partner with imine **15** providing the 1,3-*syn*-1,5-aminoalcohol product **25** in 67% yield (dr = 4:1; rr ≥ 95:5),⁹ and on cyclization, the corresponding 2,4-*trans*-disubstituted piperidine **26** in 76% yield. The stereoselectivity in this case is consistent with preferential progression through a transition state geometry that has the phenyl substituent occupying a position outside the developing *cis*-fused bicyclo[3.3.0] ring system (**F** in Figure 3).^{7, 8}

Next, coupling reactions of achiral homoallylic alcohols with chiral imines were examined. As depicted in Table 2, coupling of homoallylic alcohols with chiral imine **28** proceeded in a generally efficient manner (61–83%).¹⁰ Highest selectivities were observed in coupling reactions of the terminal- (**14**), and (*Z*)-disubstituted (**27**) olefins, and provided products with dr ≥ 20:1 (entries 1 and 2). Cyclization of these 1,5-aminoalcohols (**29** and **31**) provides the corresponding substituted piperidines **30** and **32** with good efficiency (77 and 87% yield). Coupling of the (*E*)-disubstituted olefin **18** with chiral imine **28** also led to aminoalcohol **31**, yet this product was obtained with lower levels of stereoselection (25:4:1) as that observed with the isomeric olefin **27**.

Cross coupling reactions were also examined to determine the extent to which stereochemical matching and/or mismatching might occur in double diastereodifferentiating cases.¹¹ As illustrated in Table 3, coupling of the (*Z*)-homoallylic alcohol **33** with chiral imine *ent* - **28** furnished the 1,4-*anti*-1,5-*syn*-1,5-aminoalcohol **34** in 76% yield (dr ≥ 50:1), cyclization of which provided the 2,3-*trans*-2,5-*trans*-tetrasubstituted piperidine **35** (entry 1a). Coupling of the same homoallylic alcohol (**33**) with **28** proceeded with low conversion (30%) and provided a mixture of diastereomeric products (dr = 4:1:1; entry 2a). Although the efficiency of this coupling reaction can be increased on warming (entry 2b), the dr is affected. The disparate reactivity and selectivity observed defines a clear matched (entry 1) and mismatched (entry 2) double asymmetric relationship in these coupling reactions.

The (*E*)-disubstituted homoallylic alcohol **38** displayed inverse double asymmetric behavior as compared with (*Z*)-disubstituted homoallylic alcohol **33**. As illustrated in entries 3 and 4, coupling of **38** with **28** represents a matched double asymmetric process (88% yield, dr = 35:4:1), whereas coupling of **38** with *ent* - **28** represents a mismatched double asymmetric reaction (dr = 19:12:4:1).

A preliminary empirical model to explain these double asymmetric reactions is proposed in Figure 4. The model is based on: (1) diastereoselective coordination of the pendent methyl ether to the titanium center,¹⁰ (2) intramolecular delivery of the olefin to the azametallacyclopropane,^{3–5} (3) minimization of 1,2-interactions in the developing five-membered ring, and (4) positioning of the alkyl group (R) outside of the developing cis-fused ring system. The sum of these factors may contribute to defining matched double asymmetric reactions for the synthesis of either 1,4-*anti*-1,5-*syn* (**34**) or 1,4-*syn*-1,5-*anti* (**36**) aminoalcohols via transition state geometries **G** and **H**.^{7,8} Based on this empirical model, the mismatched double asymmetric reactions (Table 3, entries 2 and 4) may suffer from the requirement of placing the alkyl substituent (R) on the inside of the developing cis-fused ring system (**I**, **J**; Figure 4).

These double asymmetric cross coupling reactions proceed with a great deal of variability in stereoselection that is dependent on the stereochemical relationship of the secondary alcohol, imine and olefin. Interestingly, matched double asymmetric reactions can provide stereoselective access to either the 1,4-*anti*-1,5-*syn*-1,5-aminoalcohol or 1,4-*syn*-1,5-*anti*-1,5-aminoalcohol products, and the corresponding trisubstituted piperidines **35** and **37**. The variability in the diastereomeric ratio of products observed in each of these matched double asymmetric coupling reactions is consistent with a stereoreinforcing¹² relationship based on olefin geometry: (*S,Z*)-homoallylic alcohol + *ent*-**28** is matched and benefits from a stereoreinforcing relationship with the (*Z*)-olefin, whereas (*S,E*)-homoallylic alcohol + **28** is also a matched double asymmetric reaction but there is a non-stereoreinforcing relationship with the (*E*)-olefin.

Overall, we report a titanium alkoxide-mediated reaction for regio- and stereoselective cross coupling of substituted olefins and imines. The reported process provides convenient access to substituted 1,5-aminoalcohols and piperidines. Our studies focusing on simple diastereoselection have shown that this reaction is stereoconvergent with respect to olefin stereochemistry – both (*E*) and (*Z*) disubstituted homoallylic alcohols provide *anti*-products in coupling reactions with achiral imines. We have studied this reaction in both single and double asymmetric modes, have defined matched and mismatched double asymmetric relationships, and have shown a reversal of stereochemical matching/mismatching as a function of the olefin geometry of the chiral homoallylic alcohol. Research focused on further understanding the nature of stereoselection in this cross coupling reaction, as well as application to the synthesis of complex heterocycles is underway.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge financial support of this work by the American Cancer Society (RSG-06-117-01), the American Chemical Society (PRF-45334-G1), the Arnold and Mabel Beckman Foundation, Boehringer Ingelheim, Eli Lilly & Co., the National Institutes of Health –NIGMS (GM80266) and Yale University.

References

1. For cross coupling of imines and alkynes, see: (promoted by Zr): (a)Buchwald SL, Watson BT, Wannamaker MW, Dewan JC. *J Am Chem Soc* 1989;111:4486–4494.4494(b)Jensen M, Livinghouse T. *J Am Chem Soc* 1989;111:4495–4496.4496(c)Brossman RB, Davis WM, Buchwald SL. *J Am Chem Soc* 1991;113:2321–2322.2322 promoted by Ti.(d)Gao Y, Harada K, Sato F. *Tetrahedron Lett* 1995;36:5913–5916.5916 catalyzed by Ni.(e)Patel SJ, Jamison TF. *Angew Chem Int Ed* 2003;42:1364–1367.1367(f)Patel SJ, Jamison TF. *Angew Chem Int Ed* 2004;43:3941–3944.3944

- coupling of conjugated alkynes with N-sulfinyl-iminoacetates catalyzed by Rh). (g) Kong JR, Cho CW, Krische MJ. *J Am Chem Soc* 2005;127:11269–11276.11276 [PubMed: 16089454]
2. For cross coupling of zirconium-imine complexes and terminal olefins, see 1(a).
 3. Ryan J, Micalizio GC. *J Am Chem Soc* 2006;128:2764–2764. [PubMed: 16506731]
 4. Reichard HA, Micalizio GC. *Angew Chem Int Ed* 2007;46:1440–1443.
 5. McLaughlin M, Takahashi M, Micalizio GC. *Angew Chem Int Ed* 2007;46in press
 6. For an example of a related PPh₃, CCl₄-promoted cyclodehydration, see: Song YC, Okamoto S, Sato F. *Tetrahedron Lett* 2002;43:8635–8637.8637
 7. The stereochemistry of the C–Ti bond in the proposed bicyclic azametallacyclopentanes is not known.
 8. The coordination number of titanium in the transition states for these carbometalation reactions is not known. The stereoselection observed can be accounted for by assuming a transition state structure for the generation of a cis-fused bicyclo[3.3.0]-metallacycle. Proposition of an azametallacyclopentane intermediate is in accord with well-known reactivity patterns of azametallacyclopropanes.
 9. For a nickel-catalyzed coupling of substituted 1,3-dienes to imines for the synthesis of related 1,3-syn-functionalized products (i.e. 25), see: Kimura M, Miyachi A, Kojima K, Tanaka S, Tamaru Y. *J Am Chem Soc* 2004;126:14360–14361.14361 [PubMed: 15521748]
 10. For examples of stereoselective alkyne–imine cross coupling reactions with chiral imine 28, see: Fukuhara K, Okamoto S, Sato F. *Org Lett* 2003;5:2145–2148.2148 [PubMed: 12790550] (b) ref. 5
 11. Masamune S, Choy W, Petersen JS, Sita LR. *Angew Chem Int Ed Engl* 1985;24:1–30.30 For an example of triple asymmetric synthesis, see Duplantier AJ, Nantz MH, Roberts JC, Short RP, Somfai P, Masamune S. *Tetrahedron Lett* 1989;30:7357–7360.7360
 12. Evans DA, Dart MJ, Duffy JL, Yang MG. *J Am Chem Soc* 1996;118:4322–4343.




Figure 1. Bimolecular coupling of imines and substituted olefins via formation of azametallacyclopentanes.




Figure 2. Cross coupling of substituted olefins and imines – a potential pathway for the synthesis of aminoalcohols and substituted heterocycles.




With (Z)-disubstituted olefins:**With (E)-disubstituted olefins:****With 1,1-disubstituted olefins:**

Figure 3.
A model for stereoselection in imine–alkene cross coupling.^{7, 8}




Figure 4. A model for stereoselection in the double asymmetric cross coupling reactions of disubstituted olefins with chiral imines.

NIH-PA Author Manuscript

NIH-PA Author Manuscript

NIH-PA Author Manuscript

Table 1

piperidine^b (yield)

dr

rr


yield (%)

n/a

≥95:5


76


J Am Chem Soc. Author manuscript; available in PMC 2009 August 26.




piperidine^b (yield)

yield (%)	rr	dr
73	≥95.5	≥95.5



piperidine^b (yield)

yield (%)	rr	dr
63	≥95.5	≥95.5

piperidine^b (yield)


yield (%)	rr	dr
67	≥95:5	4:1

piperidine^b (yield)

dr

rr

yield (%)





piperidine^b (yield)

dr


24:1

yield (%)

83



J Am Chem Soc. Author manuscript; available in PMC 2009 August 26.



piperidine^b (yield)

dr


yield (%)

20:1

75



J Am Chem Soc. Author manuscript; available in PMC 2009 August 26.


piperidine^b (yield)

dr

yield (%)

25:4:1

61



piperidine^b (yield)

dr


yield (%)

76

≥50:1



35 (79%)

piperidine^b (yield)

dr


yield (%)

4:1:1

30

6:3:1

85


piperidine^b (yield)

dr


yield (%)

35:4:1

88



37 (78%)

piperidine^b (yield)

dr

yield (%)

19:12:4:1

76

J Am Chem Soc. Author manuscript; available in PMC 2009 August 26.

