

Cardiopulmonary exercise testing and its application

K Albouaini, M Egred, A Alahmar, D J Wright

Postgrad Med J 2007;**83**:675–682. doi: 10.1136/hrt.2007.121558

Cardiopulmonary exercise testing (CPET) has become an important clinical tool to evaluate exercise capacity and predict outcome in patients with heart failure and other cardiac conditions. It provides assessment of the integrative exercise responses involving the pulmonary, cardiovascular and skeletal muscle systems, which are not adequately reflected through the measurement of individual organ system function. CPET is being used increasingly in a wide spectrum of clinical applications for evaluation of undiagnosed exercise intolerance and for objective determination of functional capacity and impairment. This review focuses on the exercise physiology and physiological basis for functional exercise testing and discusses the methodology, indications, contraindications and interpretation of CPET in normal people and in patients with heart failure.

Cardiopulmonary exercise testing (CPET) provides assessment of the integrative exercise responses involving the pulmonary, cardiovascular, haematopoietic, neuropsychological, and skeletal muscle systems, which are not adequately reflected through the measurement of individual organ system function. This non-invasive, dynamic physiological overview permits the evaluation of both submaximal and peak exercise responses, providing the doctor with relevant information for clinical decision making. CPET is increasingly being used in a wide spectrum of clinical applications for the evaluation of undiagnosed exercise intolerance and for the objective determination of functional capacity and impairment. Its use in patient management is increasing with the understanding that resting pulmonary and cardiac function testing cannot reliably predict exercise performance and functional capacity and that overall health status correlates better with exercise tolerance than with resting measurements.¹

CPET involves measurements of respiratory oxygen uptake ($\dot{V}O_2$), carbon dioxide production ($\dot{V}CO_2$), and ventilatory measures during a symptom-limited exercise test.

PHYSIOLOGY OF EXERCISE

Peak exercise capacity is defined as "the maximum ability of the cardiovascular system to deliver oxygen to exercising skeletal muscle and of the exercising muscle to extract oxygen from the blood".² Consequently, exercise tolerance is determined by three factors: pulmonary gas exchange; cardiovascular performance, including the peripheral vascular tree; and skeletal muscle metabolism.

The Fick equation

Understanding the Fick equation is of paramount importance for appreciating the utility of functional exercise testing. At rest, the Fick equation states that oxygen uptake ($\dot{V}O_2$) equals cardiac output times the arterial minus mixed venous oxygen content:

$$\dot{V}O_2 = (SV \times HR) \times (CaO_2 - Cvo_2)$$

where SV is the stroke volume, HR is the heart rate, CaO_2 is the arterial oxygen content, and Cvo_2 is the mixed venous oxygen content. Oxygen uptake is often normalised for body weight and expressed in units of ml O_2 /kg/min. One metabolic equivalent (MET) is the resting oxygen uptake in a sitting position and equals 3.5 ml/kg/min.

At maximal exercise, the Fick equation is expressed as follows:

$$\dot{V}O_{2max} = (SV_{max} \times HR_{max}) \times (CaO_{2max} - Cvo_{2max})$$

This reflects the maximal ability of a person to take in, transport and use oxygen. It defines that person's functional aerobic capacity. $\dot{V}O_{2max}$ has become the preferred laboratory measure of cardiorespiratory fitness and is the most important measurement during functional exercise testing. In healthy people, a $\dot{V}O_2$ plateau occurs at near maximal exercise. This plateau in $\dot{V}O_2$ has traditionally been used as the best evidence of $\dot{V}O_{2max}$. It represents the maximal achievable level of oxidative metabolism involving large muscle groups. However, in clinical testing, a clear plateau may not be achieved before symptom limitation of exercise. Consequently, peak $\dot{V}O_2$ ($P\dot{V}O_2$) is often used as an estimate of $\dot{V}O_{2max}$.³

Exercise intolerance

Exercise intolerance (functional aerobic impairment) is defined as an abnormally low $\dot{V}O_{2max}$. This can occur with any factor that affects one or more of the four variables of the Fick equation that determine $\dot{V}O_{2max}$: a reduction in maximal heart rate, maximal stroke volume, or maximal CaO_2 ; or an increase in rest Cvo_2 . As an example, one of the factors limiting $\dot{V}O_{2max}$ in patients with congestive heart failure (CHF) is the marked reduction in stroke volume response to exercise with smaller reductions in maximal heart rate and maximal arterial minus mixed venous oxygen content ($CaO_{2max} - Cvo_{2max}$).^{4–6} On the other hand, diseases of the lungs, skeletal muscles and

Abbreviations: CHF, congestive heart failure; CPET, cardiopulmonary exercise testing; HF, heart failure; MET, metabolic equivalent; NYHA, New York Heart Association; $P\dot{V}O_2$, peak $\dot{V}O_2$; RER, respiratory exchange ratio; RQ, respiratory quotient; VAT, ventilatory anaerobic threshold

This is a reprint of a paper that appeared in *Heart*, October 2007, Volume 93, pages 1285–92. Reprinted with kind permission of the authors and publisher.

See end of article for authors' affiliations

Correspondence to:
Dr K Albouaini,
Cardiothoracic Centre,
Thomas Drive, Liverpool
L14 3PE, UK; Albouaini@
aol.com

Accepted 29 May 2007



Figure 1 Cardiopulmonary exercise testing machine using the cycle ergometry. A non-rebreathing valve is connected to a mouthpiece with continuous ECG and blood pressure monitoring. Resuscitation equipment on hand. Informed consent was obtained for publication of this figure.

haematological system often have a profound effect on VO_2max by affecting arterial or mixed venous oxygen content.

Exercise training in healthy people

Several important changes occur in the Fick equation as a healthy person goes from rest to maximal exercise before and after exercise training⁷:

- The VO_2max response to exercise is linear until maximal VO_2 is achieved. In many people, a plateau is reached at near maximal exercise, beyond which the VO_2 does not change. Exercise training enables the person to achieve a greater maximal workload and a higher VO_2max .
- After training, the heart rate is lower at rest and at each stage of exercise, but the maximal heart rate does not change. This approximates “220 bpm – age”.⁷
- The stroke volume response is curvilinear, increasing early in exercise with little change thereafter. The training effect increases the resting stroke volume and the stroke volume at each workload.
- The a–v O_2 content difference widens as the mixed venous O_2 content falls since arterial O_2 content does not change in normal subjects. The maximal a–v O_2 content difference increases after training.

CARDIOPULMONARY EXERCISE TESTING

Several different methods exist for measuring ventilation and respiratory gas parameters during exercise. Most clinical

Table 1 Exercise equipment: cycle ergometry vs treadmill

Variable	Cycle	Treadmill
Peak oxygen content (PVO_2)	Lower	Higher
Work rate measurement	Yes	No
Blood gas collection	Easier	More difficult
Noise and artefacts	Less	More
Safety	Safer	Less safe?
Weight bearing in obese subjects	Less	More
Degree of leg muscle training	Less	More
More appropriate for	Patients	Active normal subjects

Adapted from ATS/ACCP Statement on Cardiopulmonary Exercise Testing.¹

systems rely on breath-by-breath analysis techniques because they provide the best measures of the metabolic response to exercise. A non-rebreathing valve is connected to a mouthpiece to prevent mixing of inspired and expired air (fig 1). Oxygen and carbon dioxide gas analysers are usually incorporated in a “metabolic cart” designed specifically for functional testing. Respiratory volumes are computed by integrating the air flow signals over the time of inspiration and expiration. Average minute volumes are derived from the breath-by-breath data multiplied by the respiratory rate.

CPET protocols

Many different protocols are used for functional testing. The purpose of the test and the functional capabilities of the patient determine the choice of protocol. In evaluating patients with CHF, both bicycle and treadmill protocols have been used. The rate of workload progression is somewhat arbitrary, although it has been suggested that optimal exercise duration for functional assessment on the bicycle is between 8 and 17 minutes.⁸ Bicycle work is quantified in watts (W) or in kilopod metres/min (kpm/min; 1 W = \sim 6 kpm/min). The initial workload for patients with CHF is usually 20–25 W and increased by 15–25 W every 2 minutes until maximal exertion is reached. Alternatively, the workload can be computer controlled for electronically braked bicycle ergometers, and a ramp protocol (eg, 10 W/min) is often used. The modified Naughton protocol is recommended for treadmill exercise testing in patients with heart failure.⁹ This protocol is designed to increase the workload by approximately 1 MET (3.5 ml O_2 /kg/min) for each 2-minute stage.

Treadmill exercise testing has several advantages over cycle ergometry (table 1),¹⁰ and for most people, treadmill walking is a more familiar activity than cycling. It involves a larger muscle mass and more work against gravity. Consequently, PVO_2 is, on average, 5–10% higher on the treadmill than on a cycle ergometer. Holding onto the treadmill handrails usually decreases the metabolic cost of treadmill walking and should be discouraged if possible. A cycle ergometer is less prone to induce noise artefacts with better quantification of the metabolic cost. Generally it is less expensive and requires less space than the treadmill.

Patients with heart disease require continuous ECG monitoring and frequent blood pressure measurements during exercise testing. Since verbal communication is usually not possible with the mouthpiece apparatus, hand signals are usually used by the patient during exercise.




Figure 2 Cardiopulmonary exercise testing in a healthy 49-year-old man using the Bruce protocol. The progressive linear increase in VO_2 is noted, reaching a steady state after 2 minutes in each of the first two stages. Peak VO_2 (PVO_2) was 3.09 l/min.

Table 2 Normal cardiopulmonary exercise testing variables

Variables	Normal value
Peak oxygen content (PVO ₂)	>84% Predicted
Ventilatory anaerobic threshold (VAT)	>40% PVO ₂ (40–80%)
Maximum heart rate (HRmax)	>90% Age predicted
Heart rate reserve (HRR)	<15 Beats/min
Blood pressure (BP)	<220/90
O ₂ pulse (VO ₂ /HR)	>80%
Ventilatory reserve (VR)	MVV–VE _{max} >11 litres or VE _{max} /MVV × 100 <85%
Respiratory rate (RR)	<60 Breaths/min
Minute ventilation/carbon dioxide output ratio (VE/VCO ₂) at VAT	<34

MVV, maximal voluntary ventilation; VE, expired ventilation.
Adapted from ATS/ACCP Statement on Cardiopulmonary Exercise Testing.¹

Oxygen uptake and peak oxygen uptake

Oxygen uptake (VO₂) is determined by cellular O₂ demand up to some level that equates to maximal rate of O₂ transport, which then is determined by that maximal rate of transport. As VO₂ increases with increasing external work, one or more of the determinants of VO₂ approach limitations (eg, stroke volume, heart rate, or tissue extraction), and VO₂ versus work rate may begin to plateau. This plateau in VO₂ has traditionally been used as the best evidence of VO_{2max}.

The main determinants of a normal PVO₂ are genetic factors, quantity of exercising muscle, age, sex and body size. It can also be affected by training and patient motivation. PVO₂ should be expressed in absolute values (l/min) and as a percentage of the predicted value. VO₂ can increase from a resting value of about 3.5 ml/kg/min (about 250 ml/min in an average person) to PVO₂ values about 15 times the resting value (30–50 ml/kg/min). Athletes may attain values over 20 times their resting values (up to 80 ml/kg/min).

Figure 2 is an example of a normal CPET using the Bruce protocol. Table 2 shows the normal values of variables derived from CPET.¹

Respiratory exchange ratio

The ratio of carbon dioxide output/oxygen uptake (VCO₂/VO₂) is called the gas exchange ratio or respiratory exchange ratio (RER). Under steady state conditions, the RER equals the respiratory quotient (RQ). The RQ value is determined by the




Figure 3 Determination of the ventilatory anaerobic threshold (VAT) by the ventilatory equivalents method. The VAT is the VO₂ (1400 ml) at which the ventilatory equivalent for O₂ (VE/VO₂ ratio) begins to increase systematically without an immediate increase in the ventilatory equivalent for CO₂ (VE/VCO₂).

fuels used for metabolic processes. An RQ of 1 indicates metabolism primarily of carbohydrates, whereas an RQ of <1 indicates a mixture of carbohydrates with fat (RQ about 0.7) or protein (RQ about 0.8). The term “RQ” is often reserved for expressing events at the tissue level which are difficult to measure. The term “RER” is usually measured by gas exchange at the mouth and in true steady state, the blood and gas transport systems keep pace with tissue metabolism; thus, the RER can be used as a rough index of metabolic events (RQ). RER increases during exercise owing to either the buffered lactic acid or hyperventilation (usually towards the end of exercise).

Ventilatory anaerobic threshold

The ventilatory anaerobic threshold (VAT), formerly referred to as the anaerobic threshold, is an index used to estimate exercise capacity. During the initial (aerobic) phase of CPET, which lasts until 50–60% of VO_{2max} is reached, expired ventilation (VE) increases linearly with VO₂ and reflects aerobically produced CO₂ in the muscles. Blood lactate levels do not change substantially during this phase, since muscle lactic acid production is minimal.

During the latter half of exercise, anaerobic metabolism occurs because oxygen supply cannot keep up with the increasing metabolic requirements of exercising muscles. At this time, there is a significant increase in lactic acid production in the muscles and in the blood lactate concentration. The VO₂ at the onset of blood lactate accumulation is called the lactate threshold or the VAT. The VAT is also defined as the point at which minute ventilation increases disproportionately relative to VO₂, a response that is generally seen at 60–70% of VO_{2max}.

The VAT is a useful measure as work below this level encompasses most daily living activities. The ability to achieve the VAT can help distinguish cardiac and non-cardiac (pulmonary or musculoskeletal) causes of exercise limitation, since patients who fatigue before reaching VAT are likely to have a non-cardiac problem.^{11–12} This is not universally true, as patients with mitral stenosis, for instance, often stop exercising before reaching VAT, while on the other hand patients with chronic obstructive pulmonary disease commonly pass the VAT.

Several methods are available to determine the VAT, and these can be classified as either invasive (measurement of lactic acid or bicarbonate) or non-invasive. Invasive estimation of the VAT is usually done by direct blood sampling and is rarely used in a clinical setting since determination of the point of rise in lactate (or the reciprocal drop in bicarbonate) requires multiple blood samples. When this method is used, the “VAT” is defined as the VO₂ at which blood lactate level increases or, alternatively, when the bicarbonate starts to drop. However, the precise definition remains controversial.¹ Non-invasive methods require observing the pattern of change in VCO₂ and VE relative to VO₂ as exercise intensity increases. The two main non-invasive methods for estimating the VAT are:¹³

- The ventilatory equivalents method (fig 3): the VAT is the VO₂ at which the ventilatory equivalent for O₂ (VE/VO₂ ratio) and end-tidal oxygen tension (PET O₂) begin to increase systematically without an immediate increase in the ventilatory equivalent for CO₂ (VE/VCO₂) and end-tidal CO₂ tension (PET CO₂).
- The V-slope method (fig 4): the VAT is defined as the VO₂ at which the rate of increase in VCO₂ relative to VO₂ increases in the absence of hyperventilation.¹⁴ The VAT determined by this method is a more reproducible estimate.

Unfortunately, there is considerable inter- and intraobserver variability in the visual detection of the onset of anaerobic metabolism from the breath-by-breath data.¹⁵




Figure 4 Determination of the ventilatory anaerobic threshold (VAT) by the V-slope method in a healthy 49-year-old man. The VAT is the point at which the slope of the relative rate of increase in V_{CO_2} relative to VO_2 changes. In this patient, it occurred at a VO_2 of 1.5 l/min, or 49% of PVO_2 .

INDICATIONS AND CONTRAINDICATIONS OF CPET

The ACC/AHA Update of Practice Guidelines for Exercise Testing, published in 2002, listed the indications for ordering a functional VO_2 exercise test (table 3).^{16, 17} Contraindications for CPET are listed in (table 4). Although CPET provides excellent risk stratification in primary pulmonary hypertension, caution is advised when exercising these patients owing to the increased risk of sudden death as pulmonary pressure increases with exercise.¹⁸

INDICATIONS FOR TERMINATING CPET

Symptoms at maximal exercise that result in test termination include muscle fatigue, exhaustion, extreme dyspnoea, and light-headedness. Cardiac arrhythmias are usually not an indication to stop the test unless sustained tachyarrhythmias develop.

A decrease in systolic blood pressure below the resting pressure is a sign of insufficient increase in cardiac output to compensate for the exercise-induced systemic vasodilatation, and is an indication to stop the test. Other indications for CPET termination are given in (table 5).

CPET AND HEART FAILURE (HF)

Exercise intolerance is one of the main manifestations of HF, varying directly with the severity of the disease. Thus, decreased maximal exercise capacity is associated with higher New York Heart Association (NYHA) functional class, worse symptoms, poor quality of life and decreased patient survival. Exercise training may increase functional capacity and therefore

improve the quality of life. Its effect on survival is unclear at present.¹⁹

Exercise capacity is reduced even in mild HF. The cardiac output may be relatively normal at rest, but usually does not increase adequately with even mild exertion.²⁰ As in normal subjects, PVO_2 in HF is directly related to cardiac output and muscle blood flow at peak exercise. However, the inability to appropriately increase cardiac output results in insufficient increase in perfusion to exercising muscles, which can cause early anaerobic metabolism, muscle fatigue, and eventually can contribute to muscle wasting.²¹ Patients with HF typically do not achieve a true VO_{2max} , therefore the term "peak VO_2 " is more appropriate and is more often used. Figure 5 shows a typical example of CPET in a patient with HF.

CPET has an important role in the diagnosis, quantification of symptoms, prognosis and assessment of the success of therapeutic interventions in patients with HF.^{22, 23}

Factors contributing to exercise intolerance in HF

Several factors may contribute to exercise intolerance in patients with HF:²⁴

- The inotropic and chronotropic response to catecholamines is reduced; this defect is due, at least in part, to down-regulation of β -receptors in the presence of chronically raised circulating catecholamine levels.
- Augmentation of stroke volume by the Starling mechanism may be limited by diastolic dysfunction or pericardial constraint.
- In contrast with normal subjects, exercise is associated with a higher increase in the pulmonary wedge pressure. This can exacerbate pulmonary congestion, thereby causing dyspnoea and limiting exercise capacity.
- The presence of pulmonary hypertension and increased pulmonary vascular resistance can reduce the cardiac output response to exercise, leading to impaired exercise capacity.²⁵
- Concurrent mitral regurgitation.
- Peripheral factors affecting skeletal muscles are often present, limiting the metabolic capacity of these muscles. This includes abnormalities of the endothelial function, ergoreflex activation, vasodilatory capacity and distribution of cardiac output.²⁶

Use of CPET in assessing HF severity

The CPET is a global test of a patient's cardiorespiratory capacity, since it reflects the entire oxygen transport system

Table 3 American College of Cardiology/American Heart Association guidelines for cardiopulmonary exercise testing

Class	Indication
I (indicated)	1 Evaluation of exercise capacity and response to treatment in patients with heart failure who are being considered for heart transplantation 2 Assistance in the differentiation of cardiac versus pulmonary limitations as a cause of exercise-induced dyspnoea or impaired exercise capacity when the cause is uncertain
IIa (good supportive evidence)	Evaluation of exercise capacity when indicated for medical reasons in patients for whom the estimates of exercise capacity from exercise test time or work rate are unreliable
IIb (weak supportive evidence)	1 Evaluation of the patient's response to specific therapeutic interventions in which improvement of exercise tolerance is an important goal or end point 2 Determination of the intensity for exercise training as part of comprehensive cardiac rehabilitation
III (not indicated)	Routine use to evaluate exercise capacity

Adapted from ATS/ACCP Statement on Cardiopulmonary Exercise Testing.¹

Table 4 Absolute and relative contraindications for cardiopulmonary exercise testing

Absolute	Relative
Acute myocardial infarction (3–5 days)	Left main coronary stenosis or its equivalent
Unstable angina	Moderate stenotic valvular heart disease
Uncontrolled arrhythmias causing symptoms or haemodynamic compromise	Severe untreated arterial hypertension at rest or haemodynamic compromise (>200 mm Hg systolic, >120 mm Hg diastolic)
Syncope	Tachyarrhythmias or bradyarrhythmias
Active endocarditis	High-degree atrioventricular block
Acute myocarditis or pericarditis	Hypertrophic cardiomyopathy
Symptomatic severe aortic stenosis	Significant pulmonary hypertension
Uncontrolled heart failure	Advanced or complicated pregnancy
Acute pulmonary embolus or pulmonary infarction	Electrolyte abnormalities
Thrombosis of lower extremities	Orthopaedic impairment that compromises exercise performance
Suspected dissecting aneurysm	
Uncontrolled asthma	
Pulmonary oedema	
Room air desaturation at rest $\leq 85\%$ *	
Respiratory failure	
Acute non-cardiopulmonary disorder that may affect exercise performance or be aggravated by exercise (ie, infection, renal failure, thyrotoxicosis)	
Mental impairment leading to inability to cooperate	

Adapted from ATS/ACCP Statement on Cardiopulmonary Exercise Testing.¹

beginning with the lungs and ending with the skeletal muscles. This objective assessment offers advantages over other methods in assessing the severity of HF:

- The NYHA classification of functional impairment in HF is not always accurate since it is subjective rather than objective.
- Resting central haemodynamics do not always correlate well with functional impairment measured during exercise testing.
- The symptoms of exercise intolerance in HF, such as dyspnoea, fatigue, or both, result from a complex interplay of mechanisms originating from both the central and peripheral components of the oxygen transport system. These symptoms are non-specific and may also be due to the side effects of the drugs or to other coexisting conditions that may or may not be related to the underlying heart disease.

Therefore, CPET is often helpful for classifying disease severity for treatment decisions and in the differential diagnosis of exercise intolerance and symptoms of dyspnoea and fatigue (table 6).¹

CPET in predicting the prognosis in HF

The identification of patients at high risk of HF is of utmost importance in order to guide their pharmacological and non-pharmacological treatment (device-related treatments, heart transplantation).^{27–29} CPET, when properly performed, provides

Table 5 Indications for exercise termination

Chest pain suggestive of ischaemia
Ischaemic ECG changes
Complex ectopy
Second- or third-degree heart block
Fall in systolic pressure >20 mm Hg from the highest value during the test
Hypertension (>250 mm Hg systolic; >120 mm Hg diastolic)
Severe desaturation: $SpO_2 \leq 80\%$ when accompanied by symptoms and signs of severe hypoxaemia
Sudden pallor
Loss of coordination
Dizziness or faintness
Signs of respiratory failure
Mental confusion

Adapted from ATS/ACCP Statement on Cardiopulmonary Exercise Testing.¹

the best objective measurement of peak functional capacity, and has become an important clinical tool in defining the severity of HF and in predicting outcome.³⁰ The most relevant measures derived from CPET which have direct impact on decision making and the management plan in patients with HF are as follows:³¹

Peak oxygen consumption (PVO_2)

There is extensive evidence supporting the use of PVO_2 in stratifying the risk in patients with chronic HF. The study by Mancini *et al* is considered to be the cornerstone of the documentation of the prognostic power of PVO_2 in this group of patients.³² In the study 116 consecutive male patients with CHF referred for heart transplantation were divided into the following three groups:

1. Patients with PVO_2 of <14 ml/kg/min who had been accepted for heart transplantation.
2. Patients with PVO_2 of ≥ 14 ml/kg/min who were considered too well for transplantation.




Figure 5 Cardiopulmonary exercise testing in a 46-year-old man with cardiomyopathy in NYHA class III. A modified Bruce protocol was used. The patient reached a PVO_2 of 14 ml/kg/min (4 METs), 42% of predicted for age, gender, and weight. The ventilatory anaerobic threshold occurred at a $V̇O_2$ of 955 ml/min (slopes intersection point). This blunted cardiopulmonary response is typical in severe cardiomyopathy. The patient also had blunting in his heart rate response, with an HRmax of 120 bpm. He was listed for cardiac transplantation. HR, heart rate; METs, metabolic equivalents.

Table 6 American Thoracic Society/American College of Chest Physicians: usual cardiopulmonary exercise response patterns

Measurement	Heart failure	COPD	ILD	Pulmonary vascular disease	Obesity	Deconditioned
PV _{O₂}	↓	↓	↓	↓	↓ for actual, N for ideal weight	↓
VAT	↓	N / ↓ / indeterminate	N or ↓	↓	N	N or ↓
Peak HR	Variable, N in mild	↓, N in mild	↓	N / slightly ↓	N / slightly ↓	N / slightly ↓
O ₂ Pulse	↓	N or ↓	N or ↓	↓	N	↓
VE/MVV × 100	N or ↓	↑	N or ↑	N	N or ↑	N
VE/V _{CO₂} at VAT	↑	↑	↑	↑	N	N
VD/VAT	↑	↑	↑	↑	N	N
PaO ₂	N	Variable	↓	↓	N / may ↑	N
P(A-a)O ₂	Usually N	Variable, usually ↑	↑	↑	May ↓	N

COPD, chronic obstructive pulmonary disease; HR, heart rate; ILD, interstitial lung disease; MVV, maximum voluntary ventilation; N, normal; P(A-a)O₂, alveolar-arterial difference for oxygen pressure; PV_{O₂}, peak oxygen uptake; VAT, ventilatory anaerobic threshold; VD/VAT, ratio of physiological dead space to tidal volume; VE, minute ventilation; V_{CO₂}, carbon dioxide output.

Adapted from ATS/ACCP Statement on Cardiopulmonary Exercise Testing.¹

3. Patients with PV_{O₂} of <14 ml/kg/min but with significant comorbidity that precluded heart transplantation.

The 1-year survival rates in groups 1, 2 and 3 were 48%, 94% and 47%, respectively. Of note, a PV_{O₂} of <10 ml/kg/min was associated with significantly poorer predicted survival. The results of this study were adopted by the AHA/ACC consensus statement on the selection and treatment of candidates for heart transplantation.³³ This consensus statement emphasised that once maximal treatment has been instituted and maintained, PV_{O₂} has a predictive role and that for patients with PV_{O₂} <14 ml/kg/min, it is important to prove that exercise testing was truly maximal by documenting achievement of the VAT at approximately 50–70% of PV_{O₂}.

As the value of PV_{O₂} can be influenced by many factors other than the severity of CHF, the PV_{O₂} value adjusted for these factors should theoretically improve the predictive accuracy. Stelken *et al*³⁴ retrospectively studied 181 patients with CHF in order to compare the percentage achieved of the predicted PV_{O₂}, taking into account age, gender, and weight, with the traditionally used absolute PV_{O₂} measured in ml/kg/min. During the 12-month follow-up, non-survivors achieved a lower percentage of the predicted PV_{O₂} and a lower absolute PV_{O₂} than survivors, and multivariate analysis showed that a value <50% of the predicted PV_{O₂} was the strongest predictor of cardiac events, better than the cut-off absolute value of PV_{O₂} of <14 ml/kg/min. Aaronson and Mancini refuted these results in another study.³⁵ Osman *et al* have documented that the adjustment of PV_{O₂} to lean body weight provides greater prognostic value than the traditional standard PV_{O₂}. Lean PV_{O₂}, either as a continuous variable or using a cut-off value of ≤19 ml/kg/min, was a better predictor of outcome than unadjusted PV_{O₂} in 225 patients with chronic HF.³⁶

Ventilatory anaerobic threshold

The VAT has been proposed as a submaximal index of exercise capacity, independent of the patient's motivation. A failure to reach the VAT strongly suggests poor motivation or non-cardiovascular limitation of exercise tolerance. The VAT is undetectable in a large proportion of patients with CHF, especially in those with reduced exercise tolerance, restricting the prognostic and decisional value of PV_{O₂}.³⁷

Opasich *et al* evaluated the predictive role of PV_{O₂} in 505 male patients with CHF in whom VAT had and had not been identified.³⁸ When the VAT was detected, the cardiac event rates were:

- 59% in patients with PV_{O₂} values of ≤10 ml/kg/min
- 32% in patients with PV_{O₂} values of >10 to ≤14 ml/kg/min

- 32% in patients with PV_{O₂} values of >14 to ≤18 ml/kg/min
- 15% in patients with PV_{O₂} values of >18 ml/kg/min.

Therefore, when VAT is detected, patients with a PV_{O₂} of <10 ml/kg/min have a high event rate, whereas those with a PV_{O₂} of >18 ml/kg/min have a good prognosis. Patients with intermediate functional capacity (PV_{O₂} 10–18 ml/kg/min) fall into a range in which PV_{O₂} does not provide predictive or decisional information.

When VAT was not detected, the event rate was 46% in patients with PV_{O₂} of ≤10 ml/kg/min, and was almost identical among those with PV_{O₂} values of >10 to ≤14 ml/kg/min, >14 to ≤18 ml/kg/min, and >18 ml/kg/min (29%, 23% and 22%, respectively). Thus, when the VAT is undetectable, patients with PV_{O₂} of <10 ml/kg/min have a high risk of events; and for those with PV_{O₂} >10 ml/kg/min, the risk stratification is considered to be inconclusive and the replication of CPET is recommended.

Ventilatory expired gas parameters (VE/V_{CO₂}slope)

Recently, the VE/V_{CO₂} slope has been proposed as an alternative prognostic indicator in patients with CHF. This dimensionless ratio indicates how many litres of air are being breathed to eliminate 1 litre of CO₂. An abnormally high relationship between minute ventilation (VE) and carbon dioxide output (V_{CO₂}) (VE/V_{CO₂} slope) is associated with a poor outcome. Chua *et al* reported that a VE/V_{CO₂} slope of >34 was associated with worse prognosis in 173 patients with chronic HF,³⁹ and Kleber *et al* selected a VE/V_{CO₂} slope that was >130% of the age adjusted and sex-adjusted value as the best predictive cut-off point in 142 patients.⁴⁰ MacGowan *et al* substantiated that the combination of a peak VE/V_{CO₂} slope of >50 and a peak V_{O₂} of ≤15 ml/kg/min was associated with an 82% mortality rate in 104 patients with chronic HF.⁴¹ Finally, Robbins *et al* found that a VE/V_{CO₂} slope of ≥44.7 at peak exercise was better than a peak V_{O₂} of ≤13.9 ml/kg/min in predicting 18-month survival.⁴²

Recommended risk stratification algorithm

Taking into account the above mentioned measures derived from CPET to stratify risk for patients with HF and results from other trials,^{43–45} Corra and Mezzani proposed a new risk stratification algorithm (fig 6).³¹ This algorithm is structured on a multiparametric decoding scrutiny employing the stepwise introduction of PV_{O₂}, VE/V_{CO₂}slope, and peak RER. They divided patients with CHF into the following three groups:

- PV_{O₂} of ≤10 ml/kg/min identifies high-risk patients. The peak RER can correctly discriminate outcome, as those reaching an RER of at least 1.15 at peak effort have a higher risk, whereas those who do not attain a peak RER of 1.15




Figure 6 A new prognostic algorithm that is structured on a multiparametric decoding scrutiny employing the stepwise introduction of PV_{O_2} , VE/V_{CO_2} slope, and peak respiratory exchange ratio (RER). The arrow indicates that patients with intermediate exercise capacity (ie, PV_{O_2} of >10 and <18 ml/kg/min) and excessive ventilatory response (ie, VE/V_{CO_2} slope of ≥ 35) have a total mortality rate that is comparable to that detected with a PV_{O_2} of ≤ 10 ml/kg/min (whole population). (Adapted from Corra and Mezzani³¹.)

have a prognosis comparable to that of patients with a better functional capacity.

- PV_{O_2} of ≥ 18 ml/kg/min is compatible with a fairly good long-term prognosis.
- PV_{O_2} between 10 and 18 ml/kg/min indicates a moderate risk of cardiac events. A VE/V_{CO_2} slope of ≥ 35 allows the identification of those patients with worse outcomes.

The VE/V_{CO_2} slope yields an efficient predictive contribution for almost one-quarter of patients with moderate chronic HF. The attainment of a peak RER of ≥ 1.15 allows the identification of nearly half of patients with severe exercise intolerance with “true” low peak aerobic power, who are thus at high risk. This stepwise process can assist doctors in clinical decision making by describing a reliable risk for the individual patient.

CPET DATA AND ITS REPORTING

The reporting of CPET should clearly show the indication as well as the beginning and end of exercise. Table 7 summarises the information that should be included in the final report, and an example of a CPET data report is available online as supplementary material (see <http://heart.bmj.com/supplemental>).

CONCLUSIONS

CPET is a global test of the cardiorespiratory capacity that reflects the entire oxygen transport system starting with the lungs and ending with the skeletal muscles. Interpretation of

CPET can be a complex task; however, PV_{O_2} derived from CPET is a strong and independent factor in determining the prognosis of patients with CHF. In the past few years, it has been discovered that a single arbitrary cut-off point for PV_{O_2} may be unlikely to describe the true risk of events, and that a stepwise approach may be more helpful. Specific research is still needed to confirm the importance of a stepwise multiparametric interpretation of CPET. Finally, CPET has an important role as part of a holistic approach to the assessment and management of cardiac patients.



Supplementary material available online at <http://heart.bmj.com/supplemental>

.....

Authors' affiliations

Khaled Albouaini, Mohaned Egred, Albert Alahmar, David Justin Wright, Cardiothoracic Centre, Liverpool, UK

Conflict of interest: None declared.

REFERENCES

- 1 **ATS/ACCP statement on cardiopulmonary exercise testing.** *Am J Respir Care Med* 2003;**167**:211–77.
- 2 **Dennis C.** Rehabilitation of patients with coronary artery disease. In: Braunwald E, ed. *Heart disease, a textbook of cardiovascular medicine*, 4th ed. Philadelphia: Saunders, 1992:1382.
- 3 **Brooks GA.** Anaerobic threshold: review of the concept and directions for future research. *Med Sci Sports Exerc* 1985;**17**:22–34.
- 4 **Simonton CA, Higginbotham MB, Cobb FR.** The ventilatory threshold: quantitative analysis of reproducibility and relation to arterial lactate concentrations in normal subjects and in patients in congestive heart failure. *Am J Cardiol* 1988;**62**:100–7.
- 5 **Matsumura N, Nishijima H, Kojima S et al.** Determination of an anaerobic threshold for assessment of functional state in patients with congestive heart failure. *Circulation* 1983;**68**:360–7.
- 6 **Neuberg GW, Friedman SH, Weiss MB, et al.** Cardiopulmonary exercise testing. The clinical value of gas exchange data. *Arch Intern Med* 1988;**148**:2221–6.
- 7 **Mitchell JH, Blomqvist G.** Maximal oxygen uptake. *N Engl J Med* 1971;**284**:1018–22.
- 8 **Buchfuhrer MJ, Hansen JE, Robinson TE, et al.** Optimizing the exercise protocol for cardiopulmonary assessment. *J Appl Physiol* 1983;**55**:1558–64.
- 9 **Naughton J, Sevelius G, Balke B.** Physiological responses of normal and pathologic subjects to a modified work capacity test. *J Sports Med* 1963;**31**:201–7.
- 10 **Wright DJ, Khan KM, Gossage EM, et al.** Assessment of a low-intensity cardiac rehabilitation programme using the six-minute walk test. *Clin Rehabil* 2001;**15**:119–24.
- 11 **Davis JA.** Anaerobic threshold: review of the concept and directions for future research. *Med Sci Sports Exerc* 1985;**17**:6–21.
- 12 **Jennings GL, Esler MD.** Circulatory regulation at rest and exercise and the functional assessment of patients with congestive heart failure. *Circulation* 1990;**81**:115–13.
- 13 **Wasserman K, Hansen JE, Sue DY, et al.** *Principles of exercise testing and interpretation*. Philadelphia: Lea and Febiger, 1987.
- 14 **Beaver WL, Wasserman K, Whipp BJ.** A new method for detecting the anaerobic threshold by gas exchange. *J Appl Physiol* 1986;**60**:2020–7.
- 15 **Yeh MP, Gardner RM, Adams TD, et al.** “Anaerobic threshold”: problems of determination and validation. *J Appl Physiol* 1983;**55**:1178–86.

Table 7 Summary of the data that should be included in the cardiopulmonary exercise testing (CPET) final report

Pretest information

- Reason for test
- General information: age, height, weight
- Pretest diagnoses, drugs, resting ECG and BP

Exercise modality and equipment

- Treadmill vs cycle ergometer
- Exercise protocol: ramp, incremental
- Modality of gas sampling: mask vs mouthpiece

Observations during exercise

- Reason(s) for termination
- Symptoms and complications
- Subjective assessment of effort
- Gas exchange and ventilatory data at peak and at VAT(if determined): absolute and percentage relative to reference
- HR, BP, and ECG changes

CPET interpretation

- Presence and severity of functional impairment
- Probable cause of functional impairment
- Comparison with previous functional evaluation if available

Adapted from reference⁴⁶.

BP, blood pressure; HR, heart rate; VAT, ventilatory anaerobic threshold.

- 16 **Gibbons RJ**, Balady GJ, Bricker JT, *et al*. ACC/AHA 2002 guideline update for exercise testing: summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1997 Exercise Testing Guidelines). *Circulation* 2002;**106**:1883–92.
- 17 **Gibbons RJ**, Balady GJ, Beasley JW, *et al*. ACC/AHA guidelines for exercise testing: A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Exercise Testing). *J Am Coll Cardiol* 1997;**30**:260–311.
- 18 **Rochmis P**, Blackburn H. Exercise tests: a survey of procedures, safety, and litigation experience in approximately 170,000 tests. *JAMA* 1971;**217**:1061–6.
- 19 **Pina IL**, Apstein CS, Balady GJ, *et al*. Exercise and heart failure: a statement from the American Heart Association Committee on exercise, rehabilitation, and prevention. *Circulation* 2003;**107**:1210–25.
- 20 **Reddy HK**, Weber KT, Janicki JS, *et al*. Hemodynamic, ventilatory and metabolic effects of light isometric exercise in patients with chronic heart failure. *J Am Coll Cardiol* 1988;**12**:353–8.
- 21 **Harrington D**, Anker SD, Chua TP, *et al*. Skeletal muscle function and its relation to exercise tolerance in chronic heart failure. *J Am Coll Cardiol* 1997;**30**:1758–64.
- 22 **Wright DJ**, Tan LB. The role of exercise testing in the evaluation and management of heart failure. *Postgrad Med J* 1999;**75**:453–8.
- 23 **Ghali JK**, Cooper R, Ford E. Trends in hospitalisation rates for heart failure in the US 1973–86. *Arch Intern Med* 1990;**150**:769–73.
- 24 **Colucci WS**. In vivo studies of myocardial beta-adrenergic receptor pharmacology in patients with congestive heart failure. *Circulation* 1990;**82**:144–51.
- 25 **Butler J**, Chomsky DB, Wilson JR. Pulmonary hypertension and exercise intolerance in patients with heart failure. *J Am Coll Cardiol* 1999;**34**:1802–6.
- 26 **Pina IL**, Apstein CS, Balady GJ, *et al*. Exercise and heart failure: a statement from the American Heart Association Committee on exercise, rehabilitation, and prevention. *Circulation* 2003;**107**:1210–25.
- 27 **McMurray J**, Pfeffer MA. New therapeutic options in congestive heart failure: part I. *Circulation* 2002;**105**:2099–106.
- 28 **McMurray J**, Pfeffer MA. New therapeutic options in congestive heart failure: part II. *Circulation* 2002;**105**:2223–8.
- 29 **Abraham WT**, Fisher WG, Smith AL, *et al*. Cardiac resynchronization in chronic heart failure. *N Engl J Med* 2002;**346**:1845–53.
- 30 **Benjaminovitz A**, Mancini DM. The role of exercise-based prognosticating algorithms in the selection of patients for heart transplantation. *Curr Opin Cardiol* 1999;**4**:114–20.
- 31 **Corra U**, Mezzani A, Bosimini E, *et al*. Cardiopulmonary exercise testing and prognosis in chronic heart failure. *Chest* 2004;**126**:942–50.
- 32 **Mancini DM**, Eisen H, Kusmaul W, *et al*. Value of peak oxygen consumption for optimal timing of cardiac transplantation in ambulatory patients with heart failure. *Circulation* 1991;**83**:778–86.
- 33 **Costanzo MR**, Augustine S, Bourge R, *et al*. Selection and treatment of candidates for heart transplantation. *Circulation* 1995;**92**:3595–612.
- 34 **Stelken AM**, Younis LT, Jennison SH, *et al*. Prognostic value of cardiopulmonary exercise testing using percent achieved of predicted peak oxygen uptake for patients with ischemic and dilated cardiomyopathy. *J Am Coll Cardiol* 1996;**27**:345–52.
- 35 **Aaronson KD**, Mancini DM. Is percentage of predicted maximal exercise oxygen consumption a better predictor of survival than peak exercise oxygen consumption for patients with severe heart failure. *J Heart Lung Transplant* 1995;**14**:981–9.
- 36 **Osman AF**, Mehra MR, Lavie CJ, *et al*. The incremental prognostic importance of body fat adjusted peak oxygen consumption in chronic heart failure. *J Am Coll Cardiol* 2000;**36**:2126–31.
- 37 **Cohen-Solal A**, Zannad F, Kayanakis JG, *et al*. Multicenter determination of the oxygen uptake and ventilatory threshold. *Eur Heart J* 1991;**12**:1055–63.
- 38 **Opasich C**, Pinna GD, Bobbio M, *et al*. Peak oxygen consumption in chronic heart failure: toward efficient use in the individual patient. *J Am Coll Cardiol* 1998;**31**:766–75.
- 39 **Chua TP**, Ponikowski P, Harrington D, *et al*. Clinical correlates and prognostic significance of the ventilatory response to exercise in CHF. *J Am Coll Cardiol* 1997;**29**:1585–90.
- 40 **Kleber FX**, Vietzke G, Wernecke KD, *et al*. Impairment of ventilatory efficiency in heart failure: prognostic impact. *Circulation* 2000;**101**:2803–9.
- 41 **MacGowan GA**, Janosko K, Cecchetti A, *et al*. Exercise related ventilatory abnormalities and survival in congestive heart failure. *Am J Cardiol* 1997;**79**:1264–6.
- 42 **Robbins M**, Francis G, Pashkow FJ, *et al*. Ventilatory and heart rate response to exercise: better predictors of heart failure mortality than peak oxygen consumption. *Circulation* 1999;**100**:2411–7.
- 43 **Corra U**, Mezzani A, Bosimini E, *et al*. Ventilatory response to exercise improves risk stratification in chronic heart failure patients with intermediate functional capacity. *Am Heart J*, **143**:418–26.
- 44 **Mezzani A**, Corra U, Baroffio C, *et al*. Habitual activities and peak aerobic capacity in patients with asymptomatic and symptomatic left ventricular dysfunction: use of a new physical activity scoring system. *Chest* 2000;**117**:1291–9.
- 45 **Mezzani A**, Corra U, Bosimini E, *et al*. Contribution of peak respiratory exchange ratio to peak VO₂ prognostic reliability in patients with chronic heart failure and severely reduced exercise capacity. *Am Heart J* 2003;**145**:1102–7.
- 46 **Task Force of the Italian Working Group on Cardiac Rehabilitation and Prevention** (Gruppo Italiano di Cardiologia Riabilitativa e Prevenzione, GICR); Working Group on Cardiac Rehabilitation and Exercise Physiology of the European Society of Cardiology, Piepoli MF, *et al*. Statement on cardiopulmonary exercise testing in chronic heart failure due to left ventricular dysfunction: recommendations for performance and interpretation Part II: How to perform cardiopulmonary exercise testing in chronic heart failure. *Eur J Cardiovasc Prev Rehabil* 2006;**13**:300–11.