



Published in final edited form as:

J Org Chem. 2008 December 19; 73(24): 9610–9618. doi:10.1021/jo801410s.

Synthesis of a 7-Azaindole by Chichibabin Cyclization: Reversible Base-Mediated Dimerization of 3-Picolines

Yun Ma, Sean Breslin, Ivan Keresztes, Emil Lobkovsky, and David B. Collum*

Contribution from the Department of Chemistry and Chemical Biology Baker Laboratory, Cornell University, Ithaca, New York 14853-1301


Abstract

The lithium diisopropylamide (LDA)-mediated condensation of 2-fluoro-3-picoline and benzonitrile to form 2-phenyl-7-azaindole via a Chichibabin cyclization is described. Facile dimerization of the picoline via a 1,4-addition of the incipient benzyllithium to the picoline starting material and fast 1,2-addition of LDA to benzonitrile cause the reaction to be complex. Both adducts are shown to reenter the reaction coordinate to produce the desired 7-azaindole. The solution structures of the key intermediates and the underlying reaction mechanisms are studied by a combination of IR and NMR spectroscopies.

Introduction

7-Azaindoles have been found to be therapeutically active for an array of maladies.¹ The primary synthetic challenge presented by azaindoles reduces to one of heteroannulation--appending a pyrrole to a pyridine or vice versa.^{1,2} In connection with a program at sanofi-aventis to synthesize polycyclic pyrrole derivatives to be tested for the treatment of asthma,^{2c} the conversion of 2-fluoro-3-picoline (**1**) to 7-azaindole **2** was examined (Scheme 1). The procedure was modeled after that of Wakefield and coworkers in which metalation of 3-picoline and subsequent addition of benzonitrile (PhCN) affords 7-azaindole **2** in one pot.³ The key cyclization of putative intermediate **4** to **2** corresponds to a Chichibabin reaction--the nucleophilic addition of an alkali metal amide to a pyridine.^{4,5} Substitution of the fluoro moiety precludes an air oxidation required by standard Chichibabin protocols. The fluorine does not improve the yield, but ¹⁹F NMR spectroscopy proves to be particularly useful to study the reaction.⁶

The Cornell group was charged with studying the underlying structural and mechanistic organolithium chemistry. The results are particularly odd owing to *non-fatal* condensations of benzyllithium **3** with picoline **1** and of LDA with PhCN (eq 1), which leave *none* of the original reagents intact. We examined how adducts **5** and **6** form and reenter the reaction coordinate to provide azaindole **2** in high yield. The results are summarized at the beginning of the discussion section for the benefit of the non-specialist. The importance of dimerizations during lithiations of heterocycles is discussed in a broader context.



(1)

Results

Azaindole Synthesis

The conversion of fluoropicoline **1** to azaindole **2** proceeds smoothly by sequential addition of picoline **1** to 2.1 equiv of LDA in THF at $-40\text{ }^{\circ}\text{C}$. After 60 min, addition of 1.2 equiv of PhCN to the blood-red reaction with stirring for an additional 2.0 h at $-40\text{ }^{\circ}\text{C}$ affords **2** in 80% yield. Alternatively, reversing the order of addition--adding 1.05 equiv PhCN to 2.1 equiv LDA at $-40\text{ }^{\circ}\text{C}$ followed by 1.0 equiv fluoropicoline **1**--affords azaindole **2** in 82% yield. An inferior (15–20%) yield of **2** is obtained if only 1.05 equiv of LDA is used. The second equivalent may be required for a tautomerization following the cyclization of **4**. No ketone derived from hydrolysis of putative intermediate **4** could be observed despite prompt aqueous quenches at low temperature.

Byproducts Derived from Picoline Dimerization


A number of byproducts emerged during the optimizations that attest to picoline dimerizations.⁷ The byproducts, prepared in optimized yields as illustrated in Scheme 2, foreshadow an underlying mechanistic complexity. They were characterized by a combination of ^1H , ^{13}C , ^{19}F , $[\text{H},^{13}\text{C}]$ HMBC, $[\text{H},^{13}\text{C}]$ HSQC, $[\text{H},\text{H}]$ COSY, and $[\text{H},^{19}\text{F}]$ *J*-resolved spectroscopies, IR spectroscopy, and mass spectrometry (supporting information).

Lithiating **1** at $-40\text{ }^{\circ}\text{C}$ followed by quenching with wet THF at $-78\text{ }^{\circ}\text{C}$ affords **7**, the product of 1,4-addition of benzyllithium **3** to picoline **1**, as a transiently stable intermediate.⁸ Dihydropyridine **7** displays vinyl ^1H resonances at δ 5.87 and 4.06 ppm characteristic of the enamino moiety as well as a 1:1 pair of ^{19}F resonances at δ -70.1 and -114.8 ppm. Whereas the ^{19}F resonance at δ -70.1 ppm is consistent with a fluoropyridine moiety,⁹ the resonance at δ -114.8 ppm is decidedly upfield. Aqueous workup of **7** affords a 3:1 mixture of lactams **8a** and **8b** isolated in 89% combined yield along with traces of **9** (11%). Lactams **8a** and **8b** were separated, but their stereochemistries were not assigned. Presumably **8a** and **8b** result from hydrolysis of fluorodihydropyridine **7** whereas **9** derives from its air oxidation. The yield of **9** increases to 41% by adding O_2 at $-40\text{ }^{\circ}\text{C}$ prior to the aqueous quench.¹⁰ If solutions of **1** and LDA are warmed to $0\text{ }^{\circ}\text{C}$ prior to an aqueous quench, product **10** resulting from dimerization by a 1,2-addition is isolated in 88% yield.^{11–13}

Origins of 1,4-Adducts

Metalation of **1** by 1.0 equiv of $[\text{Li},^{15}\text{N}]$ LDA affords lithiated dihydropyridine **5** (eq 2) of unknown aggregation number. ^6Li NMR spectroscopy reveals the resonance corresponding to unreacted LDA dimer **11**^{14,15} along with a single resonance displaying no ^6Li - ^{15}N coupling, initially presumed to derive from lithiated picoline **3**. The ^{19}F NMR spectra, however, display *two* resonances in a 1:1 proportion at markedly different chemical shifts (δ -93.5 and -70.5


ppm).¹⁶ A combination of ¹⁹F and ¹H NMR spectroscopies showed coupling patterns fully consistent with **5** (eq 2); vinyl resonances at 5.83 and 3.80 ppm are especially characteristic of the lithioimine moiety. At no point did we observe resonances corresponding to **9** (or a lithiated counterpart¹⁷), confirming that **9** forms during workup.



(2)

The underlying mechanism of the dimerization is shown to involve rate-limiting picoline metalation with a subsequent rapid dimerization; details of the rate studies are described in the experimental section.¹⁸ The idealized¹⁹ rate law (eq 3), in conjunction with the assignment of LDA/THF as a disolvated dimer **11**,^{14,15} is consistent with lithiation via a tetrasolvated monomer-based transition structure, which we offer **12**. The third-order dependence on the THF concentration (measured 2.9 ± 0.2) was surprising. Precedent suggests that the THF order could include secondary-shell solvation effects.²⁰ Nonetheless, using 2,2,5,5-Me₄THF as a polar, but chemically inert cosolvent afforded no reduction in the THF order.²⁰ Evidence of high coordinate lithium support such a highly solvated transition structure as possible.²¹ The depicted Li-F interaction is supported by evidence that such Li-F interactions are highly stabilizing.²² With that said, the solvation number and existence of an Li-F interaction in **12** should be viewed with a healthy skepticism.


$$-d[1]/dt = k[1][\text{LDA}]^{1/2}[\text{THF}]^3 \quad (3)$$



Reversible formation of **5** was demonstrated by a double-crossover study summarized in Scheme 3.²³ Mixing independently prepared samples of **5-d₀** and **5-d₅** at $-78\text{ }^{\circ}\text{C}$ and quenching after 10 min afforded a mixture of **9-d₀** and **9-d₅** contaminated with only traces (<5%) of crossover products **9-d₂** and **9-d₃** as shown by GC-MS analysis. (The low levels of crossover are probably elicited by inadvertent warming.) Warming to $-40\text{ }^{\circ}\text{C}$ affords an approximate statistical distribution.²⁴ Therefore, the 1,4-addition to form **5** is reversible under the conditions of the metalation of picoline **1** ($-40\text{ }^{\circ}\text{C}$) and the rearrangement to form **10** ($0\text{ }^{\circ}\text{C}$), but *not* at low temperatures. This temperature dependence proves to be important (*vide infra*).²⁵

Origins of 1,2-Adduct **10**

Warming solutions of lithiated dihydropyridine **5** to $0\text{ }^{\circ}\text{C}$ --conditions affording 1,2-adduct **10** in high yield (eq 4)--causes the ^{19}F resonances of **5** to disappear via a first-order decay.^{26, 27} A single ^{19}F resonance at $\delta -77.9$ ppm corresponding to a lithiated form of **10** (presumably **13**) appears concomitantly. Quenching affords **10** ($\delta -70.9$ ppm). Treating a purified sample of **10** with LDA causes the resonance of **13** at $\delta -77.9$ ppm to reappear. The approximate first-order decay of **5** and the crossover experiments are congruent with formation of **10** via benzyllithium **3**.²⁸



(4)

Condensation of Lithiated Dihydropyridine **5** with PhCN


Monitoring the solutions containing PhCN, **5** and residual LDA (0.5 equiv relative to starting picoline **1**) using ^{19}F NMR spectroscopy shows the disappearance of picoline **1** at $-40\text{ }^{\circ}\text{C}$. Adduct **2** (or its lithium salt) is not detectable by ^{19}F NMR spectroscopy. An absence of fluorine-containing intermediates suggests that cyclization of imidolithium **4** to afford azaindole **2** is very fast.²⁹

Condensation of LDA with PhCN

Monitoring the reaction described above using *in situ* IR spectroscopy³⁰ shows that PhCN disappears with a 3 min half life at $-78\text{ }^{\circ}\text{C}$ ($\ll 1$ min at $-40\text{ }^{\circ}\text{C}$.) This is substantially faster than the loss of **5**. This loss of PhCN was traced to the condensation of the remaining LDA (0.5–1.5 equiv; not shown) to form amidinolithium **6** as the first step of a cascade of reactions leading to PhCN trimer **15** (Scheme 4).^{31–33} Indeed, warming a mixture of **6** and PhCN to $0\text{ }^{\circ}\text{C}$ affords an intensely purple solution (appearing black), which affords PhCN-derived trimer **15** in 83% yield. Base-mediated trimerizations of aryl nitriles have been reported previously.^{31,32}

The condensation of LDA and PhCN is important because it affords **6** *substantially faster than the metalation of 1 by LDA*. We examined the kinetics by monitoring the loss of PhCN (2230 cm^{-1}) and the formation of amidinolithium **6** (1627 cm^{-1}) as described in the experimental section. The first term in the resulting rate law in eq 5 is consistent with open dimer-based transition structures **16** or **17**.^{18,34} (The reaction rate at low THF concentrations was too high to measure the LDA order in the second term of eq 5.)

$$-d[\text{PhCN}]/dt = k[\text{PhCN}][\text{LDA}][\text{THF}]^0 + k'[\text{PhCN}][\text{LDA}]^n[\text{THF}]^{-1} \quad (5)$$




(6)


Chichibabin Cyclization using Amidinolithium 6

Solutions containing amidinolithium **6** at $-40\text{ }^\circ\text{C}$ --conditions affording Chichibabin cyclization--also cause further oligomerization of **6** (actually homodimer **19**) to a new species with a slightly downfield-shifted ^6Li resonance that we attribute to oligomer(s) **14**. (Possible intermediates in PhCN trimerizations have been discussed by Wheatley and coworkers.³²) Monitoring Chichibabin cyclizations by ^6Li NMR spectroscopy reveal that **6** and **14** form but then reenter the reaction coordinate (although more reluctantly for **14**).


The most logical explanation for the success of the protocol in eq 1 is that that **6** and **14** form reversibly (eq 7). We did, however, consider alternative models. In principle, **6** could condense with **1** via 1,2-addition to form **20** with subsequent LDA-mediated cyclization (eq 8). Nonetheless, addition of picoline **1** to a nearly LDA-free solution of **6** (prepared in situ) shows very little loss of starting material and <20% yield of azaindole **2** (eq 9). Amidinolithium **6** does not react with the relatively unhindered 2-fluoropyridine at $-40\text{ }^\circ\text{C}$. Therefore, adduct **20** is not likely to be an intermediate en route to **2**. Alternatively, the cyclization shown in eq 10³⁸ and nucleophilic additions of alkylolithiums to imidolithiums of general structure $\text{ArN}=\text{C}(\text{OLi})\text{R}$ ³⁹ suggest that **6** could, in theory, be the key electrophile. Nonetheless, **6** certainly does not have the trappings of a good electrophile.



(7)




(8)



no excess LDA: 10-20% yield
 1.0 equiv excess LDA: 82% yield

(9)



(10)

Chichibabin Cyclization Using 3-Picoline

To ascertain the role of the 2-fluoro moiety in **1** we briefly investigated the metalation of 3-picolone (**21**). As described by Wakefield and coworkers,³ treatment of **21** with 3.3 equiv of LDA at 0 °C followed by addition of PhCN with subsequent warming to 40 °C affords azaindole **2** in 90% yield (Scheme 5). Cursory rate studies using IR spectroscopy following the loss of **21** (1575 cm⁻¹) show that **21** is at least 20-fold less reactive than **1**, lithiating smoothly only

on warming to 10 °C. ¹H NMR spectroscopic analysis was thwarted by precipitation. Metalation of **21** in the absence of PhCN affords **22** as a bright yellow solid that could be isolated, but resisted spectroscopic characterization due to insolubility in THF.⁴⁰ Quenching the yellow solid with THF-*d*₈/H₂O (40:1) afforded dihydropyridine **23** in 20% spectroscopic purity that was surprisingly stable to further hydrolysis. Alternatively, treatment of the yellow solid suspended in THF with O₂ followed by standard aqueous workup afforded aromatized dimer **24** (70% yield) previously detected by Goto and coworkers.⁴¹

Discussion

In the context of studies of 7-azaindoles and related bicyclic pyrrole-based heterocycles being carried out at sanofi-aventis,^{2c} we investigated the Chichibabin cyclization illustrated in Scheme 1. Although the design is a straightforward adaptation of a protocol used by Wakefield and coworkers,³ the underlying organolithium chemistry is peculiar, and the potential ramifications are consequential. The results are summarized in the context of Scheme 2 and Scheme 6 as follows.

Summary

Sequential treatment of fluoropicoline **1** with LDA (2.0 equiv) and PhCN affords azaindole **2** in excellent (80–85%) yield. The order of mixing has no effect on the yield as long as the reaction is kept at ≤−40 °C. Products resulting from the condensation of benzyllithium **3** with fluoropicoline **1** offer a glimpse of underlying complexity (Scheme 2). Quenching solutions of LDA and **1** with H₂O affords transiently stable (but fully characterized spectroscopically) dihydropyridine **7**, which is converted to lactams **8a,b** and traces of air oxidized dimer **9** on workup. Warming prior to quenching affords net 1,2-adduct **10** in high yield. Monitoring the reaction by ¹⁹F NMR spectroscopy reveals lithiated dihydropyridine **5** in >95% purity to the exclusion of benzyllithium **3**. Rate studies show that **5** forms by a rate limiting metalation of **1** via highly solvated LDA monomer (see **12**). Adding PhCN to solutions of **5** shows no fluorine-containing intermediates. The extruded 0.5 equiv of **1** reenters the reaction coordinate, as required by the high yields of **2**.


Concurrent with the dimerization of picoline **1**, LDA undergoes reversible condensation with PhCN to form amidinolithium **6**, which exists as hetero- and homodimers **18** and **19**. Rate studies reveal an LDA dimer-based mechanism via **16** or **17**.¹⁵ If adduct **6** stands for several hours or is warmed above −40 °C, further oligomerization occurs, eventually affording trimer **15** (Scheme 4).^{31,32} Provided the cascade leading to **15** is not allowed to occur, formation of amidinolithium **6** does *not* preclude formation of azaindole **2**.

The peculiarity of the Chichibabin cyclization is underscored by an inverse addition in which LDA and PhCN are mixed prior to addition of picoline **1** (eq 1): *None of the original reagents are present in the vessel*. Yet, somehow **5** and **6** are readily converted to azaindole **2**. The simplest explanation is that **5** and **6** form reversibly. The conversion of 1,4-adduct **5** to 1,2-adduct **10** certainly hints at reversible addition (although **10** could, in principle, arise from an accelerated 1,3-sigmatropic rearrangement.)⁴² Crossover studies show that **5** does indeed form reversibly at −40 °C—conditions of the Chichibabin cyclization.


Ascertaining how amidinolithium **6** re-enters the reaction coordinate proved challenging. Crossover studies showed that *i*-Pr₂N moiety of **6** exchanges with free LDA. However, the conclusion that **6** extrudes free PhCN is assailable. The conditions that elicit exchange of LDA with amidinolithium **6** are relatively forcing and the extent of crossover limited. Moreover, the LDA exchange could, at least in principle, proceed by a nucleophilic addition rather than by dissociation of PhCN. We considered the possibility that **6** acts as a nucleophile toward 2-fluoropyridino moieties (eq 8) or even as an electrophile, but found no support.

Consequences of Picoline Dimerization

Optimized yields for metalation and functionalization of the nitrogen-based heterocycles can be highly substrate and electrophile dependent.⁵ In the case of the Chichibabin cyclization in Scheme 1, self-condensation of picoline **1** to form **5** ties up 50% of the starting picoline in the form of a lithiated dihydropyridino moiety, which also leaves residual LDA free to form adduct **6**. Fortunately, neither of these chemical cul-de-sacs proves fatal. We got lucky. If one had used a different electrophile, however, the formation of **5** might not be benign. As a simple example, using H_3O^+ (or D_3O^+) as an electrophile affords dihydropyridine **7**. If the goal had been to deuterate the benzylic position of **1**, formation of **5** causes total failure.⁴³ The dichotomous behavior of PhCN and D_2O may be important to those interested in metalating picolines, pyridines, and related nitrogen-based heteroaromatics.⁵ One wonders, for example, how many reactions relying on ortho or benzylic metalations of aromatic heterocycles provide poor yields or completely fail because of self-condensations. There are anecdotal reports of 1,2- and 1,4-adducts as byproducts in metalations^{44–51} such as in pyridines (eq 11)^{48,49} and pyrimidines (eq 12).⁵¹ We do not know how many dimer-derived byproducts are buried in the enormous literature of heterocycles beyond our gaze. There may be many unpublished examples in pharmaceutical research labs and many more that have simply gone undetected.



(11)



(12)

We do not fully understand the consequences of dimerization. The self-condensation of 3-picoline (**21**, Scheme 5) is a particularly disturbing case in point. Wakefield³ and others⁵² obtained excellent results metalating and functionalizing **21**. It seems unlikely that they suspected the intermediacy of 1,4-adduct **22**. We believe that some attempts to functionalize **21**—there may very well be many unreported—have been frustrated by either the failure of electrophiles to react with **22** or by the incompatibility of the electrophiles with excess base lurking in solution.⁵³ One group attributes such a failure to the high pK_A of **21**⁵⁴ whereas another group reports the pK_A of **21** without realizing that **21** had dimerized.⁵⁵ Are picoline dimers **5** and **22** as well as other putative dimers such as **26** and **27** problematic side products, or are they key intermediates en route to desired functionalizations^{49,56}

Conclusions

Nearly half of the top twenty largest selling pharmaceutical agents contain a pyridine or pyridine-like ring.⁵⁷ One of us (DBC) once spent an entire day at a major pharmaceutical company discussing exclusively a number of problematic pyridine metalations. There is little doubt that the metalations and subsequent functionalizations of pyridines present important and confounding challenges.

Despite self condensation of **1** to form **5** and consequent condensation of leftover LDA with PhCN to form amidinolithium **6**, the Chichibabin cyclization in eq 1 works remarkably well. A colleague astutely noted that **5** is, in essence, a protected benzylolithium, in which half of the starting picoline **1** is consumed as the protecting group. Thus, although one might view base-mediated self condensations of heteroaromatics as annoying side reactions to be avoided, maybe they should be embraced and exploited.

Experimental Section

Reagents and Solvents

THF, hexane, and pentane were distilled from blue or purple solutions containing sodium benzophenone ketyl. The hydrocarbon stills contained 1% tetraglyme to dissolve the ketyl. LDA, [⁶Li]LDA, and [⁶Li,¹⁵N]LDA³⁵ were purified as crystalline solids using recrystallized *n*-BuLi.⁵⁸ Air- and moisture sensitive materials were manipulated under argon or nitrogen using standard glove box, vacuum line, and syringe techniques. Solutions of *n*-BuLi and LDA were titrated using a literature method.⁵⁹

NMR Spectroscopic Analyses

Samples for monitoring reactions using NMR spectroscopy were prepared by layering stock solutions at $-196\text{ }^{\circ}\text{C}$, sealed under partial vacuum, and warmed to $-78\text{ }^{\circ}\text{C}$. The samples were vigorously shaken in a $-78\text{ }^{\circ}\text{C}$ bath for about 30 s and then transferred to the NMR probe for acquisition of spectra. Standard ⁶Li, ¹³C, ¹⁵N, and ¹⁹F NMR spectra were recorded on a 500 MHz spectrometer at 76.73, 125.79, 50.66 and 470.35 MHz (respectively). The ⁶Li, ¹³C, ¹⁵N, and ¹⁹F resonances are referenced to 0.30 M [⁶Li]LiCl/MeOH at $-90\text{ }^{\circ}\text{C}$ (δ 0.0 ppm), the CH₂O resonance of THF at $-90\text{ }^{\circ}\text{C}$ (δ 67.57 ppm), neat Me₂NEt at $-90\text{ }^{\circ}\text{C}$ (δ 25.7 ppm), and C₆F₆ in neat THF at $-90\text{ }^{\circ}\text{C}$ (δ -162.5 ppm), respectively.

In situ preparation of dimer 5

A 1.6 M solution of *n*-butyllithium (1.33 mL, 2.1 mmol) in hexanes was added via syringe to dry THF (20.0 mL) at $-40\text{ }^{\circ}\text{C}$ under inert Ar atmosphere. Dry diisopropylamine (310 μL , 224 mg, 2.1 mmol) was added to the solution via syringe. After stirring for 5 min at $-40\text{ }^{\circ}\text{C}$, 3-methyl-2-fluoropyridine **1** (200 μL , 220 mg, 2.0 mmol) was added to the LDA solution. The solution became a bright red as *N*-lithiodihydropyridine **5** forms over the course of 1.0 h. Carrying out an analogous procedure in an NMR tube using THF-*d*₈ affords the following spectroscopic data: ¹H NMR (THF-*d*₈, $-78\text{ }^{\circ}\text{C}$) δ 1.60 (d, $J_{\text{H,F}}=1.4\text{ Hz}$, 3H), 2.43 (dd, $J_{\text{H,H}}=12.4, 7.2\text{ Hz}$, 1H), 2.77 (dd, $J_{\text{H,H}}=12.4, 3.5\text{ Hz}$, 1H), 3.50 (ddd, $J_{\text{H,F}}=3.7, J_{\text{H,H}}=6.7, 3.9\text{ Hz}$, 1H), 3.69 (dddd, $J_{\text{H,F}}=6.8, J_{\text{H,H}}=7.2, 3.4, 3.4\text{ Hz}$, 1H), 5.84 (dd, $J_{\text{H,F}}=2.0, J_{\text{H,H}}=6.7\text{ Hz}$, 1H), 7.15 (dddd, $J_{\text{H,F}}=2.1, J_{\text{H,H}}=7.0, 4.7, 0.4\text{ Hz}$, 1H), 7.63 (ddd, $J_{\text{H,F}}=9.5, J_{\text{H,H}}=7.0, 2.1\text{ Hz}$, 1H), 7.96 (ddm, $J_{\text{H,H}}=4.7, 2.1\text{ Hz}$, 1H); ¹³C NMR (THF-*d*₈, $-78\text{ }^{\circ}\text{C}$) δ 20.1 (d, $J_{\text{C,F}}=22.3\text{ Hz}$), 47.0 (d, $J_{\text{C,F}}=6.6\text{ Hz}$), 69.7 (d, $J_{\text{C,F}}=29.0\text{ Hz}$), 80.1 (d, $J_{\text{C,F}}=7.0\text{ Hz}$), 91.4, 121.6, 123.4 (d, $J_{\text{C,F}}=31.7\text{ Hz}$), 139.3 (d, $J_{\text{C,F}}=21.8\text{ Hz}$), 143.4 (d, $J_{\text{C,F}}=6.6\text{ Hz}$), 144.7 (d, $J_{\text{C,F}}=15.0\text{ Hz}$), 163.0 (d, $J_{\text{C,F}}=215.0\text{ Hz}$), 163.3 (d, $J_{\text{C,F}}=235.0\text{ Hz}$); ¹⁹F NMR (THF-*d*₈, $-78\text{ }^{\circ}\text{C}$) δ -93.5, -70.5; ⁶Li NMR (THF-*d*₈, $-78\text{ }^{\circ}\text{C}$) δ -0.01, 0.76.

Synthesis of Dihydropyridine 7

A solution of *N*-lithiodihydropyridine **5** (0.05 M) at $-40\text{ }^{\circ}\text{C}$ prepared in THF- d_8 as described above was quenched by THF- d_8 -H₂O at $-78\text{ }^{\circ}\text{C}$ to afford dihydropyridine **7**. The N-H proton is not observable at $0\text{ }^{\circ}\text{C}$; spectra were recorded on **7** in situ at $-70\text{ }^{\circ}\text{C}$: ¹H NMR (THF- d_8) δ 1.61 (d, $J=2.3\text{ Hz}$, 3H), 2.54 (m, 1H), 2.87 (m, 1H), 3.30 (m, 1H), 4.06 (dd, $J=8.0, 2.3\text{ Hz}$, 1H), 5.87 (dt, $J=8.0, 2.0\text{ Hz}$, 1H), 7.21 (tt, $J=7.0, 1.7\text{ Hz}$, 1H), 7.40 (br, 1H), 7.67 (dt, $J=7.0, 2.0\text{ Hz}$, 1H), 8.03 (m, 1H); ¹³C NMR (THF- d_8) δ 11.5, 35.2, 41.5 (dd, $J=4.0, 1.9\text{ Hz}$), 80.7, 98.3, 121.8, 122.0, 126.2, 142.8, 145.3, 149.6, 162.8.

Synthesis of Azaindole 2

To a solution of *N*-lithiodihydropyridine **5** (0.05 M) at $-40\text{ }^{\circ}\text{C}$ prepared as described above was added benzonitrile (250 μL , 253 mg, 2.45 mmol). After stirring at $-40\text{ }^{\circ}\text{C}$ for 2 h the vessel was warmed to $0\text{ }^{\circ}\text{C}$ for 30 min, the reaction was quenched with wet THF. The solvent was evaporated under reduced pressure. The resulting yellow solid was redissolved in EtOAc (15 mL), and the organic layer was washed with aqueous NaHCO₃ (3 \times 10 mL) and aqueous NaCl (3 \times 10 mL). The organic layer was dried with Na₂SO₄, filtered, and evaporated to dryness. The bright yellow solid was recrystallized from 1:4 ethyl acetate:hexane or, alternatively, flash chromatographed eluting with 2:1 ethyl acetate:hexanes. Compound **2** (310 mg, 1.60 mmol) was isolated as an off-white solid in 80% yield. ¹H NMR (DMSO- d_6) δ 6.90 (s, 1H), 7.05 (dd, $J=7.8, 4.7\text{ Hz}$, 1H), 7.33 (tt, $J=7.3, 1.2\text{ Hz}$, 1H), 7.45 (t, $J=7.8\text{ Hz}$, 2H), 7.91 (dd, $J=7.9, 1.5\text{ Hz}$, 1H), 7.96 (dd, $J=7.8, 1.2\text{ Hz}$, 2H), 8.24 (dd, $J=1.6, 4.7\text{ Hz}$, 1H), 12.23 (s, 1H); ¹³C NMR (DMSO- d_6) δ 97.1, 116.0, 121.0, 125.4, 127.8, 128.0, 128.9, 131.7, 138.3, 142.9, 149.8. HRMS [C₁₃H₁₀N₂] requires m/z 194.0844. Found: 194.0839.

Synthesis of Azaindole 2: Inverse Addition

A 1.6 M solution of *n*-butyllithium (2.66 mL, 4.2 mmol) in hexanes was added via syringe to dry THF (20.0 mL) at $-40\text{ }^{\circ}\text{C}$ under inert Ar atmosphere. Dry diisopropylamine (620 μL , 448 mg, 4.2 mmol) was added to the solution via syringe. After stirring for 5 min at $-40\text{ }^{\circ}\text{C}$, benzonitrile (215 μL , 217 mg, 2.1 mmol) was added. After stirring at $-40\text{ }^{\circ}\text{C}$ for 2 h, and fluoropicoline **1** (200 μL , 220 mg, 2.0 mmol) was added and stirring was continued for 2 more hours. Workup and purification as described above afford **2** (320 mg, 1.65 mmol) as an off-white solid in 82% yield.

Synthesis of 8a and 8b

A solution of *N*-lithiodihydropyridine **5** (0.05 M) at $-40\text{ }^{\circ}\text{C}$ prepared as described above was cooled to $-78\text{ }^{\circ}\text{C}$ and quenched with THF/water (40:1, 10.3 mL) dropwise via syringe. After warming to rt, the solvent was evaporated in vacuo. The resulting residue was redissolved in EtOAc (15 mL), and the organic layer was washed with aqueous NaHCO₃ (3 \times 10 mL) and aqueous NaCl (3 \times 10 mL). The organic extract was dried with Na₂SO₄, filtered, and evaporated to dryness. Flash chromatography (2:1 ethyl acetate:hexanes) afforded lactams **8a,b** as a colorless liquid (196 mg, 0.89 mmol) in 89% yield. HPLC separation (degassed 5:1 pentane:diethyl ether) afforded analytically pure **8a** and **8b** for spectroscopic analysis. **8a**: ¹H NMR (acetone- d_6) δ 1.20 (d, $J=7.2\text{ Hz}$, 3H), 2.49 (dd, $J=13.0, 10.4\text{ Hz}$, 1H), 2.63 (d, $J=6.7\text{ Hz}$, 1H), 2.75–2.81 (d, $J=5.1\text{ Hz}$, 1H), 2.82–2.86 (m, 1H), 2.86 (br s, 1H), 4.80 (dd, $J=8.0, 4.9\text{ Hz}$, 1H), 6.18 (dd, $J=7.7, 4.4\text{ Hz}$, 1H), 7.25 (tdd, $J=6.1, 2.1, 1.8\text{ Hz}$, 1H), 7.77–7.84 (m, $J=8.8, 2.4, 1.7, 1\text{ Hz}$), 8.05–8.09 (m, $J=5.2, 1\text{ Hz}$); ¹³C NMR (acetone- d_6) δ 11.5, 28.9 ($J=3.1$); 37.5 ($J=1.1$), 40.1, 107.5, 122.6 ($J=4.3$), 122.8 ($J=31.1$), 126.6, 143 ($J=5.6\text{ Hz}$), 146.33 ($J=15.4\text{ Hz}$), 163 ($J=235.4\text{ Hz}$), 173.27; ¹⁹F NMR (acetone- d_6) δ -74.5 (d, $J=9.4$); HRMS [C₁₂H₁₃N₂OF] requires m/z 220.1012. Found: 220.1003; IR (film, cm⁻¹): 1698, 1661, 1605, 1578, 1438. **8b**: ¹H NMR (acetone- d_6) δ 1.19 (d, $J=7.2\text{ Hz}$, 3H), 2.30 (d, $J=7.0\text{ Hz}$, 1H), 2.45–2.52 (m, 1H), 2.64–2.71 (m, 1H), 2.76–2.82 (m, 1H), 2.84 (br s, 1H), 4.84 (dd, $J=7.8, 4.8\text{ Hz}$,

1H), 6.18 (dd, $J=7.7, 4.4$ Hz, 1H), 7.25 (tdd, $J=6.1, 2.1, 1.8$ Hz, 1H), 7.77–7.84 (m, 1H), 8.05–8.09 (d, $J=5.2$ Hz, 1H); ^{13}C NMR (acetone- d_6) δ 11.5, 28.7 ($J=3.1$), 37.5 ($J=1.1$ Hz), 40.1, 107.4, 122.6 ($J=4.3$ Hz), 122.8 ($J=31.1$ Hz), 126.5, 142.9 ($J=5.6$ Hz), 146.33 ($J=15.4$ Hz), 162.97 ($J=235.4$ Hz), 173.3; ^{19}F NMR (acetone- d_6) δ -73.9 (d, $J=9.1$ Hz); IR (film, cm^{-1}): 1698, 1661, 1605, 1578, 1438. HRMS [$\text{C}_{12}\text{H}_{13}\text{N}_2\text{OF}$] requires m/z 220.1012. Found: 220.1028.

Synthesis of 9

A solution of *N*-lithiodihydropyridine **5** (0.05 M) at -40 °C prepared as described above was cooled to -78 °C and purged with gaseous O_2 . After stirring at -78 °C for 4 h, the reaction was quenched with a THF/water mixture (40:1, 10.3 mL). Water (250 μL , 250 mg) in THF (10 mL) was added to the reaction dropwise via syringe. After warming to rt, the solvent was evaporated in vacuo. The resulting residue was redissolved in EtOAc (15 mL), and the organic layer was washed with aqueous NaHCO_3 (3×10 mL) and aqueous NaCl (3×10 mL). The organic extract was dried with Na_2SO_4 , filtered, and evaporated to dryness. Flash chromatography (2:1 ethyl acetate:hexanes) afforded **9** as a white solid (90 mg, 0.41 mmol) in 41% yield. mp 78.5–80.5 °C; ^1H NMR (acetone- d_6) δ 2.24 (s, 3H), 4.14 (s, 2H), 7.01 (d, $J=5.1$ Hz, 1H), 7.29 (dt, $J=5.5, 2.0, 1.8$ Hz, 1H), 7.71 (dt, $J=9.8, 2.0, 1.2$ Hz, 1H), 7.97 (d, $J=5.1$ Hz, 1H), 8.14 (d, $J=5.5$ Hz, 1H); ^{13}C NMR (acetone- d_6) δ 10.04, 31.20 (dd, $J=4.0, 1.9$ Hz), 118.4 (d, $J=32.8$ Hz), 120.7 (d, $J=32.6$ Hz), 122.7 (d, $J=4.2$ Hz), 123.5 (d, $J=4.2$ Hz), 141.8 (d, $J=5.6$ Hz), 144.6 (d, $J=15.9$ Hz), 146.3 (d, $J=15.3$ Hz), 151.5 (d, $J=5.0$ Hz), 161.4 (d, $J=96.4$ Hz), 162.7 (d, $J=101.8$ Hz); ^{19}F NMR (acetone- d_6) δ -69.59 ($J=9.1$), -71.59; HRMS [$\text{C}_{12}\text{H}_{10}\text{N}_2\text{F}_2$] requires m/z 220.0812. Found: 220.0819. An x-ray crystal structure of **9** is reported in supporting information.

Synthesis of 10

A solution of *N*-lithiodihydropyridine **5** (0.05 M) at -40 °C prepared as described above was warmed to 0 °C in a water/ice bath. Stirring was continued for 15 min. The reaction was quenched with a THF/water mixture (40:1, 10.3 mL). After warming to rt, the solvent was evaporated in vacuo. The resulting residue was redissolved in EtOAc (15 mL), and the organic layer was washed with aqueous NaHCO_3 (3×10 mL) and aqueous NaCl (3×10 mL). The organic extract was dried with Na_2SO_4 , filtered, and evaporated to dryness. Flash chromatography (2:1 ethyl acetate:hexanes) afforded 1,2-adduct **10** as a colorless liquid (178 mg, 0.88 mmol) in 88% yield. ^1H NMR (acetone- d_6) δ 2.34 (s, 3H), 4.16 (s, 2H), 7.14 (dd, $J=7.6, 4.9$ Hz, 1H), 7.22 (dt, $J=6.1, 2.0, 0.9$ Hz, 1H), 7.56 (d, $J=7.6$ Hz, 1H), 7.66 (dt, $J=8.6, 2.4, 1.8$ Hz, 1H), 8.05–8.09 (m, 1H), 8.28 (dd, $J=4.91, 1.2$ Hz, 1H); ^{13}C NMR (acetone- d_6) δ 18.7 (d, $J=1.9$ Hz), 34.8 (d, $J=1.9$ Hz), 122.3, 122.7 (d, $J=3.2$ Hz), 122.8 (d, $J=3.8$ Hz), 132.3, 138.5, 142.6 (d, $J=5.8$ Hz), 146.2 (d, $J=15.6$ Hz), 147.6 (d, $J=1.5$ Hz), 157.6, 162.9 (d, $J=236.2$ Hz); ^{19}F NMR (acetone- d_6) δ -70.2 (d, $J=8.6$); HRMS [$\text{C}_{12}\text{H}_{11}\text{N}_2\text{F}$] requires m/z 202.0906. Found: 202.0906.

Synthesis of Trimer 15

A 1.6 M solution of *n*-butyllithium (1.25 mL, 2.0 mmol) in hexanes was added via syringe to dry THF (10.0 mL) at -40 °C under inert Ar atmosphere. Dry diisopropylamine (280 μL , 202 mg, 2.0 mmol) was added to the solution via syringe. After stirring for 5 min at -40 °C, PhCN (408 μL , 512 mg, 4.0 mmol) was added to the LDA solution. The solution became bright yellow as amidinolithium **6** formed over the course of 15 min. After stirring for 4 h, quenching with a THF/water mixture (40:1, 10.3 mL) dropwise via syringe, and warming to rt, the solvent was evaporated in vacuo. The resulting residue was redissolved in copious amounts of EtOAc (100 mL) due to limited solubility, and the organic layer was washed with aqueous NaHCO_3 (3×10 mL) and aqueous NaCl (3×10 mL). The organic extract was dried with Na_2SO_4 , filtered, and

evaporated to dryness. The off-white solid was recrystallized from neat ethyl acetate. Known³¹ trimer **15** (340 mg, 1.10 mmol) was isolated as a white solid in 83% yield. ¹H NMR (DMSO-*d*₆) δ, 8.74–8.78 (6H, m).

Preparation of Lithiated Picoline Dimer **22**

A solution of 3-picoline (**21**, 1.50 mL, 1.44 g, 15.5 mmol) in 3.0 mL THF was added via syringe to LDA (800 mg, 7.5 mmol) in THF (5.0 mL) and hexanes (60 mL) at –78 °C under Ar. Warming to 0 °C and stirring for 1.0 h resulted in a bright yellow precipitate. The suspension was cooled by dry ice/acetone for 1 hr, and the solvent was removed via syringe. The yellow solid was washed by THF/hexanes (1:10) and vacuumed for 3 h to afford a yellow solid (yield 87%). The resulting stable, bright yellow solid was too insoluble for spectroscopic analysis but was characterized by quenching to form **23** as described below.

Preparation of Picoline Dimer **23**

An NMR tube charged with lithiated 3-dimer **22** as a yellow solid (9.6 mg, 0.05 mmol), cooled to –78 °C, and treated with THF-*d*₈/H₂O (40:1) via syringe. The yellow solid became a colorless solution containing dihydropyridine **23** in greater than 90% spectroscopic purity: ¹H NMR (THF-*d*₈) δ 1.59 (s), 2.54 (ddm, *J*=13.1, 7.4 Hz, 1H), 2.74 (ddm, *J*=13.1, 4.2 Hz, 1H), 3.19 (m, 1H), 3.97 (ddd, *J*=7.7, 4.2, 1.5 Hz, 1H), 5.76 (ddd, *J*=4.7, 1.5, 1.4 Hz, 1H); 5.90 (ddd, *J*=7.7, 4.7, 1.2 Hz, 1H), 6.18 (br s 1H), 7.17 (ddd, *J*=7.7, 4.8, 0.9 Hz, 1H), 7.49 (m, 1H), 8.32 (m, 2H); ¹³C NMR (THF-*d*₈) δ 18.5, 39.7, 40.8, 96.3, 123.2, 123.8, 127.5, 35.4, 137.5, 146.9, 150.5.

1,4-Adduct **5**: Crossover Studies

A solution of LDA (0.275 M, 100 μL, 0.030 mmol) in neat THF was added to one side of a double-well vessel (Figure 1) under inert Ar atmosphere at –40 °C. A solution of **1-d**₃ (0.250 M, 100 μL, 0.025 mmol) in THF was injected into the reaction vessel and allowed to stir at –40 °C for 24 h. To the other well of the vessel was added an aliquot of LDA solution (0.275 M, 100 μL, 0.030 mmol) followed by a solution of **1-d**₀ (0.250 M, 100 μL, 0.025 mmol) in THF. After 1.5 h, the reaction vessel was tilted in the bath to fully mix both wells. The reaction was allowed to stir for the desired amount of time and quenched with a THF/water mixture (40:1, 500 μL). The contents of the vessel were transferred to a 1.5 mL GC vial and analyzed via GC/MS.

Amidinolithium **6**: Crossover Studies

A solution of [⁶Li, ¹⁵N]LDA (0.30 M, 400 μL, 0.12 mmol) in neat THF was added to an NMR tube at –78 °C under Ar atmosphere capped by two septa. To the NMR tube was added benzonitrile solution (1.56 M, 100 μL, 0.16 mmol) in THF. The reaction was allowed to stand at –78 °C for 1 hour. Once conversion to **6** was complete as indicated by ⁶Li NMR, a solution of [⁶Li]LDA (1.20 M, 100 μL, 0.12 mmol) was added and warmed to –40 °C. Crossover was allowed to occur over the course of 12 h with periodic cooling to –90 °C to acquire spectra.

Kinetics of 3-Fluoropicoline Metalation

IR spectra were recorded using an in situ IR spectrometer fitted with a 30-bounce, silicon-tipped probe. The spectra were acquired in 16 scans at a gain of 1 and a resolution of 8 cm^{–1}. A representative reaction was carried using the reaction of LDA and PhCN emblematically as follows: The IR probe was inserted through a nylon adapter and FETFE O-ring seal into an oven-dried, cylindrical flask fitted with a magnetic stir bar and a T-joint. The T-joint was capped by a septum for injections and an argon line. Pseudo-first-order conditions were established by maintaining low concentrations of **1** (0.005 M) at high, yet adjustable, concentrations of LDA (0.10–0.75 M) and THF (3.9–12.3 M)⁶⁰ with pentane as the cosolvent.

It was most convenient to follow the formation of **5** (1667 cm^{-1}) by IR spectroscopy, which displays a first-order behavior to ≥ 5 half-lives.^{30,61} No evidence of LDA-picoline complexation was noted. The resulting pseudo-first-order rate constants (k_{obsd}) are independent of picoline concentration (0.004 – 0.04 M), confirming the first-order substrate dependence. Re-establishing the IR baseline and monitoring a second injection reveals no significant change in the rate constant, showing that conversion-dependent autocatalysis or autoinhibition are unimportant under these conditions. Comparisons of **1** versus **1-d₃**, provided $k_{\text{H}}/k_{\text{D}} = 29 \pm 1$, confirming a rate-limiting proton transfer. Unusually large isotope effects have been observed for other lithiations.⁶² Plots of k_{obsd} versus THF concentration and k_{obsd} versus LDA concentration (at 4.1 M THF affords the rate law described by eq 3. Data are included in supporting information.

Kinetics of LDA-PhCN Condensation

Pseudo-first-order conditions were established by maintaining the concentration of PhCN low (0.01 M) and the concentrations of LDA (0.10–0.30 M) and THF (2.0–12.3 M)⁶⁰ high (yet adjustable) with pentane as the cosolvent. The loss of PhCN (1667 cm^{-1}) by IR spectroscopy display a first-order behavior to ≥ 5 half-lives.^{30,61} No evidence of LDA-PhCN complexation was noted. Plots of k_{obsd} versus LDA concentration (at 10.0 M THF) and k_{obsd} versus THF concentration (supporting information) afford the reaction orders indicated in eq 5.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The Cornell group thanks the National Science Foundation and sanofi-aventis for support.

References and Footnotes

1. (a) Popowycz F, Routier S, Joseph B, Merour J-Y. *Tetrahedron* 2007;63:1031. (b) Fang Y-Q, Yuen J, Lautens M. *J. Org. Chem* 2007;72:5152. [PubMed: 17559273]
2. (a) Merour J-Y, Joseph B. *Current Org. Chem* 2001;5:471. (b) Song JJ, Reeves JT, Gallou F, Tan Z, Yee N, Senanayake CH. *Chem. Soc. Rev* 2007;36:1120. [PubMed: 17576479] (c) Hopkins CR, Collar N. *Tetrahedron Lett* 2004;45:8087.
3. Davis ML, Wakefield BJ, Wardell JA. *Tetrahedron* 1992;48:939.
4. (a) Vorbrugenn H. *Adv. Heterocycl. Chem* 1990;49:117. (b) McGill CK, Rappa A. *Adv. Heterocycl. Chem* 1988;44:1.
5. (a) Queguiner G, Marsais F, Snieckus, Epszajn J. *Adv. Heterocycl. Chem* 1991;52:187. (b) Mongin F, Queguiner G. *Tetrahedron* 2001;57:5897. (c) Mongin F, Queguiner G. *Tetrahedron* 2001;57:4059. (d) Merino P. *Progr Heterocycl. Chem* 1999;11:21. Clayden, J. *The Chemistry of Organolithium Compounds*. Rappoport, Z.; Marek, I., editors. Vol. 1. New York: Wiley; 2004. p. 495. Caubere, P. *Reviews of Heteroatom Chemistry*. Vol. 4. Tokyo: MYU; 1991. p. 78-139. (g) Caubere P. *Chem. Rev* 1993;93:2317. (h) Marsais F, Queguiner G. *Tetrahedron* 1983;39:2009. (i) Collins I. *Perkin* 1 2000:2845. (j) Schlosser M, Mongin F. *Chem. Soc. Rev* 2007;36:1161. [PubMed: 17576483]
6. For a review of structural studies using ^{19}F NMR spectroscopy, see: (a) Gakh YG, Gakh AA, Gronenborn AM. *Magn. Reson. Chem* 2000;38:551. (b) McGill CA, Nordon A, Littlejohn D. *J. Process Analyt. Chem* 2001;6:36.
7. Scheme 2 does not represent a comprehensive profile of byproducts.
8. Stout DM, Meyers AI. *Chem. Rev* 1982;82:223.
9. Nanney JR, Mahaffy CAL. *J. Fluorine Chem* 1994;68:181.

10. Treating **5** with MnO₂ prior to the aqueous quench affords **9** in 50%. For a related MnO₂ rearomatization, see: (a) Godard A, Jacquelin JM, Queguiner G. *J. Organomet. Chem* 1988;354:273. (b) Eynde JJ, Delfosse F, Mayence A, Haverbeke V. *Tetrahedron* 1995;51:6511.
11. For the 1,2-addition of an aryllithium to **1**, see: DuPriest MT, Schmidt CL, Kuzmich D, Williams SB. *J. Org. Chem* 1986;51:2021.
12. Dimer **9** was also characterized by x-ray crystallography. The product of oxidative coupling (**i**) was also characterized as a rogue crystal by x-ray crystallography. We were unable to intentionally produce detectable **i** by introducing oxygen to an LDA/**1** mixture.
13. We find fluoropyridines to be sensitive to traces of acid, including the residual HCl in chlorinated solvents.
14. Galiano-Roth AS, Collum DB. *J. Am. Chem. Soc* 1989;111:6772.
15. (a) Collum DB. *Acc. Chem. Res* 1993;26:227. (b) Schleyer, PvR; Snaith, R. *Adv. Inorganic. Chem* 1991;37:47. (c) Mulvey RE. *Chem. Soc. Rev* 1991;20:167.
16. Traces (approx 5%) of **13** (δ -77.9 ppm) are also observed.
17. Lithiation of an isolated sample of **9** with LDA affords new ¹⁹F resonances at δ -78.9 and -75.0 ppm that we attribute to **ii**. These resonances were *not* observed in the reaction mixture.
18. For a review of rate studies of LDA-mediated reactions, see: Collum DB, McNeil AJ, Ramirez A. *Angew. Chem., Int. Ed* 2007;46:3002.
19. We define the idealized rate law as that obtained by rounding the observed reaction orders to the nearest rational order.
20. Ma Y, Collum DB. *J. Am. Chem. Soc* 2007;129:14818. [PubMed: 17985891] and references cited therein.
21. For leading references and discussion of high-coordinate lithium amides, see ref 20.
22. (a) van Eikema Hommes NJR, Schleyer PvR. *Angew. Chem., Int. Ed. Engl* 1992;31:755. (b) Stalke D, Klingebiel U, Sheldrick GM. *Chem. Ber* 1988;121:1457. (c) Armstrong DR, Khandelwal AH, Kerr LC, Peasey S, Raithby PR, Shields GP, Snaith R, Wright DS. *Chem. Commun* 1998:1011. (d) Plenio H, Diodone R. *J. Am. Chem. Soc* 1996;118:356. (e) Henderson KW, Dorigo AE, Liu Q-Y, Williard PG. *J. Am. Chem. Soc* 1997;119:11855. (f) Chadwick ST, Rennels RA, Rutherford JL, Collum DB. *J. Am. Chem. Soc* 2000;122:8640. (g) Chadwick ST, Ramirez A, Gupta L, Collum DB. *J. Am. Chem. Soc* 2007;129:2259. [PubMed: 17269777] (h) Williard PG, Liu Q-Y. *J. Org. Chem* 1994;59:1596.
23. Carpenter, BK. *Determination of Organic Reaction Mechanisms*. New York: Wiley; 1984.
24. A deviation from statistical stems from difficulty in obtaining precisely equal proportions of **5-d₀** and **5-d₅** due to the especially large kinetic isotope effect on the metalation.
25. It has been suggested that the *n*-BuLi adduct of pyridine can reduce pyridine by extruding lithium hydride: Clegg W, Dunbar L, Horsburgh L, Mulvey RE. *Angew. Chem., Int. Ed. Engl* 1996;35:753.
26. (a) The fit to $[A] = [A_0]\exp(k_{\text{obsd}}t)$ showed some discrepancies (supporting information), yet a fit to a non-linear Noyes equation (ref 26b) to obtain the order by best-fit methods revealed an order of 1.0. Briggs TF, Winemiller MD, Collum DB, Parsons RL Jr, Davulcu AK, Harris GD, Fortunak JD, Confalone PN. *J. Am. Chem. Soc* 2004;126:6291.
27. k_{obsd} for the decay of **5** shows a minor (50%) rise with a three-fold decrease in THF concentration.
28. The crossover described in Scheme 3 occurs at substantially lower temperatures than the rearrangement to provide **10**. As expected, analogous crossover is observed in **10**.
29. We have never observed [⁶Li]LiF by NMR spectroscopy. This may derive from a very low solubility.
30. Rein AJ, Donahue SM, Pavlosky MA. *Curr. Opin. Drug Discovery Dev* 2000;3:734.
31. (a) Compagnon, P; Louis, G; Gasquez, F.; Kimny, T. *Bull. Soc. Chim. Belges* 1986;95:57. (b) Chen X, Bai S-D, Wang L, Liu D-S. *Heterocycles* 2005;65:1425.
32. (a) Davies RP, Raithby PR, Shields GP, Snaith R, Wheatley AEH. *Organometallics* 1997;16:2223. (b) Armstrong DR, Clegg W, MacGregor M, Mulvey RE, O'Neil PA. *J. Chem. Soc, Chem. Commun* 1993:608. (c) Armstrong DR, Henderson KW, MacGregor M, Mulvey RE, Ross MJ, Clegg W, O'Neil PA. *J. Organomet. Chem* 1995;486:79.
33. Crystal structures of amidinolithiums: Aharonovich S, Kapon M, Botoshanski M, Eisen MS. *Organometallics* 2008;27:1869.

34. For calculated open dimer-based additions of methyllithium to nitriles, see: Uchiyama M, Matsumoto Y, Usui S, Hashimoto Y, Morokuma K. *Angew. Chem., Int. Ed* 2007;46:926.
35. Romesberg FE, Gilchrist JH, Harrison AT, Fuller DJ, Collum DB. *J. Am. Chem. Soc* 1991;113:5751.
36. The nuclear spins of ^6Li and ^{15}N are 1 and 1/2, respectively.
37. (a) Armstrong DR, Barr D, Snaith R, Clegg W, Mulvey RE, Wade K, Reed D. *J. Chem. Soc., Dalton Trans* 1987:1071. (b) Barr D, Snaith R, Mulvey RE, Wade K, Reed D. *Mag. Reson. Chem* 1986;24:713. (c) Reed D, Barr D, Mulvey RE, Snaith R. *J. Chem. Soc., Dalton Trans* 1986:557. (d) Barr B, Clegg W, Mulvey RE, Snaith R, Wade K. *J. Chem. Soc., Chem. Commun* 1986:295.
38. (a) Turner JA. *J. Org. Chem* 1983;48:3401. (b) Hands D, Bishop B, Cameron M, Edwards JS, Cottrell IF, Wright SHB. *Synthesis* 1996:877. (c) Houlihan WJ, Parrino VA, Uike Y. *J. Org. Chem* 1981;46:4511.
39. Smith K, El-Hiti GA, Hamilton A. *J. Chem. Soc., Perkin Trans. 1* 1998;24:4041.
40. Dissolving the yellow solid with added *N,N,N',N'*-tetramethylethylenediamine (TMEDA) resulted in very complex mixtures. We suspect that TMEDA mediates reversal of the 1,4-addition, allowing other reactions to occur.
41. Tagawa Y, Hama K, Goto Y. *Heterocycles* 1992;34:1605.
42. (a) Zoeckler MT, Carpenter BK. *J. Am. Chem. Soc* 1981;103:7661. (b) Carpenter BK. *Tetrahedron* 1978;34:1877.
43. By contrast, the relatively weak acid hexamethyldisilazane converts dimer **5** to picoline **1**.
44. (a) Knauss E, Ondrus T, Giam C. *J. Heterocycl. Chem* 1976;13:789. (b) Thomas EW. *J. Org. Chem* 1986;51:2184. (c) Giam CA, Stout JL. *J. Chem. Soc., Dalton Trans* 1970:478. (d) Ludt RE, Crowther GP, Hauser CR. *J. Org. Chem* 1970;35:1288. (e) Francis RF, Crews CD, Scott BS. *J. Org. Chem* 1978;43:3227. (f) Marsais F, Granger P, Queguiner G. *J. Org. Chem* 1981;46:4494. (g) Gronowitz S, Roe J. *Acta Chem. Scand* 1965;19:1741. (h) Marsais F, Le Nard G, Queguiner G. *Synthesis* 1982:235. (i) Also, see ref 9.
45. (a) Abramovitch RA, Poulton GA. *J. Chem. Soc. (B)* 1969:901. (b) Abramovitch RA, Giam C-S, Poulton GA. *J. Chem. Soc. (C)* 1970:128. (c) Abramovitch RA, Giam C-S. *Can. J. Chem* 1964;42:1627. (d) Abramovitch RA, Giam C-S. *Can. J. Chem* 1963;41:3127. (e) Clegg W, Dunbar L, Horsburgh L, Mulvey RE. *Angew. Chem., Int. Ed. Engl* 1996;35:753.
46. (a) Abarca B, Hayles DJ, Jones G, Sliskovic DR. *J. Chem. Res., Synop* 1983:181. (b) Postigo A, Rossi RA. *Org. Lett* 2001;3:1197. [PubMed: 11348193] (c) Clayden J, Turnbull R, Pinto I. *Org. Lett* 2004;6:609. [PubMed: 14961635] (d) Overberger CG, Heron LP. *J. Org. Chem* 1962;27:2423. (e) Giam CS, Knaus EE. *Tetrahedron Lett* 1971:4961. (f) Gros P, Fort Y. *J. Chem. Soc., Perkin I* 1998:3515. (g) Zhang L, Tan Z. *Tetrahedron Lett* 2000;41:3025. (h) Kauffmann T. *Angew. Chem., Int. Ed. Engl* 1979;18:1. and references cited therein.
47. Mixed condensation of heteroaryllithiums with heteroarenes: Gros P, Fort Y. *J. Chem. Soc., Perkin I* 1998:3515. Kauffmann T. *Angew. Chem., Int. Ed. Engl* 1979;18:1. and references cited therein.
48. (a) Gros P, Fort Y, Caubere P. *J. Chem. Soc., Perkin I Trans* 1997:3597. (b) Gros P, Fort Y, Caubere P. *J. Chem. Soc., Perkin I Trans* 1997:3071. (c) Gros P, Fort Y. *J. Org. Chem* 2003;68:2028. [PubMed: 12608830] (d) Kaufmann T. *Angew. Chem., Int. Ed. Engl* 1979;18:1. (e) Martens RJ, Den Hertzog HJ, van Ammers M. *Tetrahedron Lett* 1964:3207. (f) Marais RJ, Breant P, Ginguene A, Queguiner G. *J. Organomet. Chem* 1981;216:139. (g) Newkome GR, Hager DC. *J. Org. Chem* 1982;47:599. (h) Clarke AJ, McNamara S, Meth-Cohn O. *Tetrahedron Lett* 1974:2373. (i) Wiessenfels M, Ulrici B. *Z. Chem* 1978;18:382. (j) Cottet F, Schlosser M. *Eur. J. Org. Chem* 2004:3793.
49. An alternative radical mechanism for the dimerization of pyridine has been proposed (ref 48g).
50. (a) Kress KJ. *J. Org. Chem* 1979;44:2081. (b) Plé N, Turck A, Couture K, Queguiner G. *J. Org. Chem* 1995;60:3781.
51. (a) Kaufmann T, Greving B, König J, Mitschker A, Wotermann A. *Angew. Chem., Int. Ed. Engl* 1975;14:713. (b) Seggio A, Chevalier F, Vaultier M, Mongin F. *J. Org. Chem* 2007;72:6602. [PubMed: 17655361]
52. Alkylations of 3-picoline proceed in high yield. See, for example: (a) Gedig T, Dettner K, Seifert K. *Tetrahedron* 2007;63:2670. (b) Baldwin JE, Melman A, Lee V, Firkin CR, Whitehead RC. *J. Am. Chem. Soc* 1998;120:8559. (c) Romeril SP, Lee V, Baldwin JE, Claridge TDW. *Tetrahedron* 2005;61:1127. (d) Perez-Balado C, Nebbioso A, Rodriguez-Grana P, Minichiello A, Miceli M,

- Altucci L, de Lera AR. *J. Med. Chem* 2007;50:2497. [PubMed: 17447750] (e) Davies-Coleman MT, Faulkner JD, Dubowchik GM, Roth GP, Polson C, Fairchild C. *J. Org. Chem* 1993;58:5925.
53. (a) Kaiser EM, Petty JD. *Synthesis* 1975:705. (b) Salas M, Joule JA. *J. Chem. Res., Synop* 1990:98. (c) Reynolds S, Levine R. *J. Am. Chem. Soc* 1960;82:472.
54. Fraser RR, Mansour TS, Savard S. *J. Org. Chem* 1985;50:3232.
55. Mamane V, Aubert E, Fort Y. *J. Org. Chem* 2007;72:7294. [PubMed: 17696549]
56. There are discussions of reversible radical-based dimerizations of heteroaromatics: Shimoishimi H, Tero-Kubota S, Ikegami Y. *Bull. Chem. Soc. Japan* 1985;58:553.
57. <http://www.drugtopics.com/Top+200+Drugs><http://www.chem.cornell.edu/jn96/outreach.html>
58. (a) Kottke T, Stalke D. *Angew. Chem., Int. Ed. Engl* 1993;32:580. (b) Rennels RA, Maliakal AJ, Collum DB. *J. Am. Chem. Soc* 1998;120:421.
59. Kofron WG, Baclawski LM. *J. Org. Chem* 1976;41:1879.
60. Precipitation occurs below 3.9 M THF.
61. Following product formation ([P]) and fitting the data to $[P]=[P^\infty][1-\exp(-k_{\text{obsd}}t)]$ affords the pseudo-first-order rate constant, k_{obsd} . The quality of the fit, however, is *not* diagnostic of first-order behavior because this function also fits second-order data reasonably well. The problem stems from determining the concentration of P at $t = \infty$ ($[P^\infty]$) as an adjustable parameter. The first order dependence was confirmed by (1) monitoring the value of k_{obsd} at different initial concentrations of starting material, and (2) monitoring the loss of picoline **1** ($\delta -70.2$ ppm) by ^{19}F NMR spectroscopy and showing adequate fit to $[A]=[A_0]\exp(-k_{\text{obsd}}t)$.
62. (a) Singh KJ, Collum DB. *J. Am. Chem. Soc* 2006;128:13753. [PubMed: 17044703] (b) Rennels RA, Rutherford JL, Collum DB. *J. Am. Chem. Soc* 2000;122:8640. and references cited therein. (c) Anderson DR, Faibish NC, Beak P. *J. Am. Chem. Soc* 1999;121:7553. (d) Meyers AI, Mihelich ED. *J. Org. Chem* 1975;40:3158.







Figure 1.
Vessel used for double crossover experiment.




Scheme 1.




Scheme 2.





Scheme 3.



Scheme 4.



Scheme 5.



Scheme 6.