

The Fission Yeast HIRA Histone Chaperone Is Required for Promoter Silencing and the Suppression of Cryptic Antisense Transcripts^{∇†}

Holly E. Anderson,^{1‡} Josephine Wardle,^{1‡} Şenay Vural Korkut,^{1§} Heather E. Murton,¹
Luis López-Maury,² Jürg Bähler,² and Simon K. Whitehall^{1*}

*Institute for Cell & Molecular Biosciences, Newcastle University, Newcastle upon Tyne NE2 4HH, United Kingdom,¹ and
Department of Genetics, Evolution and Environment and UCL Cancer Institute, University College London,
London WC1E 6BT, United Kingdom²*

Received 29 May 2009/Returned for modification 22 June 2009/Accepted 11 July 2009

The assembly of nucleosomes by histone chaperones is an important component of transcriptional regulation. Here, we have assessed the global roles of the HIRA histone chaperone in *Schizosaccharomyces pombe*. Microarray analysis indicates that inactivation of the HIRA complex results in increased expression of at least 4% of fission yeast genes. HIRA-regulated genes overlap with those which are normally repressed in vegetatively growing cells, such as targets of the Clr6 histone deacetylase and silenced genes located in subtelomeric regions. HIRA is also required for silencing of all 13 intact copies of the Tf2 long terminal repeat (LTR) retrotransposon. However, the role of HIRA is not restricted to bona fide promoters, because HIRA also suppresses noncoding transcripts from solo LTR elements and spurious antisense transcripts from cryptic promoters associated with transcribed regions. Furthermore, the HIRA complex is essential in the absence of the quality control provided by nuclear exosome-mediated degradation of illegitimate transcripts. This suggests that HIRA restricts genomic accessibility, and consistent with this, the chromosomes of cells lacking HIRA are more susceptible to genotoxic agents that cause double-strand breaks. Thus, the HIRA histone chaperone is required to maintain the protective functions of chromatin.

Chromatin restricts the access of the transcription machinery to the DNA template, and as a result, modulation of chromatin structure is central to eukaryotic transcriptional control. Two major mechanisms by which changes to chromatin structures can be achieved are posttranslational modification of histones (42) and ATP-dependent nucleosome remodeling (6). In addition, it is becoming increasingly apparent that chromatin structures are also modified by a diverse group of proteins known as histone chaperones that facilitate the assembly and, in some cases, the disassembly of nucleosomes (12).

The bulk of nucleosome assembly is coupled to DNA replication via the action of replication-dependent histone chaperones, such as CAF-1 (43). However, other DNA-dependent processes, such as transcription initiation and elongation, can result in nucleosome displacement, and so the requirement for nucleosome assembly is not restricted to S phase. Accordingly, eukaryotic cells possess a repertoire of replication-independent histone chaperones, which include an evolutionarily conserved class of WD repeat protein called HIRA (or HIR) (12). Humans have a single HIRA protein (23), whereas budding and fission yeast cells have two related proteins that function in concert (Hir1 and Hir2 in *Saccharomyces cerevisiae* and Hip1

and Slm9 in *Schizosaccharomyces pombe*) (4, 19, 44). HIRA proteins are components of multisubunit complexes and these complexes purified from a number of cell types have been shown to mediate replication-independent nucleosome deposition in vitro (15, 35, 38, 46).

Histone chaperone-dependent removal and replacement of nucleosomes have emerged as important components of transcriptional control (48), and accordingly, HIRA complexes have been linked to transcription. Indeed, *S. cerevisiae* Hir1 and Hir2 were originally identified as factors that repress the transcription of histone genes outside S phase (44). Subsequently, it has been demonstrated that nucleosomes assembled by the *S. cerevisiae* HIR complex are refractory to remodeling by the SWI/SNF complex in vitro (35), indicating that this chaperone is capable of organization of repressive forms of chromatin. Consistent with this, HIRA proteins have also been implicated in assembly of heterochromatin and silencing in a range of organisms. Loss of either HIRA protein in fission yeast leads to defective pericentric and *mat* locus heterochromatin (4), and mutations in *S. cerevisiae* HIR genes exacerbate the silencing defects associated with inactivation of CAF-1 (20, 37, 40). The assembly of senescence-associated heterochromatin in human cells is also dependent upon HIRA (49), and HIRA is required for maintenance of *knox* gene silencing during organogenesis in *Arabidopsis* (34). Despite these findings, other evidence is consistent with positive roles for HIRA in transcription. Experiments with chicken DT40 cell lines suggest that the N-terminal domain of HIRA can function as an activator (2), and HIRA is implicated in the replication-independent deposition of the histone variant H3.3, which marks actively transcribed genes (3). Furthermore, genetic interactions have also suggested that HIRA complexes are involved in transcription elongation (13).

* Corresponding author. Mailing address: Institute for Cell & Molecular Biosciences, Newcastle University, Framlington Place, Newcastle upon Tyne NE2 4HH, United Kingdom. Phone: 44 (0)191 222 5989. Fax: 44 (0)191 222 7424. E-mail: S.K.Whitehall@ncl.ac.uk.

† Supplemental material for this article may be found at <http://mcb.asm.org/>.

‡ These authors contributed equally to this study.

§ Present address: Yildiz Technical University, Faculty of Arts and Sciences, Department of Biology, 34210 Esenler/Istanbul, Turkey.

∇ Published ahead of print on 20 July 2009.

Here, we have assessed the role of the *S. pombe* HIRA complex in the global regulation of transcription. We find that deletion of *hip1*⁺ or *slm9*⁺ leads to upregulation of mRNA from a large number of genes, indicating that the HIRA complex plays an important role in suppression of transcription. HIRA targets overlap with silenced genes, and consistent with this, we demonstrate that HIRA represses the expression of all intact copies of the Tf2 long terminal repeat (LTR) retrotransposable element. Furthermore, HIRA also suppresses the production of noncoding transcripts from solo LTR elements and spurious antisense transcripts originating from cryptic promoters. In addition, we also find that the HIRA complex protects cells from agents that cause double-strand breaks. Thus, the HIRA complex is required for promoter silencing, suppression of antisense transcription, and maintenance of genomic integrity.

MATERIALS AND METHODS

Plasmids and strains. Culture of *S. pombe* and general genetic methods were performed as described previously (29). A strain with a *lacZ* reporter integrated into Tf2-1 was constructed as follows. A DNA fragment was amplified by PCR using primers Tf2REPC (5'-GCATAGGAATCTAGTACATCGCTATTCCAC CAG-3') and Tf2REPD (5'-GCATTGGGATCCAATTGCTTTGTCGCTTG TAG-3') and cloned into the EcoRI and BamHI sites of pSPI356 (22) to give pSPI356Tf2CD. A second DNA fragment, containing the Tf2-1 5' LTR, was amplified using primers Tf2REPA (5'-GCATAGGGATCCTGTGCAAGT ACTACACTACG-3') and Tf2REPB (5'-GCATAGCTGCAGGGAGTAAT TCTTGCCATGTAAG-3') and cloned into the PstI and BamHI sites of pSPI356Tf2CD. Following confirmation of the sequence, the resulting plasmid was linearized with BamHI and transformed into wild-type (NT5) *S. pombe* cells. Integration at the correct locus was confirmed by PCR. The Tf2-10 element was tagged in the same way. The remaining Tf2 elements were tagged using the same approach, except that primer Tf2REPA was replaced with a primer complementary to a sequence located 300 to 500 bp upstream of the 5' LTR of the appropriate Tf2 element.

A Tf2-1 *lacZ* reporter was also integrated into an ectopic locus on chromosome II (base pairs 1877855 to 1878381). This was achieved by amplifying two DNA fragments, the first with ChrmIIA (5'-TATGTCGAGGATCCTGTAACC GTTTAGATTGCAGC-3') and ChrmIIB (5'-GTACTACTGCAGTTGTTACT GTTGTGTAGAGCC-3') and the second with ChrmIIC (5'-CCTAGAGAATT CCAACTAACCGTAATACATCGG-3') and ChrmIID (5'-CGGTTACAGGA TCCTCGACATAACACTTGCAGGTG-3'). These fragments were used as the template in an overlapping PCR with primers ChrmIIC and ChrmIIB, and the resulting DNA fragment was cloned into the EcoRI and PstI sites of pSPI356 to give pSPI356ChrmIICDAB. A fragment of Tf2-1 was amplified using Tf2REPB and Tf2(A)PstI (5'-CGATAGCTGCAGGTGCAAGTACTACACTACG-3'), cleaved with PstI, and cloned into the PstI site of pSPI356ChrmIICDAB. The resulting plasmid was linearized with BamHI and transformed into wild-type *S. pombe* cells. Integration at the correct locus was confirmed by PCR. A strain carrying a solo LTR-*lacZ* reporter was constructed using the same approach, except that the Tf2REPB primer was replaced with LTR3'del349 (5'-CGAAT CCTGCAGGCATTGTAAGCTACGCAGTTTGGA-3'). All *lacZ* reporters were introduced into the *hip1Δ* background by using standard genetic crosses. The other strains used in this study were 972 (*h*⁻), SW577 (*h*⁻ *hip1::ura4*⁺), SW578 (*h*⁻ *slm9::ura4*⁺), NT5 (*h*⁻ *ade6-M216 leu1-32 ura4-D18*), SW137 (*h*⁻ *ade6-M210 leu1-32 ura4-D18 hip1::ura4*⁺), SW138 (*h*⁺ *ade6-M210 leu1-32 ura4-D18 hip1::ura4*⁺), JK2246 (*h*⁻ *leu1-32 ura4-D18 slm9::ura4*⁺), SW318 (*h*⁻ *ade6-M210 leu1-32 ura4-D18 hip3::ura4*⁺), yYH7a (*h*⁻ *ade6 leu1-32 ura4-D18 rrp6::kanMX6*), SW524 (*h*⁺ *ade6-M216 leu1-32 ura4-D18 clr6-1*), and SW611 (*h*⁻ *ade6-M210 leu1-32 ura4-D18 clr6-1 hip1::ura4*⁺).

Microarray analysis. We used DNA microarrays displaying probes for >99% of all known and predicted genes of *S. pombe* spotted in duplicate onto glass slides. RNA extraction, hybridization, and initial data processing and normalization were performed as previously described (24). Four independent biological experiments were performed, including two dye swaps, for each mutant. The data were visualized and analyzed using GeneSpring GX7.3 (Agilent). The significance of overlaps between different gene lists was calculated with GeneSpring by using a standard Fisher exact test, and *P* values were adjusted with a Bonferroni multiple-testing correction. An average mutant/wild-type ratio was calculated for each gene, and cutoff values of 1.5-fold were used for comparisons between

mutants. Gene annotations were downloaded from the *S. pombe* GeneDB (<http://www.genedb.org/genedb/pombe/>). Clustering along chromosomes of genes with induced expression was analyzed using an in-house Perl script which compares clustered genes to a random distribution. *P* values were adjusted for multiple testing using the Benjamini-Hochberg false-discovery-rate method.

Telomere length. Telomere length measurements were performed as previously described (10). A probe for telomere repeat sequences was made with a SacI-PstI fragment from pJCF1640 (27). A probe that was used as a loading control was made by PCR amplifying a fragment by using primers ApaIF (5'-C GTGATAGTGCATTGACGATC-3') and ApaIR (5'-CCATCTTGCATGGCA CGCTTC-3').

RT-PCR. RNA was purified as described for microarray analysis and subjected to reverse transcription-PCR (RT-PCR) using a one-step RT-PCR kit (Qiagen). For strand-specific RT-PCR, one primer complementary to the sense or antisense transcript was added during first-strand cDNA synthesis while the second primer was added prior to the PCR amplification steps. The primers were as previously described (30). cDNA for quantitative (real-time) RT-PCR was made using a Superscript II kit (Invitrogen). Real-time PCRs were performed using a LightCycler 2.0 PCR system (Roche) and SYBR green mixture (Molecular Probes), using the appropriate primers. Reactions were normalized using primers specific to *act1*⁺.

Pulsed-field gel electrophoresis (PFGE). DNA plugs were prepared as described previously (36). DNA was fractionated on 1% ultrapure DNA grade agarose (Bio-Rad) gel prepared in 1× Tris-acetate-EDTA by using a CHEF-DR III system (Bio-Rad). Gels were run for 48 h at 14°C at a voltage of 2 V cm⁻¹, with an included angle of 106° and initial and final switching times of 1,200 s and 1,800 s, respectively.

Microarray data accession number. The microarray data set is available at Array Express (E-MTAB-130).

RESULTS

The HIRA complex is a global suppressor of transcription.

We have previously demonstrated that the *S. pombe* HIRA proteins Hip1 and Slm9 interact with a structurally unrelated protein called Hip3 to form a complex that is required for heterochromatic transcriptional silencing (16). In order to determine whether the HIRA complex regulates transcription in euchromatin, the RNA expression profiles of *hip1Δ* and *slm9Δ* cells were compared with those of wild-type cells by using microarrays detecting >99% of all known or predicted open reading frames and other genomic elements (24). As expected, the genes that were found to be differentially expressed in *hip1Δ* and *slm9Δ* cells were highly correlated ($P < 3 \times 10^{-243}$) (Fig. 1A; see also Table S1 in the supplemental material). A total of 195 mRNAs were found to be commonly upregulated ≥ 1.5 -fold in *hip1Δ* and *slm9Δ* mutants, suggesting that the HIRA complex suppresses transcription from a large number of RNA polymerase II promoters. We observed a significant overlap ($P < 6 \times 10^{-33}$) between genes upregulated in *hip1Δ* and *slm9Δ* mutants and genes known to be derepressed in a *clr6-1* background, which harbors a point mutation in an essential class I histone deacetylase (HDAC) (17). Clr6 functions as the catalytic core of an *S. pombe* HDAC-Sin3 corepressor complex, which is targeted to the promoters of numerous genes (30).

The functional overlap with the Clr6 HDAC suggested that the HIRA complex regulates genes whose expression is repressed under normal growth conditions. Indeed, the transcriptomes of HIRA mutants were significantly enriched for mRNAs with no measurable microarray signal in vegetative cells ($P < 1 \times 10^{-8}$) (unpublished data) and for the 10% most lowly expressed mRNAs ($P < 1 \times 10^{-9}$) (21). Consistent with these data, transcripts from Tf2 LTR retrotransposons (which are largely suppressed in wild-type cells) were greatly increased




FIG. 1. Transcriptomes of HIRA mutants. (A) Venn diagrams showing overlap between genes upregulated (≥ 1.5 -fold) in HIRA mutants with genes upregulated under the indicated condition. The P values indicate the probability that the observed overlap happened by chance. (B) Graphs showing mutant versus wild-type (wt) expression levels of subtelomeric genes of the right (R) and left (L) arms of chromosomes I and II. Lines indicating the 1.5-fold threshold are shown. Genes whose expression was confirmed by quantitative RT-PCR are marked by asterisks.

in *hip1* Δ and *slm9* Δ backgrounds. Also, many of the silenced genes located in subtelomeric regions of chromosomes I and II (47) were upregulated in the absence of HIRA (Fig. 1B), and analysis of the location of HIRA-repressed genes confirmed that they are enriched for subtelomeric genes ($P < 1 \times 10^{-7}$). Overall, our results indicate that the HIRA complex is important for the global maintenance of transcriptional silencing.

Further examination of gene expression patterns revealed a significant overlap ($P < 3 \times 10^{-49}$) between genes upregulated in *hip1* Δ and *slm9* Δ mutants and those upregulated in cells deleted for telomerase (26) (Fig. 1A). Many of the genes that have been shown to be upregulated as cells undergo telomere crisis are core environmental stress response genes (9), and

this is also true of those upregulated in response to loss of the HIRA complex ($P < 2 \times 10^{-22}$). These findings prompted us to examine whether or not the HIRA complex influences telomere length regulation, particularly as *clr6-1* mutants are known to have elongated telomeres (17). However, Southern blotting revealed that loss of the HIRA complex did not result in increased telomere length; indeed, *hip1* Δ and *slm9* Δ mutants had telomeres that were slightly shorter than those of wild-type cells. Furthermore, deletion of *hip1* Δ also partially reversed the elongated telomere phenotype that is associated with the *clr6-1* allele (Fig. 2).

Although loss of the HIRA complex predominantly led to increased mRNA levels, the expression levels of 38 genes were




FIG. 2. Influence of HIRA on telomere length. DNA samples purified from the indicated strains were digested with *Apa*I and analyzed by Southern blotting with a telomere repeat probe (lower panel) and loading control probe (upper panel) as described in Materials and Methods. w.t., wild type.

reduced (≤ 1.5 -fold) (see Table S1 in the supplemental material). Prominent among these genes were *feo1*⁺, *frp1*⁺, *str1*⁺, *str3*⁺, and *fip1*⁺, whose expression is regulated in response to intracellular iron status by the Fep1 repressor ($P < 6 \times 10^{-5}$) (32, 33). The expression of these genes is also downregulated in other mutants that are defective in silencing, such as *clr1-5* and *clr3-735*, which lack the activity of the SHREC complex (17, 45). However, this effect is thought to be indirect, as in the *clr1-5* and *clr3-735* mutant backgrounds, downregulation of the iron regulon is correlated with increased levels of mRNA encoding the Fep1 repressor (17). In contrast, microarray and quantitative RT-PCR analyses indicated that mutation of *hip1*⁺ or *slm9*⁺ did not result in an increased abundance of *fep1*⁺ mRNA (see Table S1 in the supplemental material; also unpublished data). Furthermore, loss of HIRA did not result in increased transcript levels of components of the SWI/SNF complex, which has also recently been shown to repress the iron regulon (28) or alter intracellular iron levels (see Table S1 in the supplemental material; also unpublished data). Thus, it is possible that regulation of the iron regulon may be direct and some genes may require the HIRA complex for full expression.

HIRA represses transcription of all 13 Tf2 LTR retrotransposons. Previous analysis has implicated the HIRA complex in the silencing of Tf2 LTR retrotransposons (16), and as outlined above, the microarray analysis confirmed that the levels of Tf2 mRNA were highly increased in the *hip1Δ* and *slm9Δ* backgrounds. The fission yeast laboratory strain (972) contains 13 full-length copies of this element (Tf2-1 to Tf2-13), and probes for nine of these elements were present on the microarray. However these elements are highly homogeneous having a pairwise identity of greater than 99% (5) and cross-hybridization between the Tf2 probes is expected. Therefore, it was not possible to determine whether the global increase in Tf2 mRNA resulted from derepression of a single element, a subset of elements, or all of the 13 Tf2 elements.

In order to address this issue, a series of 13 strains was constructed, each containing an integrated *lacZ* reporter gene under the control of a specific Tf2 element (Fig. 3A). β -Galactosidase assays of the resulting reporter strains indicated that all 13 elements are expressed at similar low levels, which is consistent with these elements being silenced (Fig. 3B). To determine the effect of HIRA on the expression of individual Tf2 elements, the *lacZ* reporter strains were then crossed into a *hip1Δ* background. Expression levels of all 13 Tf2-*lacZ* reporters were found to be increased in the absence of Hip1, indicating that the HIRA complex represses the expression of all of the Tf2 elements (Fig. 3C). Notably, the level of derepression of the Tf2-11 reporter was reproducibly lower than that of the other reporters. The Tf2-11 element has a 5' LTR which is a hybrid of a Tf2 element and a Tf1 element, a related retrotransposon that is now extinct in the 972 genome (5). As a result, Tf2-11, unlike the other elements, is not induced in response to limitation of oxygen by the SREBP homologue Sre1 (39). Nonetheless, our results indicate that Tf2-11 is subject to HIRA-mediated repression.

We next determined whether HIRA-dependent repression is maintained when the Tf2 element is moved to a new region of the genome that does not naturally contain a retrotransposon. Therefore, we integrated a Tf2-*lacZ* reporter into a large intergenic region of chromosome II in between a tRNA gene (*SPBTRNAPRO.05*) and *SPBC3H7.03c* (Fig. 3D). This region was chosen as there is no evidence to suggest that HIRA complexes regulate the expression of RNA polymerase III-transcribed genes and our microarray experiments indicated that HIRA does not affect the expression of *SPBC3H7.03c* (see Table S1 in the supplemental material). β -Galactosidase assays revealed that the level of expression of this reporter was low in wild-type cells and was dramatically increased in a *hip1Δ* background, indicating that HIRA-dependent silencing is retained at a novel site in the genome. In addition to the 5' LTR, the Tf2 reporter also contains the 5' untranslated region and DNA encoding the first 333 bp of the Tf2 *gag* gene. In order to determine whether these sequences were important for HIRA-mediated repression, a reporter in which the *lacZ* gene is solely controlled by the 5' LTR from the Tf2-1 element was constructed. Following integration into the ectopic locus on chromosome II, we again found that the level of expression of this reporter was low in wild-type cells and was dramatically increased in a *hip1Δ* background (Fig. 3D). This indicates that a Tf2 LTR element alone is a functional promoter that is repressed by the HIRA complex.

This finding is important because, in addition to the 13 full-length elements, the genome contains an extensive, but heterogeneous, population of (~ 250) solo LTR elements that have arisen through LTR-LTR recombination and so mark the position of previous retrotransposon insertion events (5). In order to determine whether or not the HIRA complex is required to silence naturally occurring solo LTRs, expression from a selection of 11 of these elements from all three chromosomes was analyzed using quantitative RT-PCR. Using this approach, we were unable to detect expression from four LTRs either in wild-type or in mutant backgrounds. However, non-coding transcripts were detected for seven elements, and in all these cases, transcript levels were substantially increased in the absence of the HIRA complex (Fig. 4). Importantly, we found




FIG. 3. HIRA silences the expression of all 13 Tf2 LTR retrotransposons. (A) Schematic of the *lacZ* reporter used to tag individual Tf2 elements. UTR, untranslated region. (B) Expression of Tf2-*lacZ* reporters in wild-type cells. Strains harboring a tagged Tf2 element were grown in rich medium (YE5S) at 30°C to mid-log phase. Cells were then harvested and processed for liquid β -galactosidase assays. Shown are the mean values of results from three experiments. Error bars indicate standard deviations. (C) Expression of Tf2-*lacZ* reporters in a *hip1* Δ background. Strains were processed as described for panel B. (D) HIRA-dependent repression is maintained at a novel site in the genome. Wild-type or *hip1* Δ strains harboring either the Tf2-1 or the LTR *lacZ* reporter integrated into chromosome II were processed as described for panel B.

examples of both Tf2- and Tf1-type LTRs that were regulated in this way. This suggests that the *S. pombe* HIRA complex plays a widespread role in the suppression of cryptic transcripts from solo LTRs.

HIRA represses spurious antisense transcription. As the HIRA complex functions to limit spurious transcripts from solo LTR elements, we investigated whether it is also required to suppress other illegitimate transcripts. Recent data indicate that mutations that impair the integrity of chromatin result in

the accumulation of antisense transcripts from cryptic promoters in coding regions (30). We therefore used strand-specific RT-PCR to examine several loci where defects in chromatin are known to result in increased levels of antisense transcripts. Consistent with previous findings (30), antisense transcripts originating at the *zer1⁺* locus were not detectable in wild-type cells, but in contrast, mutation in any one of the genes encoding subunits of the HIRA complex (*hip1⁺*, *slm9⁺*, or *hip3⁺*) resulted in a large increase in the levels of these antisense




FIG. 4. HIRA represses expression from Solo LTRs. RNA was prepared from wild-type (wt), *hip1Δ*, and *slm9Δ* cells. Quantitative real-time RT-PCR was then used to quantify the levels of noncoding transcripts ~100 bp downstream of selected LTR elements. Change in expression (mutant versus wild type) is shown. Error bars indicate standard deviations. For each element, the location, type, and homology to the Tf2-1 LTR is shown.

transcripts (Fig. 5A). Furthermore, deletion of the HIRA complex had a similar effect on antisense transcripts from *hrp1* and *sfc3* loci (Fig. 5A). Thus, in addition to suppressing transcripts from numerous gene promoters and LTRs, the HIRA complex is required to suppress spurious antisense transcripts from the coding regions of RNA polymerase II-transcribed genes.

The HDAC Clr6 also plays a key role in the suppression of antisense transcripts (30). Clr6 is present in at least two distinct HDAC-Sin3 corepressor complexes (complexes I and II). While complex I predominantly represses transcription from gene promoters, complex II appears to be the functional equivalent of the *S. cerevisiae* Rpd3S complex and targets tran-

scribed regions (30). We therefore investigated the genetic relationship between the HIRA complex and Clr6 with respect to antisense transcription. Introduction of the *clr6-1* allele into the *hip1Δ* background did not result in further increase in antisense transcript levels at the *hrp1* and *zer1* loci (Fig. 5B), suggesting that the HIRA complex and Clr6 complex II function in the same pathway to suppress antisense transcription.

The nuclear exosome recognizes and degrades spurious antisense transcripts, preventing their accumulation (30). Consistent with this, combining mutations in *rrp6* (which encodes a 3' exonuclease) (18) with mutations in genes encoding components of Clr6 complex II leads to a synergistic increase in




FIG. 5. Loss of HIRA leads to the accumulation of spurious antisense transcripts. (A) RNA was purified from wild-type (w.t.), *hip1Δ*, *slm9Δ*, and *hip3Δ* cells and analyzed by strand-specific RT-PCR. One primer, complementary to either the forward or the reverse transcripts, was included during the reverse transcription step, and the second primer was then added during PCR amplification. Control reactions omitting the reverse transcription step (-RT) were included to demonstrate the absence of DNA. (B) RNA from wild-type, *hip1Δ*, *rrp6Δ*, *clr6-1*, and *hip1Δ clr6-1* strains was analyzed as described for panel A. (C) The nuclear exosome is essential in the absence of HIRA. The left hand panel shows an example of a tetrad resulting from a cross between *hip1Δ* and *rrp6Δ* strains, and the right hand panel shows the terminal morphology of the *hip1Δ rrp6Δ* double mutant.




FIG. 6. Antisense transcription is correlated with sense transcription in wild-type but not HIRA mutant cells. (A) Schematic diagram showing regulation of *thi4*⁺ expression. (B) Wild-type and *hip1*Δ cells were grown to log phase in minimal (EMM) medium, thiamine was then added to give a final concentration of 2 μM, and samples of cells were harvested at the indicated time points. RNA, prepared from the resulting cell pellets, was analyzed by strand-specific RT-PCR as described in the legend to Fig. 5. –RT, control reactions omitting the reverse transcription step. (C) Antisense transcripts persist under conditions of constant *thi4*⁺ repression. *slm9*Δ cells were grown to log phase either in EMM medium or in EMM medium supplemented with 2 μM thiamine. RNA preparation and RT-PCR analysis were performed as described in the legend to Fig. 5.

antisense transcripts (30). On the basis of this finding, we predicted that there would also be a genetic interaction between the nuclear exosome and HIRA. Indeed, we consistently failed to recover a viable double mutant from genetic crosses of *hip1*Δ and *rrp6*Δ strains. The same was also true for crosses of *hip3*Δ and *rrp6*Δ cells. Microscopic examination revealed that double mutant strains were able to germinate but failed to progress past six or seven cell divisions (Fig. 5C). Thus, the nuclear exosome is essential in the absence of the HIRA complex, suggesting that this histone chaperone plays a major role in the suppression of spurious antisense transcripts.

Sense transcription is required for antisense transcription. Accumulating evidence indicates that passage of RNA polymerase II can disrupt chromatin (48), suggesting that it is transcription in the sense direction that allows access to cryptic promoters. As such, sense transcription may be a prerequisite for antisense transcription. In order to examine the interdependence of sense and antisense transcriptions, we examined the *thi4*⁺ gene, whose expression is regulated in response to the availability of thiamine, being very highly expressed in its absence and repressed in its presence (50). Strand-specific RT-PCR analysis confirmed that in wild-type cells, the production of sense transcripts was reduced by the addition of thiamine to the medium (Fig. 6). Further analysis revealed that in the absence of thiamine (when high levels of sense transcription were occurring), *thi4* antisense transcripts were clearly detectable. However, the addition of thiamine and the inhibition of sense transcription resulted in a concomitant suppression of antisense transcripts (Fig. 6). Thus, in wild-type cells, the levels of sense and antisense transcripts are correlated and the production of *thi4* antisense transcripts is dependent upon high levels of transcription in the sense direction. Notably, the addition of thiamine to the medium also inhibited sense transcription in *hip1*Δ cells, but the level of antisense transcripts was not reduced by this treatment (Fig. 6). Furthermore, in a HIRA mutant background, antisense transcripts were detectable even when cells were cultured under conditions of constant *thi4*⁺ repression (Fig.

6C). Thus, in the absence of the HIRA histone chaperone, access to cryptic promoters is maintained even when sense transcription is inhibited.

HIRA is required for resistance to DNA damage. It has been proposed that increased levels of cryptic antisense transcripts are indicative of an open chromatin conformation (30). Consistent with this hypothesis, the genomes of *clr6-1* mutants show an increased sensitivity to DNA-damaging agents (30). We therefore investigated the sensitivity of cells lacking the HIRA complex to a range of genotoxic agents, such as the alkylating agent methyl methanesulfonate, the radiomimetic zeocin, the topoisomerase I poison camptothecin, and hydroxyurea, which inhibits ribonucleotide reductase and causes replication fork stalling. Serial dilution assays revealed that mutation in any one of the genes encoding components of the HIRA complex (*hip1*Δ, *slm9*Δ, or *hip3*Δ) results in a marked increase in sensitivity to methyl methanesulfonate, zeocin, and camptothecin (Fig. 7A). In contrast, HIRA mutants exhibited only a very subtle increase in sensitivity to hydroxyurea. Thus, the HIRA complex is important for protection against agents that cause DNA double-strand breaks but not against a replication stress.

In order to determine whether HIRA mutants have increased susceptibility to chromosomal breakage, cells were first exposed to bleomycin and then chromosome integrity was analyzed using PFGE (Fig. 7B). As previously reported (30), the *clr6-1* mutant exhibited enhanced levels of bleomycin-induced chromosome breakage as evidenced by the appearance of DNA fragments and the loss of intact chromosomes. Furthermore, the chromosomes of *hip1*Δ cells were also more susceptible to breakage than those of wild-type cells. This finding suggests that HIRA-mediated nucleosome assembly is required to maintain the global protective functions of chromatin. Interestingly, analysis of a *clr6-1 hip1*Δ double mutant indicated that this strain was more sensitive to genotoxic agents than either of the parental strains (Fig. 7C), indicating that the HIRA complex and Clr6 do not have identical roles in the response to DNA double-strand breaks.




FIG. 7. HIRA is required for protection against genotoxic agents. (A) The indicated strains were grown to log phase in YE5S medium, subjected to fivefold serial dilutions, and spotted on YE5S agar and YE5S agar supplemented with the indicated genotoxic agent. wt, wild type; HU, hydroxyurea; MMS, methyl methanesulfonate. (B) The indicated strains were treated with bleomycin at the indicated concentrations for 90 min. Chromosomal DNA was then analyzed using PFGE. (C) The indicated strains were grown to log phase and treated as described for panel A.

DISCUSSION

Previous evidence has demonstrated that the HIRA histone chaperone is required for transcriptional silencing at heterochromatic loci (16). Here, we demonstrate that its function is not limited to heterochromatin, because HIRA represses transcripts from numerous promoters distributed throughout the genome, including subtelomeric genes, LTR retrotransposons, and their remnants, and it also limits the levels of cryptic antisense transcripts. Furthermore, loss of HIRA leads to increased access of genotoxic agents to the genome, indicating that HIRA is required for maintenance of the protective functions of chromatin.

Our data revealed a functional overlap between the HIRA complex and the class I HDAC Clr6 with respect to both promoter silencing and prevention of cryptic transcription. The simplest explanation of this finding is that HIRA is required to reassemble or repair nucleosomes that are subsequently modified by Clr6-Sin3 complexes. Interestingly, class I HDACs have been linked to HIRA function in vertebrate cells, as HDAC-2 stably interacts with chicken HIRA through an LXXLL motif located in the HIRA C-terminal region (1). However, neither Hip1 nor Slm9 has LXXLL motifs, and there is no evidence of a stable interaction between the fission yeast HIRA complex and Clr6. Indeed, large-scale affinity purification of Clr6-interacting subunits did not identify any of the components of the HIRA complex (30). Furthermore, a large-scale purification of Prw1, a homologue of RbAp48 that is present in all fission yeast Clr6-Sin3 complexes, also did not identify any subunits of the HIRA complex (unpublished data).

Analysis of the expression of individual Tf2 elements indicated that the high levels of Tf2 mRNA present in *hip1Δ* and *slm9Δ* cells result from increased expression levels of all 13 copies of this element. It is possible that Tf2 elements are located within regions of the genome that are associated with

HIRA-dependent forms of chromatin. Arguing against this, HIRA-dependent repression of a Tf2 reporter was also maintained when it was moved to a novel site in the genome, suggesting that these elements contain sequences that limit their own expression.

Packaging of retrotransposons into repressive chromatin structures is known to occur in many cell types and is thought to limit the potentially harmful spread of these elements (25). The chromatin structures that suppress the expression of Tf2 retrotransposons appear to be dependent upon the HIRA complex. It is possible that the interaction of HIRA with these elements is transient because, as yet, we have been unable to detect binding of HIRA to Tf2 LTRs by chromatin immunoprecipitation. Global silencing of Tf2 elements is also dependent upon HDACs (Clr6, Clr3, and Hst4) and Cenp-B homologues (7, 11, 17, 45). While the influence of these factors on the expression of individual elements has not been formally investigated, chromatin immunoprecipitation analysis has indicated that Clr3 and Cenp-B proteins are located at multiple Tf2 elements (7, 45), implying that they are required for silencing of all Tf2 elements.

The involvement of HIRA proteins in the regulation of retrotransposons is not limited to fission yeast, because Hir1 and Hir2 have been implicated in the regulation of Ty1/Copia elements in *S. cerevisiae*. Indeed, *hir* mutants were found to suppress the deleterious effects of the insertion of a Ty LTR (δ element) into the *HIS4* locus (41). Furthermore, mutation of *HIR* genes in combination with *CAC* genes also increases Ty1 transposition frequency, although this increased transposition was not associated with increased levels of Ty1 mRNA (37). It is important to note that the Tf2 elements of *S. pombe* are members of the Gypsy group of LTR retrotransposons and as such are only distantly related to the Ty1/Copia elements of *S. cerevisiae* (5). The finding that HIRA complexes regulate

such distinct elements suggests that this chaperone regulates other classes of LTR retrotransposons in other eukaryotic organisms.

The substantial population of solo LTR elements present in the *S. pombe* genome have been generated by recombination between two LTRs resulting in the removal of the internal retrotransposon coding sequences. Thus, solo LTRs are the remnants of retrotransposons and mark the positions of previous insertion events. Our data, along with those of others (7, 30, 45), are consistent with these elements being assembled into silent chromatin in order to limit the production of spurious noncoding transcripts. While in many cases expression from solo LTRs would not be advantageous, there are an increasing number of examples of cells exploiting retrotransposable elements to regulate gene expression (14). Indeed, in *S. pombe* a set of solo LTRs that are closely related to the LTRs associated with full-length Tf2 elements are known to function as oxygen-responsive promoters (39). Some of these confer oxygen-dependent expression to neighboring genes. Our data indicate that HIRA-dependent repression is not restricted to elements that are induced in response to low oxygen levels, as all but one of the solo LTR elements examined in this work lack a consensus SRE element and we also identified Tf1-type LTR elements that were repressed by Hip1 and Slm9.

The finding that HIRA mutants have high levels of spurious antisense transcription is consistent with this histone chaperone being required to maintain the integrity of chromatin in transcribed regions. Furthermore, mutation of *hip1* being synthetic lethal with *rrp6Δ* is consistent with HIRA having a major role in prevention of such antisense transcription in fission yeast. This result would not necessarily have been predicted from studies of *S. cerevisiae* where mutation of *HIR* genes did not result in high levels of internal initiation at the synthetic *FLO8-HIS3* reporter gene, although it did increase the level of spurious sense transcripts observed in a *spt2Δ* background (31). Nonetheless, our analysis is the first report that HIRA proteins are required to prevent cryptic antisense transcripts at naturally occurring genes.

Passage of RNA polymerase II through chromatin has been proposed to result in partial or complete disassembly of nucleosomes (48). That antisense transcription (and by implication access to cryptic promoters) at the *thi4⁺* gene is dependent upon high levels of sense transcription is entirely consistent with this notion. Notably, loss of the HIRA complex apparently leads to continued access to these promoters even in the absence of high levels of sense transcription. We propose that the HIRA complex restores chromatin structure in the wake of RNA polymerase II passage. Consistent with this idea, Hir1 and Hir2 have been shown to localize to transcribed regions in *S. cerevisiae* (31). In higher cells, HIRA is linked to deposition of variant histone H3.3 at actively transcribed genes (3), and so it will be interesting to determine whether this contributes to the prevention of spurious transcription initiation.

We also show that the HIRA complex plays an important role in protecting cells against double-strand breaks. The loss of HIRA led to an increased susceptibility to bleomycin-induced chromosomal breaks, suggesting that HIRA is required to maintain the protective functions of chromatin. This may not be the only role for HIRA in the response to DNA dam-

age, as other histone chaperones, such as *S. cerevisiae* Asf1, have been shown to be required for restoration of chromatin structures following DNA repair (8). Thus, it will be important to determine whether or not HIRA is also required for removal or replacement of nucleosomes during repair of double-strand breaks.

ACKNOWLEDGMENTS

We thank Richard Maraia and Julie Cooper for strains and plasmids and Janet Partridge, Elizabeth Veal, and Brian Morgan for comments on the manuscript.

This work was funded by a BBSRC project grant (BB/E014445/1) to S.K.W. and by a Cancer Research UK program grant (C9546/A6517) to J.B. H.E.A. was the recipient of a BBSRC DTG award.

REFERENCES

- Ahmad, A., Y. Takami, and T. Nakayama. 2004. WD dipeptide motifs and LXXLL motif of chicken HIRA are essential for interactions with the p48 subunit of chromatin assembly factor-1 and histone deacetylase-2 in vitro and in vivo. *Gene* **342**:125–136.
- Ahmad, A., Y. Takami, and T. Nakayama. 2003. WD dipeptide motifs and LXXLL motif of chicken HIRA are necessary for transcription repression and the latter motif is essential for interaction with histone deacetylase-2 in vivo. *Biochem. Biophys. Res. Commun.* **312**:1266–1272.
- Ahmad, K., and S. Henikoff. 2002. The histone variant H3.3 marks active chromatin by replication-independent nucleosome assembly. *Mol. Cell* **9**:1191–1200.
- Blackwell, C., K. A. Martin, A. Greenall, A. Pidoux, R. C. Allshire, and S. K. Whitehall. 2004. The *Schizosaccharomyces pombe* HIRA-like protein Hip1 is required for the periodic expression of histone genes and contributes to the function of complex centromeres. *Mol. Cell. Biol.* **24**:4309–4320.
- Bowen, N. J., I. K. Jordan, J. A. Epstein, V. Wood, and H. L. Levin. 2003. Retrotransposons and their recognition of pol II promoters: a comprehensive survey of the transposable elements from the complete genome sequence of *Schizosaccharomyces pombe*. *Genome Res.* **13**:1984–1997.
- Cairns, B. R. 2005. Chromatin remodeling complexes: strength in diversity, precision through specialization. *Curr. Opin. Genet. Dev.* **15**:185–190.
- Cam, H. P., K. Noma, H. Ebina, H. L. Levin, and S. I. Grewal. 2008. Host genome surveillance for retrotransposons by transposon-derived proteins. *Nature* **451**:431–436.
- Chen, C. C., J. J. Carson, J. Feser, B. Tamburini, S. Zabaronick, J. Linger, and J. K. Tyler. 2008. Acetylated lysine 56 on histone H3 drives chromatin assembly after repair and signals for the completion of repair. *Cell* **134**:231–243.
- Chen, D., W. M. Toone, J. Mata, R. Lyne, G. Burns, K. Kivinen, A. Brazma, N. Jones, and J. Bahler. 2003. Global transcriptional responses of fission yeast to environmental stress. *Mol. Biol. Cell* **14**:214–229.
- Cooper, J. P., E. R. Nimmo, R. C. Allshire, and T. R. Cech. 1997. Regulation of telomere length and function by a Myb-domain protein in fission yeast. *Nature* **385**:744–747.
- Durand-Dubief, M., I. Sinha, F. Fagerstrom-Billai, C. Bonilla, A. Wright, M. Grunstein, and K. Ekwall. 2007. Specific functions for the fission yeast Sirtuins Hst2 and Hst4 in gene regulation and retrotransposon silencing. *EMBO J.* **26**:2477–2488.
- Eitoku, M., L. Sato, T. Senda, and M. Horikoshi. 2008. Histone chaperones: 30 years from isolation to elucidation of the mechanisms of nucleosome assembly and disassembly. *Cell. Mol. Life Sci.* **65**:414–444.
- Formosa, T., S. Ruone, M. D. Adams, A. E. Olsen, P. Eriksson, Y. Yu, A. R. Rhoades, P. D. Kaufman, and D. J. Stillman. 2002. Defects in *SPT16* or *POB3* (yFACT) in *Saccharomyces cerevisiae* cause dependence on the Hir/Hpc pathway: polymerase passage may degrade chromatin structure. *Genetics* **162**:1557–1571.
- Goodier, J. L., and H. H. Kazazian, Jr. 2008. Retrotransposons revisited: the restraint and rehabilitation of parasites. *Cell* **135**:23–35.
- Green, E. M., A. J. Antczak, A. O. Bailey, A. A. Franco, K. J. Wu, J. R. Yates III, and P. D. Kaufman. 2005. Replication-independent histone deposition by the HIR complex and Asf1. *Curr. Biol.* **15**:2044–2049.
- Greenall, A., E. S. Williams, K. A. Martin, J. M. Palmer, J. Gray, C. Liu, and S. K. Whitehall. 2006. Hip3 interacts with the HIRA proteins Hip1 and Slm9 and is required for transcriptional silencing and accurate chromosome segregation. *J. Biol. Chem.* **281**:8732–8739.
- Hansen, K. R., G. Burns, J. Mata, T. A. Volpe, R. A. Martienssen, J. Bahler, and G. Thon. 2005. Global effects on gene expression in fission yeast by silencing and RNA interference machineries. *Mol. Cell. Biol.* **25**:590–601.
- Huang, Y., M. A. Bayfield, R. V. Intine, and R. J. Maraia. 2006. Separate RNA-binding surfaces on the multifunctional La protein mediate distinguishable activities in tRNA maturation. *Nat. Struct. Mol. Biol.* **13**:611–618.

19. Kano, J., and P. Russell. 2000. Slm9, a novel nuclear protein involved in mitotic control in fission yeast. *Genetics* **155**:623–631.
20. Kaufman, P. D., J. L. Cohen, and M. A. Osley. 1998. Hir proteins are required for position-dependent gene silencing in *Saccharomyces cerevisiae* in the absence of chromatin assembly factor I. *Mol. Cell. Biol.* **18**:4793–4806.
21. Lackner, D. H., T. H. Beilharz, S. Marguerat, J. Mata, S. Watt, F. Schubert, T. Preiss, and J. Bahler. 2007. A network of multiple regulatory layers shapes gene expression in fission yeast. *Mol. Cell* **26**:145–155.
22. Lafuente, M. J., T. Petit, and C. Gancedo. 1997. A series of vectors to construct *lacZ* fusions for the study of gene expression in *Schizosaccharomyces pombe*. *FEBS Lett.* **420**:39–42.
23. Lamour, V., Y. Lecluse, C. Desmaze, M. Spector, M. Bodescot, A. Aurias, M. A. Osley, and M. Lipinski. 1995. A human homolog of the *S. cerevisiae* HIR1 and HIR2 transcriptional repressors cloned from the DiGeorge syndrome critical region. *Hum. Mol. Genet.* **4**:791–799.
24. Lyne, R., G. Burns, J. Mata, C. J. Penkett, G. Rustici, D. Chen, C. Langford, D. Vetric, and J. Bahler. 2003. Whole-genome microarrays of fission yeast: characteristics, accuracy, reproducibility, and processing of array data. *BMC Genomics* **4**:27.
25. Maksakova, I. A., D. L. Mager, and D. Reiss. 2008. Keeping active endogenous retroviral-like elements in check: the epigenetic perspective. *Cell. Mol. Life Sci.* **65**:3329–3347.
26. Mandell, J. G., J. Bahler, T. A. Volpe, R. A. Martienssen, and T. R. Cech. 2005. Global expression changes resulting from loss of telomeric DNA in fission yeast. *Genome Biol.* **6**:R1.
27. Miller, K. M., O. Rog, and J. P. Cooper. 2006. Semi-conservative DNA replication through telomeres requires Taz1. *Nature* **440**:824–828.
28. Monahan, B. J., J. Villen, S. Marguerat, J. Bahler, S. P. Gygi, and F. Winston. 2008. Fission yeast SWI/SNF and RSC complexes show compositional and functional differences from budding yeast. *Nat. Struct. Mol. Biol.* **15**:873–880.
29. Moreno, S., A. Klar, and P. Nurse. 1991. Molecular genetic analysis of fission yeast *Schizosaccharomyces pombe*. *Methods Enzymol.* **194**:795–823.
30. Nicolas, E., T. Yamada, H. P. Cam, P. C. Fitzgerald, R. Kobayashi, and S. I. Grewal. 2007. Distinct roles of HDAC complexes in promoter silencing, antisense suppression and DNA damage protection. *Nat. Struct. Mol. Biol.* **14**:372–380.
31. Nourani, A., F. Robert, and F. Winston. 2006. Evidence that Spt2/Sin1, an HMG-like factor, plays roles in transcription elongation, chromatin structure, and genome stability in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **26**:1496–1509.
32. Pelletier, B., J. Beaudoin, Y. Mukai, and S. Labbe. 2002. Fep1, an iron sensor regulating iron transporter gene expression in *Schizosaccharomyces pombe*. *J. Biol. Chem.* **277**:22950–22958.
33. Pelletier, B., J. Beaudoin, C. C. Philpott, and S. Labbe. 2003. Fep1 represses expression of the fission yeast *Schizosaccharomyces pombe* siderophore-iron transport system. *Nucleic Acids Res.* **31**:4332–4344.
34. Phelps-Durr, T. L., J. Thomas, P. Vahab, and M. C. Timmermans. 2005. Maize rough sheath2 and its *Arabidopsis* orthologue ASYMMETRIC LEAVES1 interact with HIRA, a predicted histone chaperone, to maintain *knox* gene silencing and determinacy during organogenesis. *Plant Cell* **17**:2886–2898.
35. Prochasson, P., L. Florens, S. K. Swanson, M. P. Washburn, and J. L. Workman. 2005. The HIR corepressor complex binds to nucleosomes generating a distinct protein/DNA complex resistant to remodeling by SWI/SNF. *Genes Dev.* **19**:2534–2539.
36. Prudden, J., J. S. Evans, S. P. Hussey, B. Deans, P. O'Neill, J. Thacker, and T. Humphrey. 2003. Pathway utilization in response to a site-specific DNA double-strand break in fission yeast. *EMBO J.* **22**:1419–1430.
37. Qian, Z., H. Huang, J. Y. Hong, C. L. Burck, S. D. Johnston, J. Berman, A. Carol, and S. W. Liebman. 1998. Yeast Ty1 retrotransposition is stimulated by a synergistic interaction between mutations in chromatin assembly factor I and histone regulatory proteins. *Mol. Cell. Biol.* **18**:4783–4792.
38. Ray-Gallet, D., J. P. Quivy, C. Scamps, E. M. Martini, M. Lipinski, and G. Almouzni. 2002. HIRA is critical for a nucleosome assembly pathway independent of DNA synthesis. *Mol. Cell* **9**:1091–1100.
39. Sehgal, A., C. Y. Lee, and P. J. Espenshade. 2007. SREBP controls oxygen-dependent mobilization of retrotransposons in fission yeast. *PLoS Genet.* **3**:e131.
40. Sharp, J. A., E. T. Fouts, D. C. Krawitz, and P. D. Kaufman. 2001. Yeast histone deposition protein Asf1p requires Hir proteins and PCNA for heterochromatic silencing. *Curr. Biol.* **11**:463–473.
41. Sherwood, P. W., and M. A. Osley. 1991. Histone regulatory (*hir*) mutations suppress delta insertion alleles in *Saccharomyces cerevisiae*. *Genetics* **128**:729–738.
42. Sims, R. J., III, and D. Reinberg. 2008. Is there a code embedded in proteins that is based on post-translational modifications? *Nat. Rev. Mol. Cell Biol.* **9**:815–820.
43. Smith, S., and B. Stillman. 1989. Purification and characterization of CAF-I, a human cell factor required for chromatin assembly during DNA replication in vitro. *Cell* **58**:15–25.
44. Spector, M. S., A. Raff, H. DeSilva, K. Lee, and M. A. Osley. 1997. Hir1p and Hir2p function as transcriptional corepressors to regulate histone gene transcription in the *Saccharomyces cerevisiae* cell cycle. *Mol. Cell. Biol.* **17**:545–552.
45. Sugiyama, T., H. P. Cam, R. Sugiyama, K. Noma, M. Zofall, R. Kobayashi, and S. I. Grewal. 2007. SHREC, an effector complex for heterochromatic transcriptional silencing. *Cell* **128**:491–504.
46. Tagami, H., D. Ray-Gallet, G. Almouzni, and Y. Nakatani. 2004. Histone H3.1 and H3.3 complexes mediate nucleosome assembly pathways dependent or independent of DNA synthesis. *Cell* **116**:51–61.
47. Wang, S. W., A. L. Stevenson, S. E. Kearsey, S. Watt, and J. Bahler. 2008. Global role for polyadenylation-assisted nuclear RNA degradation in post-transcriptional gene silencing. *Mol. Cell. Biol.* **28**:656–665.
48. Williams, S. K., and J. K. Tyler. 2007. Transcriptional regulation by chromatin disassembly and reassembly. *Curr. Opin. Genet. Dev.* **17**:88–93.
49. Zhang, R., M. V. Poustovoitov, X. Ye, H. A. Santos, W. Chen, S. M. Daganzo, J. P. Erzberger, I. G. Serebriiskii, A. A. Canutescu, R. L. Dunbrack, J. R. Pehrson, J. M. Berger, P. D. Kaufman, and P. D. Adams. 2005. Formation of MacroH2A-containing senescence-associated heterochromatin foci and senescence driven by ASF1a and HIRA. *Dev. Cell* **8**:19–30.
50. Zurlinden, A., and M. E. Schweingruber. 1994. Cloning, nucleotide sequence, and regulation of *Schizosaccharomyces pombe thi4*, a thiamine biosynthetic gene. *J. Bacteriol.* **176**:6631–6635.