



Published in final edited form as:

J Am Chem Soc. 2009 September 9; 131(35): 12576–12578. doi:10.1021/ja9058926.

Improved Dienophilicity of Nitrocycloalkenes: Prospects for the Development of a *trans*-Diels-Alder Paradigm

Woo Han Kim[†], Jun Hee Lee[‡], and Samuel J. Danishefsky^{*,†,‡}

[†]Laboratory for Bioorganic Chemistry, Sloan-Kettering Institute for Cancer Research, 1275 York Avenue, New York, NY 10065

[‡]Department of Chemistry, Columbia University, Havemeyer Hall, 3000 Broadway, New York, NY 10027

Abstract

The development of an efficient and stereoselective *trans*-Diels-Alder paradigm is described. A central element of this transformation is the introduction of a temporary dienophilic functionality (A), which serves both to activate the substrate for Diels-Alder cycloaddition and, through its subsequent removal, to facilitate conversion of the *cis*-fused cycloadducts to the *trans*-fused series.

The importance of the Diels-Alder reaction can hardly be exaggerated. Its many features have been widely discussed.¹ The ability to reach *cis*-fused bicyclic systems by a Diels-Alder pathway has been a bulwark of complex target-oriented synthesis.² We well recognize that strategy in chemical synthesis is primarily beholden to capabilities arising from advances in methodology. This said, strategy also involves an intrinsic cognitive element. It connects a particular target to the huge database of organic chemistry, seeking to apply its most relevant and salient features to the challenge at hand. Central in this regard has been the logic of retrosynthetic analysis, which prioritizes various bond disconnections in an inverse sequential manner.³ The attainments resulting from strategic bond retrosynthetic analysis are legion.

Complementary to the powerful thought process of bond disconnections, we have been entertaining a different approach which we term “pattern analysis.”⁴ Although these approaches have some commonality, pattern analysis emphasizes a holistic view of the target structure, seeking connectivity between its key substructural characteristics (sometimes obvious but sometimes quite subtle) and established or prospectively implementable pathways. For instance, in pattern analysis, the possibility of a Diels-Alder (DA) application is provoked by the recognition of a *cis* junction. By contrast, *trans* junctions tend to teach away⁵ from a Diels-Alder universe.

It goes without saying that the scope of pattern analysis would be dramatically expanded if targets containing *trans* junctions could also be encompassed in the general Diels-Alder logic. The simplest scenario would contemplate an antarafacial cycloaddition.⁶ Not having any constructive suggestions in this regard, we turned to the next best thing. As outlined in Scheme 1, we envisioned equipping an otherwise unreactive dienophile (such as cyclohexene) with a temporary, readily removable activating group (A). Following cycloaddition, a product of the type **2** would be obtained. The resultant *cis*-fused substructure would be diverted to the *trans* series (**1**) through excision of the activating moiety and its controlled replacement. The traceless

s-danishefsky@ski.mskcc.org.

Supporting Information Available: Experimental procedures, copies of spectral data, and characterization (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>

function (A) would first serve the purpose of activating the dienophile for cycloaddition, thereby affording a *cis* junction. Subsequent chemistry would provide entry to the *trans*-fused series. We selected a nitro group as (A), based on its dienophile activating properties,⁷ its known success in controlling DA regioselectivity,⁸ and its ability to generate free radical intermediates.⁹ We describe herein the use of nitrocycloalkenes as dienophiles, pointing toward an emerging *trans*-Diels-Alder paradigm.

We began by studying the dienophilicity of compound **3**. At the outset, there had been only one evaluation of it as a dienophile in a Diels-Alder reaction.¹⁰ In the event, exploitation of its reactivity in cycloadditions with hydrocarbon dienes (see entry 1-4, Table 1) proved to be a particularly challenging problem. At temperatures below ~120-130 °C, there was little or no indication of cycloaddition. At temperatures above 140 °C there seemed to be very serious deterioration of the dienophile, resulting in extremely low yields of cycloadduct. We were, however, able to identify a narrow temperature range in which marginally useful cycloaddition could be achieved. Still, in hydrocarbon solvents such as toluene or on THF, the yields of isolated cycloadducts were quite low, apparently reflecting competition between the innate destruction of **3**, versus its cycloaddition.

During the course of these studies, we did note some improvement of cycloaddition yield in alcoholic solvents¹¹ (see entry 5, Table 1). Fortunately a more substantial increase in yield was realized when we turned to 2,2,2-trifluoroethanol as solvent, particularly when thermolysis was conducted under microwave conditions (entry 7). We emphasize, however, that the hydroxylic solvent effect, including 2,2,2-trifluoroethanol, apparently does not arise from catalysis (through hydrogen bonding or via other means). Thus, we see no indication that cycloaddition occurs at lower temperatures with 2,2,2-trifluoroethanol. Rather, the differences in yield seem to reflect a more favorable distribution of cycloaddition relative to decomposition. While the basis of this effect remains to be explored, it was very helpful to our program. Reasonable yields of cycloaddition from compound **3** could now be achieved. Happily, the protocol which was worked out for 2,3-dimethyl-1,3-butadiene was extendable to several other dienes (see Table 1). Moreover, we prepared 1-nitrocyclopentene (**4**)¹⁰ and evaluated its dienophilicity with the same dienes. Yields for these reactions are also included in Table 1. Here, we note that the temperatures required for cycloaddition were substantially lower (ca 100-110 °C) for **4** than for **3**, though, for the moment, the isolated yields tend to be similar. Table 1 summarizes yields for various conditions under which cycloaddition of dienophiles **3** and **4** with various acyclic dienes could be achieved.

Having substantially advanced the practicality of cycloadditions with nitrocycloalkene dienophiles, we next faced the central issue of denitration. We well recognized that if reductive denitration would lead to a *cis* junction, the nitrocycloalkene dienophile would have functioned as an equivalent of otherwise unreactive cyclohexene or cyclopentene.¹² However, if the excision-replacement sequence would afford a *trans* junction, an overall *trans*-Diels-Alder type paradigm would have been demonstrated.

In the event, we began by treating compound **5a** with tri-*n*-butyltin hydride in the presence of AIBN. We noted an 8:1 ratio of *trans*-**6a** to *cis*-**7a**, as established by comparison with authentic reference compounds prepared in multistep sequences.¹³ In a similar way, the isoprene adduct **5b** also gave rise to an 8:1 ratio of *trans*-**6b** to *cis*-**7b**. Other examples are shown in Table 2. Thus, while one could hope for even stronger stereoselectivity, these data encourage application of Diels-Alder logic to the synthesis of *trans* octalins.

Perhaps, this selectivity pattern represents a preference for conformer type **A** relative to conformer **B** thereby accounting for the selective formation of *trans* product (Figure 1).

However, in the hydrindene series, the ratio of the denitration products, was much closer to unity, presumably reflecting altered *trans:cis* preferences.

In the next step in our exploration, we evaluated the consequences of using a more functionalized diene. In that way, additional exploitable functionality would be delivered to the eventual denitration product. We began by studying the cycloaddition reaction of compound **3** with diene **8**.¹⁴ Fortunately, our conditions described above for cycloaddition with acyclic hydrocarbon dienes work quite well with diene **8**. In 2,2,2-trifluoroethanol at 80 °C, a 61% yield of adduct **9** was obtained.¹⁵ With 1-nitrocyclopentene (**4**), cycloaddition under comparable conditions afforded adduct **12** (Scheme 2).

We then turned to denitration reactions. For each compound, this process was studied in two different sequences. In one arm, the silyl enol ethers were hydrolyzed and the subsequent ketones were denitrated. In this case, a 5:1 ratio of *trans* compound **10** to *cis* compound **11** was obtained. A very similar result pertained when we reversed the order of steps, wherein denitration was carried out at the stage of adduct **9** and followed by hydrolysis of the silyl enol ether, to provide a 6:1 ratio of *trans*-**10** : *cis*-**11**.¹⁶

We applied the same methodology to adduct **12**, which arose from **4**. Once again, both denitration sequences were pursued. When denitration was conducted at the stage of the silyl enol ether, followed by hydrolysis, a 1.4:1 ratio of *trans*-**14**¹⁷ to *cis*-**13**¹⁸ was obtained. By contrast, when denitration was conducted at the ketone stage (i.e. after hydrolysis), a 15:1 ratio of *cis*-**13** to *trans*-**14** was produced. Thus, once again, in the octalin series a stereoselective route to *trans*-fused ketone **10** has been established. By contrast, in the hydrindane series, a stereoselective route to the *cis* fusion has been realized. Unfortunately, no corresponding protocol now available to us provides a *trans* hydrindanone junction with useful levels of stereoselection (*vide infra*).

Finally, we turned our attention to the cycloaddition reaction of diene **15**¹⁹ with 1-nitrocyclohexene (**3**). As it turned out, this reaction could not be conducted in 2,2,2-trifluoroethanol because of rapid conversion of the diene to the corresponding ketone, methoxybutenone. Accordingly, the reaction was conducted in xylene under reflux, as shown in Scheme 3. This treatment gave rise to a mixture of *endo* and *exo* Diels-Alder product, indicated as **16**. Again, we examined the stereochemical outcome of the denitration reaction when conducted at one of the two different stages. Thus, when reduction was conducted at the stage of the ketone **16** (obtained by prior hydrolysis of the silyl enol ether function), a 2:1 ratio of *trans:cis* octalones was obtained. In contrast, when the denitration was conducted at the silyl enol ether stage and the denitration product subjected to acid hydrolysis of the silyl enol ether, an 8:1 ratio of *trans*-**17** to *cis*-**18** was produced.²⁰ Once again, we see how the chemistry described above points the way to introduction of *trans* ring junctions in Diels-Alder driven constructions.

In summary then, while many provocative possibilities remain to be explored, the chemistry described above already holds promise for application to two types of situations. In the hydrindane series, denitration can be used to generate a *cis* fusion, thereby establishing a Diels-Alder equivalency for the otherwise inert cyclopentene. By contrast in the octalin series, it is possible to take advantage of this chemistry to produce, selectively, *trans* junctions. In that sense, the nitrocyclohexene will have served in a Diels-Alder context as an equivalent of the otherwise unavailable dienophile, *E*-cyclohexene. Studies addressing a menu of follow-up possibilities suggested by these findings are in progress.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Support was provided by the NIH (HL25848 and CA103823 to SJD). W.H.K. is grateful for a Korea Research Foundation Grant funded by the Korean government (KRF-2007-357-c00060). We thank Rebecca Wilson and Dana Ryan for assistance with the preparation of the manuscript and Prof. W. F. Berkowitz and Dr. Pavel Nagorny for helpful discussions. We also thank Dr. George Sukenick, Ms. Hui Fang, Sylvi Rusli (NMR Core Facility, Sloan-Kettering Institute) and Dr. Yasuhiro Itagaki (Mass Spectral Core Facility, Columbia University) for mass spectral and NMR spectroscopic analysis.

References

1. (a) Corey EJ. *Angew Chem Int Ed* 2009;48:2100–2117. (b) Corey EJ. *Angew Chem Int Ed* 2002;41:1650–1667. (c) Bear BR, Sparks SM, Shea KJ. *Angew Chem Int Ed* 2001;40:820–849. (d) Williams RM, Stocking EM. *Angew Chem Int Ed* 2003;42:3078–3115. (e) Jørgensen KA. *Angew Chem Int Ed* 2000;39:3558–3588.
2. (a) Nicolaou KC, Snyder SA, Montagnon T, Vassilikogiannakis G. *Angew Chem Int Ed* 2002;41:1668–1698. (b) Danishefsky S. *Acc Chem Res* 1981;14:400–406.
3. (a) Corey EJ. *Pure Appl Chem* 1967:19–37. (b) Corey EJ, Wipke WT. *Science* 1969;166:178–192. [PubMed: 17731475] (c) Corey EJ. *Angew Chem Int Ed* 1991;30:455–612. Corey, EJ.; Chelg, X. *The Logic of Chemical Synthesis*. Wiley-VCH; New York: 1995.
4. (a) Wilson RM, Danishefsky SJ. *Acc Chem Res* 2006;39:539–549. [PubMed: 16906750] (b) Wilson RM, Danishefsky SJ. *J Org Chem* 2007;72:4293–4305. [PubMed: 17539594]
5. The possibility of epimerizing a *cis* junction, obtained from a Diels Alder reaction, to the *trans* series was well recognized and practiced by Woodward and co-workers in their total synthesis of steroids. Other instances of epimerization are also cited herein. (a) Woodward RB, Sondheimer F, Taub D, Heusler K, McLamore WM. *J Am Chem Soc* 1952;74:4223–4251.4251 (b) Stevens RV, Angle SR, Kloc K, Mak KF, Trueblood KN, Liu YX. *J Org Chem* 1986;51:4347–4353.4353 and references therein. (c) Mukhopadhyay A, Ali SM, Husain M, Suryawanshi SN, Bhakuni DS. *Tetrahedron Lett* 1989;30:1853–1856.1856 Of course, this approach requires a vicinal keto group to enable epimerization and the capacity to apply strict thermodynamic or kinetic controls to govern the junction stereochemistry.
6. Woodward RB, Hoffmann R. *Angew Chem Int Ed* 1969;8:781–932.
7. Ono, N. *The Nitro Group in Organic Synthesis*. Wiley-VCH; New York: 2001. and pertinent references cited therein
8. (a) Danishefsky S, Prisbylla MP, Hiner S. *J Am Chem Soc* 1978;100:2918–2920. (b) Corey EJ, Estreicher H. *Tetrahedron Lett* 1981;22:603–606.
9. (a) Ono N, Miyake H, Tamura R, Kaji A. *Tetrahedron Lett* 1981;22:1705–1708. (b) Ono N, Miyake H, Kamimura A, Hamamoto I, Tamura R, Kaji A. *Tetrahedron* 1985;41:4013–4023. (c) Ono N, Miyake H, Kaji A. *J Chem Soc Chem Commun* 1982:33–34.
10. Corey EJ, Estreicher H. *J Am Chem Soc* 1978;100:6294–6295.
11. Breslow R, Guo TJ. *Am Chem Soc* 1988;110:5613–5617.
12. Following completion of this work, we noted that in footnote (18) in reference (10), Professor Corey futuristically envisioned potential value of reductive denitration at the junction. However no results are provided. Nonetheless the kernel of the idea can be found in Corey's comment.
13. Lee JH, Kim WH, Danishefsky SJ. *Tetrahedron Lett*. 2009;10.1016/j.tetlet.2009.07.068 In Press
14. Jung ME, McCombs CA. *Tetrahedron Lett* 1976;17:2935–2938.
15. Cycloaddition of compound 3 with several 2-silyloxy-1,3-butadienes as well as diene 8 in toluene under the variety of conditions did not proceed at all.
16. (a) Oritani T, Yamashita K. *J Org Chem* 1984;49:3689–3694. (b) Rao HSP, Reddy KS. *Tetrahedron Lett* 1994;35:171–174. (c) Augustine RL. *J Org Chem* 1958;23:1853–1856. (d) Macdonald TL, Mahalingam S. *J Am Chem Soc* 1980;102:2113–2115.
17. Krawczyk AR, Jones JB. *J Org Chem* 1989;54:1795–1801.
18. Okuda Y, Morizawa Y, Oshima K, Nozaki H. *Tetrahedron Lett* 1984;25:2483–2486. and pertinent references cited therein
19. Danishefsky S, Kitahara T. *J Am Chem Soc* 1974;96:7807–7808.

20. (a) Corey EJ, Boger DL. *Tetrahedron Lett* 1978;19:4597–4600. (b) Angell EC, Fringuelli F, Pizzo F, Taticchi A, Wenkert E. *J Org Chem* 1988;53:1424–1426.







Figure 1.
Structures of the tertiary bridge head radical.



Scheme 1.
Trans-Diels-Alder Paradigm.

**Scheme 2a.**


^aKey: (a) CF₃CH₂OH, 80 °C, MW, 12 h, 61%. (b) HF, CH₃CN, rt, 30 min. (c) *n*Bu₃SnH, AIBN, benzene, reflux, 2 h. (d) CF₃CH₂OH, 80 °C, MW, 6 h, 72%.

**Scheme 3a.**

^aKey: (a) xylene, reflux, 36 h. (b) 0.05 M HCl, THF, 2 h, 67% for 2 steps. (c) *n*Bu₃SnH, AIBN, benzene, reflux, 2 h, 89%. (d) TFAA, benzene, Dean-Stark, 24 h, 60%. (e) toluene, 130 °C, 36 h, 90%. (f) *n*Bu₃SnH, AIBN, benzene, reflux, 2 h. (g) HF, CH₃CN, rt, 10 min, 53% for 2 steps.

Diels-Alder Reaction of Nitrocycloalkenes.

Table 1



entry	5	n	diene	solvent	T (°C)	T (h)	yield (%) ^d
1	a	1	R ¹ =R ² =CH ₃	toluene ^b	110	12	trace
2	a	1	R ¹ =R ² =CH ₃	toluene ^b	140	36	18
3	a	1	R ¹ =R ² =CH ₃	toluene ^b	150	36	trace
4	a	1	R ¹ =R ² =CH ₃	THF	130 MW ^c	12	17
5	a	1	R ¹ =R ² =CH ₃	CH ₃ CH ₂ OH	130 MW ^c	14	36
6	a	1	R ¹ =R ² =CH ₃	CF ₃ CH ₂ OH	130	24	70
7	a	1	R ¹ =R ² =CH ₃	CF ₃ CH ₂ OH	130 MW ^c	12	78
8	b	1	R ¹ =H, R ² =CH ₃	CF ₃ CH ₂ OH	125 MW ^c	14	57 ^{d,e}
9	c	1	R ¹ =R ² =(CH ₂) ₄	CF ₃ CH ₂ OH	130 MW ^c	10	75
10	d	1	R ¹ =R ² =Ph	toluene ^{b,f}	150	48	45
11	e	0	R ¹ =R ² =CH ₃	CF ₃ CH ₂ OH	100	12	78
12	f	0	R ¹ =H, R ² =CH ₃	CF ₃ CH ₂ OH	110	12	77 ^g

^(d) Isolated yield.^(b) In the presence of 2,6-di-*tert*-butylresol (5 mol%).^(c) MW=microwave.^(d) *p*-directed/*m*-directed=15:1.^(e) The reaction in toluene gave a 1:1 mixture of regioisomers as a 5% yield

(f) The diene was insoluble in $\text{CF}_3\text{CH}_2\text{OH}$.

(g) *p*-directed/*m*-directed=11:1.

Radical Denitration.

Table 2

trans = **6a-f**
cis = **7a-f**

entry	6,7	n	diene	yield (%) ^d	6:7 ^b
1	a	1	R ¹ =R ² =CH ₃	67	8:1
2	b	1	R ¹ =H, R ² =CH ₃	53	8:1
3	c	1	R ¹ =R ² =(CH ₂) ₄	81	12:1 ^c
4 ^d	d	1	R ¹ =R ² =Ph	83	7:1
5	e	0	R ¹ =R ² =CH ₃	61	1.3:1
6	f	0	R ¹ =H, R ² =CH ₃	35 ^d	1.9:1

^(a) Isolated yield.^(b) Determined by ¹H-NMR.^(c) Determined by GC.^(d) The compound is volatile.