

New Medium for the Isolation of *Streptococcus mutans* and Its Differentiation from Other Oral Streptococci

HARALD A. B. LINKE

Department of Microbiology, New York University Dental Center, New York, New York 10010

Received for publication 14 February 1977

A new, improved agar medium for the isolation of *Streptococcus mutans*, the etiological agent of dental caries, was developed. In contrast to mitis-salivarius agar, this medium not only recovers a greater number of *S. mutans* strains from most oral specimens but, because of its mannitol and sorbitol content, it also facilitates the differentiation of *S. mutans* from other oral streptococci, e.g., *S. salivarius*, *S. mitis*, and *S. sanguis*, which do not grow or produce scanty growth only after 10 days of incubation. The medium is easy to prepare because of its simple and unique composition, is characterized by the presence of an acid indicator, and can be utilized under aerobic and anaerobic conditions as well. The medium cannot be used to distinguish among the eight serotypes, *a* to *g* and SL-1, of *S. mutans*. Mannitol-utilizing bacteria such as streptococci (e.g., *S. faecalis*) and other microorganisms (e.g., *Staphylococcus aureus*) are able to grow on this medium and can be distinguished from *S. mutans* by their unique colony morphology.

Streptococcus mutans routinely is isolated together with other oral streptococci on mitis-salivarius (MS) agar. *S. mutans* is differentiated from the other oral streptococci by its ability to ferment mannitol and sorbitol and to adhere to smooth surfaces in the presence of sucrose (17, 19). The recovery of *S. mutans* from MS agar is never quantitative (5) and depends on the origin of MS agar (12, 18). In the following study we utilized the unique ability of *S. mutans* among oral streptococci to ferment mannitol and sorbitol to develop a new isolation and differentiation medium specifically for this *Streptococcus* species.

MATERIALS AND METHODS

Bacterial strains and isolation specimen. For comparative growth studies, 28 defined *Streptococcus* strains were utilized: 13 *S. mutans* strains (Table 1), representing the eight serotypes, *a* to *g* and SL-1, according to Bratthall (1) and Perch et al. (14); and 15 other streptococcal strains (Table 1) typical of viridans (2 *S. sanguis*, 3 *S. salivarius*, and 3 *S. mitis* strains), enterococcus (2 *S. faecalis*, 1 *S. faecalis* subsp. *liquefaciens*, and 1 *S. durans* strain), pyogenic (1 *S. pyogenes* and 1 *S. agalactiae* strain) and lactic groups (1 *S. lactis* strain). For enumeration of unknown microbial strains, a specimen was used that was composed of 3 ml of saliva mixed with 100 mg of plaque material (from a human source) diluted with 9 ml of distilled water.

Media and chemicals. MSFA agar, used for isolation and differentiation of *S. mutans*, was composed of: D-sorbitol (Sigma Chemical Co., St. Louis, Mo.),

10 g; D-mannitol (Sigma), 10 g; yeast extract (Difco Laboratories, Detroit, Mich.), 20 g; sodium azide (Fisher Scientific Co., Fair Lawn, N.J.), 100 mg; basic fuchsin (Allied Chemical Co., Morristown, N.J.), 5 mg; CaCO₃ (precipitated; Merck & Co., Inc., Rahway, N.J.), 10 g; agar (Fisher Scientific Co.), 15 g; and distilled water, 1,000 ml; the pH was adjusted to 7.0. MS agar (Difco) was prepared according to the manufacturer's recommendation. Blood agar was prepared by mixing tryptic soy agar (Difco), prepared according to the manufacturer's recommendation, with 50 ml of sheep blood per 1,000 ml. Trypan blue for growth inhibition studies was obtained from Matheson Coleman & Bell, Norwood, Ohio, and crystal violet was obtained from Difco. The *Streptococcus* strains were maintained in screw-cap test tubes containing 9 ml of a medium containing 2% glucose (dextrose, Difco) and 2% yeast extract (Difco) at pH 6.5, sterilized in an autoclave for 10 min at 20 lb/in² and 126°C. The tubes were supplemented with 10% inoculum and incubated at 28°C. The cultures were transferred every 5 to 6 days.

Growth and enumeration of bacterial strains. The agar plates were inoculated with each of the 28 *Streptococcus* strains by streaking 0.1 ml of a suspension onto the plate, cutting into the agar surface at the last 0.5 inch (ca. 1.27 cm) of the streak. The plates were incubated aerobically at 36°C for up to 2 weeks. Growth was observed at intervals of 4 days. For anaerobic growth conditions, the plates were incubated in an anaerobic jar (BBL, Cockeysville, Md.) in the presence of a gas-generating envelope (GasPak no. 70304, BBL) and an anaerobic indicator strip (GasPak no. 70404, BBL) for 36°C for up to 2 weeks. For enumeration of the unknown *Streptococcus* strains, a dilution series of the aqueous saliva-

TABLE 1. *The 28 Streptococcus species studied*

| Strain | Serotype | Origin | Source |
|---|------------------|---------|--|
| <i>S. mutans</i> | | | |
| NCTC 10449 | <i>c</i> | Human | H. D. Slade, Chicago, Ill. |
| AHT | <i>a</i> | Human | |
| B-13 | <i>d</i> | Human | |
| B-14 | <i>e</i> | Human | |
| FA-1 | <i>b</i> | Rat | |
| SL-1 | SL-1/ <i>d/g</i> | Human | |
| OMZ-176 | <i>d</i> | Human | B. Guggenheim, Zürich, Switzerland |
| OMZ-65 | <i>g</i> | Human | |
| OMZ-175 | <i>f</i> | Human | |
| BHT | <i>b</i> | Human | D. D. Zinner, Miami, Fla. |
| HS-6 | <i>a</i> | Hamster | R. J. Fitzgerald, Miami, Fla. |
| P-4 | <i>e</i> | Human | S. Edwardsson, Malmö, Sweden |
| B-2 | <i>e</i> | Human | |
| <i>S. mitis</i> ATCC 15909 | | | American Type Culture Collection, Rockville, Md. |
| <i>S. mitis</i> ATCC 15910 | | | |
| <i>S. salivarius</i> ATCC 9759 | | | |
| <i>S. salivarius</i> ATCC 13419 | | | |
| <i>S. sanguis</i> ATCC 10556 | | | |
| <i>S. sanguis</i> ATCC 10557 | | | |
| <i>S. faecalis</i> ATCC 19433 | | | |
| <i>S. faecalis</i> subsp. <i>liquefaciens</i> | | | Midwest Culture Service, Terre Haute, Ind. |
| <i>S. pyogenes</i> | | | |
| <i>S. durans</i> | | | |
| <i>S. agalactiae</i> | | | |
| <i>S. lactis</i> | | | |
| <i>S. salivarius</i> | | | |
| <i>S. mitis</i> | | | |
| <i>S. faecalis</i> | | | |

plaque material mixture was prepared, and 0.2-ml portions were evenly distributed, with the aid of a bent glass rod, on top of the agar plates, which were incubated at 36°C for up to 2 weeks. Then the colonies were counted with the aid of a colony counter (model C-110; New Brunswick Scientific Co., New Brunswick, N.J.).

Growth inhibitions studies. Stock solutions of sodium azide, basic fuchsin, crystal violet, and trypan blue were prepared and sterilized in an autoclave; the solutions were then diluted in sterile water to yield the desired inhibitor concentrations. For each test, 1.0 ml of the appropriate inhibitor dilution and 0.5 ml of sterile water were added to 5.0 ml of a 2% glucose-2% yeast extract medium at pH 6.5 in screw-cap test tubes. The final concentrations of sodium azide, basic fuchsin, crystal violet, and trypan blue in the medium were 20 to 1,000, 1 to 500, 1 to 500, and 1 to 500 µg/ml, respectively. Each tube was then inoculated with 0.5 ml of a 48-h culture of the appropriate *Streptococcus* strain to yield a total volume of 7.0 ml. Inoculated tubes with 1.5 ml of sterile water added to the glucose-yeast extract medium served as controls. The test tubes were incubated in a gyratory water bath shaker (model G76, New Brunswick

Scientific Co.) at 36°C. After 24 h the optical density, as a measure of growth, was determined at 546 nm with a Bausch & Lomb Spectronic 20 spectrophotometer. For each inhibitor dilution, a blank tube was prepared to aid adjustment of the spectrophotometer's transmission.

RESULTS

Growth inhibition studies. Most *Streptococcus* species are relatively resistant to sodium azide treatment (Fig. 1); some strains are still able to survive the presence of 1,000 µg/ml. At a concentration of 200 µg/ml, on the average, a 50% growth inhibition was obtained with all tested strains.

A significant growth inhibition of *S. mutans*, but not of *S. faecalis*, by basic fuchsin was obtained at a concentration of 10 µg/ml (Fig. 2); concentrations lower than 5 µg/ml had no or little effect on streptococcal growth.

At a concentration of 10 µg/ml, crystal violet inhibited most of the tested *Streptococcus* strains (Fig. 3); even 1 µg of this compound per




FIG. 1. Growth inhibition (optical density [OD] at 546 nm) of glucose-grown *Streptococcus* strains in the presence of sodium azide.




FIG. 2. Growth inhibition (optical density [OD] at 546 nm) of glucose-grown *Streptococcus* strains in the presence of basic fuchsin.




FIG. 3. Growth inhibition (optical density [OD] at 546 nm) of glucose-grown *Streptococcus* strains in the presence of crystal violet.

ml inhibited the growth of all tested strains up to 25%.

Growth of the *Streptococcus* strains in the presence of trypan blue concentrations higher

than 50 µg/ml could not be determined under the experimental conditions. Three of the tested *S. mutans* strains, P-4, HS-6, and NCTC 10449, were almost completely growth inhibited by a concentration of 50 µg/ml (Fig. 4); therefore, a trypan blue concentration of 75 µg/ml, as used in MS agar preparations, is definitely not desirable for isolation and growth of many *S. mutans* strains. A significant reduction in streptococcal growth, up to 35%, was obtained at a trypan blue concentration of 10 µg/ml (Fig. 4).

Growth on MSFA agar and MS agar. Most of the studied *S. mutans* strains grew much better on MSFA agar than on MS agar (Table 2). The other *Streptococcus* species that grew well on MS agar, e.g., *S. salivarius*, *S. mitis*, and *S. sanguis*, did not grow or produced only scanty growth on MSFA agar. On MSFA agar most streptococci colonies were pink to red, some having a purple to blue tint. The colonies were often pinpoint to small and were shiny or had a silky to frosted-glass appearance; some appeared to be raspberry-like. *S. faecalis* grew in large mat colonies, often with a center indentation. Good growth on MSFA agar developed usually after 1 week of incubation, but, in the cases of *S. salivarius*, *S. mitis*, and *S. sanguis*, scanty growth appeared after an incubation period of 10 days to 2 weeks.

Total viable counts of bacteria on various growth media. A dilution series of the saliva-plaque material specimen was prepared, and portions of the specimen dilutions were plated on the three different media: blood agar, MS agar, and MSFA agar. Results in Table 3 represent readings from plates that received inoculum from the 10⁻⁴ dilution. Blood agar produced the highest viable count of bacteria (=100%), followed by MS agar, which recovered 84% of




FIG. 4. Growth inhibition (optical density [OD] at 546 nm) of glucose-grown *Streptococcus* strains in the presence of trypan blue.

TABLE 2. Growth of 28 *Streptococcus* strains on the new isolation medium (MSFA agar) and on MS agar

| Strain | Growth ^a on MSFA agar | Colony characteristics | Growth ^a on MS agar |
|---|----------------------------------|----------------------------|--------------------------------|
| <i>S. mutans</i> | | | |
| OMZ-175 | ++ | Pink to red, frosted glass | + |
| OMZ-176 | ++ | Pink, frosted glass | ++ |
| OMZ-65 | ++ | Pink, raspberry-like | + |
| SL-1 | +++ | Pink, smooth | + |
| HS-6 | ++ | Pink to red, frosted glass | + |
| B-13 | ++ | Pink, silky | (+) |
| B-14 | +++ | Red, silky | + |
| BHT | ++ | Pink, mat | ++ |
| AHT | +++ | Pink to red, frosted glass | ++ |
| NCTC 10449 | +++ | Blue-purple, silky | ++ |
| P-4 | ++ | Red, silky | ++ |
| FA-1 | ++ | Pink, silky | + |
| B-2 | ++ | Red, silky | + |
| <i>S. mitis</i> ATCC 15909 | (+) | Pink-red, mat | + |
| <i>S. mitis</i> ATCC 15910 | (+) | Pink-red, silky | +++ |
| <i>S. mitis</i> | + | Pink-red, mat | ++ |
| <i>S. salivarius</i> ATCC 9759 | - | | ++++ |
| <i>S. salivarius</i> ATCC 13419 | (+) | Pink-blue, smooth | +++++ |
| <i>S. salivarius</i> | + | Pink-blue, smooth | ++ |
| <i>S. sanguis</i> ATCC 10556 | - | | ++ |
| <i>S. sanguis</i> ATCC 10557 | (+) | Pink-blue, mat | + |
| <i>S. faecalis</i> ATCC 19433 | +++ | Pink-blue, indented | ++ |
| <i>S. faecalis</i> | ++ | Light pink, mat | ++ |
| <i>S. faecalis</i> subsp. <i>liquefaciens</i> | +++ | Pink, mat | +++ |
| <i>S. pyogenes</i> | (+) | Light pink, silky | - |
| <i>S. durans</i> | + | Pink, mat | ++ |
| <i>S. agalactiae</i> | + | Pink-red, mat | + |
| <i>S. lactis</i> | +++ | Pink-blue, silky | +++ |

^a Growth after 10 days of aerobic incubation at 36°C; colonies usually appeared circular and convex. Symbols: -, no growth; (+), scanty growth (haze of growth); +, light growth (pinpoint colonies); ++, moderate growth (colony size ca. 1 mm); +++, very good growth (colony size ca. 2 mm); +++++, excellent growth (colony size ca. 3 mm); ++++++, luxuriant growth (colony size > 3 mm).

TABLE 3. Comparison of total viable counts of bacteria from a saliva-plaque material sample (diluted 10⁻⁴) on various growth media

| Agar medium | No. of colonies/ml | Recovery (%) |
|-------------|--------------------|--------------|
| Blood | 155 | 100 |
| MS | 130 | 84 |
| MSFA | 120 | 77 |

the viable cells. Many colonies on MS resembled *S. salivarius* according to their large size and smooth convex appearance. On MSFA agar the recovery of colonies was 77% as compared with that on blood agar, or 92% as compared with that on MS agar. It was difficult to isolate *S. salivarius*, *S. mitis*, and *S. sanguis* colonies from MSFA agar because of scanty growth.

Growth of other microorganisms besides *Streptococcus* species on MSFA agar. Besides the *Streptococcus* species already mentioned, several other mannitol-utilizing microorganisms were able to grow on MSFA agar. *Lactobacillus casei* produced good growth in pink-blue mat colonies; *L. acidophilus* produced very

scanty growth. Under semianaerobic conditions, *Actinomyces israelii* grew well in spreading, tan, mat-scaly colonies; *Actinomyces viscosus* produced pink to red mat colonies under aerobic conditions. *Staphylococcus aureus* exhibited fair growth in mucoid, pink-blue, silky colonies on MSFA agar, and also *Escherichia coli* ATCC 25922, which grew in pink-red mat colonies. Furthermore, *Neisseria sicca* exhibited scanty growth in pink-red mat colonies with raspberry-like centers. Occasionally, diphtheroid colonies from saliva specimens could be recovered on MSFA agar.

DISCUSSION

S. mutans, the etiological agent of dental caries, was first discovered and described in 1924 (3). Since that time several investigators have isolated, from carious teeth of humans and animals, numerous *S. mutans* strains by using various complex growth media (7, 10, 20) including MS medium (11). MS agar is now commonly used to isolate *S. mutans* as well as other *Streptococcus* species from natural

sources. MS agar was modified to be more specific for the isolation of *S. mutans* by the addition of either sulfonamide (2), polymyxin (6), or bacitracin (9). By comparison of *S. mutans* growth on various selective media, different rates of recovery were observed (5). Different MS agar preparations from different manufacturers (BBL; Difco) recovered different numbers of *S. mutans* strains from the same specimen (12), and some *S. mutans* strains were even inhibited by these different MS agar preparations (18). This growth inhibition of MS agar preparations to certain strains of *S. mutans* was probably caused by the high (75 µg/ml) trypan blue content of the medium. We found that this large amount of trypan blue in MS agar definitely exerted a growth inhibitory effect on most *S. mutans* strains.

Modified MS agar preparations are also not very useful in the differentiation of *S. mutans* from other oral streptococci, e.g., *S. salivarius*, *S. sanguis*, and *S. mitis*; this differentiation is usually achieved after isolation of several *Streptococcus* spp. from MS agar and utilizes the ability of *S. mutans* to ferment mannitol and sorbitol and to adhere to smooth surfaces in the presence of sucrose (17, 19). Recently, it was found that some strains of *S. mutans* are not able to ferment sorbitol (4). *S. salivarius*, *S. sanguis*, and *S. mitis* ferment neither mannitol nor sorbitol.

On this basis we developed an isolation and differentiation medium (MSFA agar) for *S. mutans*, which contains the carbohydrates mannitol and sorbitol as carbon sources and yeast extract mainly as a nitrogen source. The addition of sodium azide and basic fuchsin provides some advantages over other isolation media (8); the former suppresses the growth of gram-negative microorganisms, including *Proteus*, and the latter reduces the development of undesirable gram-positive organisms. In addition, basic fuchsin has the advantage of producing colored colonies. Since *S. mutans* and other *Streptococcus* species are high acid producers (13, 15), some precipitated calcium carbonate was added to the medium to neutralize produced acids, which, in turn, enhanced the microbial growth on the agar surface. Due to the calcium carbonate content of the medium, acid-producing colonies appeared to be surrounded by a clear zone, a convenient visual check. Therefore, the new MSFA agar in comparison with MS medium is more useful for the isolation of *S. mutans* strains and its differentiation from other oral streptococci. In addition, the new medium is easy to prepare because of its simple and unique composition, is chemically stable, and has a long shelf life. The medium can be

utilized under both aerobic and anaerobic conditions.

MSFA agar allows growth of most mannitol-utilizing microorganisms, e.g., *S. faecalis*, *S. faecalis* subsp. *liquefaciens*, *Staphylococcus aureus*, and *Actinomyces* spp. The new medium cannot be utilized for biochemical determination of the eight serological groups, *a* to *g* and SL-1, of *S. mutans* according to Bratthall (1) and Perch et al. (14); previously a biochemical scheme was developed for the separation of *S. mutans* into five biotypes that correlated with the serotypes *a* to *e*, based on the utilization of mannitol with and without bacitracin and other biochemical tests (16).

ACKNOWLEDGMENTS

We express our gratitude to Donald A. M. Mackay of Life Savers, Inc., New York, for a grant-in-aid to support this study and to Heiner Hoffman for helpful discussions and valuable suggestions. The technical assistance of Byron Baker and Mariann Renna is gratefully appreciated, and we thank Sherry Johanas for preparation of the manuscript.

LITERATURE CITED

1. Bratthall, D. 1970. Demonstration of five serological groups of streptococcal strains resembling *Streptococcus mutans*. *Odontol. Revy* 21:143-152.
2. Carlsson, J. 1967. A medium for isolation of *Streptococcus mutans*. *Arch. Oral Biol.* 12:1657-1658.
3. Clarke, J. K. 1924. On the bacterial factor in the aetiology of dental caries. *Br. J. Exp. Pathol.* 5:141-147.
4. Edwardsson, S. 1970. The caries-inducing property of variants of *Streptococcus mutans*. *Odontol. Revy* 21:153-157.
5. Emilson, C. G., and D. Bratthall. 1976. Growth of *Streptococcus mutans* on various selective media. *J. Clin. Microbiol.* 4:95-98.
6. Fitzgerald, R. J., and B. O. Adams. 1975. Increased selectivity of Mitis-Salivarius agar containing polymyxin. *J. Clin. Microbiol.* 1:239-240.
7. Fitzgerald, R. J., H. V. Jordan, and H. R. Stanley. 1960. Experimental caries and gingival pathologic changes in the gnotobiotic rat. *J. Dent. Res.* 39:923-935.
8. Fitzgerald, R. J., and P. H. Keyes. 1960. Demonstration of the etiologic role of streptococci in experimental caries in the hamster. *J. Am. Dent. Assoc.* 61:9-19.
9. Gold, O. G., H. V. Jordan, and J. van Houte. 1973. A selective medium for *Streptococcus mutans*. *Arch. Oral Biol.* 18:1357-1364.
10. Guggenheim, B., K. G. König, and H. R. Mühlemann. 1965. Modifications of the oral bacterial flora and their influence on dental caries in the rat. I. The effects of inoculating 4 labelled strains of streptococci. *Helv. Odontol. Acta* 9:121-129.
11. Jordan, H. V., B. Krasse, and A. Möller. 1968. A method of sampling human dental plaque for certain "caries-inducing" streptococci. *Arch. Oral Biol.* 13:919-927.
12. Liljemark, W. F., D. H. Okrent, and C. G. Bloomquist. 1976. Differential recovery of *Streptococcus mutans* from various mitis-salivarius agar preparations. *J. Clin. Microbiol.* 4:108-109.
13. Linke, H. A. B., and C. A. Chang. 1976. Physiological effects of sucrose substitutes and artificial sweeteners on growth pattern and acid production of glucose-

- grown *Streptococcus mutans* strains *in vitro*. Z. Naturforsch. 31c:245-251.
14. Perch, B., E. Kjems, and T. Ravn. 1974. Biochemical and serological properties of *Streptococcus mutans* from various human and animal sources. Acta Pathol. Microbiol. Scand. Sect. B 82:357-370.
 15. Robrish, S. A., and M. I. Krichevsky. 1972. Acid production from glucose and sucrose by growing cultures of caries-conductive streptococci. J. Dent. Res. 51:734-739.
 16. Shklair, I. L., and H. J. Keene. 1974. A biochemical scheme for the separation of the five varieties of *Streptococcus mutans*. Arch. Oral Biol. 19:1079-1081.
 17. Shklair, I. L., H. J. Keene, and P. Cullen. 1974. The distribution of *Streptococcus mutans* on the teeth of two groups of naval recruits. Arch. Oral Biol. 19:199-202.
 18. Staat, R. H. 1976. Inhibition of *Streptococcus mutans* strains by different mitis-salivarius agar preparations. J. Clin. Microbiol. 3:378-380.
 19. Syed, S. A., and W. J. Loesche. 1973. Efficiency of various growth media in recovering oral bacterial flora from human dental plaque. Appl. Microbiol. 26:459-465.
 20. Zinner, D. D., J. M. Jablon, A. P. Aran, and M. S. Saslaw. 1965. Experimental caries induced in animals by streptococci of human origin. Proc. Soc. Exp. Biol. Med. 118:766-770.