



Published in final edited form as:

*J Org Chem.* 2009 September 4; 74(17): 6784–6791. doi:10.1021/jo901100m.

## Efficient Enantio- and Diastereo-divergent Synthesis of Poison-frog Alkaloids **251O** and *trans*-**223B**

Naoki Toyooka<sup>\*,1</sup>, Dejun Zhou<sup>1</sup>, Hideo Nemoto<sup>1</sup>, Yasuhiro Tezuka<sup>2</sup>, Shigetoshi Kadota<sup>2</sup>, Nirina R. Andriamaharavo<sup>3</sup>, H. Martin Garraffo<sup>3</sup>, Thomas F. Spande<sup>3</sup>, and John W. Daly<sup>3,#</sup>

<sup>1</sup>Graduate School of Medicine and Pharmaceutical Sciences, University of Toyama, Sugitani 2630, Toyama 930-0194, Japan

<sup>2</sup>Institute of Natural Medicine, University of Toyama, Sugitani 2630, Toyama 930-0194, Japan

<sup>3</sup>Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, DHHS, Bethesda, MD 20892, USA

### Abstract

An efficient and flexible synthesis of poison-frog alkaloids **251O** and *trans*-**223B** has been achieved using for both alkaloids an enantiodivergent process starting from the common lactam **1**. The relative stereochemistry of **251O** and *trans*-**223B** was determined to be **7** (R = *n*-C<sub>7</sub>H<sub>15</sub>, R' = *n*-Pr) and **14** by the present enantioselective synthesis.

### Keywords

poison-frog alkaloids; **251O**; *trans*-**223B**

### Introduction

A variety of lipophilic alkaloids have been detected in skin extracts of neotropical poison-frogs and over 800 alkaloids from 24 classes have been detected to date.<sup>1</sup> These alkaloids serve as a chemical defense against predation and some of them also exhibit significant inhibitory effects on nicotinic acetylcholine receptors (nAChRs).<sup>2</sup>

Very interestingly, most of these alkaloids appear to be sequestered from dietary arthropods.<sup>3</sup> Thus, the 3,5-disubstituted pyrrolizidine **251O**, detected in the poison-frogs *Mantella haraldmeieri*, *M. bernhardi* and *M. baroni* of Madagascar, has also been found in Malagasy ants of the genus *Anochetus grandidieri*.<sup>4</sup> So far, several syntheses of *trans*, *trans*-type of 3,5-disubstituted pyrrolizidines such as xenovenine have been reported,<sup>5</sup> whereas no synthesis of the *trans*, *cis*-type of 3,5-disubstituted pyrrolizidine alkaloids has been reported. Natural xenovenine is found in frogs as *cis*-**223H**. It has the 3*R*, 5*S*, 8*S* absolute configuration as shown in Figure 1 (unpublished work, HMG, TFS and JWD). Alkaloid **251O** has a *trans*, *cis*-type of pyrrolizidine ring system; however, both the relative and absolute stereochemistry are still unknown (Figure 1). As part of a program directed at studying the synthesis of biologically

toyooka@pha.u-toyama.ac.jp.

<sup>#</sup>Deceased on March 5, 2008.

#### Supporting Information Available

Experimental details for compounds ent-**4**, -**5**, -**6**, -**7**, -**11**, -**12**, -**13**, and -**14**, characterization data for all new synthetic compounds, including <sup>1</sup>H and <sup>13</sup>C NMR spectra, the GC-MS of coinjection data, and FTIR spectra of synthetic stereoisomers of **251O** and natural **251O** are available free of charge via the internet at <http://pubs.acs.org>.

active alkaloids,<sup>6</sup> we report herein, the efficient enantio- and diastereo-divergent synthesis of **251O** starting from a common lactam, **1**.<sup>7</sup>

## Results and Discussion

The known olefin **2**,<sup>8</sup> derived from **1**, was converted to a Boc-imide, which was treated with *n*-Pr or *n*-C<sub>7</sub>H<sub>15</sub> magnesium bromide according to Martin's reaction conditions<sup>9</sup> to provide the ketones **3** (R = *n*-Pr or *n*-C<sub>7</sub>H<sub>15</sub>) in high yield. Both ketones **3** were subjected to cyclization and subsequent reduction of the resulting iminium ion with Ph<sub>3</sub>SiH<sup>9</sup> to give rise to the *cis*-substituted pyrrolidines **4** (R = *n*-Pr or *n*-C<sub>7</sub>H<sub>15</sub>). The cross-metathesis reaction of **4** (R = *n*-Pr or *n*-C<sub>7</sub>H<sub>15</sub>) with ethyl acrylate using Grubbs' second-generation catalyst<sup>10</sup> afforded the unsaturated esters **5** (R = *n*-Pr or *n*-C<sub>7</sub>H<sub>15</sub>). The key pyrrolizidine ring-closure reaction of **5** (R = *n*-Pr or *n*-C<sub>7</sub>H<sub>15</sub>) resulted after treatment with 2 equivalents of AlCl<sub>3</sub> in CH<sub>2</sub>Cl<sub>2</sub> followed by K<sub>2</sub>CO<sub>3</sub> in the desired pyrrolizidines **6** (R = Et or *n*-C<sub>6</sub>H<sub>13</sub>) in excellent yield with high diastereoselectivity (>30:1). The stereochemistry of **6** (R = Et) was determined to be as shown based on an NOE experiment. In a difference NOE experiment, an NOE enhancement (ca. 2.3%) was observed on one of the methylene protons of the side chain at C-5 upon irradiation of the methine proton at C-3.

The highly selective formation of the desired **6** (R = Et or *n*-C<sub>6</sub>H<sub>13</sub>) is explained by invoking kinetic control as shown in Figure 2.<sup>11</sup> Finally, half-reduction of the ester moiety with diisobutylaluminum hydride, Wittig olefination of the resulting aldehyde followed by hydrogenation of the corresponding olefin furnished **7** (R = *n*-Pr, R' = *n*-C<sub>7</sub>H<sub>15</sub> or R = *n*-C<sub>7</sub>H<sub>15</sub>, R' = *n*-Pr). (Scheme 1)

### Reagents and conditions

(a) Boc<sub>2</sub>O, DMAP, MeCN, rt (95%); (b) *n*-C<sub>3</sub>H<sub>7</sub>MgBr or *n*-C<sub>7</sub>H<sub>15</sub>MgBr, TMEDA, THF, -78 °C (88% for R = *n*-Pr, 78% for R = *n*-C<sub>7</sub>H<sub>15</sub>); (c) B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>, Ph<sub>3</sub>SiH, CH<sub>2</sub>Cl<sub>2</sub>, -78 °C to rt (85% for R = *n*-Pr, 89% for R = *n*-C<sub>7</sub>H<sub>15</sub>); (d) ethyl acrylate, Grubbs' 2nd catalyst, CH<sub>2</sub>Cl<sub>2</sub>, reflux (96% for R = *n*-C<sub>3</sub>H<sub>7</sub>, 96% for R = *n*-C<sub>7</sub>H<sub>15</sub>); (e) AlCl<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, rt then K<sub>2</sub>CO<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, rt (93% for R = Et, 88% for R = *n*-C<sub>6</sub>H<sub>13</sub>); (f) DIBAL, CH<sub>2</sub>Cl<sub>2</sub>, -78 °C; (g) *n*-C<sub>5</sub>H<sub>11</sub>P<sup>+</sup>Ph<sub>3</sub>Br<sup>-</sup>, *n*-BuLi or MeP<sup>+</sup>Ph<sub>3</sub>I<sup>-</sup>, *n*-BuLi, THF, 0 °C to rt (46% for R = *n*-Pr, R' = C<sub>7</sub>H<sub>13</sub>, 49% for R = *n*-C<sub>7</sub>H<sub>15</sub>, R' = allyl); (h) 10% Pd/C, H<sub>2</sub>, EtOAc, 1 atm (quant for R = *n*-Pr, R' = *n*-C<sub>7</sub>H<sub>15</sub>, 95% for R = *n*-C<sub>7</sub>H<sub>15</sub>, R' = *n*-Pr)

Ent-**7** (R = *n*-Pr, R' = *n*-C<sub>7</sub>H<sub>15</sub> or R = *n*-C<sub>7</sub>H<sub>15</sub>, R' = *n*-Pr) was synthesized from **1** in a similar way as shown in Scheme 2.

### Reagents and conditions

(a) EtMgBr or *n*-C<sub>6</sub>H<sub>13</sub>MgBr, THF, -35 °C (52% for R = *n*-Pr, 50% for R = *n*-C<sub>7</sub>H<sub>15</sub>); (b) Boc<sub>2</sub>O, DMAP, MeCN, rt (94% for R = *n*-Pr, 93% for R = *n*-C<sub>7</sub>H<sub>15</sub>); (c) 4-butenylMgBr, TMEDA, THF, -78 °C (95% for R = *n*-Pr, 98% for R = *n*-C<sub>7</sub>H<sub>15</sub>); (d) B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>, Ph<sub>3</sub>SiH, CH<sub>2</sub>Cl<sub>2</sub>, -78 °C to rt (73% for R = *n*-Pr, 86% for R = *n*-C<sub>7</sub>H<sub>15</sub>); (e) ethyl acrylate, Grubbs' 2nd catalyst, CH<sub>2</sub>Cl<sub>2</sub>, reflux (96% for R = *n*-Pr, 95% for R = *n*-C<sub>7</sub>H<sub>15</sub>); (f) AlCl<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, rt then K<sub>2</sub>CO<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, rt (93% for R = *n*-Pr, 91% for R = *n*-C<sub>7</sub>H<sub>15</sub>); (g) DIBAL, CH<sub>2</sub>Cl<sub>2</sub>, -50 °C; (h) *n*-C<sub>5</sub>H<sub>11</sub>P<sup>+</sup>Ph<sub>3</sub>Br<sup>-</sup>, *n*-BuLi or MeP<sup>+</sup>Ph<sub>3</sub>I<sup>-</sup>, *n*-BuLi, THF, 0 °C to rt (48% for R = *n*-Pr, R' = C<sub>7</sub>H<sub>13</sub>, 52% for R = *n*-C<sub>7</sub>H<sub>15</sub>, R' = allyl); (h) 10% Pd/C, H<sub>2</sub>, EtOAc, 1 atm (96% for R = *n*-Pr, R' = *n*-C<sub>7</sub>H<sub>15</sub>, quant for R = *n*-C<sub>7</sub>H<sub>15</sub>, R' = *n*-Pr).

The enantiodivergent synthesis of *trans*-**223B** was also achieved starting from common lactam **2** as shown in Scheme 3

## Reagents and conditions

(a) Boc<sub>2</sub>O, DMAP, MeCN, rt (95%); (b) *n*-BuMgBr, TMEDA, THF, -78 °C (98%); (c) B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>, Ph<sub>3</sub>SiH, CH<sub>2</sub>Cl<sub>2</sub>, -78 °C to rt (73%); (d) ethyl acrylate, Grubbs' 2nd catalyst, CH<sub>2</sub>Cl<sub>2</sub>, reflux (96%); (e) AlCl<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, rt then K<sub>2</sub>CO<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, rt (92%); (f) DIBAL, CH<sub>2</sub>Cl<sub>2</sub>, -50 °C; (g) EtP<sup>+</sup>Ph<sub>3</sub>Br<sup>-</sup>, *n*-BuLi, THF, 0 °C to rt (50%); (h) 10% Pd/C, H<sub>2</sub>, EtOAc, 1 atm (quant); (i) Boc<sub>2</sub>O, DMAP, MeCN, rt (95%); (j) 10% Pd/C, H<sub>2</sub>, EtOAc, 1 atm, then 4-butenylMgBr, TMEDA, THF, -78 °C (96%); (k) B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>, Ph<sub>3</sub>SiH, CH<sub>2</sub>Cl<sub>2</sub>, -78 °C to rt (72%); (l) ethyl acrylate, Grubbs' 2nd catalyst, CH<sub>2</sub>Cl<sub>2</sub>, reflux (84%); (m) AlCl<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, rt then K<sub>2</sub>CO<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, rt (92%); (n) DIBAL, CH<sub>2</sub>Cl<sub>2</sub>, -50 °C; (o) EtP<sup>+</sup>Ph<sub>3</sub>Br<sup>-</sup>, *n*-BuLi, THF, 0 °C to rt (53%); (p) 10% Pd/C, H<sub>2</sub>, EtOAc, 1 atm (quant).

The GC-MS and GC-FTIR spectra of synthetic **7** (R = *n*-C<sub>7</sub>H<sub>15</sub>, R' = *n*-Pr) were identical with those for natural **2510** present in skin extracts of three Madagascan frogs (*Mantella haraldmeieri* collected in 2004, *M. bernhardi* collected in 1997, and *M. baroni* collected in 2004, unpublished results; see supporting information). Furthermore, the GC-MS and GC-FTIR spectra of synthetic **14** were also identical with those for natural *trans*-**223B** present in a skin extract of the Argentinean frog *Melanophryniscus stelzneri*.<sup>12</sup> The mass spec evidence, with a strong loss of propyl from the molecular ion at m/z 208 (251-43) indicates a pseudo-axial position for the propyl group: a group that is trans-antiparallel to the radical ion on the N atom should cleave preferentially, all other things being equal. The heptyl group, being larger cleaves better, but the comparison of the MS on the Supplemental Information, between *trans*-stereoisomers with a Pr-group pseudo-axial or a Pr-group pseudo-equatorial, clearly indicates a bigger fragmentation of the propyl group for the one in the pseudo-axial position. This MS evidence indicates that natural **2510** is (3S, 5S, 8R) or the enantiomer (3R, 5R, 8S) same as **7** (R = *n*-C<sub>7</sub>H<sub>15</sub>, R' = *n*-Pr) or ent-**7** (R = *n*-C<sub>7</sub>H<sub>15</sub>, R' = *n*-Pr). Unfortunately, we could not achieve the separation of both enantiomers of synthetic **7** by GC analysis using two different columns. For the enantiomers of synthetic **14**, we anticipate even more difficulty in the separation since the two side-chains are identical; however, this separation was not attempted.

In conclusion, we have achieved the first synthesis of alkaloids **2510** and *trans*-**223B** in enantio- and diastereo-divergent fashion in 8 or 9 steps in 25~29% overall yield from the known lactam **1**, respectively. Furthermore, the relative stereochemistries of natural **2510** and *trans*-**223B** were determined to be **7** (R = *n*-C<sub>7</sub>H<sub>15</sub>, R' = *n*-Pr) and **14**, respectively, by the present efficient syntheses and comparisons with the natural compounds. It is difficult to separate the enantiomers of **7** and **14**, but further work on the GC separation of the enantiomers of **7** and **14** for the determination of the absolute stereochemistry of natural **2510** and *trans*-**223B** is in progress.<sup>13</sup>

## Experimental Section

### (2R)-(-)-2-(But-3-enyl)-5-oxopyrrolidine-1-carboxylic acid *tert*-butyl ester

To a stirred solution of **2**<sup>8</sup> (337 mg, 2.42 mmol) in MeCN (20 mL) was added DMAP (326 mg, 2.67 mmol) at 0 °C, and the reaction mixture was stirred at 0 °C for 30 min. To the reaction mixture was added Boc<sub>2</sub>O (633 mg, 2.90 mmol) at 0 °C, and the resulting mixture was stirred at room temperature for 45 h. The volatiles were evaporated, and the residue was chromatographed on silica gel (20 g, hexane/acetone=15:1) to give the Boc-imide (550 mg, 95%) as a colorless oil.

IR (neat) 3078, 1785, 1750, 1714, 1308, 1153 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 1.52 (9H, s), 1.59 (1H, m), 1.78 (1H, m), 1.90 (1H, m), 2.03–2.19 (3H, br m), 2.43 (1H, ddd, *J* = 9.0, 8.5, 2.1 Hz), 2.57 (1H, dd, *J* = 9.0, 8.5 Hz), 4.12 (1H, m), 5.01 (1H, d, *J* = 10.7 Hz), 5.04 (1H, d, *J* = 15.0 Hz), 5.80 (1H, m); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 21.9 (t), 27.7 (q), 29.6 (t), 31.0 (t),

32.4 (t), 57.1 (d), 82.2 (s), 115.0 (t), 136.7 (d), 149.3 (s), 173.7 (s); MS 182 ( $M^+$ -57), 84 (100); HRMS Calcd for  $C_9H_{12}O_3N$  182.0817, Found 182.0830;  $[\alpha]_D^{26}$  -56.68 (*c* 2.34,  $CHCl_3$ ).

**(1*R*)-(+)-[1-(3-Oxohexyl)pent-4-enyl]carbamic acid *tert*-butyl ester (**3**, R = *n*-Pr)**

To a stirred solution of above Boc-imide (239 mg, 1.00 mmol) in THF (5 mL) was added a solution of *n*-PrMgBr, prepared from *n*-PrBr (0.27 mL, 3.00 mmol) and Mg (72 mg, 3.00 mmol) in THF (10 mL) at reflux, and TMEDA (0.48 mL, 3.00 mmol) at -78 °C, and the resulting mixture was stirred at the same temperature for 1.5 h. The reaction was quenched with *i*-PrOH (5 mL), and the reaction mixture was diluted with Et<sub>2</sub>O. The organic layer was washed with 10% HCl (aq) solution, dried over MgSO<sub>4</sub>, and evaporated to give pale yellow oil, which was chromatographed on silica gel (20 g, hexane/acetone=50:1-30:1) to give **3** (R = *n*-Pr, 250 mg, 88%) as a colorless solid (mp 41-43 °C). IR (KBr) 3349, 3081, 1708, 1689, 1528, 1175  $cm^{-1}$ ; <sup>1</sup>H NMR (500 MHz,  $CDCl_3$ ) δ 0.90 (3H, t, *J* = 7.2 Hz), 1.42 (9H, s), 1.48-1.63 (5H, br m), 1.79 (1H, m), 2.08 (2H, q-like, *J* = 7.2 Hz), 2.37 (2H, t, *J* = 7.2 Hz), 2.46 (2H, t, *J* = 7.2 Hz), 3.53 (1H, br), 4.25 (1H, br d, *J* = 10.2 Hz), 4.95 (1H, d, *J* = 9.4 Hz), 5.01 (1H, d, *J* = 14.8 Hz), 5.78 (1H, m); <sup>13</sup>C NMR (75 MHz,  $CDCl_3$ ) δ 13.6 (q), 17.1 (t), 28.2 (q), 29.0 (t), 30.1 (t), 35.1 (t), 39.1 (t), 44.6 (t), 49.9 (d), 78.5 (s), 114.5 (t), 137.6 (d), 155.4 (s), 210.3 (s); MS 226 ( $M^+$ -57), 57 (100); HRMS Calcd for  $C_{12}H_{20}O_3N$  226.1443, Found 226.1457;  $[\alpha]_D^{26}$  +1.17 (*c* 1.21,  $CHCl_3$ ).

**(1*R*)-(-)-[1-(3-Oxodecyl)pent-4-enyl]carbamic acid *tert*-butyl ester (**3**, R = *n*-C<sub>7</sub>H<sub>15</sub>)**

To a stirred solution of above Boc-imide (239 mg, 1.00 mmol) in THF (5 mL) was added a solution of *n*-C<sub>7</sub>H<sub>15</sub>MgBr, prepared from *n*-C<sub>7</sub>H<sub>15</sub>Br (0.47 mL, 3.00 mmol) and Mg (72 mg, 3.00 mmol) in THF (10 mL) at reflux, and TMEDA (0.48 mL, 3.00 mmol) at -78 °C, and the resulting mixture was stirred at the same temperature for 1.5 h. The reaction was quenched with *i*-PrOH (5 mL), and the reaction mixture was diluted with Et<sub>2</sub>O. The organic layer was washed with 10% HCl (aq) solution, dried over MgSO<sub>4</sub>, and evaporated to give a pale yellow oil, which was chromatographed on silica gel (20 g, hexane/acetone=50:1-30:1) to give **3** (R = *n*-C<sub>7</sub>H<sub>15</sub>, 265 mg, 78%) as a colorless solid (mp 60-62 °C). IR (KBr) 3347, 3080, 1709, 1684, 1525, 1173  $cm^{-1}$ ; <sup>1</sup>H NMR (500 MHz,  $CDCl_3$ ) δ 0.85 (3H, t, *J* = 7.2 Hz), 1.26 (8H, br), 1.43 (9H, s), 1.46-1.61 (5H, br m), 1.81 (1H, m), 2.13 (2H, m), 2.39 (2H, t, *J* = 7.2 Hz), 2.47 (2H, t, *J* = 7.2 Hz), 3.53 (1H, br), 4.24 (1H, br d, *J* = 9.6 Hz), 4.95 (1H, d, *J* = 9.6 Hz), 5.01 (1H, d, *J* = 15.0 Hz), 5.78 (1H, m); <sup>13</sup>C NMR (75 MHz,  $CDCl_3$ ) δ 14.0 (q), 22.5 (t), 23.7 (t), 28.3 (q), 29.0 (t), 29.1 (t), 30.1 (t), 31.6 (t), 35.2 (t), 39.2 (t), 42.9 (t), 50.0 (d), 78.7 (s), 114.6 (t), 137.7 (d), 155.4 (s), 210.6 (s); MS 282 ( $M^+$ -57), 57 (100); HRMS Calcd for  $C_{16}H_{28}O_3N$  282.2069, Found 282.2091;  $[\alpha]_D^{26}$  -1.56 (*c* 0.74,  $CHCl_3$ ).

**(2*R*,5*S*)-(-)-2-But-3-enyl-5-propylpyrrolidine-1-carboxylic acid *tert*-butyl ester (**4**, R = *n*-Pr)**

To a stirred solution of **3** (R = *n*-Pr, 226 mg, 0.80 mmol) in  $CH_2Cl_2$  (5 mL) was added a solution of  $(C_6F_5)_3B$  (82 mg, 0.16 mmol) and  $Ph_3SiH$  (415 mg, 1.59 mmol) in  $CH_2Cl_2$  (5 mL) at -78 °C, and the reaction mixture was stirred at -78 °C for 30 min, and then at room temperature for 20 h. The reaction was quenched with Et<sub>3</sub>N (0.6 mL), and the resulting mixture was stirred at room temperature for 20 min. The mixture was diluted with Et<sub>2</sub>O, and the organic layer was washed successively with 10% AcOH (aq) solution and satd. NaHCO<sub>3</sub> (aq) solution, dried over MgSO<sub>4</sub>, and evaporated to give a pale yellow oil, which was chromatographed on silica gel (20 g, hexane/acetone=200:1-150:1) to give **4** (R = *n*-Pr, 182 mg, 85%) as a colorless oil.

IR (neat) 3073, 1694, 1389  $cm^{-1}$ ; <sup>1</sup>H NMR (500 MHz,  $CDCl_3$ ) δ 0.92 (3H, t, *J* = 7.2 Hz), 1.28 (6H, br m), 1.46 (9H, s), 1.61 (2H, m), 1.90 (2H, m), 2.04 (2H, m), 3.76 (2H, br), 4.94 (1H, d, *J* = 9.6 Hz), 5.03 (1H, d, *J* = 15.6 Hz), 5.81 (1H, m); <sup>13</sup>C NMR (75 MHz,  $CDCl_3$ ) δ 14.2 (q), 19.6 (t), 28.5 (q), 29.4 (t), 30.7 (t), 35.1 (t), 38.2 (t), 57.8 (d), 58.1 (d), 78.7 (s), 114.2 (t), 138.3

(d), 154.7 (s); MS 267 (M<sup>+</sup>), 57 (100); HRMS Calcd for C<sub>16</sub>H<sub>29</sub>O<sub>2</sub>N 267.2198, Found 267.2215; [ $\alpha$ ]<sub>D</sub><sup>26</sup> -3.23 (c 1.10, CHCl<sub>3</sub>).

**(2R,5S)-(-)-2-But-3-enyl-5-heptylpyrrolidine-1-carboxylic acid *tert*-butyl ester (4, R = *n*-C<sub>7</sub>H<sub>15</sub>)**

To a stirred solution of **3** (R = *n*-C<sub>7</sub>H<sub>15</sub>, 250 mg, 0.77 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added a solution of (C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>B (78 mg, 0.15 mmol) and Ph<sub>3</sub>SiH (401 mg, 1.54 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) at -78 °C, and the reaction mixture was stirred at -78 °C for 30 min, and then at room temperature for 20 h. The reaction was quenched with Et<sub>3</sub>N (0.6 mL), and the resulting mixture was stirred at room temperature for 20 min. The mixture was diluted with Et<sub>2</sub>O, and the organic layer was washed successively with 10% AcOH (aq) solution and satd. NaHCO<sub>3</sub> (aq) solution, dried over MgSO<sub>4</sub>, and evaporated to give a pale yellow oil, which was chromatographed on silica gel (20 g, hexane/acetone=200:1–150:1) to give **4** (R = *n*-C<sub>7</sub>H<sub>15</sub>, 222 mg, 89%) as a colorless oil.

IR (neat) 3075, 1695, 1390, 1174, 1107 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  0.88 (3H, t, *J* = 7.2 Hz), 1.27 (14H, br), 1.46 (9H, s), 1.62 (2H, m), 1.90 (2H, m), 2.07 (2H, br), 3.75 (2H, br), 4.95 (1H, d, *J* = 10.2 Hz), 5.02 (1H, d, *J* = 15.8 Hz), 5.84 (1H, m); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  14.1 (q), 22.7 (t), 26.4 (t), 28.5 (q), 29.3 (t), 29.6 (t), 30.8 (t), 31.8 (t), 35.1 (t), 35.9 (t), 57.8 (d), 58.4 (d), 78.7 (s), 114.2 (t), 138.3 (d), 154.7 (s); MS 323 (M<sup>+</sup>), 168 (100); HRMS Calcd for C<sub>20</sub>H<sub>37</sub>O<sub>2</sub>N 323.2824, Found 323.2847; [ $\alpha$ ]<sub>D</sub><sup>26</sup> -0.97 (c 0.55, CHCl<sub>3</sub>).

**(2R,5S)-(-)-2-(4-Ethoxycarbonylbut-3-enyl)-5-propylpyrrolidine-1-carboxylic acid *tert*-butyl ester (5, R = *n*-Pr)**

To a stirred solution of **4** (R = *n*-Pr, 117 mg, 0.44 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (8 mL) were added Grubbs 2<sup>nd</sup> catalyst (15 mg, 0.018 mmol) and ethyl acrylate (0.24 mL, 2.20 mmol), and the resulting mixture was refluxed for 5.5 h. After cooling, the solvent was removed under reduced pressure, and the residue was chromatographed on silica gel (20 g, hexane/acetone=100:1–60:1) to give **5** (R = *n*-Pr, 143 mg, 96%) as a pale yellow oil.

IR (neat) 1721, 1693, 1390, 1174 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  0.91 (3H, t, *J* = 7.2 Hz), 1.20–1.30 (8H, br, including at  $\delta$  1.27, 3H, t, *J* = 7.2 Hz), 1.45 (9H, s), 1.61 (2H, m), 1.90 (3H, m), 2.19 (2H, br), 3.76 (2H, br), 4.16 (2H, q, *J* = 7.2 Hz), 5.85 (1H, d, *J* = 15.8 Hz), 6.96 (1H, dt, *J* = 15.8, 6.8 Hz); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  14.1 (q), 14.2 (q), 19.5 (t), 28.5 (q), 29.1 (t), 29.5 (t), 34.2 (t), 38.2 (t), 57.6 (d), 58.1 (d), 60.0 (t), 78.8 (s), 121.1 (d), 148.4 (d), 154.6 (s), 166.2 (s); MS 339 (M<sup>+</sup>), 196 (100); HRMS Calcd for C<sub>19</sub>H<sub>33</sub>O<sub>4</sub>N 339.2410, Found 339.2392; [ $\alpha$ ]<sub>D</sub><sup>26</sup> -6.81 (c 0.66, CHCl<sub>3</sub>).

**(2R,5S)-(-)-2-(4-Ethoxycarbonylbut-3-enyl)-5-heptylpyrrolidine-1-carboxylic acid *tert*-butyl ester (5, R = *n*-C<sub>7</sub>H<sub>15</sub>)**

To a stirred solution of **4** (R = *n*-C<sub>7</sub>H<sub>15</sub>, 134 mg, 0.41 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (7 mL) were added Grubbs 2<sup>nd</sup> catalyst (14 mg, 0.016 mmol) and ethyl acrylate (0.23 mL, 2.07 mmol), and the resulting mixture was refluxed for 5.5 h. After cooling, the solvent was removed under reduced pressure, and the residue was chromatographed on silica gel (20 g, hexane/acetone=100:1–60:1) to give **5** (R = *n*-C<sub>7</sub>H<sub>15</sub>, 157 mg, 96%) as a pale yellow oil.

IR (neat) 1716, 1696, 1387, 1174 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  0.87 (3H, t, *J* = 7.2 Hz), 1.17–1.33 (13H, br, including at  $\delta$  1.27, 3H, t, *J* = 7.2 Hz), 1.45 (9H, s), 1.60 (2H, m), 1.92 (2H, m), 2.20 (2H, m), 3.76 (2H, br), 4.16 (2H, q, *J* = 7.2 Hz), 5.81 (1H, d, *J* = 15.6 Hz), 6.97 (1H, dt, *J* = 15.6, 6.8 Hz); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  13.9 (q), 14.1 (q), 22.4 (t), 26.2 (t), 28.3 (q), 29.0 (t), 29.1 (t), 29.4 (t), 31.6 (t), 34.1 (t), 35.8 (t), 57.4 (d), 58.2 (d), 59.7 (t),

78.6 (s), 121.0 (d), 148.2 (d), 154.4 (s), 165.9 (s); MS 338 ( $M^+$ -57), 57 (100); HRMS Calcd for  $C_{19}H_{32}O_4N$  338.2331, Found 338.2338;  $[\alpha]_D^{26}$  -1.75 (*c* 0.88,  $CHCl_3$ ).

**(3*R*,5*S*,8*S*)-(+)-(5-Propylhexahydropyrrolizin-3-yl)acetic acid ethyl ester (6, R = Et)**

To a stirred solution of **5** (R = *n*-Pr, 123 mg, 0.36 mmol) in  $CH_2Cl_2$  (10 mL) was added  $AlCl_3$  (106 mg, 0.80 mmol) at 0 °C, and the resulting suspension was stirred at room temperature for 24 h. The reaction was quenched with satd.  $NaHCO_3$  (aq) solution, and the organic layer was separated. The aqueous layer was extracted with  $CHCl_3$  (10 mL×5), and the organic layer and extracts were combined, dried over  $K_2CO_3$ , and evaporated to give the residue. To a stirred solution of this residue in  $CH_2Cl_2$  (10 mL) was added  $K_2CO_3$  (100 mg, 0.72 mmol), and the resulting suspension was stirred at room temperature for 48 h. The insoluble material was filtered off, and washed with  $CH_2Cl_2$ . The filtrate was evaporated to afford the residue, which was chromatographed on silica gel (20 g, hexane/acetone=20:1–12:1) to give **6** (R = Et, 80 mg, 93%) as a pale yellow oil, and the stereoisomer at the 3-position (all *cis*-pyrrolizidine, 2.5 mg, 3%) also as a pale yellow oil.

IR (neat) 1732, 1178  $cm^{-1}$ ;  $^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$  0.94 (3H, t, *J* = 7.2 Hz), 1.26 (3H, t, *J* = 7.2 Hz), 1.31 (1H, m), 1.33–1.44 (4H, br m), 1.52 (1H, m), 1.74–1.81 (4H, br m), 2.03 (2H, m), 2.28 (1H, dd, *J* = 15.1, 9.8 Hz), 2.57 (1H, dd, *J* = 15.1, 3.9 Hz), 3.01 (1H, br), 3.40 (1H, m), 3.57 (1H, br), 4.12 (2H, q, *J* = 7.2 Hz);  $^{13}C$  NMR (75 MHz,  $CDCl_3$ )  $\delta$  14.2 (q), 14.4 (q), 21.4 (t), 30.1 (t), 30.8 (t), 31.3 (t), 32.4 (t), 33.5 (t), 43.2 (t), 54.2 (d), 60.0 (t), 63.5 (d), 65.6 (d), 172.1 (s); MS 239 ( $M^+$ ), 196 (100); HRMS Calcd for  $C_{14}H_{25}O_2N$  239.1885, Found 239.1869;  $[\alpha]_D^{26}$  +25.94 (*c* 1.95,  $CHCl_3$ ).

Stereoisomer at the 3-position: IR (neat) 1736, 1174  $cm^{-1}$ ;  $^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$  0.90 (3H, t, *J* = 7.2 Hz), 1.26 (3H, t, *J* = 7.2 Hz), 1.22–1.37 (2H, m), 1.38–1.52 (5H, m), 1.91–1.98 (2H, m), 2.01–2.04 (2H, m), 2.28 (1H, dd, *J* = 15.0, 8.6 Hz), 2.53 (1H, dd, *J* = 15.0, 3.6 Hz), 2.67 (1H, m), 3.15 (1H, m), 3.57 (1H, br), 4.12 (2H, q, *J* = 7.2 Hz); MS 239 ( $M^+$ ), 83 (100).

**(3*R*,5*S*,8*S*)-(+)-(5-Heptylhexahydropyrrolizin-3-yl)acetic acid ethyl ester (6, R = *n*- $C_6H_{13}$ )**

To a stirred solution of **5** (R = *n*- $C_7H_{15}$ , 270 mg, 0.68 mmol) in  $CH_2Cl_2$  (20 mL) was added  $AlCl_3$  (200 mg, 1.50 mmol) at 0 °C, and the resulting suspension was stirred at room temperature for 24 h. The reaction was quenched with satd.  $NaHCO_3$  (aq) solution, and the organic layer was separated. The aqueous layer was extracted with  $CHCl_3$  (20 mL×5), and the organic layer and extracts were combined, dried over  $K_2CO_3$ , and evaporated to give the residue. To a stirred solution of this residue in  $CH_2Cl_2$  (20 mL) was added  $K_2CO_3$  (189 mg, 1.37 mmol), and the resulting suspension was stirred at room temperature for 48 h. The insoluble material was filtered off, and washed with  $CH_2Cl_2$ . The filtrate was evaporated to afford the residue, which was chromatographed on silica gel (30 g, hexane/acetone=20:1–12:1) to give **6** (R = *n*- $C_6H_{13}$ , 178 mg, 88%) as a pale yellow oil.

IR (neat) 1731, 1176  $cm^{-1}$ ;  $^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$  0.87 (3H, t, *J* = 7.2 Hz), 1.25 (3H, t, *J* = 7.2 Hz), 1.17–2.41 (14H, br m), 1.51 (1H, m), 1.76 (3H, m), 2.01 (2H, m), 2.27 (1H, dd, *J* = 14.8, 9.6 Hz), 2.56 (1H, dd, *J* = 14.8, 4.1 Hz), 2.99 (1H, br), 3.38 (1H, m), 3.56 (1H, m), 4.10 (2H, q, *J* = 7.2 Hz);  $^{13}C$  NMR (75 MHz,  $CDCl_3$ )  $\delta$  13.9 (q), 14.1 (q), 22.5 (t), 28.1 (t), 29.1 (t), 29.7 (t), 30.1 (t), 30.6 (t), 31.1 (t), 31.2 (t), 31.6 (t), 32.2 (t), 43.1 (t), 54.0 (d), 59.8 (t), 63.6 (d), 65.5 (d), 171.8 (s); MS 295 ( $M^+$ ), 196 (100); HRMS Calcd for  $C_{18}H_{33}O_2N$  295.2510, Found 295.2501;  $[\alpha]_D^{26}$  +18.82 (*c* 1.14,  $CHCl_3$ ).

**(3*S*,5*S*,8*S*)-(+)-3-Heptyl-5-propylhexahydropyrrolizine (7, R = *n*-Pr, R' = *n*- $C_7H_{15}$ )**

To a stirred solution of **6** (R = Et, 79 mg, 0.32 mmol) in  $CH_2Cl_2$  (7 mL) was added a solution of DIBAL (0.98 M in hexane, 0.36 mL, 0.35 mmol) at -50 °C, and the reaction mixture was

stirred at  $-50\text{ }^{\circ}\text{C}$  for 30 min. The reaction was quenched with MeOH, and satd. Rochelle (aq) solution, and the organic layer was separated. The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  (5 mL $\times$ 3), and the organic layer and extracts were combined, dried over  $\text{K}_2\text{CO}_3$ , and evaporated to give a pale yellow oil, which was used directly in the next step.

To a stirred suspension of  $n\text{-C}_5\text{H}_{11}\text{P}^+\text{Ph}_3\text{Br}^-$  (529 mg, 1.28 mmol) in THF (10 mL) was added a solution of  $n\text{-BuLi}$  (1.6 M in hexane, 0.7 mL, 1.12 mmol) at  $0\text{ }^{\circ}\text{C}$ , and the resulting orange suspension was stirred at  $0\text{ }^{\circ}\text{C}$  for 10 min. To the suspension was added a solution of the above aldehyde in THF (3 mL) at  $0\text{ }^{\circ}\text{C}$ , and the resulting suspension was stirred at room temperature for 22 h. The reaction was quenched with  $\text{H}_2\text{O}$ , and the aqueous mixture was extracted with  $\text{Et}_2\text{O}$  (15 mL $\times$ 4). The organic extracts were combined, dried over  $\text{K}_2\text{CO}_3$ , and evaporated to give a residue, that was chromatographed on silica gel (20 g, hexane/acetone=25:1–10:1) to give the corresponding olefin (37 mg, 46%) as a mixture of *E*- and *Z*-isomers.

$^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.89 (3H, t,  $J = 7.2$  Hz), 0.94 (3H, t,  $J = 7.1$  Hz), 1.21–1.53 (12H, br m), 1.72–1.88 (3H, m), 1.89–1.94 (1H, m), 1.96–2.07 (3H, m), 2.30–2.41 (1H, m), 2.93–3.00 (1H, m), 3.02–3.16 (1H, m), 3.62 (1H, br), 5.32–5.47 (2H, m).

To a stirred solution of the above olefin (20 mg, 0.08 mmol) in EtOAc (3 mL) was added 10% Pd/C (10 mg), and the resulting suspension was stirred under a hydrogen atmosphere at 1 atm for 40 h. The catalyst was removed by filtration and the filtrate was evaporated to give **7** (R = *n*-Pr, R' = *n*-C<sub>7</sub>H<sub>15</sub>, 20 mg, quant) as a pale yellow oil.

IR (neat) 2926, 2869, 1457  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.88 (3H, t,  $J = 7.1$  Hz), 0.94 (3H, t,  $J = 7.1$  Hz), 1.27–1.43 (17H, br m), 1.54 (1H, br), 1.77 (3H, m), 1.97 (3H, m), 2.90 (1H, br), 3.05 (1H, br), 3.57 (1H, br);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  14.2 (q), 14.6 (q), 21.6 (t), 22.8 (t), 27.2 (t), 29.4 (t), 29.9 (t), 30.2 (t), 30.9 (t), 31.7 (t), 31.9 (t), 32.1 (t), 33.8 (t), 38.6 (t), 57.7 (d), 63.8 (d), 65.8 (d); MS 251 ( $\text{M}^+$ ), 208 (100); HRMS Calcd for  $\text{C}_{17}\text{H}_{33}\text{N}$  251.2613, Found 251.2601;  $[\alpha]_{\text{D}}^{26} +36.81$  ( $c$ 0.44,  $\text{CHCl}_3$ ).

### (3*R*,5*S*,8*S*)-(+)-3-Allyl-5-heptylhexahydropyrrolizine

To a stirred solution of **6** (R = *n*-C<sub>6</sub>H<sub>13</sub>, 91 mg, 0.31 mmol) in  $\text{CH}_2\text{Cl}_2$  (7 mL) was added a solution of DIBAL (0.98 M in hexane, 0.35 mL, 0.34 mmol) at  $-50\text{ }^{\circ}\text{C}$ , and the reaction mixture was stirred at  $-50\text{ }^{\circ}\text{C}$  for 30 min. The reaction was quenched with MeOH, and then satd. Rochelle (aq) solution, and the organic layer was separated. The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  (5 mL $\times$ 3), and the organic layer and extracts were combined, dried over  $\text{K}_2\text{CO}_3$ , and evaporated to give a pale yellow oil, that was used directly in the next step.

To a stirred suspension of  $\text{MeP}^+\text{Ph}_3\text{I}^-$  (501 mg, 1.24 mmol) in THF (10 mL) was added a solution of  $n\text{-BuLi}$  (1.6 M in hexane, 0.68 mL, 1.09 mmol) at  $0\text{ }^{\circ}\text{C}$ , and the resulting orange suspension was stirred at  $0\text{ }^{\circ}\text{C}$  for 10 min. To the suspension was added a solution of the above aldehyde in THF (3 mL) at  $0\text{ }^{\circ}\text{C}$ , and the resulting suspension was stirred at room temperature for 27 h. The reaction was quenched with  $\text{H}_2\text{O}$ , and the aqueous mixture was extracted with  $\text{Et}_2\text{O}$  (15 mL  $\times$  4). The organic extracts were combined, dried over  $\text{K}_2\text{CO}_3$ , and evaporated to give a residue, that was chromatographed on silica gel (20 g, hexane/acetone=20:1–10:1) to give the corresponding olefin (38 mg, 49%) as a pale yellow oil.

IR (neat) 3074, 2953, 2927, 2857, 1465, 909  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.87 (3H, t,  $J = 7.2$  Hz), 1.20–1.41 (14H, br m), 1.50 (1H, m), 1.73–1.83 (3H, m), 1.91 (1H, m), 1.97–2.04 (2H, m), 2.36 (1H, m), 2.95–3.03 (2H, m), 3.57 (1H, m), 4.98 (1H, d,  $J = 10.2$  Hz), 5.02 (1H, d,  $J = 15.0$  Hz), 5.98 (1H, m);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  14.2 (q), 22.8 (t), 28.4 (t), 29.3 (t), 30.0 (t), 30.2 (t), 31.0 (t), 31.4 (t), 31.5 (t), 31.6 (t), 31.9 (t), 43.0 (t), 57.1 (d), 64.0

(d), 66.0 (d), 115.8 (t), 136.6 (d); MS 249 (M<sup>+</sup>), 208 (100); HRMS Calcd for C<sub>17</sub>H<sub>31</sub>N 249.2455, Found 249.2473; [ $\alpha$ ]<sub>D</sub><sup>26</sup> +36.81 (c 0.44, CHCl<sub>3</sub>).

**(3S,5S,8R)-(+)-3-Heptyl-5-propylhexahydropyrrolizine (7, R = *n*-C<sub>7</sub>H<sub>15</sub>, R' = *n*-Pr)**

To a stirred solution of the above olefin (25 mg, 0.10 mmol) in EtOAc (3 mL) was added 10% Pd/C (13 mg), and the resulting suspension was stirred under a hydrogen atmosphere at 1 atm for 44 h. The catalyst was removed by filtration and the filtrate was evaporated to give **7** (R = *n*-C<sub>7</sub>H<sub>15</sub>, R' = *n*-Pr, 24 mg, 95%) as a pale yellow oil.

IR (neat) 2966, 2927, 2851, 1458 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  0.87 (3H, t, *J* = 7.1 Hz), 0.90 (3H, t, *J* = 6.9 Hz), 1.40–1.50 (19H, br m), 1.56 (1H, m), 1.79 (2H, m), 1.98 (2H, m), 2.92 (1H, m), 3.08 (1H, br), 3.64 (1H, br); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  14.2 (q), 14.4 (q), 20.4 (t), 22.7 (t), 28.3 (t), 29.3 (t), 30.0 (t), 30.1 (t), 30.7 (t), 31.3 (t), 31.7 (t), 31.9 (t), 32.1 (t), 40.4 (t), 57.8 (d), 64.0 (d), 66.0 (d); MS 251 (M<sup>+</sup>), 208 (100); HRMS Calcd for C<sub>17</sub>H<sub>33</sub>N 251.2613, Found 251.2620; [ $\alpha$ ]<sub>D</sub><sup>26</sup> +25.48 (c 0.47, CHCl<sub>3</sub>).

**(5S)-(-)-5-Propyl-2-oxopyrrolidine (8, R = *n*-Pr)**

To a stirred suspension of CuI (3.8 g, 20 mmol) in THF (30 mL) was added a solution of EtMgBr (0.96 M in THF, 42 mL, 40 mmol) at -35 °C, and the resulting suspension was stirred at the same temperature for 30 min. To the suspension was added a solution of **1** (1.8 g, 8 mmol) in THF (15 mL) at -35 °C, and then the reaction mixture was stirred at -35 °C for 18 h. The reaction was quenched with satd. NH<sub>4</sub>Cl (aq) solution, and the insoluble material was filtered off and washed with CHCl<sub>3</sub>. The organic layer was separated, and the aqueous layer was extracted with CHCl<sub>3</sub> (30 mL  $\times$  2). The organic layer and extracts were combined, dried over MgSO<sub>4</sub>, and evaporated to give a residue, that was chromatographed on silica gel (50 g, hexane/acetone=20:1–2:1) to give **8** (R = *n*-Pr, 525 mg, 52%) as a pale yellow oil.

IR (neat) 3193, 1699, 1286 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  0.93 (3H, t, *J* = 7.2 Hz), 1.36 (2H, m), 1.44 (1H, m), 1.51 (1H, m), 1.69 (1H, m), 2.25 (1H, m), 2.32 (2H, m), 3.63 (1H, quint, *J* = 6.8 Hz), 6.42 (1H, br); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  13.6 (q), 18.7 (t), 26.8 (t), 30.2 (t), 38.6 (t), 54.3 (d), 178.3 (s); MS 127 (M<sup>+</sup>), 84 (100); HRMS Calcd for C<sub>7</sub>H<sub>13</sub>ON 127.0997, Found 127.1001; [ $\alpha$ ]<sub>D</sub><sup>26</sup> -9.20 (c 1.19, CHCl<sub>3</sub>).

**(2S)-(+)-2-Heptyl-5-oxopyrrolidine (8, R = *n*-C<sub>7</sub>H<sub>15</sub>)**

To a stirred suspension of CuI (4.52 g, 23.8 mmol) in THF (30 mL) was added a solution of *n*-C<sub>7</sub>H<sub>15</sub>MgBr, prepared from *n*-C<sub>7</sub>H<sub>15</sub>Br (7.34 mL, 52.31 mmol) and Mg (1.26 g, 52.31 mmol) in THF (50 mL) at reflux, at -35 °C, and the resulting suspension was stirred at same temperature for 30 min. To the suspension was added a solution of **1** (2.14 g, 9.51 mmol) in THF (15 mL) at -35 °C, and then the reaction mixture was stirred at -35 °C for 18 h. The reaction was quenched with satd. NH<sub>4</sub>Cl (aq) solution, and the insoluble material was filtered off and washed with CHCl<sub>3</sub>. The organic layer was separated, and the aqueous layer was extracted with CHCl<sub>3</sub> (30 mL  $\times$  2). The organic layer and extracts were combined, dried over MgSO<sub>4</sub>, and evaporated to give the residue, that was chromatographed on silica gel (50 g, hexane/acetone=20:1–5:1) to give **8** (R = *n*-C<sub>7</sub>H<sub>15</sub>, 870 mg, 50%) as a pale yellow oil.

IR (neat) 3209, 1698, 1284 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>)  $\delta$  0.88 (3H, t, *J* = 7.2 Hz), 1.30 (9H, m), 1.43–1.54 (2H, m), 1.68–1.76 (2H, m), 2.23 (1H, m), 2.33 (2H, m), 3.62 (1H, quint, *J* = 6.9 Hz), 5.77 (1H, br); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  13.84 (q), 22.4 (t), 25.5 (t), 26.9 (t), 28.9 (t), 29.2 (t), 30.3 (t), 31.5 (t), 36.5 (t), 54.6 (d), 178.3 (s); MS 183 (M<sup>+</sup>), 84 (100); HRMS Calcd for C<sub>11</sub>H<sub>21</sub>ON 183.1623, Found 183.1608; [ $\alpha$ ]<sub>D</sub><sup>26</sup> +8.30 (c 0.89, CH<sub>2</sub>Cl<sub>2</sub>), Lit. [ $\alpha$ ]<sub>D</sub><sup>26</sup> +9.0 (c 2.0, CH<sub>2</sub>Cl<sub>2</sub>).



**(2R)-(-)-5-Propyl-2-oxopyrrolidine-1-carboxylic acid *tert*-butyl ester**

To a stirred solution of **8** (R = *n*-Pr, 247 mg, 1.94 mmol) in MeCN (10 mL) was added DMAP (261 mg, 2.14 mmol) at 0 °C, and the reaction mixture was stirred at 0 °C for 30 min. To the reaction mixture was added Boc<sub>2</sub>O (508 mg, 2.33 mmol) at 0 °C, and then the resulting mixture was stirred at room temperature for 43 h. The solvent was removed under reduced pressure, and the residue was chromatographed on silica gel (20 g, hexane/acetone=30:1–15:1) to give the title Boc-imide (414 mg, 94%) as a pale yellow oil.

IR (neat) 1786, 1749, 1714, 1306, 1153 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.95 (3H, t, *J* = 7.2 Hz), 1.34 (1H, m), 1.39 (1H, m), 1.52 (9H, s), 1.69–1.78 (2H, m), 2.42 (1H, m), 2.57 (1H, m), 4.10 (1H, m); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 13.6 (q), 18.5 (t), 22.1 (t), 27.6 (q), 31.0 (t), 31.5 (t), 35.4 (t), 57.4 (d), 82.0 (s), 149.3 (s), 173.8 (s); MS 227 (M<sup>+</sup>), 84 (100); HRMS Calcd for C<sub>12</sub>H<sub>21</sub>O<sub>3</sub>N 227.1522, Found 227.1527; [α]<sub>D</sub><sup>26</sup> –62.86 (c 0.87, CHCl<sub>3</sub>).

**(2R)-(-)-2-Heptyl-5-oxopyrrolidine-1-carboxylic acid *tert*-butyl ester**

To a stirred solution of **8** (R = *n*-C<sub>7</sub>H<sub>15</sub>, 602 mg, 3.29 mmol) in MeCN (20 mL) was added DMAP (442 mg, 3.62 mmol) at 0 °C, and the reaction mixture was stirred at 0 °C for 30 min. To the reaction mixture was added Boc<sub>2</sub>O (862 mg, 3.95 mmol) at 0 °C, and then the resulting mixture was stirred at room temperature for 48 h. The solvent was removed under reduced pressure, and the residue was chromatographed on silica gel (40 g, hexane/acetone=30:1–15:1) to give the desired Boc-imide (866 mg, 93%) as a pale yellow oil.

IR (neat) 1788, 1750, 1714, 1308, 1153 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.88 (3H, t, *J* = 7.2 Hz), 1.28 (10H, m), 1.46 (1H, br), 1.52 (9H, s), 1.76 (2H, m), 2.08 (1H, m), 2.41 (1H, m), 2.53 (1H, m), 4.09 (1H, m); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 13.9 (q), 22.3 (t), 22.4 (t), 25.4 (t), 27.8 (q), 29.0 (t), 31.2 (t), 31.5 (t), 33.5 (t), 57.8 (d), 82.2 (s), 149.5 (s), 174.0 (s); MS 283 (M<sup>+</sup>), 57 (100); HRMS Calcd for C<sub>16</sub>H<sub>29</sub>O<sub>3</sub>N 283.2148, Found 283.2128; [α]<sub>D</sub><sup>26</sup> –57.34 (c 1.21, CHCl<sub>3</sub>).

**(1R)-(+)-[1-(3-Oxohept-7-enyl)butyl]carbamic acid *tert*-butyl ester (**9**, R = *n*-Pr)**

To a stirred solution of the above Boc-imide (1.08 g, 4.75 mmol) in THF (15 mL) was added a solution of 4-butenylMgBr, prepared from 1-bromo-4-butene (1.45 mL, 14.27 mmol) and Mg (342 mg, 14.27 mmol) in THF (60 mL) at reflux, and TMEDA (2.27 mL, 14.27 mmol) at –78 °C, and the resulting mixture was stirred at the same temperature for 1.5 h. The reaction was quenched with *i*-PrOH (5 mL), and the reaction mixture was diluted with Et<sub>2</sub>O. The organic layer was washed with 10% HCl (aq) solution, dried over MgSO<sub>4</sub>, and evaporated to give a pale yellow oil, that was chromatographed on silica gel (40 g, hexane/acetone=50:1–40:1) to give **9** (R = *n*-Pr, 1.28 g, 95%) as a colorless solid (mp 64–66 °C).

IR (KBr) 3349, 3083, 1707, 1685, 1528, 1174 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.89 (3H, t, *J* = 7.2 Hz), 1.26–1.38 (5H, br m), 1.42 (9H, s), 1.50 (1H, m), 1.72 (1H, m), 2.30 (2H, q, *J* = 7.2 Hz), 2.48 (3H, q, *J* = 7.2 Hz), 3.51 (1H, br), 4.23 (1H, br d, *J* = 9.1 Hz), 4.94 (1H, d, *J* = 9.5 Hz), 5.01 (1H, d, *J* = 15.0 Hz), 5.79 (1H, m); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 13.7 (q), 18.9 (t), 27.5 (t), 28.1 (q), 29.0 (t), 37.9 (t), 39.1 (t), 41.6 (t), 49.8 (d), 78.3 (s), 114.7 (t), 136.6 (d), 155.4 (s), 209.2 (s); MS 283 (M<sup>+</sup>), 57 (100); HRMS Calcd for C<sub>16</sub>H<sub>29</sub>O<sub>3</sub>N 283.2148, Found 283.2142; [α]<sub>D</sub><sup>26</sup> +5.10 (c 1.15, CHCl<sub>3</sub>).

**(1R)-(-)-[1-(3-Oxohept-7-enyl)octyl]carbamic acid *tert*-butyl ester (**9**, R = *n*-C<sub>7</sub>H<sub>15</sub>)**

To a stirred solution of the above Boc-imide (724 mg, 2.56 mmol) in THF (10 mL) was added a solution of 4-butenylMgBr, prepared from 1-bromo-4-butene (0.78 mL, 7.68 mmol) and Mg (185 mg, 7.68 mmol) in THF (40 mL) at reflux, and TMEDA (1.16 mL, 7.68 mmol) at –78 °C, and the resulting mixture was stirred at the same temperature for 1.5 h. The reaction was

quenched with *i*-PrOH (5 mL), and the reaction mixture was diluted with Et<sub>2</sub>O. The organic layer was washed with 10% HCl (aq) solution, dried over MgSO<sub>4</sub>, and evaporated to give a pale yellow oil, that was chromatographed on silica gel (30 g, hexane/acetone=50:1–30:1) to give **9** (R = *n*-C<sub>7</sub>H<sub>15</sub>, 850 mg, 98%) as a colorless solid (mp 48~50 °C).

IR (KBr) 3348, 3080, 1707, 1685, 1531, 1173 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.87 (3H, t, *J* = 7.2 Hz), 1.25 (12H, br), 1.43 (9H, s), 1.49 (1H, br), 1.77 (1H, m), 2.31 (2H, q-like, *J* = 7.2 Hz), 2.49 (4H, m), 3.50 (1H, br), 4.20 (1H, br d, *J* = 9.4 Hz), 4.95 (1H, d, *J* = 9.6 Hz), 5.01 (1H, d, *J* = 15.0 Hz), 5.79 (1H, m); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 14.0 (q), 22.5 (t), 25.8 (t), 27.6 (t), 28.3 (q), 29.1 (t), 29.2 (t), 29.3 (t), 31.6 (t), 35.9 (t), 39.3 (t), 41.8 (t), 50.3 (d), 78.6 (s), 114.8 (t), 136.8 (d), 155.5 (s), 209.5 (s); MS 339 (M<sup>+</sup>), 57 (100); HRMS Calcd for C<sub>16</sub>H<sub>28</sub>O<sub>3</sub>N 282.2068, Found 282.2091; [α]<sub>D</sub><sup>26</sup> -1.04 (*c* 1.05, CHCl<sub>3</sub>).

#### (1R)-(-)-[1-(3-Oxoheptyl)pent-4-enyl]carbamic acid *tert*-butyl ester (**10**)

To a stirred solution of the Boc-imide (239 mg, 1.00 mmol), prepared from **2** (as described in the preparation of **3**), in THF (10 mL) was added a solution of *n*-PrMgBr, prepared from *n*-BuBr (0.32 mL, 3.00 mmol) and Mg (72 mg, 3.00 mmol) in THF (10 mL) at reflux, and TMEDA (0.48 mL, 3.00 mmol) at -78 °C, and the resulting mixture was stirred at the same temperature for 1.5 h. The reaction was quenched with *i*-PrOH (5 mL), and the reaction mixture was diluted with Et<sub>2</sub>O. The organic layer was washed with 10% HCl (aq) solution, dried over MgSO<sub>4</sub>, and evaporated to give a pale yellow oil, that was chromatographed on silica gel (20 g, hexane/acetone=50:1–30:1) to give **10** (292 mg, 98%) as a colorless solid (mp 42~43 °C).

IR (KBr) 3351, 3080, 1688, 1530, 1174 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.88 (3H, t, *J* = 7.2 Hz), 1.28 (2H, sext, *J* = 7.2 Hz), 1.42 (9H, s), 1.47–1.61 (5H, br m), 1.77 (1H, m), 2.08 (2H, q-like, *J* = 7.2 Hz), 2.38 (2H, t, *J* = 7.2 Hz), 2.46 (2H, t, *J* = 7.2 Hz), 3.50 (1H, br), 4.26 (1H, br d, *J* = 9.1 Hz), 4.94 (1H, d, *J* = 10.2 Hz), 5.00 (1H, d, *J* = 15.0 Hz), 5.78 (1H, m); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 13.7 (q), 22.1 (t), 25.7 (t), 28.2 (q), 29.0 (t), 30.0 (t), 35.0 (t), 39.1 (t), 42.4 (t), 49.8 (d), 78.4 (s), 114.5 (t), 137.6 (d), 155.4 (s), 210.4 (s); MS 240 (M<sup>+</sup>-57), 57 (100); HRMS Calcd for C<sub>13</sub>H<sub>22</sub>O<sub>3</sub>N 240.1600, Found 240.1616; [α]<sub>D</sub><sup>26</sup> -0.47 (*c* 1.20, CHCl<sub>3</sub>).

#### (2R,5S)-(-)-2-But-3-enyl-5-butylpyrrolidine-1-carboxylic acid *tert*-butyl ester (**11**)

To a stirred solution of **10** (279 mg, 0.94 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (7 mL) was added a solution of (C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>B (96 mg, 0.19 mmol) and Ph<sub>3</sub>SiH (490 mg, 1.88 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) at -78 °C, and the reaction mixture was stirred at -78 °C for 30 min, and then at room temperature for 24 h. The reaction was quenched with Et<sub>3</sub>N (1.0 mL), and the resulting mixture was stirred at room temperature for 20 min. The mixture was diluted with Et<sub>2</sub>O, and the organic layer was washed successively with 10% AcOH (aq) solution and satd. NaHCO<sub>3</sub> (aq) solution, dried over MgSO<sub>4</sub>, and evaporated to give a pale yellow oil, that was chromatographed on silica gel (20 g, hexane/acetone=200:1–150:1) to give **11** (194 mg, 73%) as a colorless oil.

IR (neat) 3080, 1696, 1388 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.90 (3H, t, *J* = 7.2 Hz), 1.24–1.40 (6H, br m), 1.46 (9H, s), 1.61 (2H, br), 1.93 (4H, br m), 2.05 (2H, br m), 3.75 (2H, br), 4.80 (1H, d, *J* = 10.5 Hz), 5.03 (1H, d, *J* = 16.0 Hz), 5.82 (1H, m); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 14.1 (q), 22.7 (t), 28.5 (q), 28.6 (t), 29.4 (t), 30.7 (t), 35.1 (t), 35.6 (t), 57.8 (d), 58.3 (d), 78.6 (s), 114.2 (t), 138.2 (d), 154.6 (s); MS 281 (M<sup>+</sup>), 170 (100); HRMS Calcd for C<sub>17</sub>H<sub>31</sub>O<sub>2</sub>N 281.2355, Found 281.2355; [α]<sub>D</sub><sup>26</sup> -2.72 (*c* 1.15, CHCl<sub>3</sub>).

**(2R,5S)-(-)-2-(4-Ethoxycarbonylbut-3-enyl)-5-butylpyrrolidine-1-carboxylic acid *tert*-butyl ester (12)**

To a stirred solution of **11** (155 mg, 0.55 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) were added Grubbs 2<sup>nd</sup> catalyst (19 mg, 0.022 mmol) and ethyl acrylate (0.30 mL, 2.76 mmol), and the resulting mixture was refluxed for 5 h. After cooling, the solvent was removed under reduced pressure, and the residue was chromatographed on silica gel (20 g, hexane/acetone=100:1–60:1) to give **12** (186 mg, 96%) as a pale yellow oil.

IR (neat) 1718, 1696, 1389, 1174 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.89 (3H, t, *J* = 7.2 Hz), 1.20–1.37 (10H, br, including at δ 1.28, 3H, t, *J* = 7.2 Hz), 1.45 (9H, s), 1.60 (2H, br), 1.91 (3H, br), 2.20 (2H, br), 3.77 (2H, br), 4.17 (2H, q, *J* = 7.2 Hz), 5.82 (1H, d, *J* = 15.7 Hz), 6.96 (1H, dt, *J* = 15.7, 6.8 Hz); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 14.0 (q), 14.1 (q), 22.6 (t), 28.4 (q), 28.5 (t), 29.1 (t), 29.4 (t), 34.2 (t), 35.6 (t), 57.6 (d), 58.3 (d), 59.9 (t), 78.8 (s), 121.1 (d), 148.3 (d), 154.6 (s), 166.1 (s); MS 353 (M<sup>+</sup>), 252 (100); HRMS Calcd for C<sub>20</sub>H<sub>35</sub>O<sub>4</sub>N 353.2564, Found 353.2544; [α]<sub>D</sub><sup>26</sup> -2.89 (*c* 0.75, CHCl<sub>3</sub>).

**(3R,5S,8S)-(+)-(5-Butylhexahydropyrrolizin-3-yl)acetic acid ethyl ester (13)**

To a stirred solution of **12** (100 mg, 0.28 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (7 mL) was added AlCl<sub>3</sub> (83 mg, 0.62 mmol) at 0 °C, and the resulting suspension was stirred at room temperature for 24 h. The reaction was quenched with satd. NaHCO<sub>3</sub> (aq) solution, and the organic layer was separated. The aqueous layer was extracted with CHCl<sub>3</sub> (10 mL×5), and the organic layer and extracts were combined, dried over K<sub>2</sub>CO<sub>3</sub>, and evaporated to give a residue. To a stirred solution of this residue in CH<sub>2</sub>Cl<sub>2</sub> (7 mL) was added K<sub>2</sub>CO<sub>3</sub> (78 mg, 0.57 mmol), and the resulting suspension was stirred at room temperature for 48 h. The insoluble material was filtered off, and washed with CH<sub>2</sub>Cl<sub>2</sub>. The filtrate was evaporated to afford a residue, that was chromatographed on silica gel (20 g, hexane/acetone=20:1–8:1) to give **13** (66 mg, 92%) as a pale yellow oil.

IR (neat) 1733, 1181 cm<sup>-1</sup>; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.90 (3H, t, *J* = 7.2 Hz), 1.26 (3H, t, *J* = 7.2 Hz), 1.28–1.43 (7H, br m), 1.52 (1H, m), 1.69 (1H, br), 1.78 (3H, m), 2.28 (1H, dd, *J* = 15.0, 9.8 Hz), 2.57 (1H, dd, *J* = 15.0, 4.3 Hz), 3.00 (1H, br), 3.40 (1H, m), 3.57 (1H, br), 4.11 (2H, q, *J* = 7.2 Hz); <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 14.1 (q), 14.2 (q), 22.9 (t), 30.2 (t), 30.4 (t), 30.8 (t), 31.0 (t), 31.3 (t), 32.4 (t), 43.2 (t), 54.2 (d), 60.0 (t), 63.7 (d), 65.6 (d), 172.1 (s); MS 253 (M<sup>+</sup>), 196 (100); HRMS Calcd for C<sub>15</sub>H<sub>27</sub>O<sub>2</sub>N 253.2042, Found 253.2058; [α]<sub>D</sub><sup>26</sup> +21.06 (*c* 0.83, CHCl<sub>3</sub>).

**(3S,5S,8S)-(+)-3,5-Dibutylhexahydropyrrolizine (14)**

To a stirred solution of **13** (70 mg, 0.28 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (7 mL) was added a solution of DIBAL (0.98 M in hexane, 0.31 mL, 0.31 mmol) at -50 °C, and the reaction mixture was stirred at -50 °C for 30 min. The reaction was quenched with MeOH, and satd. Rochelle (aq) solution, and the organic layer was separated. The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (5 mL × 3), and the organic layer and extracts were combined, dried over K<sub>2</sub>CO<sub>3</sub>, and evaporated to give a pale yellow oil, that was used directly in the next step.

To a stirred suspension of EtP<sup>+</sup>Ph<sub>3</sub>Br<sup>-</sup> (410 mg, 1.12 mmol) in THF (7 mL) was added a solution of *n*-BuLi (1.6 M in hexane, 0.6 mL, 0.98 mmol) at 0 °C, and the resulting orange suspension was stirred at 0 °C for 10 min. To the suspension was added a solution of the above aldehyde in THF (3 mL) at 0 °C, and the resulting suspension was stirred at room temperature for 27 h. The reaction was quenched with H<sub>2</sub>O, and the aqueous mixture was extracted with Et<sub>2</sub>O (15 mL × 4). The organic extracts were combined, dried over K<sub>2</sub>CO<sub>3</sub>, and evaporated to give a residue, that was chromatographed on silica gel (20 g, hexane/acetone=25:1–10:1) to give the corresponding olefin (31 mg, 50%) as a mixture of *E*- and *Z*-isomers.

$^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.90 (3H, t,  $J = 6.8$  Hz), 1.24–1.41 (8H, br m), 1.48 (1H, m), 1.62 (3H, d,  $J = 6.8$  Hz), 1.72–1.93 (4H, br m), 1.95–2.08 (2H, m), 2.31 (1H, br), 2.96 (1H, m), 3.04 (1H, br), 3.60 (1H, br), 5.36–5.51 (2H, br m).

To a stirred solution of the above olefin (20 mg, 0.09 mmol) in EtOAc (3 mL) was added 10% Pd/C (10 mg), and the resulting suspension was stirred under a hydrogen atmosphere at 1 atm for 45 h. The catalyst was removed by filtration and the filtrate was evaporated to give **14** (20 mg, quant) as a pale yellow oil.

IR (neat) 2928, 2858, 1457, 1099  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  0.89 (3H, t,  $J = 7.2$  Hz), 0.90 (3H, t,  $J = 6.8$  Hz), 1.20–1.51 (15H, br m), 1.60 (1H, m), 1.81 (2H, m), 1.98 (2H, m), 2.91 (1H, m), 3.10 (1H, br), 3.68 (1H, br);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  14.1 (q), 14.2 (q), 22.9 (t), 23.0 (t), 29.4 (t), 30.1 (t), 30.4 (t), 30.7 (t), 30.9 (t), 31.6 (t), 32.1 (t), 37.6 (t), 58.2 (d), 64.0 (d), 66.0 (d); MS 223 ( $\text{M}^+$ ), 55 (100); HRMS Calcd for  $\text{C}_{15}\text{H}_{29}\text{N}$  223.2300, Found 223.2312;  $[\alpha]_{\text{D}}^{26} +29.77$  ( $c$  0.40,  $\text{CHCl}_3$ ).

### (1R)-(+)-[1-(3-Oxohept-7-enyl)pentyl]carbamic acid *tert*-butyl ester (**15**)

To a stirred solution of **2**<sup>8</sup> (337 mg, 2.42 mmol) in MeCN (20 mL) was added DMAP (326 mg, 2.67 mmol) at 0 °C, and the reaction mixture was stirred at 0 °C for 30 min. To the reaction mixture was added  $\text{Boc}_2\text{O}$  (633 mg, 2.90 mmol) at 0 °C, and the resulting mixture was stirred at room temperature for 45 h. The volatiles were evaporated, and the residue was chromatographed on silica gel (20 g, hexane/acetone=15:1) to give the Boc-imide (550 mg, 95%) as a colorless oil.

IR (neat) 3078, 1785, 1750, 1714, 1308, 1153  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  1.52 (9H, s), 1.59 (1H, m), 1.78 (1H, m), 1.90 (1H, m), 2.03–2.19 (3H, br m), 2.43 (1H, ddd,  $J = 9.0, 8.5, 2.1$  Hz), 2.57 (1H, dd,  $J = 9.0, 8.5$  Hz), 4.12 (1H, m), 5.01 (1H, d,  $J = 10.7$  Hz), 5.04 (1H, d,  $J = 15.0$  Hz), 5.80 (1H, m);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  21.9 (t), 27.7 (q), 29.6 (t), 31.0 (t), 32.4 (t), 57.1 (d), 82.2 (s), 115.0 (t), 136.7 (d), 149.3 (s), 173.7 (s); MS 182 ( $\text{M}^+ - 57$ ), 84 (100); HRMS Calcd for  $\text{C}_{13}\text{H}_{21}\text{O}_3\text{N}$  239.1520, Found 239.1534;  $[\alpha]_{\text{D}}^{26} -56.68$  ( $c$  2.34,  $\text{CHCl}_3$ ).

To a stirred solution of the above Boc-imide (239 mg, 1 mmol) in EtOAc (10 mL) was added 10% Pd/C (50 mg), and the resulting suspension was stirred under a hydrogen atmosphere at 1 atm for 48 h. The catalyst was removed by filtration and the filtrate was evaporated to give the corresponding imide as a pale yellow oil, that was used directly in the next step.

To a stirred solution of the imide prepared above in THF (5 mL) was added a solution of 4-butenylMgBr, prepared from 4-bromo-1-butene (0.30 mL, 3.00 mmol) and Mg (72 mg, 3.00 mmol) in THF (8 mL) at reflux, and TMEDA (0.48 mL, 3.00 mmol) at  $-78$  °C, and the resulting mixture was stirred at the same temperature for 1.5 h. The reaction was quenched with *i*-PrOH (5 mL), and the reaction mixture was diluted with  $\text{Et}_2\text{O}$ . The organic layer was washed with 10% HCl (aq) solution, dried over  $\text{MgSO}_4$ , and evaporated to give a pale yellow oil, that was chromatographed on silica gel (20 g, hexane/acetone=50:1–30:1) to give **15** (284 mg, 96%) as a colorless solid (mp 38~39 °C).

IR (KBr) 3355, 3070, 1709, 1685, 1530, 1174  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.88 (3H, t,  $J = 7.2$  Hz), 1.30 (6H, br m), 1.43 (9H, s), 1.51 (1H, br), 1.78 (1H, m), 2.31 (2H, q-like,  $J = 6.9$  Hz), 2.49 (4H, m), 3.50 (1H, br), 4.21 (1H, br d,  $J = 9.0$  Hz), 4.96 (1H, d,  $J = 10.2$  Hz), 5.00 (1H, d,  $J = 16.0$  Hz), 5.82 (1H, m);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  13.6 (q), 22.2 (t), 27.3 (t), 27.7 (t), 28.0 (q), 28.9 (t), 35.2 (t), 39.0 (t), 41.4 (t), 49.9 (d), 78.0 (s), 114.5 (t), 136.5 (d), 155.3 (s), 209.0 (s); MS 297 ( $\text{M}^+$ ), 57 (100); HRMS Calcd for  $\text{C}_{17}\text{H}_{31}\text{O}_3\text{N}$  297.2304, Found 297.2315;  $[\alpha]_{\text{D}}^{26} +2.76$  ( $c$  1.04,  $\text{CHCl}_3$ ).

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


## Acknowledgment

This work was supported in part by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science. Work at NIH was supported by intramural funds of NIDDK.



## REFERENCES and NOTES

- (a) Daly JW. *J. Med. Chem* 2003;46:445–452. [PubMed: 12570366] (b) Daly JW, Spande TF, Garraffo HM. *J. Nat. Prod* 2005;68:1556–1575. [PubMed: 16252926]
- (a) Tsuneki H, You Y, Toyooka N, Kagawa S, Kobayashi S, Sasaoka T, Nemoto H, Kimura I, Dani JA. *Mol. Pharmacol* 2004;66:1061–1069. [PubMed: 15258256] (b) Toyooka N, Kobayashi S, Zhou D, Tsuneki H, Wada T, Sakai H, Nemoto H, Sasaoka T, Garraffo HM, Spande TF, Daly JW. *Bioorg. Med. Chem. Lett* 2007;17:5872–5875. [PubMed: 17827002] (c) Kobayashi S, Toyooka N, Zhou D, Tsuneki H, Wada T, Sasaoka T, Sakai H, Nemoto H, Garraffo HM, Spande TF, Daly JW. *Beilstein J. Org. Chem* 2007;3:30. (d) Toyooka N, Zhou D, Kobayashi S, Tsuneki H, Wada T, Sakai H, Nemoto H, Sasaoka T, Tezuka Y, Subehan, Kadota S, Garraffo HM, Spande TF, Daly JW. *Synlett* 2008:61–64.
- (a) Dumbacher JP, Beehler BM, Spande TF, Garraffo HM, Daly JW. *Science* 1992;258:799–801. [PubMed: 1439786] (b) Dumbacher JP, Spande TF, Daly JW. *Proc. Natl. Acad. Sci. USA* 2000;97:12970–12975. [PubMed: 11035772] (c) Dumbacher JP, Wako A, Derrickson SR, Samuelson A, Spande TF, Daly JW. *Proc. Natl. Acad. Sci. USA* 2004;101:15857–15860. [PubMed: 15520388]
- Clark VC, Raxworthy CJ, Rakotomalala V, Sierwald P, Fisher BL. *Proc. Natl. Acad. Sci. USA* 2005;102:11617–11622. [PubMed: 16087888]
- (a) Takano S, Otaki S, Ogasawara K. *J. Chem. Soc. Chem. Commun* 1983:1172–1174. (b) Arseniyadis S, Huang PQ, Husson HP. *Tetrahedron Lett* 1988;29:1391–1394. (c) Takahata H, Bandoh H, Momose T. *Tetrahedron: Asymmetry* 1991;2:351–352. (d) Takahata H, Bandoh H, Momose T. *J. Org. Chem* 1992;57:4401–4404. (e) Grandjean C, Rosset S, Celerier JP, Lhommet G. *Tetrahedron Lett* 1993;34:4517–4518. (f) Oppolzer W, Bochet CG, Merifield E. *Tetrahedron Lett* 1994;35:7015–7018. (g) Cuny GD, Buchwald SL. *Synlett* 1995:519–522. (h) Dhimane H, Vanucci-Bacque C, Hamon L, Lhommet G. *Eur. J. Org. Chem* 1998:1955–1963. (i) Arredondo VM, Tian S, McDonald FE, Marks TJ. *J. Am. Chem. Soc* 1999;121:3633–3639. (j) Takahata H, Takahashi S, Azer N, Eldefrawi AT, Eldefrawi ME. *Bioorg. Med. Chem. Lett* 2000;10:1293–1295. [PubMed: 10866403]
- (a) Toyooka N, Zhou D, Nemoto H. *J. Org. Chem* 2008;73:4575–4577. [PubMed: 18507441] (b) Toyooka N, Zhou D, Nemoto H, Tezuka Y, Kadota S, Jones TH, Garraffo HM, Spande TF, Daly JW. *Synlett* 2008:1894–1896. (c) Toyooka N, Tsuneki H, Kobayashi S, Zhou D, Kawasaki M, Kimura I, Sasaoka T, Nemoto. *Curr. Chem. Biol* 2007;1:97–114. (d) Toyooka N, Tsuneki H, Nemoto H. *Yuki Gosei Kagaku Kyokaishi* 2006;64:49–60. Toyooka, N.; Nemoto, H. *New Methods for the Asymmetric Synthesis of Nitrogen Heterocycles*. Vicario, JL., editor. India: Research Signpost; 2005. p. 149–163. Toyooka, N.; Nemoto, H. *Recent Research Developments in Organic Chemistry*. Pandalai, SG., editor. Vol. 6. India: TRANSWORLD RESEARCH NETWORK; 2002. p. 611–624.
- Kamimura A, Nagata Y, Kadowaki A, Uchida K, Uno H. *Tetrahedron* 2007;63:11856–11861.
- Hjelmgaard T, Sjøtofte I, Tanner D. *J. Org. Chem* 2005;70:5688–5697. [PubMed: 15989354]
- Brenneman JB, Machauer R, Martin SF. *Tetrahedron* 2004;60:7301–7314.
- Scholl M, Ding S, Lee CW, Grubbs RH. *Org. Lett* 1999;1:953–956. [PubMed: 10823227]
- Exposure of the stereoisomer at the 3-position (all *cis*-pyrrolizidine) of **6** (R = Et) to the same reaction conditions (K<sub>2</sub>CO<sub>3</sub> in CH<sub>2</sub>Cl<sub>2</sub> at room temperature for 48 h) recovered the starting material, and no formation of **6** (R = Et) was observed.
- Garraffo HM, Spande TF, Daly JW, Baldessari A, Gros EG. *J. Nat. Prod* 1993;56:357–373. [PubMed: 8482947]
- No separation of a mixture of **7** and ent-**7** (ca. 1:1) was observed with either of two beta cyclodextrin chiral columns. One column was a 30 m × 0.25 mm i.d. (0.25 μm film thickness) beta Dex 120 column



(Supelco); the other column was a 25 m × 0.22 mm i.d. (0.25 μm film thickness) permethylated beta cyclodextrin column (SGE). Both were operated at a head pressure of 20 psi and used a gas chromatograph with flame ionization detection. A 3390A recorder integrator was used and a temperature program of 100 °C to 150 °C at 1 or 2 °C/min. A separation of a mixture of **14** and ent-**14** was not attempted.




(-)-xenovenine



or


**2510**

**Figure 1.**  
Structure of xenovenine and 2510




**Figure 2.**  
Kinetic control on pyrrolizidine formation




**Scheme 1.**

Synthesis of 7 ( $R = n\text{-Pr}$ ,  $R' = n\text{-C}_7\text{H}_{15}$  or  $R = n\text{-C}_7\text{H}_{15}$ ,  $R' = n\text{-Pr}$ )

**Scheme 2.**

Synthesis of ent-7 ( $R = n\text{-Pr}$ ,  $R' = n\text{-C}_7\text{H}_{15}$  or  $R = n\text{-C}_7\text{H}_{15}$ ,  $R' = n\text{-Pr}$ )



**Scheme 3.**  
Synthesis of both enantiomers of *trans*-223B