



Published in final edited form as:

*Metabolism*. 2009 October ; 58(10): 1369–1371. doi:10.1016/j.metabol.2009.04.023.

## No association between TCF7L2 rs7903146 and euglycemic-clamp derived insulin sensitivity in a mixed-age cohort

Laura J. Rasmussen-Torvik, PhD<sup>1</sup>, James S. Pankow, PhD<sup>1</sup>, David R. Jacobs Jr, PhD<sup>1,2</sup>, and Alan R. Sinaiko, MD<sup>3,1</sup>

<sup>1</sup>Division of Epidemiology and Community Health, University of Minnesota, Minneapolis, MN

<sup>2</sup>Department of Nutrition, University of Oslo, Oslo, Norway

<sup>3</sup>Department of Pediatrics, University of Minnesota, Minneapolis, MN

### Abstract

There are conflicting reports about the significance of TCF7L2 SNP rs7903146, a SNP found to be associated with Type 2 diabetes in several genome-wide association studies, and insulin sensitivity. The association of rs7903146 and euglycemic-clamp derived insulin sensitivity was measured in a cohort of children and their parents. 470 whites (from 226 families) and 89 African Americans (from 48 families) were included in the analysis. No significant associations were seen between rs7903146 and insulin sensitivity. Adjusted genotype means were consistent across races and generational subgroups.

### Introduction

Studies have sought to determine the mechanisms underlying the well-established association between SNP rs7903146 in TCF7L2 and type 2 diabetes. Studies that have examined the association of this SNP with insulin sensitivity derived from the euglycemic clamp [1-3] or the IVGTT [4-6] have produced inconsistent results. SNP rs7903146 has been associated with insulin sensitivity in American whites and elderly Danish twins [3,5,6] but not in Hispanic Americans, Germans, African Americans, or Swedish men [1,2,4,6].

### Methods

Subjects were drawn from a longitudinal study of cardiovascular risk factors in adolescents. Details of the recruitment have been published previously [7]. The study was approved by the Human Subjects Committee of the University of Minnesota. Briefly, in 1996, Minneapolis school children were randomly selected with stratification according to sex, race (African American and white), and systolic blood pressure. Informed consent was obtained for 401 children (probands) and their parents. Probands returned for a subsequent study visit at mean age 19 at which time parents and siblings of the probands also were recruited into the study and underwent many of the same measurements as the probands. Individuals with both

---

Address correspondence to: Laura J. Rasmussen-Torvik, PhD., Division of Epidemiology and Community Health, University of Minnesota School of Public Health, 1300 S. Second St, Suite 300, Minneapolis, MN 55454-1015, rasm0218@umn.edu, Phone: 612-626-7921, Fax: 612-624-0315.

The authors have no conflicts of interest to disclose.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

rs7903146 genotypes and insulin sensitivity measurements from this visit were included in the analysis. Individuals who self-reported a diagnosis of diabetes or use of diabetes medication were excluded as were individuals with a fasting glucose measurement greater than 126 mg/dl.

Insulin sensitivity was determined from the euglycemic clamp. Euglycemic clamp studies were conducted in the University of Minnesota Clinical Research Center after a 12 hour fast as previously described [7]. Plasma glucose was measured at baseline and every five minutes during the clamp. The insulin infusion was started at time 0 and continued at 1 mU/kg/min for 3 hours. An infusion of 20% glucose was started at time 0 and adjusted, based on plasma glucose levels, to maintain plasma glucose at 100 mg/dl. Insulin sensitivity was determined from the amount of glucose administered over the final 40 minutes of the euglycemic clamp, and expressed as glucose utilization/ kg lean body mass/ minute. Percentage of body fat and lean body mass (LBM), or fat-free mass were calculated by DEXA.

SNP associations were calculated in a mixed linear regression model (SAS v.9.1, Cary, NC). A compound symmetry correlation structure was specified to account for expected correlation within families and a sandwich estimator was used to calculate the variance. All analyses were adjusted for age and sex. A 2 d.f. association test was used for SNP genotypes, except for association analyses in African Americans, where a 1 d.f. test was used because of small numbers; in this case individuals having CT or TT genotypes were pooled for analysis.

## Results

After exclusions there were 470 whites from 226 families and 89 African Americans from 48 families in the sample. Forty-seven % of the families included 1 parent, 24% included 2 parents, and 29 % included no parents. Thirty-eight % of the families included 1 child, 37% included 2 children, 16% included 3 or more children, and 9% included no children. Forty-nine % of the sample was male.

Table 1 shows the results of the regression of clamp-derived insulin sensitivity on TCF7L2 SNP rs7903146 in the total study sample, and in racial and generational subgroups. Because the patterns of genotype-specific means were similar in both races, the decision was made to combine the racial groups for analyses. In the total sample there was no significant association between insulin sensitivity and rs7903146 ( $p = .85$ ,  $F$  value = 0.16 on 2 d.f.). There was also no significant association observed in any of the subgroups, and similar patterns of genotype-specific means (a slight increase with the heterozygous or homozygous minor allele genotypes) were seen for most subgroup analyses. Additional adjustment for BMI in the total sample and in racial and generational subgroups did not materially change the results.

## Discussion

The results of this study differ from previous studies that found a significant association between rs7903146 and insulin sensitivity in whites [3,5,6]. Furthermore, the pattern of insulin sensitivity by genotype observed in this study (a slight increase with the heterozygous or homozygous minor allele genotypes) is opposite to other published studies [5,6], and not consistent with the association of the T allele and type 2 diabetes [5,6]. The inconsistencies between this study and others may be due to the use of the IVGTT, as some surrogate measures of insulin sensitivity/resistance have been shown to have different genetic determinants than clamp-derived insulin sensitivity [8], or to smaller sample sizes in previous studies which may have lead to spurious results [3,5,6]. Strengths of this study include a relatively large sample size and use of the gold-standard measure of insulin sensitivity. Weaknesses include the small number of African Americans in the sample which limited the power of analyses in this

subgroup. In conclusion, this analysis offers additional evidence that the rs7903146 SNP in TCF7L2 is not associated with insulin resistance in whites [1,2,4]. This analysis should be repeated in larger African American cohorts to verify the lack of association in this racial group.

## Acknowledgments

This study was supported by grants HL52851 and M01RR00400 from the National Institutes of Health. LJR-T is supported by NHLBI training grant T32 HL07779.

The Human Subjects Committee of the University of Minnesota approved this study.

## References

1. Dahlgren A, Zethelius B, Jensevik K, et al. Variants of the TCF7L2 gene are associated with beta cell dysfunction and confer an increased risk of type 2 diabetes mellitus in the ULSAM cohort of Swedish elderly men. *Diabetologia* 2007;50:1852–7. [PubMed: 17618413]
2. Schafer SA, Tschritter O, Machicao F, et al. Impaired glucagon-like peptide-1-induced insulin secretion in carriers of transcription factor 7-like 2 (TCF7L2) gene polymorphisms. *Diabetologia* 2007;50:2443–50. [PubMed: 17661009]
3. Wegner L, Hussain MS, Pilgaard K, et al. Impact of TCF7L2 rs7903146 on insulin secretion and action in young and elderly Danish twins. *J Clin Endocrinol Metab* 2008;93:4013–9. [PubMed: 18611970]
4. Palmer ND, Lehtinen AB, Langefeld CD, et al. Association of TCF7L2 gene polymorphisms with reduced acute insulin response in Hispanic Americans. *J Clin Endocrinol Metab* 2008;93:304–9. [PubMed: 17971425]
5. Damcott CM, Pollin TI, Reinhart LJ, et al. Polymorphisms in the transcription factor 7-like 2 (TCF7L2) gene are associated with type 2 diabetes in the Amish: replication and evidence for a role in both insulin secretion and insulin resistance. *Diabetes* 2006;55:2654–9. [PubMed: 16936218]
6. Elbein SC, Chu WS, Das SK, et al. Transcription factor 7-like 2 polymorphisms and type 2 diabetes, glucose homeostasis traits and gene expression in US participants of European and African descent. *Diabetologia* 2007;50:1621–30. [PubMed: 17579832]
7. Sinaiko AR, Jacobs DR Jr, Steinberger J, et al. Insulin resistance syndrome in childhood: associations of the euglycemic insulin clamp and fasting insulin with fatness and other risk factors. *J Pediatr* 2001;139:700–7. [PubMed: 11713450]
8. Rasmussen-Torvik LJ, Pankow JS, Jacobs DR, et al. Heritability and genetic correlations of insulin sensitivity measured by the euglycaemic clamp. *Diabet Med* 2007;24:1286–9. [PubMed: 17956454]

**Table 1**  
Adjusted\* means and confidence intervals of insulin sensitivity by rs7903146 Genotype

Population	Genotype Means (95% C.I.)** of Insulin Sensitivity in mg/kg/min		p-value <sup>†</sup>	
All	CC= 10.9 (10.6, 11.3) N = 314	CT= 11.1 (10.4, 11.7) N = 205	TT= 11.1 (9.9, 12.2) N = 40	.85
Whites	CC= 11.1 (10.6, 11.5) N = 262	CT= 11.1 (10.5, 11.7) N = 175	TT= 11.8 (10.6, 13.0) N = 33	.48
African Americans	CC= 10.1 (9.2, 11.0) N = 52	CT/TT= 10.7 (9.4, 12.0) N = 37		.40 <sup>††</sup>
Parents (ages 32-64, median = 48)	CC= 11.1 (10.4, 11.8) N = 117	CT= 11.4 (10.6, 12.3) N = 81	TT= 11.7 (9.6, 13.8) N = 14	.75
Offspring (ages 11-30, median = 18)	CC= 10.7 (10.3, 11.2) N = 197	CT= 11.0 (10.2, 11.7) N = 124	TT= 10.5 (9.2, 11.7) N = 26	.77

\* All regression analyses adjusted for age and sex

\*\* 95% confidence intervals presented in parentheses next to genotype means

<sup>†</sup> for a 2 d.f. test, except in African Americans

<sup>††</sup> where CT and TT genotypes were combined for analysis due to small numbers resulting in a 1 degree of freedom test