



Published in final edited form as:

Arterioscler Thromb Vasc Biol. 2008 September ; 28(9): 1614–1620. doi:10.1161/ATVBAHA.107.158725.

Reevaluation of the Role of VEGF-B Suggests a Restricted Role in the Revascularization of the Ischemic Myocardium

Xuri Li, Marc Tjwa, Inge Van Hove, Berndt Enholm, Elke Neven, Karri Paavonen, Michael Jeltsch, Toni Diez Juan, Richard E. Sievers, Emmanuel Chorianopoulos, Hiromichi Wada, Maarten Vanwildemeersch, Agnes Noel, Jean-Michel Foidart, Matthew L. Springer, Georges von Degenfeld, Mieke Dewerchin, Helen M. Blau, Kari Alitalo, Ulf Eriksson, Peter Carmeliet, and Lieve Moons

Vesalius Research Center (X.L., M.T., I.V.H., E.N., T.D.J., E.C., H.W., M.D., P.C., L.M.), VIB, 3000 Leuven, Belgium; Vesalius Research Center (X.L., M.T., I.V.H., E.N., T.D.J., E.C., H.W., M.D., P.C., L.M.), K.U. Leuven, 3000 Leuven, Belgium; Ludwig Institute for Cancer Research (X.L., M.V., U.E.), Karolinska Institute, Stockholm, Sweden; Molecular/Cancer Biology Laboratory (B.E., K.P., M.J., K.A.), Biomedicum Helsinki, University of Helsinki, Finland; Division of Cardiology (R.E.S., M.L.S.), University of California San Francisco; Laboratory of Biology of Tumors and Development (A.N., J.-M.F.), University of Liège, Belgium; and Baxter Laboratory in Genetic Pharmacology (M.L.S., G.v.D., H.M.B.), Stanford University, Calif. X.L. was a visiting scientist from the Ludwig Institute for Cancer Research, Stockholm, at the CTG, Leuven while performing these studies

Abstract

Objective—The endogenous role of the VEGF family member vascular endothelial growth factor-B (VEGF-B) in pathological angiogenesis remains unclear.

Methods and Results—We studied the role of VEGF-B in various models of pathological angiogenesis using mice lacking VEGF-B (VEGF-B^{-/-}) or overexpressing VEGF-B₁₆₇. After occlusion of the left coronary artery, VEGF-B deficiency impaired vessel growth in the ischemic myocardium whereas, in wild-type mice, VEGF-B₁₆₇ overexpression enhanced revascularization of the infarct and ischemic border zone. By contrast, VEGF-B deficiency did not affect vessel growth in the wounded skin, hypoxic lung, ischemic retina, or ischemic limb. Moreover, VEGF-B₁₆₇ overexpression failed to enhance vascular growth in the skin or ischemic limb.

Conclusion—VEGF-B appears to have a relatively restricted angiogenic activity in the ischemic heart. These insights might offer novel therapeutic opportunities.

Keywords

VEGF-B; angiogenesis; arteriogenesis; collateral growth; cardiac ischemia; limb ischemia

Vascular endothelial growth factor (VEGF) is a key regulator of angiogenesis in health and disease by binding to VEGF receptor-2 (VEGFR-2),¹ but the angiogenic activity of its homologue VEGF-B, which binds VEGFR-1 (Flt-1), is less defined.^{2,3} By contrast, genetic

Correspondence to Peter Carmeliet, Vesalius Research Center (VRC), Flanders Institute for Biotechnology (VIB), K.U. Leuven, Campus Gasthuisberg, Herestraat 49, B-3000, Leuven, Belgium. peter.carmeliet@med.kuleuven.be.

Current address for M.T.: Leibniz AG, Center for Molecular Medicine, University of Frankfurt, Germany.

X.L. and M.T. contributed equally to this study.

Reprints: Information about reprints can be found online at <http://www.lww.com/reprints>

Disclosures: None.

and inhibition studies revealed that Flt-1, and its other specific ligand PlGF, stimulate pathological angiogenesis only.^{4,5}

VEGF-B is produced as 2 different isoforms: a heparin-binding VEGF-B₁₆₇ and diffusible VEGF-B₁₈₆ isoform.^{3,6,7} VEGF-B is widely expressed in many tissues and cell types, including cardiac and skeletal myocytes and endothelial and mural cells.^{3,6,7} In vitro, VEGF-B stimulates endothelial cell growth and proliferation.⁸ By contrast, loss-of-function studies failed to reveal a consistent role for VEGF-B in pathological angiogenesis. Indeed, in 2 lines, loss of VEGF-B did not cause vessel defects in the embryo or healthy adult mouse, whereas recovery of coronary flow was impaired after transient occlusion *ex vivo* but, apparently, not because of reduced coronary vessel growth.^{9,10} Moreover, loss of VEGF-B failed to reduce angiogenesis in the cornea⁹ or ischemic retina.¹¹ One study proposed a role for VEGF-B in pulmonary hypertension,¹² but this finding was contested by others.¹³ The effects of VEGF-B deficiency on revascularization of the infarcted myocardium *in vivo* remain unknown. Gain-of-function studies also showed inconsistent effects. Indeed, overexpression of VEGF-B stimulates angiogenesis in skin wounds and ischemic limbs^{14–16} or promotes hypertrophy of remote myocardium after myocardial infarction,¹⁷ whereas VEGF-B had negligible effects after adenoviral gene transfer in rabbit carotid arteries or normoxic hindlimbs.^{18,19} The therapeutic potential of VEGF-B (VEGF-B₁₆₇) to promote revascularization of ischemic myocardium remains unknown. Overall, the precise role of VEGF-B in pathological angiogenesis remains to be established.

As these controversies might be attributable to differences in genetic background or phenotyping methodology, we characterized the angiogenic role of VEGF-B in different mouse models of disease. By using VEGF-B^{-/-} mice on a pure C57BL/6 inbred background, and various strategies to overexpress VEGF-B, we found that VEGF-B promotes vessel growth in the ischemic heart but not in other organs.

Methods

For detailed Methods, please see the supplemental materials (available online at <http://atvb.ahajournals.org>).

Animals, Models, Histology, Immunohistochemistry, and Morphometric Analyses

VEGF-B^{-/-} mice⁹ were back-crossed onto a C57BL/6 background for 8 generations. C57BL/6 and NMRI nu/nu mice (all 8 to 12 weeks old) were obtained from Charles River Laboratories (Les Oincs, France) and adult SCID mice were from Taconic M&B Europe (Ry, Denmark). LacZ-tagged Flt1 mice, expressing β galactosidase (β -Gal) under the control of the Flt-1 promoter were kindly provided by J. Rossant (Toronto, Ontario, Canada).²⁰ Animal experiments were approved by local committees. All experimental approaches are described in detail in the supplemental methods.

Production and Administration of VEGF-B Protein, Plasmid, and Adenovirus

Recombinant human VEGF-B₁₆₇ (rhVEGF-B₁₆₇) protein was obtained from Amrad Corporation and delivered systemically via osmotic minipumps. Adenoviruses, constructed by cloning the murine PlGF-2, or the human VEGF-B₁₆₇ or VEGF-B₁₈₆ cDNA into the pACCMVpLpA plasmid, were intradermally or intravenously injected. A plasmid expressing murine VEGF-B₁₆₇ (pcDNA3.mVEGF-B₁₆₇) or an empty pcDNA3 plasmid was administered via muscle electroporation. These experimental methods are described in more detail in the supplemental methods.

Statistics

Data (mean±SEM) were analyzed using 2-tailed Student *t* test, with $P<0.05$ considered statistically significant.

Results

Loss of VEGF-B Impairs Revascularization After Myocardial Infarction

To study the potential of VEGF-B to stimulate revascularization of the ischemic heart (infarct and border zone), we used a previously established model of acute myocardial infarction (MI) in wild-type (WT) and VEGF-B^{-/-} mice.²¹ At 7 days after MI, the density of both thrombomodulin positive (TM⁺) capillaries and smooth muscle α -actin positive (SMA⁺) covered vessels in the infarct area of VEGF-B^{-/-} mice was only 65% of that of WT mice (n=14; $P<0.05$ in all groups; Figure 1A and 1B). Revascularization of the ischemic border zone was also impaired in VEGF-B^{-/-} mice (TM⁺ vessels: 340±39 vessels/mm² in WT versus 234±38 vessels/mm² in VEGF-B^{-/-} mice; n=5; $P<0.05$). The number of macrophages, infiltrating into the infarct area, was normal (Mac3⁺ area/infarct area: 0.22±0.06% in VEGF-B^{-/-} mice versus 0.22±0.05% in WT mice; n=12; $P=NS$). Thus, loss of VEGF-B impairs angiogenesis and vessel maturation in the ischemic heart.

We next analyzed whether administration of recombinant human (rh) VEGF-B₁₆₇ protein rescued the impaired myocardial revascularization in VEGF-B^{-/-} mice, and therefore administered rhVEGF-B₁₆₇, the predominant isoform in cardiac and skeletal muscle.⁷ Continuous systemic delivery of a daily dose of 1.5 μ g rhVEGF-B₁₆₇ for 1 week to VEGF-B^{-/-} mice normalized the impaired revascularization of the infarct (n=5; $P<0.05$; Figure 1A and 1B), and increased the growth of TM⁺ vessels in the ischemic border zone by 1.5-fold ($P<0.05$, n=5).

VEGF-B₁₆₇ Therapy Enhances Ischemic Myocardial Revascularization

We then used 3 different techniques to investigate whether delivery of VEGF-B stimulated revascularization of ischemic hearts in WT mice. First, we administered VEGF-B₁₆₇ protein. Pilot studies revealed that delivery of rVEGF₁₆₇ via osmotic minipumps significantly increased the VEGF-B₁₆₇ blood plasma levels (Note I, please see supplemental materials). VEGF-B₁₆₇ protein therapy indeed increased the density of capillaries and arterioles in the infarct area (TM⁺ vessels/mm²: 206±10 after vehicle versus 285±33 after rhVEGF-B₁₆₇; SMA⁺ vessels/mm²: 47±4 after vehicle versus 62±11 after rhVEGF-B₁₆₇; n=5; $P<0.05$; Figure 1C through 1F), and stimulated vessel growth in the ischemic border by 1.5-fold and 1.3-fold, respectively ($P<0.05$; n=5).

Second, similar results were obtained after intravenous injection of an adenoviral vector encoding hVEGF-B₁₆₇ (Ad.hVEGF-B₁₆₇), known to transduce hepatocytes, which then release the transgene product into the circulation for up to 21 days (Note II, please see supplemental materials). Compared to control Ad.RR5 virus, VEGF-B₁₆₇ gene transfer increased the densities of TM⁺ and SMA⁺ vessels in the infarct and in the ischemic border (n=9; $P<0.05$ versus Ad.RR5; Figure 1G and 1H) at 7 days after MI. VEGF-B₁₆₇ gene transfer did not stimulate macrophage recruitment (Mac3⁺ area/infarct area: 0.20±0.05% after Ad.hVEGF-B₁₆₇ versus 0.19±0.06% after Ad.RR5; n=9; $P=NS$).

Third, implantation of VEGF-B₁₆₇-expressing myoblasts, but not control LacZ⁺ myoblasts, in the ischemic border zone induced vessel growth in the ischemic myocardium (supplemental Figure I). Hence, using different techniques (delivery of protein or gene transfer) and routes of administration (locally or systemically), VEGF-B₁₆₇ therapy enhanced vessel growth in ischemic hearts of WT mice.

VEGF-B Does Not Affect Vessel Growth in Skin, Lung, or Retina

To further study the role of VEGF-B in pathological angiogenesis, we analyzed skin wound healing, using a linear skin incision model. Daily measurements revealed no defects in the rate or extent of skin wound healing in VEGF-B^{-/-} mice (Figure 2A through 2C). Consistent herewith, loss of VEGF-B failed to affect the number of endothelial cell-lined and mural cell-covered vessels in the granulation tissue at 5 days after wounding (CD31⁺ vessels/mm²: 270 ±16 in WT mice versus 290±20 in VEGF-B^{-/-} mice; SMA⁺ vessels/mm²: 67±9 in WT mice versus 62±7 in VEGF-B^{-/-} mice; n=5; P=NS). Macrophage infiltration was also normal (F4/80⁺ area as % of total granulation tissue area: 4.59±0.25 in WT mice versus 4.55±0.92 in VEGF-B^{-/-} mice; n=5; P=NS). Additional studies using adenoviral vectors to locally overexpress VEGFB₁₆₇ or VEGF-B₁₈₆ in the skin confirmed that VEGF-B does not affect the skin vasculature (supplemental Figure II).

Furthermore, loss of VEGF-B failed to affect mural cell recruitment and vessel remodeling in hypoxic lungs (supplemental Figure IIIA through IIIC) or neovascularization in ischemic retinas (supplemental Figure IIIE through IIIF). Together, VEGF-B plays a negligible role in vessel growth, maturation, and remodeling in normal or wounded skin, in hypoxic lungs, or in ischemic retinas.

Loss of VEGF-B Does Not Affect Revascularization of Ischemic Limbs

We next analyzed whether VEGF-B affects revascularization of ischemic limbs.²¹ In an established mouse model of hind-limb ischemia, the ischemic gastrocnemius muscle is revascularized by capillary angiogenesis, whereas collateral vessel growth occurs in the adductor muscle.²¹ In the ischemic gastrocnemius muscle, vessel densities in the regenerating areas were comparable in VEGF-B^{-/-} mice at 7 days after ischemia (Table). In addition, laser Doppler perfusion analysis and endurance/graded treadmill exercise tests confirmed normal revascularization of ischemic limbs in VEGF-B^{-/-} mice after ischemia (Table; supplemental Figure IVA and IVB). Moreover, to exclude the possibility that VEGF-B would only act as a modifier of PlGF, we analyzed mice lacking both VEGF-B and PlGF. However, compared to WT or VEGF-B^{-/-} mice, limb reperfusion was comparably reduced in mice lacking PlGF alone or in mice lacking both VEGF-B and PlGF (supplemental Figure IVC and IVD; Table), indicating that VEGF-B is redundant for limb reperfusion even in conditions of genetically crippled limb revascularization in PlGF^{-/-} mice.²² When analyzing in the adductor muscle the number and lumen size of preexisting collateral vessels, we found, again, no genotypic differences at 7 days after ischemia (Table; supplemental Figure IVE and IVF). Macrophage recruitment around the collaterals (supplemental Figure IVG through IVI) as well as in vitro macrophage activation were also unaffected by, respectively, loss or addition of VEGF-B (supplemental Figure IVG through IVI). Thus, endogenous VEGF-B is redundant for the revascularization of ischemic limbs.

VEGF-B₁₆₇ Therapy Does Not Enhance Revascularization in Ischemic Limbs

As our findings above do not exclude the possibility that overexpression of VEGF-B₁₆₇ might enhance collateral growth or capillary angiogenesis in ischemic limbs, as suggested by others,^{15,16} we overexpressed VEGF-B₁₆₇ in ischemic hindlimbs of WT mice using various established methods. To analyze the effect of VEGF-B₁₆₇ overexpression on both collateral growth and capillary angiogenesis in the ischemic limb, we intravenously injected the adenoviral vector Ad.hVEGF-B₁₆₇, similar as done for our MI experiments and resulting in increased VEGF-B plasma levels (Note II, please see supplemental materials). At 7 days after ligation, Ad.hVEGF-B₁₆₇ gene transfer did not improve angiogenesis in the gastrocnemius muscle (Table; Figure 3A and 3B) and failed to increase the number and size of preexisting collateral vessels in the adductor muscle (Table). VEGF-B₁₆₇ therapy did not stimulate infiltration of macrophages (numbers around the main collateral vessel: 14±2 after Ad.RR5

versus 12 ± 1 after Ad.hVEGF-B₁₆₇ gene transfer, $n=5$; $P=NS$), and failed to improve limb perfusion (Table). Also, VEGF-B₁₆₇ therapy failed to stimulate ischemic limb revascularization at later time points, thus excluding a delayed effect (Note III, please see supplemental materials).

To further assess any possible effect of VEGF-B on ischemic limb revascularization (in particular capillary angiogenesis), we implanted VEGF-B₁₆₇-expressing mouse myoblasts into the anterior tibialis muscles (supplemental Figure I). In healthy (nonligated) muscle, no vessel growth was induced at the engraftment site of VEGF-B₁₆₇-expressing myoblasts, as compared to control LacZ⁺ myoblasts (Figure 3E). To assess whether VEGF-B₁₆₇ affects vessel growth in ischemic skeletal muscle, myoblasts were implanted immediately after ligation of the femoral artery. Capillary density was increased in the ischemic anterior tibialis muscle at 28 days postligation, but implantation of VEGF-B₁₆₇-expressing myoblasts did not further enhance angiogenesis (Figure 3C through 3E). To exclude a transient effect on vessel growth, legs were also harvested 7 days after ischemia induction and myoblast implantation, again not resulting in increased vessel densities (not shown).

In addition, in vivo electroporation of a plasmid expressing murine VEGF-B₁₆₇ (pmVEGF-B₁₆₇) in the adductor muscle (to analyze collateral growth) was ineffective (Note IV, please see supplemental materials; Table). Overall, using various established strategies, VEGF-B₁₆₇ therapy failed to stimulate revascularization of ischemic limbs.

Upregulation of VEGF-B₁₆₇, but not of Flt-1, in the Ischemic Myocardium

In an effort to obtain some insight in the cardio-restricted properties of VEGF-B, we analyzed, by ELISA, the levels of VEGF-B in cardiac and skeletal muscle of WT mice. Compared to skeletal muscle, the heart expressed higher levels of VEGF-B in baseline conditions (pg/mg protein: 272 ± 8 in heart versus 174 ± 6 in skeletal muscle; $n=6$; $P<0.05$). At 4 days after MI, levels of VEGF-B were increased by 1.8 ± 0.3 -fold and 2.0 ± 0.3 -fold in the infarct and infarct border zone, respectively ($n=6$; $P<0.05$). At 7 days, the fold upregulation of VEGF-B was 1.4 ± 0.1 and 1.5 ± 0.1 , in the respective areas ($n=3$; $P=NS$). By contrast, at 4 and 7 days after limb ischemia, VEGF-B protein expression did not increase in the ischemic skeletal muscle (fold increase: 0.9 ± 0.1 at day 4 and 0.8 ± 0.1 at day 7; $n=3$ to 6; $P=NS$). This cardio-restricted upregulation of VEGF-B expression was specific, as the PlGF levels were increased in both ischemic myocardium and limb (not shown). Thus, VEGF-B expression is upregulated in ischemic hearts only, consistent with the cardio-restricted phenotype of VEGF-B^{-/-} mice (for discussion, see below).

To analyze the expression of Flt-1, we performed RT-PCR using specific primers and found more abundant Flt-1 transcripts in the heart than skeletal muscle in baseline conditions (transcript levels of Flt-1 per 1000 copies of GAPDH: 1.29 ± 0.11 in the myocardium versus 0.40 ± 0.04 in skeletal muscle, $n=5$; $P<0.05$). However, Flt-1 mRNA expression did not increase in the heart or skeletal muscle at 4 and 7 days after induction of ischemia (not shown). To identify the cell types expressing Flt-1 in vivo, we subjected LacZ-tagged Flt-1 reporter mice to MI or limb ischemia. Immunostaining for β -Gal revealed a similar labeling pattern for Flt-1 in both heart and skeletal muscle in ischemic as well as in nonischemic conditions. Indeed, double immuno-labeling studies showed expression of Flt-1 in SMA⁺ SMCs and most CD31⁺ ECs of nonischemic (Figure 4A, 4C, 4E, and 4G) and ischemic heart or limb tissue at 4 (Figure 4B, 4D, 4F, and 4H) and 7 days (not shown) after induction of ischemia. In the ischemic heart and skeletal muscle, Flt-1 expression also colocalized with some of the CD45⁺ infiltrating leukocytes (not shown). Thus, the cardio-restricted activity of VEGF-B is unlikely explained by differences in Flt-1 expression.

Discussion

We studied loss- and gain-of-function of VEGF-B in pathological angiogenesis by using various genetic and experimental approaches. The principal finding is that VEGF-B has a relatively restricted role in pathological angiogenesis and, even more remarkably, predominantly in the ischemic myocardium.

The precise role and therapeutic potential of VEGF-B in the revascularization of the ischemic myocardium *in vivo* has not been studied thus far. By stimulating the ingrowth of new vessels in the ischemic infarct borders, VEGF-B resembles other angiogenic factors such as VEGF and PlGF (and others), which have, quantitatively and qualitatively, similar effects in rodent models of MI (reviewed in⁴). VEGF-B differs, however, from these agents in its selectivity to stimulate angiogenesis primarily in ischemic myocardium and not in other tissues. These loss- and gain-of-function findings in the mouse, together with findings that VEGF-B promotes compensatory hypertrophy of the remote myocardium after myocardial infarction,¹⁷ warrant further consideration of the therapeutic potential of VEGF-B for promoting functional recovery of MI.

In contrast to its role in the ischemic heart, various types of under- and overexpression studies indicate that VEGF-B is dispensable for vessel growth in the skin, lung, retina, and particularly the ischemic limb. Indeed, although others reported a minor role of VEGF-B in skin, retina, or lung,^{11–14} the negligible role of VEGF-B in ischemic limbs was unexpected, given that VEGF-B stimulates revascularization of the ischemic heart, another type of muscle. To confirm that the lack of an effect of VEGF-B was not attributable to a particular experimental condition, we used different complementary, nonoverlapping strategies. First, loss of VEGF-B did not impair revascularization of ischemic limbs, neither did it aggravate the revascularization defect in PlGF^{-/-} mice, indicating that VEGF-B was ineffective alone and as a modifier of its homologue PlGF. Gain-of-function of VEGF-B, achieved via various strategies, also failed to stimulate the revascularization of ischemic limbs. This failure was not attributable to insufficient expression, as injection of the same dose of a VEGF-B expressing adenovirus or implantation of the same engineered VEGF-B myoblasts sufficed to stimulate vessel growth in the ischemic heart. Also, each of these strategies have previously been used in ischemic limbs to demonstrate the angiogenic activity of other well-known angiogenic factors, such as VEGF and PlGF (^{21,23}; unpublished Tjwa M and Carmeliet P, 2008). In contrast to our findings, others reported that VEGF-B₁₆₇ promotes ischemic limb revascularization.^{15,16} The precise explanation for this discrepancy remains unknown, and might be attributable to experimental differences in gene transfer method, ischemic limb model or methods of analysis.

A remarkable observation is that VEGF-B has angiogenic properties, which substantially differ from those of its homologues VEGF and PlGF. Of all angiogenic members of the VEGF family, VEGF is the most widely active, affecting angiogenesis in health and disease.^{1,4} PlGF, by contrast, is redundant for embryonic vascularization and vessel maintenance in healthy conditions, but involved in the angiogenic switch in disease conditions.^{21,22} VEGF-B is not only dispensable in development and health, but also in most conditions of pathological angiogenesis, except for the ischemic heart. The selectivity and specificity of each of these VEGF family members is remarkable, when comparing it, for instance, to the largely redundant and overlapping angiogenic activity of the 24 FGF family members.²⁴ Though beyond the scope of the present study, it remains outstanding why VEGF-B has such a restricted angiogenic activity in the ischemic heart. The lack of its angiogenic activity in noncardiac tissues cannot be simply attributed to absent expression of VEGF-B or its receptor Flt-1, because both are constitutively expressed in these tissues, including in skeletal muscle.^{7,21,22} Interestingly, however, VEGF-B levels were up-regulated by ischemia in the ischemic heart but not in the ischemic muscle, which might possibly explain, at least in part, why angiogenesis was impaired

in VEGF-B^{-/-} mice in the ischemic myocardium, but not in the ischemic limb. Another possible mechanism might be that the endothelial cells in the heart have distinct tissue-specific characteristics as compared to endothelial cells in other tissues, similar to the distinct differentiation properties of endothelial cells in the nervous system, endocrine organs, lymphoid tissue, etc.²⁵ It is indeed known that coronary endothelial cells are derived from unique progenitors, ie, the epicardium-derived progenitor cells, and require distinct angiogenic signals (such as thymosin- β 4).²⁶ Perhaps, also, the intracellular signaling pathway, induced by VEGF-B, is distinct in these coronary endothelial cells. Because VEGF and PlGF transmit specific angiogenic signals through Flt-1,⁵ VEGF-B could also have a different angiogenic activity profile. Further hypothetical explanations for the selective angiogenic activity of VEGF-B in the heart might include differences in the composition of extracellular matrix proteins (which modulate the activity of VEGF-B²⁷), or in a role for Neuropilin-1, another VEGF-B receptor (reviewed in⁶).

Regardless of the mechanisms, few other tissue-selective angiogenic molecules have been identified thus far, including EG-VEGF, BDNF, and others.¹ Such tissue-specific angiogenic signals are medically relevant, as they offer attractive therapeutic opportunities. Indeed, the potent activity of VEGF and its associated risk of adverse effects (bleeding, leakage, hypotension, malignancy, etc), for instance, has precluded systemic administration of this angiogenic agent for the revascularization of ischemic tissues.^{1,4} In contrast, a molecule such as VEGF-B, which would more selectively stimulate vessel growth in the heart, and primarily in ischemic conditions, would be expected to stimulate vessels in the specific target organ without causing adverse effects. Our observations that VEGF-B did not cause any noticeable side effects of general toxicity, hemorrhage, edema, or hypotension (not shown) are consistent with such a model. Overall, these mouse genetic findings support a—therapeutically attractive—model whereby VEGF-B selectively promotes angiogenesis in the ischemic myocardium.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Sources of Funding: This work was supported by grants to PC from Bristol-Myers-Squibb, FWO (#G0125.00 & #G. 0121.02), EU (QLRT-2001-01955), Leducq foundation, Concerted Research Activities (#GOA2001/09), and IAP-P5/02, and IAP-P6/30, and by NIH (HL65572) to H.M.B. and M.L.S., and by Swedish Research Council, Novo Nordisk Foundation, Wallenbergs Foundation, and Karolinska Institute. to U.E. G.v.D. (postdoc DE 740/1-1); M.T. (postdoc IWT and FWO).

References

1. Ferrara N, Kerbel RS. Angiogenesis as a therapeutic target. *Nature* 2005;438:967–974. [PubMed: 16355214]
2. Grimmond S, Lagercrantz J, Drinkwater C, Silins G, Townson S, Pollock P, Gotley D, Carson E, Rakar S, Nordenskjold M, Ward L, Hayward NK, Weber G. Cloning and characterization of a novel human gene related to vascular endothelial growth factor. *Genome Research* 1996;6:124–131. [PubMed: 8919691]
3. Olofsson B, Pajusola K, Kaipainen A, von Euler G, Joukov V, Saksela O, Orpana A, Pettersson R, Alitalo K, Eriksson U. Vascular endothelial growth factor B, a novel growth factor for endothelial cells. *Proc Natl Acad Sci USA* 1996;93:2576–2581. [PubMed: 8637916]
4. Carmeliet P. Angiogenesis in life, disease and medicine. *Nature* 2005;438:932–936. [PubMed: 16355210]
5. Tjwa M, Luttun A, Autiero M, Carmeliet P. VEGF and PlGF: two pleiotropic growth factors with distinct roles in development and homeostasis. *Cell Tissue Res* 2003;314:5–14. [PubMed: 13680354]

6. Nash AD, Baca M, Wright C, Scotney PD. The biology of vascular endothelial growth factor-B (VEGF-B). *Pulm Pharmacol Ther* 2006;19:61–69. [PubMed: 16286239]
7. Li X, Aase K, Li H, von Euler G, Eriksson U. Isoform-specific expression of VEGF-B in normal tissues and tumors. *Growth Factors* 2001;19:49–59. [PubMed: 11678209]
8. Olofsson B, Korpelainen E, Pepper MS, Mandriota SJ, Aase K, Kumar V, Gunji Y, Jeltsch MM, Shibuya M, Alitalo K, Eriksson U. Vascular endothelial growth factor B (VEGF-B) binds to VEGF receptor-1 and regulates plasminogen activator activity in endothelial cells. *Proc Natl Acad Sci USA* 1998;95:11709–11714. [PubMed: 9751730]
9. Aase K, von Euler G, Li X, Ponten A, Thoren P, Cao R, Cao Y, Olofsson B, Gebre-Medhin S, Pekny M, Alitalo K, Betsholtz C, Eriksson U. Vascular endothelial growth factor-B-deficient mice display an atrial conduction defect. *Circulation* 2001;104:358–364. [PubMed: 11457758]
10. Bellomo D, Headrick JP, Silins GU, Paterson CA, Thomas PS, Gartside M, Mould A, Cahill MM, Tonks ID, Grimmond SM, Townson S, Wells C, Little M, Cummings MC, Hayward NK, Kay GF. Mice lacking the vascular endothelial growth factor-B gene (*Vegfb*) have smaller hearts, dysfunctional coronary vasculature, and impaired recovery from cardiac ischemia. *Circ Res* 2000;86:E29–E35. [PubMed: 10666423]
11. Reichelt M, Shi S, Hayes M, Kay G, Batch J, Gole GA, Browning J. Vascular endothelial growth factor-B and retinal vascular development in the mouse. *Clin Experiment Ophthalmol* 2003;31:61–65. [PubMed: 12580897]
12. Wanstall JC, Gambino A, Jeffery TK, Cahill MM, Bellomo D, Hayward NK, Kay GF. Vascular endothelial growth factor-B-deficient mice show impaired development of hypoxic pulmonary hypertension. *Cardiovasc Res* 2002;55:361–368. [PubMed: 12123775]
13. Louzier V, Raffestin B, Leroux A, Branellec D, Caillaud JM, Levame M, Eddahibi S, Adnot S. Role of VEGF-B in the lung during development of chronic hypoxic pulmonary hypertension. *Am J Physiol Lung Cell Mol Physiol* 2003;284:L926–L937. [PubMed: 12547729]
14. Mould AW, Greco SA, Cahill MM, Tonks ID, Bellomo D, Patterson C, Zournazi A, Nash A, Scotney P, Hayward NK, Kay GF. Transgenic overexpression of vascular endothelial growth factor-B isoforms by endothelial cells potentiates postnatal vessel growth in vivo and in vitro. *Circ Res* 2005;97:e60–e70. [PubMed: 16109918]
15. Wright CE. Effects of vascular endothelial growth factor (VEGF)A and VEGFB gene transfer on vascular reserve in a conscious rabbit hindlimb ischaemia model. *Clin Exp Pharmacol Physiol* 2002;29:1035–1039. [PubMed: 12366398]
16. Silvestre JS, Tamarat R, Ebrahimian TG, Le-Roux A, Clergue M, Emmanuel F, Duriez M, Schwartz B, Branellec D, Levy BI. Vascular endothelial growth factor-B promotes in vivo angiogenesis. *Circ Res* 2003;93:114–123. [PubMed: 12805240]
17. Tirziu D, Chorianopoulos E, Moodie KL, Palac RT, Zhuang ZW, Tjwa M, Roncal C, Eriksson U, Fu Q, Elfenbein A, Hall AE, Carmeliet P, Moons L, Simons M. Myocardial hypertrophy in the absence of external stimuli is induced by angiogenesis in mice. *J Clin Invest* 2007;117:3188–3197. [PubMed: 17975666]
18. Bhardwaj S, Roy H, Gruchala M, Viita H, Kholova I, Kokina I, Achen MG, Stacker SA, Hedman M, Alitalo K, Yla-Herttuala S. Angiogenic responses of vascular endothelial growth factors in periaortic tissue. *Hum Gene Ther* 2003;14:1451–1462. [PubMed: 14577925]
19. Rissanen TT, Markkanen JE, Gruchala M, Heikura T, Puranen A, Kettunen MI, Kholova I, Kauppinen RA, Achen MG, Stacker SA, Alitalo K, Yla-Herttuala S. VEGF-D is the strongest angiogenic and lymphangiogenic effector among VEGFs delivered into skeletal muscle via adenoviruses. *Circ Res* 2003;92:1098–1106. [PubMed: 12714562]
20. Fong GH, Klingensmith J, Wood CR, Rossant J, Breitman ML. Regulation of *flt-1* expression during mouse embryogenesis suggests a role in the establishment of vascular endothelium. *Dev Dyn* 1996;207:1–10. [PubMed: 8875071]
21. Luttun A, Tjwa M, Moons L, Wu Y, Angelillo-Scherrer A, Liao F, Nagy JA, Hooper A, Priller J, De Klerck B, Compennolle V, Daci E, Bohlen P, Dewerchin M, Herbert JM, Fava R, Matthys P, Carmeliet G, Collen D, Dvorak HF, Hicklin DJ, Carmeliet P. Revascularization of ischemic tissues by PlGF treatment, and inhibition of tumor angiogenesis, arthritis and atherosclerosis by anti-Flt1. *Nat Med* 2002;8:831–840. [PubMed: 12091877]

22. Carmeliet P, Moons L, Luttun A, Vincenti V, Compernelle V, De Mol M, Wu Y, Bono F, Devy L, Beck H, Scholz D, Acker T, DiPalma T, Dewerchin M, Noel A, Stalmans I, Barra A, Blacher S, Vandendriessche T, Ponten A, Eriksson U, Plate KH, Foidart JM, Schaper W, Charnock-Jones DS, Hicklin DJ, Herbert JM, Collen D, Persico MG. Synergism between vascular endothelial growth factor and placental growth factor contributes to angiogenesis and plasma extravasation in pathological conditions. *Nat Med* 2001;7:575–583. [PubMed: 11329059]
23. von Degenfeld G, Banfi A, Springer ML, Wagner RA, Jacobi J, Ozawa CR, Merchant MJ, Cooke JP, Blau HM. Microenvironmental VEGF distribution is critical for stable and functional vessel growth in ischemia. *Faseb J* 2006;20:2657–2659. [PubMed: 17095533]
24. Rusnati M, Presta M. Fibroblast growth factors/fibroblast growth factor receptors as targets for the development of anti-angiogenesis strategies. *Curr Pharm Des* 2007;13:2025–2044. [PubMed: 17627537]
25. Aird WC. Phenotypic heterogeneity of the endothelium: I. Structure, function, and mechanisms. *Circ Res* 2007;100:158–173. [PubMed: 17272818]
26. Smart N, Rossdeutsch A, Riley PR. Thymosin beta4 and angiogenesis: modes of action and therapeutic potential. *Angiogenesis* 2007;10:229–241. [PubMed: 17632766]
27. Ikuta T, Ariga H, Matsumoto K. Extracellular matrix tenascin-X in combination with vascular endothelial growth factor B enhances endothelial cell proliferation. *Genes Cells* 2000;5:913–927. [PubMed: 11122379]




Figure 1.

A and B, Morphometric analysis revealed a reduced number of TM⁺ and SMA⁺ vessels in the infarcted area of VEGF-B^{-/-} mice, as compared to WT animals (**P*<0.05) or to VEGF-B^{-/-} mice, treated with rhVEGF-B₁₆₇ (#*P*<0.05 vs no treatment). C–F, Immunostaining showed increased TM⁺ and SMA⁺ vessels after delivery of rhVEGF-B₁₆₇ to WT mice. G and H, Systemic injection of Ad.hVEGF-B₁₆₇ stimulates the growth of TM⁺ and SMA⁺ vessels in the infarct area and border zone of WT mice (**P*<0.05). Scale bars: 50 μm.




Figure 2.

A–C, Skin wound healing is comparable in WT and VEGF-B^{-/-} mice as analyzed by measurement of the wound width (A), and illustrated by macroscopic inspection (B and C).




Figure 3.

A and B, Immunolabeling for CD31 showed comparable vessel densities in regenerating gastrocnemius muscles of Ad.RR5 or Ad.hVEGF-B₁₆₇ injected mice. C and D, Double staining for β-Gal (green cells, marked by an asterisk) and CD31 revealed an identical vascularization pattern in gastrocnemius muscles, injected with control or VEGF-B₁₆₇ overexpressing myoblasts. E, Capillary-to-myocyte ratio is comparable in normal or ischemic limbs, injected with control or VEGF-B₁₆₇ over-expressing myoblasts. Scale bars: 50 μm.




Figure 4.

A-H, Double immunostaining of β -Gal and CD31 (A, B, E, F) or SMA (C, D, G, H) in normal and ischemic heart (A–D) or skeletal muscle (E–H) sections of LacZ-tagged Flt-1 mice revealed labeling of Flt-1 expressing cells in most CD31⁺ cells (arrowheads) and in all SMA⁺ cells (arrows). Scale bars: 50 μ m.

Table
Negligible Role of VEGF-B in Revascularization After Limb Ischemia

| | WT Mice | | | | | |
|--|-----------|----------------------------|---------------|-------------------------|-----------|---------------------------|
| | WT Mice | VEGF-B ^{-/-} Mice | Empty Plasmid | pmVEGF-B ₁₆₇ | Ad.RRS | Ad.hVEGF-B ₁₆₇ |
| Angiogenesis | | | | | | |
| Capillary-to-myocyte ratio | 0.96±0.06 | 0.97±0.06 | ND | ND | 1.16±0.06 | 1.19±0.10 |
| Total limb perfusion | | | | | | |
| Laser Doppler (% of non-ligated) | 90±5 | 90±11 | 86±9 | 82±6 | 92±5 | 94±4 |
| Treadmill exercise test | | | | | | |
| Endurance (% of baseline) | 56±6 | 50±7 | ND | ND | ND | ND |
| Grade exercise (% of baseline) | 57±4 | 52±6 | ND | ND | ND | ND |
| Collateral growth | | | | | | |
| <i>Lumen area, μm²</i> | | | | | | |
| Main collateral artery | 2810±300 | 2480±150 | 2890±190 | 2630±280 | 2294±425 | 2415±511 |
| 2nd collateral branch | 740±75 | 790±50 | 760±60 | 790±65 | 704±190 | 688±163 |
| 3rd collateral branch | 120±10 | 140±7 | 96±3 | 98±6 | 105±13 | 92±13 |
| <i>Collateral side branches (n/mm²)</i> | | | | | | |
| 2nd collateral branches | 3±0.8 | 3±0.7 | 2±0.3 | 3±0.4 | 4±1.8 | 3±1.4 |
| 3rd collateral branches | 5±1.0 | 6±1.5 | 8±1.2 | 9±1.6 | 7±1.6 | 10±3.6 |
| <i>Total perfusion area, μm²/mm²</i> | 2730±700 | 3000±800 | 3140±360 | 3460±670 | 3098±569 | 3020±524 |

Analysis was performed as described in methods. ND indicates not determined; P=NS for WT/control mice vs VEGF-B^{-/-}/VEGF-B-treated animals.