

Published in final edited form as:

Nat Genet. 2008 June ; 40(6): 776–781. doi:10.1038/ng.149.

X-linked protocadherin 19 mutations cause female-limited epilepsy and cognitive impairment

Leanne M Dibbens^{1,2,14}, Patrick S Tarpey^{3,14}, Kim Hynes^{1,4}, Marta A Bayly¹, Ingrid E Scheffer^{5,6}, Raffaella Smith³, Jamee Bomar⁷, Edwina Sutton⁴, Lucianne Vandeleur¹, Cheryl Shoubbridge¹, Sarah Edkins³, Samantha J Turner⁵, Claire Stevens³, Sarah O'Meara³, Calli Tofts³, Syd Barthorpe³, Gemma Buck³, Jennifer Cole³, Kelly Halliday³, David Jones³, Rebecca Lee³, Mark Madison³, Tatiana Mironenko³, Jennifer Varian³, Sofie West³, Sara Widaa³, Paul Wray³, John Teague³, Ed Dicks³, Adam Butler³, Andrew Menzies³, Andrew Jenkinson³, Rebecca Shepherd³, James F Gusella⁸, Zaid Afawi⁹, Aziz Mazarib⁹, Miriam Y Neufeld⁹, Sara Kivity¹⁰, Dorit Lev¹¹, Tally Lerman-Sagie¹¹, Amos D Korczyn⁸, Christopher P Derry⁵, Grant R Sutherland^{1,2,4}, Kathryn Friend¹, Marie Shaw¹, Mark Corbett¹, Hyung-Goo Kim⁸, Daniel H Geschwind⁷, Paul Thomas⁴, Eric Haan^{1,4}, Stephen Ryan¹², Shane McKee¹³, Samuel F Berkovic⁵, P Andrew Futreal³, Michael R Stratton³, John C Mulley^{1,2,4}, and Jozef Gécz^{1,2,4}

¹Department of Genetic Medicine, Level 9 Rieger Building, Women's and Children's Hospital, 72 King William Road, North Adelaide, South Australia 5006, Australia

²School of Paediatrics and Reproductive Health, University of Adelaide, Adelaide, South Australia 5005, Australia

³Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton, Cambridge CB10 1SA, UK

⁴School of Molecular & Biomedical Science, University of Adelaide, Adelaide, South Australia 5005, Australia

© 2008 Nature Publishing Group

Correspondence should be addressed to L.M.D. (leanne.dibbens@cywhs.sa.gov.au) or J.G. (jozef.gecz@adelaide.edu.au).

¹⁴These authors contributed equally to this work.

AUTHOR CONTRIBUTIONS

L.M.D. and P.S.T. contributed equally to this work. L.M.D. coordinated the project in concept and design, supervised molecular studies, managed collaborations, wrote the first draft of the manuscript and significantly edited successive manuscript drafts; P.S.T. supervised the X-chromosome gene sequencing and analysis; K. Hynes and M.A.B. carried out molecular studies. I.E.S., S.F.B., S.J.T., E.H., S.M., S.R., A. Mazarib, Z.A., M.Y.N., S.K., D.L., T.L.-S., A.D.K. and C.P.D. identified families and provided clinical information; R.S., S.E., C. Stevens, S.O., C.T., S.B., G.B., J.C., K. Halliday, D.J., T.M., J.V., S. West, S. Widaa, J.T., E.D., A.B., R.L., M.M., P.W., A. Menzies, A.J. and R.S. performed the X-chromosome gene sequencing and analysis of 737 genes. L.V. performed tissue culture work, J.B. and D.H.G. carried out and interpreted the human *in situ* hybridization analysis, K.F. performed and interpreted linkage analysis, M.S. and K. Hynes did X inactivation studies and their interpretation, and M.C. and C. Shoubbridge contributed to the supervision of molecular and cell studies. H.-G.K. and J.F.G. contributed to segregation analysis. E.S. and P.T. performed and interpreted the mouse *in situ* hybridization analysis. I.E.S. and S.F.B. contributed to the project concept and coordinated families. E.H. and G.R.S. coordinated families. P.A.F. and M.R.S. coordinated the X-chromosome gene sequencing and analysis. J.C.M. and J.G. coordinated and supervised the project in concept and design, supervised molecular studies, managed collaborations and significantly edited successive drafts. All authors contributed to the discussion of the results and the preparation of successive manuscript drafts with the opportunity to comment critically and constructively.

URLs. VEGA, <http://vega.sanger.ac.uk/index.html>.

Accession codes. GenBank: Incomplete human *PCDH19* mRNA, NM_020766; incomplete human PCDH19 protein, NP_065817; complete human *PCDH19* mRNA and protein, EF676096; human *PCDH11X* mRNA, NM_014522, NM_032967, NM_032968 and NM_032969; human PCDH11X protein, NP_055337, NP_116749, NP_116750 and NP_116751; human *PCDH11Y* mRNA, NM_032971, NM_032972 and NM_032973; human PCDH11Y protein, NP_116753, NP_116754 and NP_116755.

⁵Epilepsy Research Centre and Department of Medicine, University of Melbourne, Level 1 Neurosciences Building, Heidelberg Repatriation Hospital, Austin Health, Banksia Street, Heidelberg West, Victoria 3081, Australia

⁶Department of Paediatrics, University of Melbourne, Royal Children's Hospital, Flemington Road, Parkville, Victoria 3052, Australia

⁷Program in Neurogenetics and Neurobehavioral Genetics, Neurology Department and Semel Institute for Neuroscience and Behavior, David Geffen School of Medicine, University of California at Los Angeles, Los Angeles, California 90095-1769, USA

⁸Molecular Neurogenetics Unit, Center for Human Genetic Research, Massachusetts General Hospital, 185 Cambridge Street, Boston, Massachusetts 02114, USA and Department of Genetics, Harvard Medical School, 77 Avenue Louis Pasteur, NRB 0330, Boston, Massachusetts 02115, USA

⁹Department of Neurology, Tel Aviv Sourasky Medical Center, 6 Weizmann Street, Tel Aviv 64239, Israel

¹⁰Department of Neurology, Schneider Children's Medical Center, Petaq Tikvah 49202, Israel

¹¹Metabolic Neurogenetic Clinic, Wolfson Medical Center, 62 HaLohamim Street, Holon 58100, Israel

¹²AstraZeneca, 1800 Concord Pike, Wilmington, Delaware 19803, USA

¹³Northern Ireland Regional Genetics Service, Belfast City Hospital, Lisburn Road, Belfast BT9 7AB, Northern Ireland, UK

Abstract

Epilepsy and mental retardation limited to females (EFMR) is a disorder with an X-linked mode of inheritance and an unusual expression pattern. Disorders arising from mutations on the X chromosome are typically characterized by affected males and unaffected carrier females. In contrast, EFMR spares transmitting males and affects only carrier females. Aided by systematic resequencing of 737 X chromosome genes, we identified different protocadherin 19 (*PCDH19*) gene mutations in seven families with EFMR. Five mutations resulted in the introduction of a premature termination codon. Study of two of these demonstrated nonsense-mediated decay of *PCDH19* mRNA. The two missense mutations were predicted to affect adhesiveness of PCDH19 through impaired calcium binding. *PCDH19* is expressed in developing brains of human and mouse and is the first member of the cadherin superfamily to be directly implicated in epilepsy or mental retardation.

EFMR (MIM 300088) is a condition characterized by seizure onset in infancy or early childhood (6-36 months) and cognitive impairment. The disorder is sex-limited, with the phenotype being restricted to females. Males are apparently spared, with normal cognitive function and absence of seizures. Evidence for the trait being due to a gene on the X chromosome derives from the pattern of inheritance and from genetic linkage to Xq22 (ref. 1). EFMR is recognizable in multi-generational pedigrees by the unique sex-limited expression pattern, wherein affected females are connected through unaffected transmitting males 1-3.

We ascertained five new families carrying EFMR on the basis of their inheritance pattern, electroclinical features and gene localization to Xq22 (ref. 4 and S.M., unpublished data). To identify the underlying genetic defect, we resequenced 737 genes from the Vertebrate Genome Annotation (VEGA) database in probands from three families. We identified a single nonsense nucleotide change 2012C>G (S671X) in the *PCDH19* gene in family 3 (Fig. 1). Families 1 and 2 initially did not show any changes.

Subsequent comparative sequence analysis of the annotated *PCDH19* ORF revealed that it was incomplete, consistent with a previous report⁵. We annotated the complete 3,447-bp ORF of *PCDH19*, which consists of six exons. The full-length, processed *PCDH19* mRNA is 9,765 nt long, with exon 2 alternatively spliced (Supplementary Fig. 1 online). Sequencing of the entire *PCDH19* ORF identified a missense change 1322T>A (V441E) in family 1, a nonsense change 253C>T (Q85X) in family 2, a frameshift change 2030_2031 insT (L677fsX717) in family 4 and a frameshift change 357delC (I119fsX122) in family 5. Finally, we identified a frameshift change 1091_1092insC (P364fsX375) in the original EFMR-bearing family (family 6, ref. 1) (Fig. 1). In summary, we identified *PCDH19* nucleotide changes in all six families bearing EFMR that we studied. Another unique nucleotide change (1671C>G, N557K) was identified by screening a cohort of 87 females that included both isolated and familial cases with epilepsy and varying degrees of mental impairment. We found that the nucleotide change in this small family, with two affected girls with epilepsy and autism spectrum disorder, arose *de novo* in their carrier father on the grandmaternal X chromosome (Fig. 1, family 7).

All seven nucleotide changes (Fig. 2a) segregated with the clinical phenotype in the respective families (Fig. 1) and were not present in 250 male probands from families with putative X-linked mental retardation or in 750 control (350 male and 400 female) X chromosomes. Further, we did not detect any other potentially deleterious nucleotide changes in probands from the three families in any of the other 736 X-chromosome genes sequenced (data not shown). Taken together, the predicted loss of function of all seven *PCDH19* changes (see below), the high degree of conservation of the residues affected by the two missense changes (Fig. 2a,b) and our *PCDH19* mRNA studies (see below) strongly support *PCDH19* as the gene whose mutation causes EFMR. Moreover, *PCDH19* is located at Xq22 (Ensembl database) within the original linkage region1. EFMR locus homogeneity in these families has now been molecularly defined, and the combined lod score with markers in the Xq22 region reached 11.9 (refs. 1,4 and S.M., unpublished data). Among the seven families bearing EFMR studied, there were two (of 68) carrier females who had been classified (at the time of the original assessment) as unaffected. Therefore, the estimated penetrance of the known EFMR mutations in females was >90%.

PCDH19 was expressed predominantly in neural tissues and at different developmental stages. By RNA blot analysis using a *PCDH19*-specific probe, we detected a transcript size of approximately 9.8 kb from various areas of the male and female adult human brain (Fig. 2c and below). We also identified *PCDH19* mRNA expression in primary skin fibroblasts. This allowed us to examine the consequences of the *PCDH19* mutations 253C>T, Q85X (family 2) and 2012C>G, S671X (family 3) on the stability of their respective mRNAs. Both mutations introduce a premature termination codon (PTC) into the *PCDH19* mRNA. Such PTC-containing mRNAs are usually recognized by the nonsense-mediated decay (NMD) surveillance complexes and efficiently degraded⁶. We found that the PTC mutations in family 2 (Fig. 2d) and family 3 (data not shown) led to *PCDH19* mRNA degradation by NMD. Inhibition of translation by cycloheximide treatment of skin fibroblast cells preserved the PTC mutation-containing mRNA (Fig. 2d). The absence of mutant *PCDH19* mRNAs could not be attributed to skewing of X-inactivation. We confirmed random X-inactivation in DNA isolated from blood and skin fibroblast cultures of each affected female available (data not shown), in agreement with previous data¹.

PCDH19 encodes a 1,148-amino acid protein belonging to the protocadherin $\delta 2$ subclass of the cadherin superfamily of cell-cell adhesion molecules. *PCDH19* contains a signal sequence, six extracellular cadherin (EC) repeats, a transmembrane domain and a cytoplasmic region with conserved CM1 and CM2 domains (Fig. 2a). The biological role of the *PCDH19* protein is not known. However, members of the protocadherin family are predominantly expressed in the nervous system^{7,8} and are postulated to be involved in the establishment of neuronal connections and in signal transduction at the synaptic membrane^{9,10}. *PCDH10*, also a $\delta 2$

protocadherin, is required for growth of striatal axons and thalamocortical projections¹¹. Usher syndrome type 1, involving combined hearing loss and blindness, was the first human disorder to be associated with mutations in a *PCDH* gene (*PCDH15*)¹². Mutations in *PCDH19* have now been associated with epilepsy and mental retardation.

All mutations identified in this study were located in the large *PCDH19* extracellular domain (Fig. 2a) containing the cadherin repeats, which facilitate cell-cell interactions. We predict five of these mutations to be complete loss-of-function mutations as a consequence of NMD degradation of their respective PTC-containing mRNAs (Fig. 2d and data not shown). Given the similarity among the clinical phenotypes associated with all seven *PCDH19* mutations, it is reasonable to suggest that the remaining two mutations, missense mutations leading to V441E and N557K substitutions, also lead to loss of *PCDH19* function. The N557K mutation affects an invariant asparagine residue within the EC5 domain (Fig. 2a,b). The equivalent asparagine residue of EC1 of classical cadherins (for example, Asn100 of N-cadherin¹³) and protocadherins (for example, Asn101 of *Pcdh α 14*) is essential for calcium binding and thus for the adhesive function of the EC1 domain^{13,14}. The valine residue at position 441 (in EC4, or the equivalent of Val96 in EC1 of N-cadherin¹³ or Val97 in EC1 of *Pcdh α 14*) is also highly conserved (Fig. 2b) and in close proximity to the calcium-binding site^{13,14}. We speculate that both *PCDH19* missense variants adversely affect *PCDH19* adhesive function through impaired calcium binding.

To investigate the expression of *PCDH19* in the developing mouse and human central nervous system (CNS), we performed *in situ* hybridization analysis. In mice, *Pcdh19* had widespread expression in both the embryonic and adult brain, including the developing cortex and hippocampus (Fig. 3 and Supplementary Note online), which is consistent with our finding that mutations of this gene in humans are associated with cognitive impairment. In human tissue, in addition to *PCDH19*, we also investigated the expression pattern of *PCDH11X* and *PCDH11Y*, two paralogous protocadherin genes, whose expression and function we speculate to be relevant to EFMR (see Fig. 4 and below). Each of the three *PCDH* genes was expressed in developing cortical plate, amygdala and subcortical regions and in the ganglionic eminence (Fig. 4). However, *PCDH11X/Y* expression showed a marked sexual dimorphism in the caudate nucleus: virtually absent expression in the male and high expression in the female (Fig. 4a-c). Additionally, closer inspection of the amygdala showed that *PCDH19* was expressed in lateral nuclei, whereas *PCDH11X/Y* was more medial.

Typically, in most X-linked dominant conditions, males are more severely affected and often die prenatally. In EFMR, the transmitting males are apparently normal (that is, without epilepsy or intellectual disability), but with some obsessive traits reported anecdotally⁴. Previous hypotheses to explain the EFMR expression pattern¹ considered either a dominant negative effect of the mutant protein in carrier females (for example, similar to that of the C-terminally truncated P-cadherin in malignant melanoma¹⁵) or the presence of a compensatory or rescue factor in males. The dominant negative effect of mutant *PCDH19* is unlikely, based on our NMD results (see above), and *PCDH19* does not have a Y-chromosome paralog to provide male rescue. Sexually dimorphic expression and function^{16,17} of *PCDH19* and/or relevant compensatory genes is also a possibility.

We propose an alternative hypothesis, based on the identity of the EFMR gene, that invokes functional rescue by a related, but nonparalogous, protocadherin gene in males. The *PCDH19* gene is subject to X inactivation (Supplementary Figs. 2 and 3 online). Therefore, hemizygous transmitting males will have a homogeneous population of *PCDH19*-negative cells, whereas affected females are likely to be mosaics comprising *PCDH19*-negative and *PCDH19*-wild type cells. Such tissue mosaicism may scramble cell-cell communication, which manifests clinically as EFMR. A similar mechanism was proposed for the craniofrontonasal

syndrome CFNS (MIM 304110)¹⁸. We hypothesize that the absence of *PCDH19* function in males may potentially be compensated for by the related but nonparalogous protocadherin gene *PCDH11Y*, a Y-chromosome gene that is expressed in human brain^{19,20}. *PCDH11Y* has an X chromosome paralog, *PCDH11X*, that has strong sequence similarity¹⁹. However, the differences in brain expression patterns between these two genes may account for differential ability to compensate for absence of *PCDH19*. The potential role of *PCDH11Y* (and that of its paralog *PCDH11X*) in the EFMR phenotype will need to be addressed in the future by engineering the appropriate animal models.

Cadherins are a large family of genes with crucial functions in human brain²¹. Our data directly implicate the protocadherin gene family in epilepsy and intellectual disability. On the basis of our observations, molecular diagnosis will result in wider recognition of EFMR, especially in smaller families and single cases, with important consequences for counseling.

METHODS

Subjects

The clinical details of families 1-4 have been reported⁴. The proband in family 5 (IV.2) experienced multiple seizures in the first year of life and has severe learning disabilities, no speech, and behavioral disturbances (S.M., unpublished data). The clinical details of family 6 have been published¹⁻³. Family 7 was ascertained on the basis of one sister having infantile seizures and Asperger's syndrome and the other having epilepsy and mild intellectual disability (I.E.S. and S.F.B., unpublished data).

RNA blotting

Human brain MTN blots II and V (Clontech) were hybridized per the manufacturer's instructions with a probe containing nucleotides 2884-3257 of the human *PCDH19* ORF. The 374-bp PCR product (see Supplementary Table 1 online for PCR primer sequences) was labeled with radioactive [³²P]dCTP (Perkin Elmer) using the Megaprime DNA labeling system (GE Healthcare).

RT-PCR analyses

Total RNA was extracted from fibroblast cells with the RNeasy Mini kit (Qiagen) and treated with DNase I (Qiagen). About 2 µg of RNA was primed with 1 µg of random hexanucleotides and subjected to reverse transcription for 90 min at 42 °C using Superscript II (Invitrogen). The efficiency of the reaction was tested by PCR using primers specific to the ubiquitously expressed *ESD* gene. cDNAs were amplified with Taq DNA polymerase (Roche) and specific single-stranded DNA primers for 35 cycles (denaturation, 94 °C for 30 s; annealing, at specific melting temperature for each pair of primers for 30 s; extension, 72 °C for 30 s). PCR products were separated on an agarose gel stained with 1% ethidium bromide.

Tissue culture of primary skin fibroblast lines

A 3-mm skin biopsy excised from the upper arm was cut finely and transferred to a tissue culture flask. The biopsy was cultured in RPMI-1640, 20% FCS (supplemented with 4 mM L-glutamine, 0.017 mg/ml benzylpenicillin) and grown at 37 °C, with 5% CO₂. Once established, fibroblasts were cultured in RPMI-1640, 10% FCS (supplements as above).

Cycloheximide treatment of skin fibroblast cell lines

Primary fibroblast cells were seeded at 1×10^4 per cm² in RPMI-1640, 10% FCS, and incubated with 50 µg/ml cycloheximide (Sigma) or medium alone for 6 h. Fibroblasts were harvested

using a sterile cell scraper (TPP), then washed once in PBS before total RNA extraction and processing to cDNA as described above.

Human *in situ* hybridization analysis

Human tissue was obtained from the Brain and Tissue Bank for Developmental Disorders at the University of Maryland and was treated as previously described²². Five brains were used, ranging in gestational age from 16 to 20 weeks, with known post mortem intervals from 1 to 2 h. For probe preparation, we performed cDNA synthesis using Superscript III (Invitrogen) from RNA of the inferior frontal gyrus of a 20-gestational-week fetus (RNeasy Mini kit, Qiagen) and amplified two nonoverlapping cDNA transcripts for both *PCDH11X/Y* and *PCDH19* using Platinum Taq Polymerase (Qiagen). Two probes for *PCDH11X/Y* were designed to recognize all *PCDH11X* and *PCDH11Y* isoforms. The PCR primer sequences used for generating the two probes and the *PCDH19* probe are provided in Supplementary Table 1. *In situ* hybridization and image analysis were performed as previously described²². Nonoverlapping probe sets for both genes confirmed the same expression pattern. Sections hybridized with sense probes showed absence of specific staining.

Mouse *in situ* hybridization analysis

Embryonic day 15.5 embryonic heads and dissected P2 brains were fixed in 4% paraformaldehyde at 4 °C, cryoprotected in 30% sucrose and frozen in OCT embedding medium. *In situ* hybridization of 16- μ m sections was performed as described previously²³. The *Pcdh19* probe was a digoxigenin-labeled antisense RNA probe prepared as described²⁴. A total of three neonates and two embryos were analyzed and representative sections are shown. No signal was detected using a sense control probe. Images were taken on a Zeiss Axiophot microscope, compiled and minimally processed (adjusted for color and light/dark) using Adobe Photoshop CS.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

We thank the members of the families studied for their participation and members of the International Genetics of Learning Disability (IGOLD) study for their collaboration. This work was supported by grants from the Australian National Health and Medical Research Council Program Grant 400121 (I.E.S., S.F.B., J.C.M. and J.G.), Thyne-Reid Charitable Trusts (L.M.D.) and the Wellcome Trust. We also acknowledge support to J.F.G. from US National Institutes of Health grant GM061354 and D.H.G. from US National Institute of Mental Health U.S. grant R01 MH 64547. We are grateful for access to the tissues used in these studies from the Developmental Brain and Tissue Bank at University of Maryland funded by the US National Institutes of Health (National Institute of Child Health and Human Development contracts NO1-HD-4-3368 and NO1-HD-4-3383).

References

1. Ryan SG, et al. Epilepsy and mental retardation limited to females: an X-linked dominant disorder with male sparing. *Nat. Genet* 1997;17:92–95. [PubMed: 9288105]
2. Juberg RC, Hellman CD. A new familial form of convulsive disorder and mental retardation limited to females. *J. Pediatr* 1971;79:726–732. [PubMed: 5116697]
3. Fabisiak K, Erickson RP. A familial form of convulsive disorder with or without mental retardation limited to females: extension of a pedigree limits possible genetic mechanisms. *Clin. Genet* 1990;38:353–358. [PubMed: 2126489]
4. Scheffer IE, et al. Epilepsy and mental retardation limited to females: an under-recognised disorder. *Brain* 2008;131:918–927. [PubMed: 18234694]

5. Wolverson T, Lalande M. Identification and characterization of three members of a novel subclass of protocadherins. *Genomics* 2001;76:66–72. [PubMed: 11549318]
6. Maquat LE. Nonsense-mediated mRNA decay: splicing, translation and mRNP dynamics. *Nat. Rev. Mol. Cell Biol* 2004;5:89–99. [PubMed: 15040442]
7. Vanhalst K, Kools P, Staes K, van Roy F, Redies C. Proto-cadherins: a gene family expressed differentially in the mouse brain. *Cell. Mol. Life Sci* 2005;62:1247–1259. [PubMed: 15905963]
8. Kim SY, et al. Spatiotemporal expression pattern of non-clustered protocadherin family members in the developing rat brain. *Neuroscience* 2007;147:996–1021. [PubMed: 17614211]
9. Wu Q, Maniatis T. A striking organization of a large family of human neural cadherin-like cell adhesion genes. *Cell* 1999;97:779–790. [PubMed: 10380929]
10. Yagi T, Takeichi M. Cadherin superfamily genes: functions, genomic organization, and neurologic diversity. *Genes Dev* 2000;14:1169–1180. [PubMed: 10817752]
11. Uemura M, Nakao S, Suzuki ST, Takeichi M, Hirano S. OL-protocadherin is essential for growth of striatal axons and thalamocortical projections. *Nat. Neurosci* 2007;10:1151–1159. [PubMed: 17721516]
12. Ahmed ZM, et al. Mutations of the protocadherin gene *PCDH15* cause Usher syndrome type 1F. *Am.J.Hum. Genet* 2001;69:25–34. [PubMed: 11398101]
13. Patel SD, et al. Type II cadherin ectodomain structures: implications for classical cadherin specificity. *Cell* 2006;124:1255–1268. [PubMed: 16564015]
14. Morishita H, et al. Structure of the cadherin-related neuronal receptor/protocadherin-alpha first extracellular cadherin domain reveals diversity across cadherin families. *J. Biol. Chem* 2006;281:33650–33663. [PubMed: 16916795]
15. Bauer R, Bosserhoff AK. Functional implication of truncated P-cadherin expression in malignant melanoma. *Exp. Mol. Pathol* 2006;81:224–230. [PubMed: 16919267]
16. Arnold AP. Sex chromosomes and brain gender. *Nat. Rev. Neurosci* 2004;5:701–708. [PubMed: 15322528]
17. Yang X, et al. Tissue-specific expression and regulation of sexually dimorphic genes in mice. *Genome Res* 2006;16:995–1004. [PubMed: 16825664]
18. Wieland I, et al. Mutations of the ephrin-B1 gene cause craniofrontonasal syndrome. *Am. J. Hum. Genet* 2004;74:1209–1215. [PubMed: 15124102]
19. Blanco P, Sargent CA, Boucher CA, Mitchell M, Affara NA. Conservation of *PCDHX* in mammals; expression of human X/Y genes predominantly in brain. *Mamm. Genome* 2000;11:906–914. [PubMed: 11003707]
20. Durand CM, et al. Expression and genetic variability of *PCDH11Y*, a gene specific to *Homo sapiens* and candidate for susceptibility to psychiatric disorders. *Am. J. Med. Genet. B. Neuropsychiatr. Genet* 2006;141:67–70. [PubMed: 16331680]
21. Takeichi M. The cadherin superfamily in neuronal connections and interactions. *Nat. Rev. Neurosci* 2007;8:11–20. [PubMed: 17133224]
22. Teramitsu I, Kudo LC, London SE, Geschwind DH, White SA. Parallel Foxp1 and Foxp2 expression in songbird and human brain predicts functional interaction. *J. Neurosci* 2004;24:3152–3163. [PubMed: 15056695]
23. Wilson LD, et al. Developmentally regulated expression of the regulator of G-protein signaling gene 2 (*Rgs2*) in the embryonic mouse pituitary. *Gene Expr. Patterns* 2005;5:305–311. [PubMed: 15661635]
24. Gaitan Y, Bouchard M. Expression of the δ -protocadherin gene *Pcdh19* in the developing mouse embryo. *Gene Expr. Patterns* 2006;6:893–899. [PubMed: 16682261]




Figure 1. Mutations in *PCDH19* cause EFMR. Pedigrees of the seven EFMR families showing the characteristic inheritance pattern of affected females and transmitting males. Each of the X chromosome-encoded *PCDH19* mutations segregated with the EFMR clinical phenotype. An example of a sequence chromatogram of a *PCDH19* mutation as detected in an affected female is shown for each family.




Figure 2. Structure and expression analysis of *PCDH19*. **(a)** Schematic diagram of the *PCDH19* protein showing the signal peptide, extracellular cadherin (EC), transmembrane (TM) and cytoplasmic (CM) domains. The positions of the mutations found in the families bearing EFMR are shown. **(b)** Partial alignment of human *PCDH*s and orthologs of *PCDH19* from other species, showing the high conservation of the residues affected by the two missense mutations, V441E (top) and N557K (bottom). Asn557 is invariant and one of the essential residues for calcium ion binding 13,14. Val441 is highly conserved and in close proximity to the calcium-binding acidic residues (bracket). **(c)** RNA blot analyses of *PCDH19* and *PCDH11X/Y* in various human tissues. Asterisks, the ~9.8 kb *PCDH19* transcripts; arrowheads, the smaller, ~9.5 kb *PCDH11X/Y* mRNAs. Brackets, either nonspecific binding of the *PCDH19* probe or *PCDH19* degradation products, as there is no evidence for such alternative isoforms of *PCDH19* available from our data (data not shown) or from published data. **(d)** Nonsense-mediated RNA decay of mutant *PCDH19* transcripts. Sequence chromatogram from EFMR-affected female from family 2 showing the detection of the mutation 253C>T in genomic DNA (gDNA) (top), the absence of the mutant sequence in fibroblast cDNA (middle) and the presence of the both mutant and wild-type cDNA after the treatment of fibroblasts with cycloheximide (bottom), which inhibits the pioneer round of translation and thus NMD. The position of the mutation is boxed.




Figure 3. Expression of *Pcdh19* in the developing mouse CNS. **(a-l)** Expression at embryonic day 15.5 **(a-f)** and postnatal day 2 **(g-l)** (representative of two males and two females studied). **(a,b)** Adjacent coronal sections through the hippocampal region stained with hematoxylin and eosin or processed for *Pcdh19* *in situ*, respectively. **(c-e)** Higher magnification images of the boxed regions in **b**. Arrowheads in **c**, *Pcdh19*-expressing cells within the cortex; asterisk in **e**, dorsolateral wall of the lateral ventricle. **(f)** Coronal section through the olfactory bulb highlighting *Pcdh19* expression in the nasal epithelium. **(g,h)** Adjacent coronal sections through the mid-hippocampal region stained with hematoxylin and eosin or processed for *Pcdh19* *in situ*, respectively. **(i)** A brain section more posterior than that in **h**, highlighting

Pcdh19 expression. (**j-l**) Higher-magnification images of the regions boxed in **g** and **h**, as indicated. Arrows in **j,k**, *Pcdh19* expression within cortical layers II-IV. Cx, cortex; CxP, cortical plate; Hn, hippocampal neuroepithelium; lv, lateral ventricle; Th, thalamus; Hy, hypothalamus; icf, intercerebral fissure; Ob, olfactory bulbs; Ne, nasal epithelium. Scale bars in **a,b,f-i**, 200 μ M; in **c-e,j-l**, 50 μ M.




Figure 4.

Expression of *PCDH11X/Y* and *PCDH19* in midgestation developing human CNS. **(a,b)** Autoradiography *in situ* hybridization of female (f; panel **a**) and male (m; panel **b**) coronal sections through the basal regions of the brain using probes for *PCDH11X/Y* and *PCDH19*, showing several areas of overlap and differences, including high expression of *PCDH11X/Y* in the ganglionic eminence and sexually dimorphic expression in the caudate nucleus. Dark color denotes high expression. Both genes are highly expressed in amygdala and developing cortical plate. A cartoon key to these sections is shown at the far right and the boxed regions are shown magnified in **c**. **(c)** Micrographs from photographic emulsions of magnified regions of *PCDH11X/Y* expression show low expression in male caudate. White denotes strong expression. GE, ganglionic eminence; Put, putamen; CaNu, caudate nucleus. **(d)** Sagittal photomicrographs highlight areas of differential expression between the two *PCDH* genes, indicating high putamen expression for *PCDH11X/Y* and high hippocampal expression for *PCDH19*. We show sections from two different planes in the male, the first more lateral and the second more medial, to show that the absence of male expression is not an artifact of section plane. **(e)** Sections hybridized with sense probes show absence of specific staining.