

Published in final edited form as:

Oral Dis. 2009 September ; 15(6): 407–413. doi:10.1111/j.1601-0825.2009.01560.x.

Expression of *Six1* homeobox gene during development of the mouse submandibular salivary gland

E. L. McCoy¹, K. Kawakami², H. L. Ford^{1,3,4}, and R. D. Coletta⁵

¹ Program in Molecular Biology, University of Colorado School of Medicine, Denver, Colorado 80045

² Division of Biology, Center for Molecular Medicine, Jichi Medical University, Shimotsuke, Tochigi 329-0498, Japan

³ Department of Obstetrics and Gynecology, University of Colorado School of Medicine, Denver, Colorado 80045

⁴ Department of Biochemistry and Molecular Genetics, University of Colorado School of Medicine, Denver, Colorado 80045

⁵ Department of Oral Diagnosis, School of Dentistry, State University of Campinas, 13414-018 Piracicaba, SP, Brazil.

Abstract

BACKGROUND—Members of the Six family of homeoproteins are expressed in numerous tissues during vertebrate embryogenesis, and are critical regulators of both cell proliferation and survival. Here we report the temporal and spatial expression of *Six1* during maturation of the mouse submandibular salivary gland (SSG) from embryonic day 18.5 (E18.5) to postnatal day 28. Additionally, we examine the role of *Six1* during SSG development using *Six1*-deficient mice.

METHODS—*Six1* expression was assessed by RT-PCR, western blot, and immunofluorescence. Proliferation was measured by bromodeoxyuride (BrdU) incorporation index, and apoptosis was evaluated by TUNEL assay.

RESULTS—*Six1* mRNA and protein levels are high in the epithelial SSG cells at E18.5 and decrease progressively in the postnatal maturing SSG. Although SSGs from *Six1*^{-/-} embryos are significantly smaller than wild type SSGs, the histological structures of the SSG acini and ducts are similar. *Six1*^{-/-} salivary epithelial cells exhibit an intrinsic defect in cell proliferation accompanied by a significant reduction in the *Six1* target gene cyclin A1, previously shown to be a critical mediator of *Six1*-induced proliferation.

CONCLUSION—Our results suggest that the reduction in size of *Six1*^{-/-} SSGs is result of a decrease in cell proliferation during development/maturation.

Keywords

submandibular salivary gland; *Six1*; development; maturation; proliferation; cyclin A1

Correspondence to: Dr. Ricardo D. Coletta, Department of Oral Diagnosis, School of Dentistry, State University of Campinas, Caixa Postal 52, 13414-018 Piracicaba, São Paulo, Brazil. E-mail: coletta@fop.unicamp.br. FAX: +55-19-21065218.

Author contributions Dr. McCoy and Dr. Coletta were responsible for the experiments and manuscript preparation. Dr. Ford and Kawakami were responsible for manuscript revisions.

Conflict of interest There is no conflict of interest.

Introduction

Homeobox genes encode a large and diverse group of DNA binding proteins that act as “master regulators” of development, performing critical functions in specifying cell proliferation, differentiation, survival, and migration (Pearson et al., 2005). The Six family of homeoproteins (Six1-6) are characterized structurally by a divergent homeodomain that is involved in DNA binding, and an N-terminally localized Six domain, that confers cooperative interactions with co-factors (Kawakami et al., 2000, Christensen et al., 2008). Animal studies demonstrate that the *Six* family members play critical roles in organogenesis via regulating cell growth and survival, as well as tissue specification (Christensen et al., 2008). Importantly, mutations in *Six* family members are found in numerous human genetic disorders, underscoring their importance in human embryonic development (Christensen et al., 2008).

Recently, *Six1*^{-/-} mice were generated, which exhibit perinatal death due to severe muscle hypoplasia that results in inadequate diaphragm muscle development and suffocation at birth. In addition, these animals exhibit hypoplastic or completely lacking kidneys and thymus, craniofacial structure malformations, and defects in neurogenesis (Laclef et al., 2003, Xu et al., 2003, Zheng et al., 2003, Ozaki et al., 2004, Li et al., 2003, Zou et al., 2004, Ikeda et al., 2007). Interestingly, cells in the *Six1*^{-/-} affected organs exhibit an increase in apoptosis and a decrease in proliferation (Li et al., 2003, Xu et al., 2003, Ozaki et al., 2004), suggesting that *Six1* is important for the expansion of tissue-specific progenitor cell populations in early development. This expansion is believed to occur because Six1 homeoprotein directly activates numerous cell cycle regulators, including *gdnf*, *c-myc*, and *cyclin D1* during development (Li et al., 2003, Yu et al., 2006). Recently, our group identified the tissue-restricted *cyclin A1* as a transcriptional target of Six1, uncovering yet another mechanism by which Six1 promotes cell cycle progression (Ford et al., 1998, Coletta et al., 2004a).

Salivary gland development is a dynamic process in which cellular proliferation and survival are carefully controlled during branching morphogenesis, directed, in part, by transcriptional regulation (Melnick & Jaskoll, 2000). Although a reduction in salivary gland size has been observed in Six1-deficient mice (Laclef et al., 2003), the mechanism by which Six1 influences submandibular salivary gland (SSG) development/maturation is unknown. To address this question, we have analyzed the expression profile of the *Six1* gene during normal SSG development/maturation, and have characterized the effect of *Six1* deficiency in this gland.

Material and methods

Animals

Generation and characterization of Six1 null mice have previously been described (Ozaki et al., 2004). The mice were housed at the Center for Laboratory Animal Care at the University of Colorado Denver (UCD) and treated in accordance with the NIH Guide to Humane Use of Animals in Research. All animal protocols were approved by the UCD-Institutional Animal Care and Use Committee (IACUC). To determine the normal temporal expression pattern of Six1, wild type C57Bl6/J mice were used. For Six1, *cyclin A1* and *cyclin A2* expression analysis, at each timepoint, SSG from 3 animals were sampled.

Reverse transcription-polymerase chain reaction (RT-PCR)

RNA was isolated from SSGs according to the manufacturer's protocol for Trizol reagent (Invitrogen, Carlsbad, CA, USA). Before the RT reactions, all RNA samples were treated with DNaseI for 10 min at room temperature in order to eliminate genomic DNA

contamination. Two micrograms of total RNA per sample were used to generate cDNA using random primers and Superscript II RNase H-reverse transcriptase (Invitrogen, Carlsbad, CA, USA). The resulting cDNAs were subsequently amplified in a 50 µl reaction mixture containing 1 µM of each primer, 2 mM MgCl₂, 0.8 mM dNTPs, and 0.025 U/µl Taq DNA polymerase. Actin was used as a housekeeping control. Primer pairs used to amplify mouse *Six1* (WT allele) were 5' GAA TCA ACT CTC TCC TCT GG 3' and 5' TTA GGA ACC CAA GTC CAC CA 3'; EGFP (mutated allele) primers were 5' CTG GTG ACC ACC CTG ACC TAC 3' and 5' TGA TCC CGG CGG CGG TCA CGA A 3'; and actin primers were 5' TAT CCT GAC CCT GAA GTA CC 3' and 5' GGT CAG GAT CTT CAT GAG GT 3'. After denaturation for 2 min at 94°C, 30 cycles of amplification were performed using a thermocycler, followed by a final extension of 10 min at 72°C. The amplification cycling parameters were: denaturation for 30 sec at 94°C, annealing for 1 min at 55°C, and extension for 2 min at 72°C. After amplification, 20 µl of PCR products were electrophoresed on a 1% agarose gel containing 0.5 µg/ml of ethidium bromide.

Quantitative real-time PCR (qPCR)

qPCR was performed using a model 7000 instrument (Applied Biosystems, Foster City, CA, USA). Amplicons were detected using Taqman fluorescence probes as described elsewhere (Lie & Petropoulos, 1998). The primers and probes used for this study were as follows: for *Six1* 5' AAC TGC AGC AGC TGT GGC T 3', 5' GTC GGC CGC GAA GTT TC 3', and 5' AAA GCG CAC TAC GTG GAG GCC G 3' (probe), and for cyclin A1 5' TTT CCC CAA TGC TGG TTG A 3', 5' AAC CAA AAT CCG TTG CTT CCT 3', and 5' CCC ACC ACC CAT GCC CAG TCA 3' (probe). The cyclin A2 and 18S rRNA primers and probes were purchased as Assays-on-Demand gene expression from Applied Biosystems. Target genes were analyzed using standard curves to determine relative levels of gene expression, and individual cDNA samples were normalized according to the levels of 18S rRNA.

Western blot analysis

SSGs were washed with cold phosphate-buffered saline (PBS) and lysed in RIPA buffer (50mM Tris-HCl pH7.4, 150mM NaCl, 1mM EDTA, 0.5% NP-40, 0.5% deoxycholic acid, 0.5% SDS, 1mM PMSF, 10mM NaF, 1mM Na₃VO₄ and 1µg/ml leupeptin). After centrifugation, protein concentrations were measured using a protein assay (Bio Rad, Hercules, CA, USA) according to the manufacturer's instructions. Western blot analysis was performed using anti-*Six1* and anti-actin antibodies as described (Ford et al., 2000).

Immunofluorescence

SSGs from *Six1*^{-/-} animals at embryonic day 18.5 (E18.5) were fixed in 4% paraformaldehyde in PBS for 16 h at 4°C. Following fixation, the tissues were washed in PBS, paraffin-embedded, and sectioned at 3 µm. After dewaxing and hydrating in graded alcohol solutions, the sections were treated with 10 mM citric acid pH6.0 in a microwave for 20 min. To prevent non-specific binding, the tissues were blocked with the M.O.M. mouse Ig blocking reagent (Vector Labs, Burlingame, CA, USA) in PBS for 16 h at 4°C. The tissues were then incubated with monoclonal mouse anti-GFP antibodies (Chemicon Int., Temecula, CA), washed with PBS, and incubated with goat anti-mouse IgG conjugated with fluorescein (Calbiochem, San Diego, CA, USA). Tissues were examined under an Olympus BX51 fluorescence microscope equipped with a Penguin 600CL camera. Tissues untreated with primary antibodies were used as negative controls.

SSG measurement, histology, cell proliferation and cell death assays

Pregnant female mice at day 18.5 of gestation were intraperitoneally injected with 100 mg of bromodeoxyuridine (BrdU) per kg body weight. Embryos were collected 2 h later and the

SSG volumes were calculated using the formula: $\text{volume} = 0.5 \times \text{Length} \times \text{Width}^2$. After measurement, the SSGs from wild *Six1*^{+/+} and *Six1*^{-/-} embryos were dissected with the aid of a stereoscopic microscope, fixed in 70% ethanol, embedded in paraffin, and sectioned at 3 μm . The sections were either hematoxylin & eosin (H&E) stained or analyzed for cell proliferation or cell death. Cell proliferation was measured via BrdU incorporation using an immunohistochemical analysis kit (GE Healthcare, Piscataway, NJ, USA), whereas cell death assays were performed by terminal deoxynucleotidyl transferase-mediated dUTP nick end labeling (TUNEL) analysis using an apoptosis detection kit (Intergen, Norcross, GA).

Statistical analysis

All assays were performed at least twice. Student's T-test was used for statistical analysis, and $p \leq 0.05$ was considered to indicate statistical significance.

Results

Six1 is dynamically expressed in the maturing SSG

To elucidate the potential role of *Six1* in SSG maturation, we first evaluated the temporal and spatial expression of *Six1* in wild-type mouse SSGs from E18.5 to postnatal day 28. *Six1* mRNA levels were high in SSGs from E18.5 to postnatal day 4, after which they declined progressively (Fig. 1A and B). In accordance with *Six1* mRNA expression levels, the amount of *Six1* protein decreased steadily in the developing postnatal SSG as shown by western blot analysis (Fig. 1C and D). To determine which cell types within the salivary gland express *Six1*, immunofluorescence was performed on E18.5 SSGs isolated from *Six1* heterozygote embryos, in which enhanced green fluorescent protein (EGFP) had been knocked into the *Six1* locus (Fig. 1E and F). EGFP expression was strongly observed in the developing ductal and acinar cells, with some mesenchymal cells showing very low levels of expression (Fig. 1G).

Six1^{-/-} SSGs are significantly smaller than wild-type littermates as a consequence of decreased proliferation

Our results confirm those from previous findings (Laclef et al., 2003) revealing that *Six1*^{-/-} SSGs are significantly smaller than *Six1*^{+/+} SSGs (Fig. 2A). The SSG volume in *Six1*^{+/+} animals (n=28) ranged from 0.67 to $2.45 \times 10^3 \text{ mm}^3$, with a mean of $1.41 \pm 0.38 \times 10^3 \text{ mm}^3$, whereas the SSG volume in *Six1*^{-/-} animals (n=21) ranged from 0.24 to $0.86 \times 10^3 \text{ mm}^3$, with a mean of $0.51 \pm 0.14 \times 10^3 \text{ mm}^3$ (Fig. 2B). Although the *Six1*^{-/-} SSGs were significantly smaller than wild-type SSGs, the histological structures of the *Six1*^{-/-} SSG acini and ducts showed normal morphogenesis (Fig. 2C).

Six1 stimulates proliferation and inhibits apoptosis in both cell culture and in mouse model systems (Coletta et al., 2004a, Ozaki et al., 2004, Xu et al., 2003, Li et al., 2003, Ikeda et al., 2007), suggesting that the loss of *Six1* leads to an exit from the cell cycle and to the premature death of epithelial cells. Thus, BrdU incorporation and TUNEL assays were performed to determine whether the decreased size of *Six1*^{-/-} SSGs was the result of attenuated cell proliferation and/or increased cell elimination by apoptosis. The percentage of total BrdU-positive cells in *Six1*^{+/+} and *Six1*^{-/-} SSGs was 33.92 ± 8.05 and $7.62 \pm 4.54\%$ at E18.5, respectively, demonstrating a marked decrease in the number of proliferating epithelial cells in the *Six1*^{-/-} SSGs (Fig. 3). However, the number of TUNEL-positive cells was very low in both *Six1*^{+/+} and *Six1*^{-/-} SSG epithelial cells, and no significant difference was observed (data not shown).

***Six1* transcriptional target cyclin A1 is decreased in *Six1*^{-/-} SSGs**

To investigate the molecular mechanism underlying the proliferation defect in *Six1*^{-/-} SSG epithelial cells, expression levels of cyclin A1, a known transcriptional target of *Six1* that mediates its effects on proliferation, was analyzed. First, the temporal expression of cyclin A1 and its functionally related gene, cyclin A2, was determined throughout SSG development. The temporal expression of cyclin A1 was very similar to that of *Six1*, with high levels of expression in the embryonic SSG. In contrast, cyclin A2 was most highly expressed in the postnatal SSG, although its levels were significantly higher than cyclin A1 throughout the course of SSG development (Fig. 4A). Expression of cyclin A1 was significantly lower in *Six1*^{-/-} SSGs as compared to *Six1*^{+/+} SSGs ($p < 0.01$), whereas cyclin A2 levels were not changed in a statistically significant manner (Fig. 4B). Thus, the proliferation defect observed in *Six1*^{-/-} SSG cells was accompanied by a downregulation of the *Six1* transcriptional target cyclin A1, which is highly expressed in the SSG during late embryogenesis.

Discussion

The SSG develops through a process of branching morphogenesis, undergoing complex stages of cell proliferation and differentiation, which are tightly coordinated by transcriptional regulatory pathways (Melnick & Jaskoll, 2000). However, the molecular mechanisms underlying these events within the SSG remain largely unknown. The homeobox gene family encodes proteins with DNA-binding and transcriptional activities that are critical in normal development (Pearson et al., 2005). To date, from a family with more than 200 members, only few homeobox genes are reported to be expressed in the mouse SSG (Biben et al., 2002, Jaskoll et al., 1998, Raju et al., 1993, Tanaka et al., 2000). In this study we have identified the expression of another homeoprotein in the mouse SSG, and we have examined the dependence of SSG maturation on this important developmental molecule. Expression of *Six1* is high during late stages of embryonic development (E18.5), and its levels progressively decline during postnatal SSG development. These findings are consistent with the general role of Six family members, which function in regulating cell proliferation and in specifying cell fate in the developing embryo (Christensen et al., 2008).

Studies examining *Six1* deficiency in mice suggest that it is a key participant in development by controlling the expansion of progenitor cell populations in specific organs (Li et al., 2003, Xu et al., 2003, Ozaki et al., 2004). Inner ear development in *Six1*^{-/-} embryos arrests at the otic vesicle stage and all components of the inner ear fail to form due to an increase in apoptosis and a decrease in proliferation of the cells of the otic epithelium (Zheng et al., 2003, Ozaki et al., 2004). In the kidney, loss of *Six1* leads to a failure of ureteric bud proliferation and invasion, with subsequent apoptosis of mesenchymal cells (Xu et al., 2003). Recently, it has been shown that *Six1* stimulates proliferation and tumor growth through direct activation of cyclin A1 (Coletta et al., 2004a). In contrast to studies which demonstrated that *Six1* alters both cell proliferation and apoptosis during inner ear and kidney development, no alterations in cell death were observed in *Six1*^{-/-} SSGs. However, *Six1* deficiency in the mouse SSG resulted in a profound decrease in proliferation. Therefore, we conclude from our studies that the significant reduction in size of *Six1*^{-/-} SSGs occurs as a result of decreased proliferation. Importantly, the lack of an apoptotic effect may be due to the developmental stages analyzed in this study. During SSG development, proliferation seems to be a constant event, whereas apoptosis is rare, except for a few apoptotic cells observed in a large duct at E15 in mouse (Melnick & Jaskoll, 2000), a stage which we did not include in our analysis. Interestingly, apoptosis is associated with rat intercalated duct development (Hayashi et al., 2000, Hecht et al., 2000).

Although our data clearly demonstrate that *Six1* deficiency decreases cell proliferation, the SSG is still properly formed and no difference in the degree of maturity of the gland was observed. The absence of morphological alterations in *Six1*^{-/-} SSGs confirms that the primary role of *Six1* in the SSG is to expand the SSG cell population and is not to influence differentiation or morphogenesis *per se*. It is likely that the SSG is not completely absent in *Six1*^{-/-} mice because additional factors are involved in the proliferation of SSG cells during development, and these can, in part, compensate for the loss of *Six1*. One possibility is that other *Six* family members compensate for the loss of *Six1*. At this time, the expression pattern of the other four *Six* family members throughout SSG development has not been investigated. Given the documented redundancy between *Six* family members during development (Grifone et al., 2005, Konishi et al., 2006, Kobayashi et al., 2007), another *Six* family member might compensate for the loss of *Six1* and promote proliferation to establish the SSG. Redundancy between *Six1* targets might also compensate for the loss of *Six1*. For example, the *Six1* transcriptional target cyclin A1 is, at least in part, functionally redundant with cyclin A2 during embryogenesis (Winston, 2001). Both A-type cyclins bind to and activate the cyclin-dependent kinases 1 and 2 and are required at critical points in the cell cycle, including the progression through S phase and the G2/M transition (Yang et al., 1999, Romanowski et al., 2000, Liu et al., 2000). Here we demonstrate that cyclin A1 expression is dramatically reduced in the *Six1*^{-/-} SSG, whereas cyclin A2 levels remain unchanged. Thus, it is possible that cyclin A2 partially compensates for the loss of cyclin A1 in the *Six1*-deficient SSG. Furthermore, as no SSG size alterations have been reported in cyclin A1 null mice (Liu et al., 1998), it is likely that other cyclin A1-independent proliferative pathways important for SSG development are affected by *Six1*.

In summary, we demonstrate that *Six1* is important for SSG development by controlling the proliferation that is necessary for the expansion of SSG epithelial cells. Recent evidence suggests that *Six1* may play important roles in multiple tumor types, as its overexpression has been observed in breast (Reichenberger et al., 2005, Coletta et al., 2004b, Ford et al., 1998), ovarian (Behbakht et al., 2007), hepatocellular carcinoma (Ng et al., 2006), cervical carcinoma (Wan et al., 2008), Wilms' tumor (Li et al., 2002), and alveolar rhabdomyosarcoma (RMS) (Yu et al., 2004, Khan et al., 1999). Interestingly, in some cases, *Six1* expression is high in cancers derived from tissues where it is normally expressed and where it plays a functional role during development, including kidney and muscle (Khan et al., 1999, Li et al., 2002). As the molecular pathways involved in carcinogenesis often represent aberrations of normal processes that control embryogenesis (Abate-Shen, 2002), it is of interest to determine whether *Six1* can contribute to salivary gland tumorigenesis.

Acknowledgments

This work was supported by grants from the NIH (R01CA095277), Susan G. Komen Breast Cancer Foundation (9862), American Cancer Society/UCCC, and Avon Foundation for H.L.F. R.D.C. was supported by fellowships from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Brazil, University of Colorado Cancer Center, Colorado Cancer League, and W.M. Thorkildsen Foundation. E.L.M. was funded by a predoctoral fellowship from the Department of Defense Breast Cancer Research Program (W81XWH-06-1-0409).

References

- Abate-Shen C. Deregulated homeobox gene expression in cancer: cause or consequence? *Nat Rev Cancer*. 2002; 2:777–85. [PubMed: 12360280]
- Behbakht K, Qamar L, Aldridge CS, Coletta RD, Davidson SA, Thorburn A, Ford HL. *Six1* overexpression in ovarian carcinoma causes resistance to TRAIL-mediated apoptosis and is associated with poor survival. *Cancer Res*. 2007; 67:3036–42. [PubMed: 17409410]

- Biben C, Wang CC, Harvey RP. NK-2 class homeobox genes and pharyngeal/oral patterning: Nkx2-3 is required for salivary gland and tooth morphogenesis. *Int J Dev Biol.* 2002; 46:415–22. [PubMed: 12141427]
- Christensen KL, Patrick AN, McCoy EL, Ford HL. The Six Family of Homeobox Genes in Development and Cancer. *Adv Cancer Res.* 2008; 101
- Coletta RD, Christensen K, Reichenberger KJ, Lamb J, Micomonaco D, Huang L, Wolf DM, Müller-Tidow C, Golub TR, Kawakami K, Ford HL. The Six1 homeoprotein stimulates tumorigenesis via reactivation of cyclin A1. *Proc Natl Acad Sci U S A.* 2004a; 101:6478–83. [PubMed: 15123840]
- Coletta RD, Christensen K, Reichenberger KJ, Lamb J, Micomonaco D, Huang L, Wolf DM, Müller-Tidow C, Golub TR, Kawakami K, Ford HL. The Six1 homeoprotein stimulates tumorigenesis by reactivation of cyclin A1. *Proc Natl Acad Sci U S A.* 2004b; 101:6478–83. [PubMed: 15123840]
- Ford HL, Kabingu EN, Bump EA, Mutter GL, Pardee AB. Abrogation of the G2 cell cycle checkpoint associated with overexpression of HSIX1: a possible mechanism of breast carcinogenesis. *Proc Natl Acad Sci U S A.* 1998; 95:12608–13. [PubMed: 9770533]
- Ford HL, Landesman-Bollag E, Dacwag CS, Stukenberg PT, Pardee AB, Seldin DC. Cell cycle-regulated phosphorylation of the human SIX1 homeodomain protein. *J Biol Chem.* 2000; 275:22245–54. [PubMed: 10801845]
- Grifone R, Demignon J, Houbroun C, Souil E, Niro C, Seller MJ, Hamard G, Maire P. Six1 and Six4 homeoproteins are required for Pax3 and Mrf expression during myogenesis in the mouse embryo. *Development.* 2005; 132:2235–49. [PubMed: 15788460]
- Hayashi H, Ozono S, Watanabe K, Nagatsu I, Onozuka M. Morphological aspects of the postnatal development of submandibular glands in male rats: involvement of apoptosis. *J Histochem Cytochem.* 2000; 48:695–8. [PubMed: 10769053]
- Hecht R, Connelly M, Marchetti L, Ball WD, Hand AR. Cell death during development of intercalated ducts in the rat submandibular gland. *Anat Rec.* 2000; 258:349–58. [PubMed: 10737853]
- Ikeda K, Ookawara S, Sato S, Ando Z, Kageyama R, Kawakami K. Six1 is essential for early neurogenesis in the development of olfactory epithelium. *Dev Biol.* 2007; 311:53–68. [PubMed: 17880938]
- Jaskoll T, Luo W, Snead ML. Msx-2 expression and glucocorticoid-induced overexpression in embryonic mouse submandibular glands. *J Craniofac Genet Dev Biol.* 1998; 18:79–87. [PubMed: 9672840]
- Kawakami K, Sato S, Ozaki H, Ikeda K. Six family genes--structure and function as transcription factors and their roles in development. *Bioessays.* 2000; 22:616–26. [PubMed: 10878574]
- Khan J, Bittner ML, Saal LH, Teichmann U, Azorsa DO, Gooden GC, Pavan WJ, Trent JM, Meltzer PS. cDNA microarrays detect activation of a myogenic transcription program by the PAX3-FKHR fusion oncogene. *Proc Natl Acad Sci U S A.* 1999; 96:13264–9. [PubMed: 10557309]
- Kobayashi H, Kawakami K, Asashima M, Nishinakamura R. Six1 and Six4 are essential for Gdnf expression in the metanephric mesenchyme and ureteric bud formation, while Six1 deficiency alone causes mesonephric-tubule defects. *Mech Dev.* 2007; 124:290–303. [PubMed: 17300925]
- Konishi Y, Ikeda K, Iwakura Y, Kawakami K. Six1 and Six4 promote survival of sensory neurons during early trigeminal gangliogenesis. *Brain Res.* 2006; 1116:93–102. [PubMed: 16938278]
- Laclef C, Souil E, Demignon J, Maire P. Thymus, kidney and craniofacial abnormalities in Six 1 deficient mice. *Mech Dev.* 2003; 120:669–79. [PubMed: 12834866]
- Li CM, Guo M, Borczuk A, Powell CA, Wei M, Thaker HM, Friedman R, Klein U, Tycko B. Gene expression in Wilms' tumor mimics the earliest committed stage in the metanephric mesenchymal-epithelial transition. *Am J Pathol.* 2002; 160:2181–90. [PubMed: 12057921]
- Li X, Oghi KA, Zhang J, Krones A, Bush KT, Glass CK, Nigam SK, Aggarwal AK, Maas R, Rose DW, Rosenfeld MG. Eya protein phosphatase activity regulates Six1-Dach-Eya transcriptional effects in mammalian organogenesis. *Nature.* 2003; 426:247–54. [PubMed: 14628042]
- Lie YS, Petropoulos CJ. Advances in quantitative PCR technology: 5' nuclease assays. *Curr Opin Biotechnol.* 1998; 9:43–8. [PubMed: 9503586]
- Liu D, Liao C, Wolgemuth DJ. A role for cyclin A1 in the activation of MPF and G2-M transition during meiosis of male germ cells in mice. *Dev Biol.* 2000; 224:388–400. [PubMed: 10926775]

- Liu D, Matzuk MM, Sung WK, Guo Q, Wang P, Wolgemuth DJ. Cyclin A1 is required for meiosis in the male mouse. *Nat Genet.* 1998; 20:377–80. [PubMed: 9843212]
- Melnick M, Jaskoll T. Mouse submandibular gland morphogenesis: a paradigm for embryonic signal processing. *Crit Rev Oral Biol Med.* 2000; 11:199–215. [PubMed: 12002815]
- Ng KT, Man K, Sun CK, Lee TK, Poon RT, Lo CM, Fan ST. Clinicopathological significance of homeoprotein Six1 in hepatocellular carcinoma. *Br J Cancer.* 2006; 95:1050–5. [PubMed: 17008870]
- Ozaki H, Nakamura K, Funahashi J, Ikeda K, Yamada G, Tokano H, Okamura HO, Kitamura K, Muto S, Kotaki H, Sudo K, Horai R, Iwakura Y, Kawakami K. Six1 controls patterning of the mouse otic vesicle. *Development.* 2004; 131:551–62. [PubMed: 14695375]
- Pearson JC, Lemons D, McGinnis W. Modulating Hox gene functions during animal body patterning. *Nat Rev Genet.* 2005; 6:893–904. [PubMed: 16341070]
- Raju K, Tang S, Dube ID, Kamel-Reid S, Bryce DM, Breitman ML. Characterization and developmental expression of Tlx-1, the murine homolog of HOX11. *Mech Dev.* 1993; 44:51–64. [PubMed: 7908826]
- Reichenberger KJ, Coletta RD, Schulte AP, Varella-Garcia M, Ford HL. Gene amplification is a mechanism of Six1 overexpression in breast cancer. *Cancer Res.* 2005; 65:2668–75. [PubMed: 15805264]
- Romanowski P, Marr J, Madine MA, Rowles A, Blow JJ, Gautier J, Laskey RA. Interaction of *Xenopus Cdc2* × cyclin A1 with the origin recognition complex. *J Biol Chem.* 2000; 275:4239–43. [PubMed: 10660590]
- Tanaka M, Komuro I, Inagaki H, Jenkins NA, Copeland NG, Izumo S. Nkx3.1, a murine homolog of *Drosophila bagpipe*, regulates epithelial ductal branching and proliferation of the prostate and palatine glands. *Dev Dyn.* 2000; 219:248–60. [PubMed: 11002344]
- Wan F, Miao X, Quraishi I, Kennedy V, Creek KE, Pirisi L. Gene expression changes during HPV-mediated carcinogenesis: a comparison between an in vitro cell model and cervical cancer. *Int J Cancer.* 2008; 123:32–40. [PubMed: 18398830]
- Winston N. Regulation of early embryo development: functional redundancy between cyclin subtypes. *Reprod Fertil Dev.* 2001; 13:59–67. [PubMed: 11545166]
- Xu PX, Zheng W, Huang L, Maire P, Laclef C, Silvius D. Six1 is required for the early organogenesis of mammalian kidney. *Development.* 2003; 130:3085–94. [PubMed: 12783782]
- Yang R, Muller C, Huynh V, Fung YK, Yee AS, Koeffler HP. Functions of cyclin A1 in the cell cycle and its interactions with transcription factor E2F-1 and the Rb family of proteins. *Mol Cell Biol.* 1999; 19:2400–7. [PubMed: 10022926]
- Yu Y, Davicioni E, Triche TJ, Merlino G. The Homeoprotein Six1 Transcriptionally Activates Multiple Protumorigenic Genes but Requires Ezrin to Promote Metastasis. *Cancer Res.* 2006; 66:1982–1989. [PubMed: 16488997]
- Yu Y, Khan J, Khanna C, Helman L, Meltzer PS, Merlino G. Expression profiling identifies the cytoskeletal organizer ezrin and the developmental homeoprotein Six-1 as key metastatic regulators. *Nat Med.* 2004; 10:175–81. [PubMed: 14704789]
- Zheng W, Huang L, Wei ZB, Silvius D, Tang B, Xu PX. The role of Six1 in mammalian auditory system development. *Development.* 2003; 130:3989–4000. [PubMed: 12874121]
- Zou D, Silvius D, Fritzscht B, Xu PX. Eya1 and Six1 are essential for early steps of sensory neurogenesis in mammalian cranial placodes. *Development.* 2004; 131:5561–72. [PubMed: 15496442]




Figure 1. Temporal and spatial expression of *Six1* during SSG late embryogenesis is dynamic. **(A)** RNA isolated from wild-type SSGs was subjected to RT-PCR assays using specific primers for *Six1* (actin was used as a control). **(B)** Comparison of *Six1* expression throughout SSG development by densitometric analysis indicates that *Six1* mRNA levels progressively decrease from E18.5 to postnatal day 28. Values are expressed in arbitrary units as the ratio of the optical density of *Six1*/actin. **(C and D)** Western blot and densitometric analyses demonstrate that the *Six1* protein is present at high levels in SSG development, and levels decrease throughout the course of development. **(E)** PCR analyses used to genotype *Six1*^{+/+}, *Six1*^{+/-}, and *Six1*^{-/-} neonates. Tail DNA was isolated and PCR was performed using specific primer pairs as described. **(F)** RT-PCR analyses of SSG from *Six1*^{+/+}, *Six1*^{+/-}, and *Six1*^{-/-} at day E18.5 confirms the absence of *Six1* mRNA and the expression of EGFP in the *Six1*^{-/-} SSG. **(G)** H&E staining and immunostaining against EGFP, which was knocked in to the *Six1* locus, demonstrate the expression of *Six1* in the ductal and acinar cells of the SSG at E18.5. (Magnification, ×40)




Figure 2.

SSGs are reduced in size in *Six1*-deficient embryos. (A) Photographs of SSGs from a *Six1*^{+/+} embryo (left) and from a *Six1*^{-/-} littermate (right) at E18.5. Top panels represent SSG within the mouse, whereas bottom panels represent glands after dissection from the mouse. (B) Measurements of SSG size demonstrate that *Six1*^{-/-} SSGs are significantly smaller than *Six1*^{+/+} SSGs. (C) E18.5 SSGs from *Six1*^{-/-} mice are completely normal in structure when compared with litter-matched wild type mice. (Magnification: A, B ×25; C, D ×40)




Figure 3. Proliferation is dramatically decreased in *Six1*^{-/-} SSG epithelial cells. BrdU-staining in SSGs from *Six1*^{+/+} (A) and *Six1*^{-/-} (B) embryos at E18.5. (C) The BrdU index, expressed as the percentage of positive cells, was determined by counting 1,500 cells in five independent samples for each genotype.




Figure 4.

Six1 and cyclin A1 expression are coordinately regulated during embryogenesis, and expression of cyclin A1 is significantly reduced in *Six1*^{-/-} SSG. SSGs were dissected, RNA isolated and converted into cDNA, and Six1, cyclin A1, and cyclin A2 expression levels determined by qPCR. **(A)** Representative expression of Six1, cyclin A1, and cyclin A2 during different developmental stages of the mouse SSG. Note that Six1 and cyclin A1 have a similar temporal expression during late development of SSG. **(B)** Expression levels of Six1, cyclin A1, and cyclin A2 in the SSG from *Six1*^{+/+} and *Six1*^{-/-} embryos at day 18.5. Results represent the average \pm SD of 3 independent samples, each sample containing the pooled SSGs of 5 mice for each genotype. SSGs from *Six1*^{-/-} mice demonstrated a significant reduction in cyclin A1 expression as compared to *Six1*^{+/+}.