



Published in final edited form as:

J Invest Dermatol. 2009 January ; 129(1): 148–154. doi:10.1038/jid.2008.205.

NF- κ B Mediates Mitogen-Activated Protein Kinase Pathway-Dependent iNOS Expression in Human Melanoma

Deon G. Uffort, Elizabeth A. Grimm, and Julie A. Ellerhorst

The Department of Experimental Therapeutics, The University of Texas M. D. Anderson Cancer Center, Houston, TX, USA

Abstract

Tumor expression of inducible nitric oxide synthase (iNOS) predicts poor outcomes for melanoma patients. We have reported the regulation of melanoma iNOS by the mitogen-activated protein kinase (MAPK) pathway. In the current study, we test the hypothesis that NF- κ B mediates this regulation. Western blotting of melanoma cell lysates confirmed the constitutive expression of iNOS. Baseline levels of activated nuclear ERK and NF- κ B were also detected by western blot. Indirect immunofluorescence confirmed the presence of NF- κ B p50 and p65 in melanoma cell nuclei, with p50 being more prevalent. Electrophoretic mobility shift assay demonstrated baseline NF- κ B activity, the findings confirmed by supershift analysis. Treatment of melanoma cells with the MEK inhibitor U0126 decreased NF- κ B binding to its DNA recognition sequence, implicating the MAPK pathway in NF- κ B activation. Two specific NF- κ B inhibitors suppressed iNOS expression, demonstrating regulation of iNOS by NF- κ B. Several experiments indicated the presence of p50 homodimers, which lack a transactivation domain and rely upon the transcriptional co-activator Bcl-3 to carry out this function. Bcl-3 was detected in melanoma cells and co-immunoprecipitated with p50. These data suggest that the constitutively activated melanoma MAPK pathway stimulates activation of NF- κ B hetero- and homodimers, which, in turn, drive iNOS expression and support melanoma tumorigenesis.

Keywords

MAPK; NF- κ B; iNOS; p50 homodimer; Bcl-3

INTRODUCTION

Malignant melanoma is currently the most virulent form of skin cancer, with incidence and mortality rates steadily rising in the United States population (Jemal *et al.*, 2007). Although much improvement has been achieved regarding the treatment of early stage disease, many patients die each year from melanoma metastases as tumor cells become increasingly resistant to chemotherapy. As such, the elucidation of pathways supporting metastatic cell growth is needed to develop specific targeted therapies for advanced disease.

The p44/42 mitogen-activated protein kinase (MAPK) pathway is arguably the most critical signaling cascade supporting the uncontrolled growth of melanoma cells (Smalley, 2003). In

Address correspondence to: Julie Ellerhorst M.D., Ph.D., Department of Experimental Therapeutics, Unit 362, The University of Texas M. D. Anderson Cancer Center, 1515 Holcombe Blvd., Houston, TX 77030, Ph: 713-792-8990, Fax: 713-563-3531, jaellerh@mdanderson.org .

CONFLICT OF INTEREST

The authors state no conflict of interest.

normal cells, the MAPK pathway is cytokine inducible. Signal induced by ligand binding is transported to a series of protein kinases that eventually culminate in the phosphorylation and activation of the extracellular signal-regulated kinase (ERK). Activated ERK translocates to the nucleus and initiates the transcription of a variety of growth-related genes. In melanoma cells, the MAPK pathway is constitutively active, a condition attributed in part to somatic mutations of *NRAS* and *BRAF*, which occur in the majority of cases (Goydos *et al.*, 2005). The aberrant activation of this pathway confers a protective benefit to melanoma cells by driving their proliferation and survival. The critical nature of the MAPK pathway has generated interest in melanoma-specific targets downstream of ERK which may participate in these oncogenic processes.

One such target of particular interest to our laboratory is inducible nitric oxide synthase (iNOS). The catalytic product of iNOS is nitric oxide (NO), a diatomic free radical that mediates processes such as neurotransmission, vasodilation, and host defense in normal cells. NO has been shown to contribute to the pathogenesis of a variety of cancers, as well. (Thomsen *et al.*, 1994; Cobbs *et al.*, 1995). Specifically, in melanoma, iNOS-generated NO has been shown to protect tumor cells from apoptosis (Salvucci *et al.*, 2001; Tang and Grimm, 2004). We have recently reported that the MAPK pathway drives constitutive iNOS expression in melanoma cells (Ellerhorst *et al.*, 2006). Furthermore, we have published data demonstrating an association of tumor iNOS expression with poor prognosis in patients diagnosed with Stage III melanoma (Ekmekcioglu *et al.*, 2006).

The findings described above have prompted us to explore the molecular links between melanoma ERK and iNOS, to further clarify this tumor-promoting pathway. A particularly promising candidate protein is the transcription factor NF- κ B, the activation of which is a hallmark of many cancers, including melanoma. (Wang *et al.*, 1999; Huang *et al.*, 2000; Dhawan and Richmond, 2002; Tian *et al.*, 2006). NF- κ B consists of a family of structurally related proteins, including Rel A (p65), Rel B, c-Rel, NF- κ B1 (p50/p105), and NF- κ B2 (p52/p100). In quiescent cells, NF- κ B exists as cytoplasmic hetero- or homodimers associated with an inhibitory protein belonging to the I κ B family. Upon appropriate cytokine stimulation, I κ B kinase (IKK) induces the phosphorylation of I κ B, promoting its subsequent ubiquitination and degradation in the proteasome (Baldwin, 1996). This process allows the translocation of NF- κ B to the nucleus where it binds to target DNA elements and induces transcription of various genes involved in inflammation and survival, including *iNOS* (Taylor *et al.*, 1998). Evidence for a role of MAPK in NF- κ B activation has been reported in several non-melanoma systems, as well as in the HS294T human melanoma cell line (Kurland *et al.*, 2003; Wang *et al.*, 1999; Dhawan and Richmond, 2002; Jiang *et al.*, 2004).

We have thus hypothesized that NF- κ B serves as the regulatory link between the MAPK pathway and iNOS expression in human melanoma cells. In the current study, we explore this proposed sequence of molecular signaling events and show that, indeed, the constitutively activated melanoma MAPK pathway drives NF- κ B activation, which, in turn induces iNOS expression.

RESULTS

Human melanoma cells constitutively express activated ERK, activated NF- κ B, and iNOS

Initial experiments were performed to demonstrate the presence in melanoma cells of the three proteins included in our proposed pathway: ERK, NF- κ B, and iNOS. The human melanoma cell lines WM35, WM793, MeWo, and HS294T, as well as primary melanocytes (FMC7C and FMC14C), were used in these experiments. WM793 and WM35 carry *BRAF* T1799A mutations, whereas MeWo, HS294T, and both melanocyte lines are *BRAF* wild type. None of these lines carry *NRAS* mutations. Nuclear extracts prepared from these cells under basal

growth conditions contained both NF- κ B p50 and p65, and phosphorylated ERK (pERK), indicating the presence of these proteins in the activated form (Figure 1a). Nuclear expression varied from one cell line to the next, but tended to be higher in the melanoma cells when compared to the melanocytes (Figure 1c). A western blot of whole cell lysates from the same cells revealed the constitutive presence of iNOS protein and demonstrated the same pattern of higher levels in the melanoma cells (Figures 1b and 1c). An unanticipated observation was the presence of iNOS protein in the nuclear extracts of the melanoma cell lines and melanocytes (Figure 1b), a finding that has not been previously reported.

Indirect immunofluorescence studies were next carried out to confirm the nuclear localization of NF- κ B in these cells. In keeping with the immunoblotting data, constitutive translocation of NF- κ B p50 and p65 to the nuclei of melanoma cells was demonstrated (Figure 2). Because nuclear NF- κ B can exist as p50/p65 heterodimers or p50 homodimers, the prevalence of each of these NF- κ B subunits was examined in the immunofluorescently stained cells. Among the melanoma cell lines, there was a significant difference in the number of cells expressing nuclear p50 and p65, with some cells expressing only p50 (Figure 3). This finding was not observed in the melanocytes (FMC7C). These data suggest that melanoma cells utilize both heterodimeric and homodimeric NF- κ B forms for transcriptional activation, and that the p50 homodimer activity may be tumor specific.

The melanoma MAPK pathway regulates NF- κ B

The first segment of the proposed MAPK/NF- κ B/iNOS pathway involves the regulation of melanoma NF- κ B by MAPK. An electrophoretic mobility shift assay (EMSA), carried out with nuclear extracts from WM793 cells at baseline, revealed NF- κ B binding to its consensus oligonucleotide DNA target sequence (Figure 4a). Supershift experiments with antibodies to NF- κ B p50 and p65 revealed even greater retardation of the DNA:protein:antibody complexes. The antibody to NF- κ B p50 produced a more complete supershift than antibody to the p65 subunit, consistent with the retardation of both p50 homodimers and p50/p65 heterodimers by the anti-p50 antibody, whereas the p65 antibody recognizes only the heterodimers, leaving the homodimers unaffected (Figure 4b).

To test the hypothesis that NF- κ B is regulated by the melanoma MAPK pathway, additional EMSA experiments were performed using WM793 cells to assess the effects of the MEK inhibitor U0126 on NF- κ B nuclear binding activity. These experiments revealed diminished NF- κ B binding as early as 2 minutes, with nearly complete abrogation of activity after 5 minutes of exposure to U0126 (Figure 4c). Experiments repeated with MeWo cells yielded similar results (data not shown). These data convincingly support the activation of NF- κ B by the MAPK pathway in melanoma cells.

NF- κ B regulates iNOS expression in melanoma cells

The second segment of the proposed pathway involves the induction of melanoma iNOS expression by NF- κ B. To examine this regulatory event, melanoma cells were treated with each of two inhibitors which target different aspects of NF- κ B activation. The first, NF- κ B Activation Inhibitor, interferes with the transcriptional ability of NF- κ B, whereas the second inhibitor, IKK-2 inhibitor V, selectively blocks I κ B α phosphorylation, preventing the translocation of NF- κ B to the nucleus.

Figure 5a shows the inhibition of iNOS protein expression in MeWo cells over the course of a 48-hour incubation with NF- κ B Activation Inhibitor. Similar findings are demonstrated in WM793 cells treated with IKK-2 inhibitor V (Fig. 5b); in this experiment, the decline in iNOS levels is preceded by a reduction in phosphorylation of I κ B α . With both inhibitors, the stable, dimeric iNOS functional unit persists for the first 12 hours, but is markedly diminished

thereafter. Similar findings were seen with WM35 cells (data not shown). These results are consistent with the hypothesis that iNOS expression is induced by NF- κ B in human melanoma.

Melanoma cells express the transcriptional co-activator Bcl-3 which complexes with NF- κ B p50

Data from several of the experiments described above indicated the presence of both NF- κ B p50 homodimers and p50/p65 heterodimers in the nuclei of melanoma cells. The p50 homodimer binds DNA *in vitro*, but lacks a transcriptional activation domain and, consequently, initiates transcription weakly, if at all. In fact, this NF- κ B homodimer appears to function as transcriptional inhibitor in some cell types by competing with p50/p65 for promoter binding (Kang *et al.*, 1992; Plaksin *et al.*, 1993). Countering this inhibitory effect in a variety of transformed cell types is the oncoprotein Bcl-3, which binds NF- κ B p50 homodimers and robustly co-activates transcription (Fujita *et al.*, 1993). Experiments were thus conducted to explore a potential role of Bcl-3 in melanoma NF- κ B p50 transcriptional co-activation. Western blots of whole cell lysates and nuclear extracts from melanoma cells revealed the presence of the Bcl-3 protein in both the cytoplasmic and nuclear compartments (Figure 6a). Notably, Bcl-3 was virtually undetectable in melanocytes. Subsequently, the association of Bcl-3 and NF- κ B p50 in melanoma cells was examined. Proteins from whole cell lysates were immunoprecipitated with rabbit IgG or anti-Bcl-3 antibody, followed by western blotting with antibody to p50. Figure 6b demonstrates the presence of NF- κ B p50 in the Bcl-3 immunoprecipitates, indicating the presence of Bcl-3/p50 complexes in melanoma cells, with the potential for transcriptional activity.

DISCUSSION

The Ras/Raf/MEK/ERK pathway is a major signaling pathway involved in tumor cell growth and survival. As human melanoma is driven by constitutive activation of this pathway, elucidation of downstream effector molecules is critical to the understanding of melanoma pathobiology. We have previously reported the positive regulation of melanoma iNOS expression by the MAPK pathway and the profoundly negative clinical implications of tumor iNOS expression in this malignancy (Ekmekcioglu *et al.*, 2006; Ellerhorst *et al.*, 2006). We now show that NF- κ B mediates this regulatory event. Previous work in the literature has described the activation of NF- κ B by the MAPK pathway in the HS294T human melanoma cell line (Dhawan and Richmond, 2002). Our data now confirm and extend this finding, as we demonstrate that the inhibition of MAPK signaling with the MEK inhibitor U0126 is accompanied by diminished nuclear NF- κ B DNA binding. Furthermore, this is the first study to show that NF- κ B regulates expression of iNOS in melanoma cells. Notably, the MAPK/NF- κ B/iNOS pathway was demonstrated in both *BRAF* mutant and wild type cells, suggesting that the source of MAPK activation does not alter these downstream events.

An unexpected finding was the presence of iNOS in the nuclei of cultured melanoma cells and melanocytes. Although nuclear iNOS has been reported in neutrophils and adipocytes of rat origin (Giordano *et al.*, 2002; Saini *et al.*, 2006), this is the first description of iNOS in the nuclei of human cells. Of note, endothelial NOS (eNOS) has been detected in the nuclei of cultured human mast cells (Gilchrist *et al.*, 2004). In those cell lines, eNOS phosphorylation preceded translocation from the cytoplasm to the nucleus. The specific role of nuclear NOS and, implicitly, nuclear NO, has yet to be explored. In melanoma cells, it could be hypothesized that nuclear compartmentalization of iNOS and NO may confer a strategic advantage in terms of efficient modification of DNA and nuclear proteins by nitric oxide, in such a way as to promote growth and survival. Although nuclear iNOS was detected in melanocytes as well, these normally quiescent cells are maintained in growth media containing potent mitogens such as cholera toxin and phorbol myristate acetate, providing growth pathway stimulation similar

to that seen in tumor cells. It therefore remains to be determined whether nuclear translocation of iNOS is a tumor-specific event. Future research in our laboratory will address this issue as well as the question of iNOS phosphorylation, as has been described for eNOS.

The presence of NF- κ B p50 homodimers in the nuclei of melanoma cells is an intriguing finding. This NF- κ B subtype may play a role in the competitive transcriptional inhibition of p50/p65, a function frequently attributed to the p50 homodimer (Kang *et al.*, 1992; Plaksin *et al.*, 1993). However, the expression of the transcriptional co-activator Bcl-3 by the melanoma cell lines, and the complexing of Bcl-3 with p50, provide preliminary evidence to suggest that the p50 homodimers may be transcriptionally active. The oncoprotein Bcl-3 is a member of the I κ B family of NF- κ B -binding proteins. The *BCL3* gene, located on chromosome 19, is translocated intact and is activated in cases of t(14;19)(q32.3;q13.2) B-cell chronic lymphocytic leukemia (Ohno *et al.*, 1990). Bcl-3 has additionally been implicated in the pathogenesis of nasopharyngeal carcinoma, breast cancer, and a variety of lymphomas, all cases giving evidence for co-activation of NF- κ B p50 by Bcl-3 as the transforming event (Fujita *et al.*, 1993; Watanabe *et al.*, 1997; Cogswell *et al.*, 2000; Thornburg *et al.*, 2003; Mathas *et al.*, 2005). Bcl-3 expression and its interaction with NF- κ B p50 have not been previously reported in melanoma. It is interesting to speculate that Bcl-3/p50 complexes may play an important role in the transcription of melanoma-promoting genes, such as *INOS*.

In summary, we have further defined the pathway leading from the constitutively active melanoma MAPK pathway to iNOS expression by demonstrating that NF- κ B is an important intermediary protein. In the process, we have made the novel observations of the nuclear localization of iNOS and of Bcl-3/NF- κ B p50 complexes in cultured melanoma cells. These findings will stimulate further exploration of these regulatory molecules and pathways, always with the ultimate goal of identifying targets for future therapy.

MATERIALS AND METHODS

Cell lines

The human melanoma cell lines WM793 and WM35 were generous gifts from Dr. Robert Kerbel (Sunnybrook Health Science Center, Toronto, ON, Canada) and the human metastatic melanoma MeWo cell line was provided by Dr. David Menter (M. D. Anderson Cancer Center). The HS294T melanoma cell line was kindly provided by Dr. Ann Richmond (Vanderbilt University School of Medicine, Nashville TN). BJ fibroblasts were purchased from the American Type Culture Collection (Manassas, VA). Cells were grown in RPMI 1640 medium, supplemented with 10% fetal bovine serum (FBS). Primary melanocytes FMC7C and FMC14C were derived from neonatal foreskin and were maintained in MCDB-153 media (Sigma, St. Louis MO), supplemented with 1% FBS, 10 ng/ml phorbol-12-myristate-13-acetate (Calbiochem, San Diego CA), 1 ng/ml basic fibroblast growth factor (Invitrogen, Carlsbad, CA), 5 ug/ml transferrin (Sigma), 10 nM cholera toxin (Calbiochem), 0.1 mM 3-isobutyl-1-methylxanthine (Calbiochem), 30 ug/ml bovine pituitary extract (Sigma), and 5ug/ml insulin (Sigma). All cells were maintained at 37°C with 5% CO₂.

Reagents and antibodies

NF- κ B Activation inhibitor and IKK-2 inhibitor V were purchased from Calbiochem and U0126 from Cell Signaling Technology (Beverly, MA). Protein A agarose was obtained from Sigma. Antibodies to ERK, phosphorylated ERK, I κ B α , and phosphorylated I κ B α were purchased from Cell Signaling Technology. Antibodies to histone, iNOS, and NF- κ B p50 and p65 were obtained from Santa Cruz Biotechnology (Santa Cruz, CA). The anti Bcl-3 antibody was purchased from Abcam (Cambridge, MA).

Whole cell extracts and western blotting

Cells were washed with cold PBS and harvested into PBS with 1 mM phenylmethylsulfonyl fluoride (PMSF). Cell pellets were then treated with lysis buffer (140 mM NaCl, 25 mM Tris HCl pH 7.4, and 1% NP-40) with freshly added protease inhibitor cocktail (BD Biosciences, San Jose, CA). The supernatants were then collected after rigorous agitation and protein concentration was measured. Proteins were separated by SDS-PAGE, transferred to a nitrocellulose membrane, and blocked for 1 hour in 5% nonfat milk in PBS. Primary antibody was diluted in 5% nonfat dry milk/PBS/0.1% Tween and incubated overnight at 4°C, followed by 45 minutes incubation with horseradish peroxidase labeled secondary antibody, again diluted in 5% nonfat dry milk/PBS/0.1% Tween. Membrane development was achieved with enhanced chemiluminescence (Amersham Pharmacia Biotech, Piscataway, NJ).

Nuclear extracts

The two extraction buffers used, A and B, comprised of the following reagents: Buffer A: 10 mM Hepes pH 7.9, 2 mM MgCl₂, 10 mM KCl, 0.1 mM EDTA, 0.1 mM EGTA, 5 mM NaF, 1 mM Na₃VO₄; Buffer B: 20 mM Hepes, 0.4 M NaCl, 1 mM EDTA, 1 mM EGTA, 5 mM NaF, 1 mM Na₃VO₄. Cells were washed with cold PBS and harvested into PBS/1mM PMSF. Cells were then treated with cold buffer A, followed by the addition of protease inhibitor cocktail 1:50. After the addition of 10% NP-40 and centrifugation at 3500g, the cytoplasmic fraction was collected. The remaining pellet was washed 3 times with buffer A, followed by the addition of Buffer B and 1:50 protease cocktail inhibitor. After final centrifugation at 16,000g, the nuclear fraction was collected.

Indirect immunofluorescence

Cells grown on chamber slides were fixed with 2% paraformaldehyde on ice (30 minutes), blocked with 5% serum in PBS (30 minutes room temperature), and incubated with primary antibody diluted in blocking solution (2 hours, 4°C). This was followed by incubation with FITC-labeled secondary antibody (1 hour, room temperature). Staining was observed and imaged with a Nikon Eclipse TE 2000-U microscope equipped with a Nikon digital DXM 1200F camera.

Electrophoretic mobility shift assay

Nuclear protein extracts for EMSA were prepared as described above. The NF-κB, AP2, and SP1 target oligonucleotides and gel shift binding buffer were purchased from Promega (Madison, WI). Binding reactions were performed with 4 μg of nuclear extract and 1.75 pmol of NF-κB target oligonucleotide (unlabeled or end-labeled with ³²P). AP2 or SP1 target DNA sequences were used as nonspecific competitor oligonucleotides. The sequence of the double-stranded NF-κB target oligonucleotide is:

5'-AGTTGAGGGGACTTTCCCAGGC-3'

3'-TCAACTCCCCTGAAAGGGTCCG-5'

DNA-protein complexes were resolved on a 6% non-denaturing gel. The gel was dried and exposed to film at -70 °C. For supershift experiments, 2-4 μg of nuclear extracts were incubated with antibodies to the different subunits of NF-κB for 20 minutes prior to the addition of the radiolabeled probe.

Immunoprecipitation

Whole cell lysates were prepared as described above. A mixture of Protein A agarose, ice-cold PBS and 1μg of anti-Bcl-3 or rabbit IgG was tumble incubated overnight at 4°C. Whole cell extracts were precleared with Protein A agarose for 30 minutes at 4°C followed by high speed

centrifugation. Precleared lysates were added to the protein A agarose/antibody mixture and incubated for 2 hours at 4°C. Complexes were washed with wash buffer (0.1% Triton X-100, 50 mM Tris pH 7.4, 300 mM NaCl, and 5 mM EDTA), resuspended in SDS-PAGE loading buffer, and examined by western blotting.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

We are grateful to the following individuals for their valuable technical assistance in this project: Ms. Marilyn Johnson, Ms. Carolyn Cooke, Dr. Shyam Dang, and Dr. Eugene Walch. This work was supported by NIH P50 CA093459 (DGU, EAG, JAE) and NIH R01 CA90282 (EAG). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the National Institutes of Health.

Abbreviations

EMSA, electrophoretic mobility shift assay; ERK, extracellular signal-regulated kinase; iNOS, inducible nitric oxide synthase; MAPK, mitogen-activated protein kinase; NF- κ B, nuclear factor-kappa B; NO, nitric oxide.

REFERENCES

- Baldwin AS. The NF- κ B and I κ B proteins: New discoveries and insights. *Ann Rev Immunol* 1996;14:649–681. [PubMed: 8717528]
- Cobbs CS, Brenman JE, Aldape KD, Bredt DS, Israel MA. Expression of nitric oxide synthase in human central nervous system tumors. *Cancer Res* 1995;55:727–730. [PubMed: 7531613]
- Cogswell PC, Guttridge DC, Funkhouser WK, Baldwin AS. Selective activation of NF- κ B subunits in human breast cancer: potential roles for NF- κ B/p52 and for Bcl-3. *Oncogene* 2000;19:1123–1131. [PubMed: 10713699]
- Dhawan P, Richmond A. A novel NF- κ B-inducing kinase-MAPK signaling pathway up-regulates NF- κ B activity in melanoma cells. *J Biol Chem* 2002;277:7920–7928. [PubMed: 11773061]
- Ellerhorst JA, Ekmekcioglu S, Johnson MK, Cooke CP, Johnson MM, Grimm EA. Regulation of iNOS by the p44/42 mitogen-activated protein kinase pathway in human melanoma. *Oncogene* 2006;25:3956–3962. [PubMed: 16474847]
- Ekmekcioglu S, Ellerhorst JA, Prieto VG, Johnson MM, Broemeling LD, Grimm EA. Tumor iNOS predicts poor survival for stage III melanoma patients. *Int J Cancer* 2006;119:861–866. [PubMed: 16557582]
- Fujita T, Nolan GP, Liou HC, Scott ML, Baltimore D. The candidate proto-oncogene *bcl-3* encodes a transcriptional coactivator that activates through NF- κ B p50 homodimers. *Genes & Dev* 1993;7:1354–1363. [PubMed: 8330739]
- Gilchrist M, McCauley SD, Befus AD. Expression, localization, and regulation of NOS in human mast cell lines: effects on leukotriene production. *Blood* 2004;104:462–469. [PubMed: 15044250]
- Giordano A, Tonello C, Bulbarelli A, Cozzi V, Cinti S, Carruba MO, et al. Evidence for a functional nitric oxide synthase system in brown adipocyte nucleus. *FEBS Lett* 2002;514:135–140. [PubMed: 11943139]
- Goydos JS, Mann B, Kim HJ, Gabriel EM, Alsina J, Germino FJ, et al. Detection of *B-RAF* and *N-RAS* mutations in human melanoma. *J Am Coll Surg* 2005;200:362–370. [PubMed: 15737846]
- Huang S, DeGuzman A, Bucana CD, Fidler IJ. Nuclear factor- κ B activity correlates with growth, angiogenesis, and metastasis of human melanoma cells in nude mice. *Clin Cancer Res* 2000;6:2573–2581. [PubMed: 10873114]
- Jemal A, Siegel R, Ward E, Murray T, Xu J, Thun MJ. Cancer statistics, 2007. *CA Cancer J Clin* 2007;57:43–66. [PubMed: 17237035]

- Jiang B, Xu S, Hou X, Pimentel DR, Brecher P, Cohen RA. Temporal control of NF- κ B activation by ERK differentially regulates interleukin-1 β -induced gene expression. *J Biol Chem* 2004;279:1323–1329. [PubMed: 14581482]
- Kang SM, Tran AC, Grilli M, Lenardo MJ. NF- κ B subunit regulation in nontransformed CD4⁺ T lymphocytes. *Science* 1992;256:1452–1456. [PubMed: 1604322]
- Kurland JF, Voehringer DW, Meyn RE. The MEK/ERK pathway acts upstream of NF- κ B1 (p50) homodimer activity and Bcl-2 expression in a murine B-cell lymphoma line. *J Biol Chem* 2003;34:32465–32470. [PubMed: 12801933]
- Mathas S, Johrens K, Joos S, Lietz A, Hummel F, Janz M, et al. Elevated NF- κ B p50 complex formation and Bcl-3 expression in classical Hodgkin, anaplastic large-cell, and other peripheral T-cell lymphomas. *Blood* 2005;106:4287–4293. [PubMed: 16123212]
- Ohno H, Takimoto G, McKeithan TW. The candidate proto-oncogene *bcl-3* is related to genes implicated in cell lineage determination and cell cycle control. *Cell* 1990;60:991–997. [PubMed: 2180580]
- Plaksin D, Baeuerle PA, Eisenbach L. KBF1 (p50 NF- κ B homodimer) acts as a repressor of H-2K^b gene expression in metastatic tumor cells. *J Exp Med* 1993;177:1651–1662. [PubMed: 8496683]
- Saini R, Patel S, Saluja R, Sahasrabudhe AA, Singh MP, Habib S, et al. Nitric oxide synthase localization in the rat neutrophils: immunocytochemical, molecular, and biochemical studies. *J Leukoc Biol* 2006;79:519–528. [PubMed: 16387842]
- Salvucci O, Carsana M, Bersani I, Tragni G, Anichini A. Antiapoptotic role of endogenous nitric oxide in human melanoma cells. *Cancer Res* 2001;61:318–326. [PubMed: 11196180]
- Smalley KSM. A pivotal role for ERK in the oncogenic behaviour of malignant melanoma? *Int J Cancer* 2003;104:527–532. [PubMed: 12594806]
- Tang CH, Grimm EA. Depletion of endogenous nitric oxide enhances cisplatin-induced apoptosis in a p53-dependent manner in melanoma cell lines. *J Biol Chem* 2004;279:288–298. [PubMed: 14576150]
- Taylor BS, de Vera ME, Ganster RW, Wang Q, Shapiro RA, Morris SM, et al. Multiple NF- κ B enhancer elements regulate cytokine induction of the human inducible nitric oxide synthase gene. *J Biol Chem* 1998;273:15148–15156. [PubMed: 9614127]
- Thomsen LL, Lawton FG, Knowles RG, Beesley JE, Riveros-Moreno V, Moncada S. Nitric oxide synthase activity in human gynecological cancer. *Cancer Res* 1994;54:1352–1354. [PubMed: 7509718]
- Thornburg NJ, Pathmanathan R, Raab-Traub N. Activation of nuclear factor- κ B p50 homodimer/Bcl-3 complexes in nasopharyngeal carcinoma. *Cancer Res* 2003;63:8293–8301. [PubMed: 14678988]
- Tian F, Zang WD, Hou WH, Liu HT, Xue LX. Nuclear factor- κ B signaling pathway constitutively activated in esophageal squamous cell carcinoma cell lines and inhibition of growth of cells by small interfering RNA. *Acta Biochim Biophys Sin* 2006;38:318–326. [PubMed: 16680372]
- Wang W, Abbruzzese JL, Evans DB, Larry L, Cleary KR, Chiao PJ. The nuclear factor- κ B relA transcription factor is constitutively activated in human pancreatic adenocarcinoma cells. *Clin Cancer Res* 1999;5:119–127. [PubMed: 9918209]
- Watanabe N, Iwamura T, Shinoda T, Fujita T. Regulation of NF- κ B1 proteins by the candidate oncoprotein BCL-3: generation of NF- κ B homodimers from the cytoplasmic pool of p50-p105 and nuclear translocation. *EMBO J* 1997;16:3609–3620. [PubMed: 9218802]




Figure 1. Expression of activated NF- κ B, activated ERK, and iNOS in melanoma cell lines and melanocytes. (a) Western blotting of nuclear extracts confirms the nuclear localization of pERK, and NF- κ B p50 and p65 in these cell lines. Histone serves as a loading control. (b) Western blot of whole cell lysates demonstrates the constitutive presence of iNOS in melanoma cells and melanocytes. iNOS is present in nuclear extracts as well. (c) Normalization of the immunoblotting results in (a) and (b) to the histone or actin bands is shown, to control for differences in loading and transfer. The iNOS chart refers to cytoplasmic levels. m: *BRAF* mutant cell lines; w: *BRAF* wild type cell lines.




Figure 2. Nuclear localization of NF- κ B p50 and p65. Immunofluorescent staining reveals the presence of the NF- κ B subunits in (a) MeWo and (b) HS294T cells.




Figure 3.

Comparison of the prevalences of NF- κ B p50 and p65 signals in the nuclei of melanoma cells and melanocytes. A significantly higher number of melanoma cells express p50 relative to p65. Error bars represent the mean \pm SD. *P-values were found to be statistically significant by chi-square analysis.




Figure 4. Melanoma NF-κB is regulated by the MAPK pathway. (A) EMSA was carried out with ^{32}P -labeled NF-κB target oligonucleotide and 4 μg of nuclear protein from unstimulated WM793 cells. Binding experiments were performed in the presence of a specific competitor (unlabeled NF-κB oligonucleotide), a non-specific competitor (SP1 target oligonucleotide), and in the absence of competitor. Results demonstrate NF-κB binding activity in these lysates. (B) Supershift experiments were conducted to confirm the specificity of NF-κB binding. Four micrograms of lysate were incubated with one microgram of antibody to p50 or p65, followed by EMSA analysis. (C) Nuclear extracts of WM793 cells treated with 10 μM U0126 were prepared and examined for NF-κB activity. Reversal of NF-κB binding is observed. Sp: specific.




Figure 5.

NF- κ B regulates iNOS expression in melanoma cells. MeWo cells (a) were treated with the NF- κ B Activation Inhibitor (10 μ M) and WM793 cells (b) with IKK-2 inhibitor V (1 μ M) over a time course of up to 48 hours. Western blotting was performed for iNOS protein and, in the case of IKK-2 Inhibitor V, for total and phosphorylated I κ B α .




Figure 6.

Melanoma cells express Bcl-3/p50 complexes. (a) Whole cell lysates and nuclear extracts from melanoma cells and melanocytes were used to carry out western blotting for Bcl-3. (b) Whole cell lysates from melanoma cells were used for immunoprecipitation with anti-Bcl-3 antibody followed by western blotting with anti-p50.