

ERRATUM

Effect of Sodium Chloride and pH on Enterotoxin B Production

CONSTANTIN GENIGEORGIS AND WALTER W. SADLER

Department of Public Health, School of Veterinary Medicine, University of California, Davis, California

Volume 92, no. 5, p. 1384: the following table which should have appeared on this page was omitted.

TABLE 2. Effect of pH and salt concentration on enterotoxin B production by strain S-6^a

Initial pH	Time	Salt concn (w/v)				
		2%	4%	6%	8%	10%
6.9	12 hr	13.5 ^b	8.7	2.8	0.2	—
	48 hr	15.3	13.4	10.5	3.4	—
	6 days	NM ^c	NM	NM	4.7	0.2
	10 days	NM	NM	NM	NM	NM
6.5	12 hr	13	8.7	1.9	—	—
	48 hr	14.9	13.4	10.1	1.9	—
	6 days	NM	NM	NM	2.2	—
	10 days	NM	NM	NM	NM	—
6.0	12 hr	9.8	7.5	1.4	—	—
	48 hr	13.9	12.5	8.7	0.9	—
	6 days	NM	NM	NM	1.4	—
	10 days	NM	NM	NM	NM	—
5.5	12 hr	0.6	0.2	—	—	—
	48 hr	3.0	1.4	0.9	—	—
	6 days	4.8	3.9	1.9	0.2	—
	10 days	NM	NM	NM	NM	—
5.1	12 hr	—	—	—	—	—
	48 hr	—	—	—	—	—
	6 days	0.2	—	—	—	—
	10 days	NM	0.2	—	—	—

^a Results were negative for 12, 14, and 16%.

^b Enterotoxin-antienterotoxin band migration (mm).

^c Not measured for width of the zone, but positive for enterotoxin.