

Published in final edited form as:


J Am Chem Soc. 2009 September 2; 131(34): 12068–12069. doi:10.1021/ja904930p.

Introduction of Cationic Charge into DNA Near the Major Groove Edge of a Guanine•Cytosine Base Pair: Characterization of Oligodeoxynucleotides Substituted with 7-Aminomethyl-7-Deaza-2'-Deoxyguanosine

 Manjori Ganguly^{†,‡,¶}, Ruo-Wen Wang^{†,‡,¶}, Luis A. Marky[‡], and Barry Gold^{*,†}
[†]University of Pittsburgh, Department of Pharmaceutical Sciences, 3501 Terrace St., Pittsburgh, PA 15261

[‡]University of Nebraska Medical Center, Department of Pharmaceutical Sciences, Omaha, NE 68198-6025

Abstract



The replacement of the 7-N atom on guanine (**G**) with a C-H to give 7-deazaguanine (**c⁷G**) alters the electronic properties of the heterocyclic base and eliminates a potential major groove cation binding site, which affects the organization of salts and water in the major groove. This has a destabilizing effect on DNA. We report herein the characterization of DNA oligomers containing 7-(aminomethyl)-7-deazaguanine (**1**) residues using a variety of spectroscopic and thermodynamic approaches. **1** is an intramolecular model for the major groove binding of cations and basic amino acid residues to **G**. In contrast to **c⁷G**, the tethering of a cation in the major groove using **1** affords

goldbi@pitt.edu.
[¶]contributed equally

Supporting Information. Thermodynamic and CD data are available free (Internet at <http://pubs.acs.org>).

DNA that is as, or more, stable than the corresponding unmodified DNA. The stabilization is associated with the folding enthalpy and hydration.

The electrostatics of DNA is generally considered in terms of polyelectrolyte theory that treats DNA as a polyanionic cylinder.¹ Even at relatively low salt concentrations, the cation concentration in the vicinity of DNA is estimated to be 1 M, and diffusible cations are treated without regard to high vs. low occupancy sites. Accordingly, the release of electrostricted ions, which is a powerful entropic driving force in the equilibrium binding of molecules to DNA,¹ is generally treated as non-sequence specific.

However, there are electrostatic factors associated with the DNA grooves, in addition to the phosphate backbone. NMR studies show that Mg^{2+} and related divalent cations preferentially bind in the major groove at G runs and appear to bend DNA.² A compilation of high-resolution structures of DNA shows a consensus for localization of monovalent cations in the major groove within 3.5 Å from the O⁶- and/or N7-positions of dG.³ The 1.2 Å resolution structure of 5'-d(CGCGAATTCGCG) in the presence of Tl^+ clearly indicates that there are high occupancy monovalent cation binding sites in the major groove near G/C regions as well as in the minor groove near A/T rich regions.⁴

In order to explore the role of groove associated cations on DNA stability, structure and reactivity we previously studied the effects of (i) removing a major groove cation binding site using 7-deazaguanine (e^7G)⁵ and (ii) introducing a tethered cation into the major groove using a 5-(3-aminopropyl)uracil modified base.⁶ The e^7G substitution results in lower thermodynamic stability that is mainly derived from a reduced enthalpy contribution.^{5a} A crystal structure of DNA with e^7G shows the predicted loss of a cation from a conserved cation binding site and the NMR indicates a destabilization at the flanking base pair.⁵ The flexible cationic 3-aminopropyl sidechain shows a complex sequence dependency in its impact on stability and structure that is related to its proximity to a major groove cation binding site.⁶

To create a structural model of groove associated cations that more accurately reflects cation distribution and location observed in DNA structures (Figure 1), we synthesized 7-aminomethyl-7-deaza-dG (**1**), which has a basic primary amine appended onto e^7G via a CH_2 linker.⁷ **1** recapitulates the location of inorganic cations that are observed near the major groove edge of G: the restricted $-NH_3^+$ is approximately 2.6 Å from the major groove face of the G (Figure 1), a distance that is also similar to that seen in crystal structures for cationic groups on basic amino acid residues of DNA binding proteins.⁸ We also prepared 7-hydroxymethyl-7-deaza-dG (**2**) as a neutral isostere of **1**.⁷ The spectroscopic and thermodynamic characterization of DNA containing e^7G , **1** and **2** are reported.

The sequences of the oligomers studied along with their thermodynamic parameters from UV melting and differential scanning calorimetry (DSC) analyses for unmodified (G), and e^7G , **1** and **2** substituted DNA are shown in Table 1. The sequence used (**OL-1**) was selected because it is self-complementary and forms an intermolecular duplex rather than an intramolecular hairpin. Similar to what was previously observed in another sequence,^{5a} the e^7G substitution at G-5 (**OL-2**) decreases stability relative to **OL-1**: $\Delta\Delta G^\circ = 2.5$ and 4.1 kcal/mol at 10 and 100 mM NaCl, respectively. In contrast, the substitution of **1** (**OL-3**) makes the DNA thermodynamically more stable than **OL-1**: $\Delta\Delta G^\circ -2.2$ and -1.4 kcal/mole at low and high salt, respectively. Relative to **OL-2**, the introduction of the tethered NH_3^+ increases stability by 4.7 and 5.5 kcal/mol at low and high salt, respectively. This translates to each $CH_2NH_3^+$ appendage increasing the stability by 2.3–2.7 kcal/mol. We attribute the stabilization predominantly to the tethered NH_3^+ ion since DNA modified with **2** (**OL-4**) is thermodynamically similar to **OL-2** with e^7G . A similar trend is observed when e^7G , **1** and **2** are introduced at G-3 (Supporting Information).

The origin of the differences in ΔG for the duplexes is derived from the ΔH term for the G-5 substitutions; c^7G in **OL-2** causes a marked reduction (21.9 and 24.8 kcal/mol at low and high salt, respectively) in enthalpic stabilization, while **1** in **OL-3** is enthalpy stabilizing by -14.7 and -7.7 kcal/mol at low and high salt, respectively. DNA with **2** (**OL-4**) behaves similar to the c^7G -modified DNA. At the 3-position in **OL-1**, the ΔH term for DNA modified with **1** is similar to unmodified DNA and remains higher than for duplexes with c^7G and **2**, particularly in 100 mM NaCl (Table 1 and Supporting Information).

The differences in the ΔH term can reflect changes in base stacking and/or hydration. Therefore, the effects of the different modifications on cation and water release upon unfolding were examined by T_M dependencies on salt concentration and osmolyte molality, respectively.⁹ There is a correlation between the ΔH term and DNA hydration. **OL-1** and **OL-3** have similar Δn_w values, which are significantly higher than observed for **OL-2** and **OL-4**. Interestingly, the release of cations upon unfolding varied little between **OL-1** and **OL-3**, but was significantly lower for **OL-2** and **OL-4**. (Table 1).

CD spectra provide global DNA conformational information of the different duplexes, and some indication of changes in base stacking by analysis of the relative intensities of the negative band at ~ 250 nm.¹⁰ The CD spectra of **OLs 1–12** are consistent with a B-conformation and there is a correlation between the intensity of the negative bands and the ΔH term at 10 mM NaCl that reflects base stacking for the oligomers (Supporting Information).

In addition to the intermolecular duplex DNA, an intramolecular hairpin with a higher T_M (**OL-5**) was modified with a c^7G (**OL-6**), or a single residue of **1**. The introduction of c^7G results in a minor destabilization (Table 1), while **OL-7** and **OL-8** are very similar to the unmodified hairpin. The addition of a second **1** (**OL-9**) places the two cations as close as 4.0 Å of each other assuming a normal B-conformation. This causes a modest reduction in the thermodynamic parameters. The minimum distance between the tethered NH_3^+ ions in **OL-3** is 8.3 Å. Of note, is that substitution of **1** (**OL-7** and **OL-8**) or two residues (**OL-9**) into the hairpin removed any T_M salt dependency; Δn_{Na^+} is 0.

In summary, the tethering of cationic and neutral polar functionalities at a distance from the floor of the major groove that mimics the distance observed for diffusible cations and basic amino acid residues is reported. In DNA with a c^7G substitution, a conserved cation binding is eliminated and the DNA is thermodynamically destabilized.⁵ In the current study, it is demonstrated that the insertion of **1** into DNA, which permanently refurbishes a cation that is lost with c^7G , increases the stability of DNA relative to that of the natural sequence. Locating a polar hydroxyl group on the major groove edge of c^7G does not restore stability. The effect of the tethered cationic modification provides an insight into the stabilizing role that “diffusible” cations associated with high occupancy sites in the major groove play in maintaining DNA structure.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

This work was supported by NIH grant RO1 CA29088. We thank C. Rizzo and A. Kozekova at Vanderbilt University for the synthesis of the DNA oligomers.

References

1. (a) Manning GS. *Quart. Rev. Biophys* 1978;2:179–246. (b) Record MT Jr, Anderson CF, Lohman TM. *Quart. Rev. Biophys* 1978;2:103–179. (c) Honig B, Nicholls A. *Science* 1995;268:1144–1149. [PubMed: 7761829]
2. (a) Braunlin WH, Nordenskiöld L, Drakenberg T. *Biopolymers* 1991;31:1343–1346. [PubMed: 1777584] (b) Buckin VA, Kankiya B, Renteperis D, Marky LA. *J. Am. Chem. Soc* 1994;116:9423–9429.
3. Howerton SB, Nagpal A, Williams LD. *Biopolymers* 2003;69:87–99. [PubMed: 12717724]
4. Howerton SB, Sines CC, VanDerveer D, Williams LD. *Biochemistry* 2001;40:10023–10031. [PubMed: 11513580]
5. (a) Ganguly M, Wang F, Kaushik M, Stone MP, Marky LA, Gold B. *Nucleic Acids Res* 2007;35:6181–6195. [PubMed: 17855404] (b) Wang F, Li F, Ganguly M, Marky LA, Gold B, Egli M, Stone MP. *Biochemistry* 2008;47:7147–7157. [PubMed: 18549246]
6. (a) Soto AM, Kankia BI, Dande P, Gold B, Marky LA. *Nucleic Acids Res* 2002;30:3171–3180. [PubMed: 12136099] (b) Li Z, Huang L, Dande P, Gold B, Stone MP. *J. Amer. Chem. Soc* 2002;124:8553–8560. [PubMed: 12121096] (c) Moulaei T, Maehigashi T, Lountos G, Komeda S, Watkins D, Stone M, Marky L, Li J-S, Gold B, Williams LD. *Biochemistry* 2004;43:7458–74648.
7. Wang R-W, Gold B. *Org. Lett* 2009;11:2465–2468. [PubMed: 19422208]
8. see Protein Data Bank (<http://www.rcsb.org/pdb>. e.g., PDB id: 1GU5, 1CF7, 1DP7, 1GCC, 3BPY, 1C7U).
9. (a) Spink CH, Chaires JB. *Biochemistry* 1999;38:496–508. [PubMed: 9890933] (b) Kaushik MN, Suehl N, Marky LA. *Biophys. Chem* 2007;126:154–164. [PubMed: 16822606]
10. (a) Tinoco, I.; Sauer, K.; Wang, J.C.; Puglisi, J. *Physical Chemistry: Principles and Applications in Biological Sciences*. Vol. 4th Ed.. Prentice Hall; 2001. (b) Basse WA, Johnson WC Jr. *Nucleic Acids Res* 1979;6:797–814. [PubMed: 424316] (c) Kypr J, Kejnovska´ I, Renc´iuk D, Michaela Vorli´c´kova´ M. *Nucleic Acids Res* 2009;37:1713–1725. [PubMed: 19190094]




Figure 1. Models of: (a) cation and (b) basic amino acid sidechain associated with major groove edge of G, (c) tethered $-NH_3^+$ in 7-(aminomethyl)-7-deazaguanine (**1**), (d) 7-(hydroxymethyl)-7-deazaguanine (**2**) and (e) 7-deazaguanine (**c⁷G**) lacking cation binding site.

Table 1

Thermodynamic Parameters for DNA Formation at 20 °C.^a

OL sequence	NaCl ^b	T _m ^c	ΔG ^{°d,e}	ΔH ^{°e}	TAS ^{°e}	ΔN _{Na} ^f	Δn _w ^f	ΔΔG vs OL-1 or -5
1 5'-GAGAGCGCTCTC	10	48.7	-6.9	-78.2	-71.3	-3.35 ± 0.17	-41 ± 3	-
	100	66.1	-12.b	-92.0	-79.b	-3.61 ± 0.18	-43 ± 4	
2 5'-GAGA-c ⁷ G-CGCTCTC	10	44.7	-4.4	-56.3	-51.9	-2.14 ± 0.11	-25 ± 2	2.5
	100	62.0	-8.4	-67.2	-58.8	-2.31 ± 0.12	-27 ± 2	4.1
3 5'-GAGA-1-CGCTCTC	10	52.0	-9.1	-92.9	-83.8	-2.86 ± 0.14	-38 ± 4	-2.2
	100	67.3	-13.9	-99.7	-85.9	-2.90 ± 0.15	-36 ± 3	-1.4
4 5'-GAGA-2-CGCTCTC	10	47.2	-4.6	-54.b	-49.9	-1.63 ± 0.10	-18 ± 2	2.3
	100	63.9	-7.6	-58.2	-50.6	-1.52 ± 0.10	-17 ± 2	4.9
5 5'-CGCGTTTCGGG	10	68.4	-4.4	-31.0	-26.6	-0.26 ± 0.02	-18 ± 2	-
6 5'-CGCGTTTC-c ⁷ G-CG	10	63.7	-3.b	-27.0	-23.5	-0.21 ± 0.02	-15 ± 2	0.9
7 5'-CGCGTTTC-1-CG	10	63.0	-4.0	-31.3	-27.3	0	-14 ± 2	0.4
8 5'-GGG-1-TTTTCGGG	10	71.3	-4.3	-28.9	-24.6	0	-16 ± 2	0.1
9 5'-CGG-1-TTTTC-1-CG	10	65.6	-3.6	-27.0	-23.4	0	-13 ± 1	0.8

^a Parameters are measured from UV (T_m) and DSC melting curves in 10 mM sodium phosphate buffer (pH 7.0). The observed standard deviations are: T_m (± 0.7), ΔH_{cal} (± 3%), ΔG[°]20 (± 5%), TAS_{cal} (± 3%).

^b Salt concentration in mM.

^c °C.

^d Determined at 20 °C.

^e kcal/mol.

^f per mol DNA