

Published in final edited form as:


Org Lett. 2008 October 2; 10(19): 4311–4314. doi:10.1021/ol8017566.

Efficient and Stereoselective Synthesis of Yellow Scale Pheromone via Alkyne Haloboration, Zr-Catalyzed Asymmetric Carboalumination of Alkenes (ZACA Reaction), and Pd-Catalyzed Tandem Negishi Coupling

Zhaoqing Xu and Ei-ichi Negishi*

Herbert C. Brown Laboratories of Chemistry, Purdue University, 560 Oval Drive, West Lafayette, Indiana 47907-2084

Abstract



A Pd-catalyzed reaction of allylzincs with the 1-octyne bromoboration product gives the desired allyl-alkenyl coupling products in good yields except with $\text{H}_2\text{C}=\text{CHCH}_2\text{ZnBr}$. This reaction is suitable for converting an alkyne bromoboration product **3** into **4** with no isomerization or β -elimination. The Pd-catalyzed isoalkyl-alkenyl coupling of **4** with the isoalkylzinc reagent derived from **2** provides yellow scale pheromone (**1**) of $\geq 98\%$ isomeric purity in 34% in six steps from TBDPS-protected homoallyl alcohol.

Hydrometalation and carbometalation of alkynes followed by Pd-catalyzed cross-coupling¹ have provided a variety of highly stereoselective and efficient routes to di- and trisubstituted alkenes.² Even so, however, there still are a large number of trisubstituted alkenes that are not readily preparable by these methods, as exemplified by the sex pheromone of yellow scale (**1**),³ a pest of citrus and ornamental plants.

It was isolated and identified without the establishment of full stereochemical details by Gieselmann and his co-workers in 1979.^{3a} In the same year, the active isomer of **1** was shown to be the *5E* isomer by a deliberate synthesis of a mixture of the *5E* and *5Z* isomers by Anderson and Henrick.^{3b} Mori and his co-workers subsequently established the *3S* configuration of the active isomer through asymmetric syntheses of both enantiomers.^{3d,e} Although highly stereoselective, their syntheses based on the chiral pool protocol suffered from a lengthy sequence of over 10 steps from methyl (*R*)-(+)-citronellate in 7-10% overall yields. A very

promising, convergent, but nonasymmetric route to **1** via silylcupration⁴ was devised more recently by Millar.^{3f}

The synthetically challenging trisubstituted alkene moiety of **1** and an opportunity for applying the ZACA reaction (Zrcatalyzed asymmetric carboalumination of alkenes),⁵ especially one-pot synthesis of (2*S*)-4-*tert*-butyldiphenylsilyloxy-2-methyl-1-butanol (**2**) of 98% ee in 52% yield from TBDPS-protected 3-buten-1-ol,^{5d} prompted us to devise an efficient and regio- and stereoselective synthesis of **1**, as outlined in Scheme 1.

For construction of the trisubstituted alkene moiety, application of alkyne haloboration developed by Suzuki⁷ was considered, and stereoselective formation of **3** in ca. 80% yield by the reaction of 3-methyl-1-butyne with BBr₃ in CH₂Cl₂ was observed. This was to be followed by selective and high-yielding cross-coupling. The high propensity of the bromoboration products, such as **3**, to undergo unwanted dehaloboration together with the modest reactivity of hindered alkenyl bromide would make desired C-C bond formation via cross-coupling of **3** challenging. In fact, the only known satisfactory cross-coupling appeared to be the Pd-catalyzed organozinc reaction² employed by Suzuki.⁷ It then occurred to us that, whereas the Pd-catalyzed alkenylmetal-allyl electrophile (alkenyl-allyl, hereafter) coupling had been extensively investigated, generally high-yielding, and often highly regio- and stereospecific with retention with respect to both alkenyl and allyl groups,⁸ little had been known about the corresponding allylmetal-alkenyl electrophile (allyl-alkenyl, hereafter) coupling.^{8b} Since the Pd-catalyzed alkenyl-allyl coupling was not a viable option for the task at hand, however, the Pd-catalyzed allyl-alkenyl coupling of (*Z*)-(2-bromo-1-octenyl)dibromoborane, generated by treating 1-octyne with BBr₃ in CH₂Cl₂, with allylzinc bromide and its variously carbon-substituted derivatives generated by treating the corresponding allylic bromides with Zn dust was examined.⁹ The results summarized in Table 1 indicate the following: (1) The Pd-catalyzed allyl-alkenyl coupling with the parent allylzinc bromide does not occur under the conditions used, but all of the other cases including those involving 2-methylallylzinc bromide gave the desired products in good yields. The sterically least demanding parent allylzinc bromide may act as a catalyst poison, although this point needs to be further investigated. (2) As anticipated, the Pd-catalyzed allyl-alkenyl coupling is more readily prone to stereoisomerization. Thus, the allylic organozinc reagents derived from both geranyl and neryl bromides led to the formation of essentially identical mixtures of the 4*E* and 4*Z* isomers (*E/Z*) 2.5/1) without producing detectable amounts of regioisomers. These results are in sharp contrast with the corresponding Pd-catalyzed alkenyl-allyl coupling proceeding with essentially complete retention of both stereo- and regiochemical identities.⁸ Despite these limitations, the desired case of the conversion of **3** to **4** via Pd-catalyzed cross-coupling of **3** with 3-methyl-2-butenylzinc bromide in the presence of 1 mol % of Pd(PPh₃)₂Cl₂ followed by iodinolysis with I₂ (2 equiv) and NaOAc (1.5 equiv) proceeded cleanly with no detectable sign of isomerization to give **4** (≥98% *E*) in 77% yield from 3-methyl-1-butyne.

For the final assembly of the full carbon skeleton of the yellow scale pheromone (**1**), **2** was iodinated in 97% yield by its treatment with I₂ (1.2 equiv), PPh₃ (1.15 equiv), and imidazole (1.3 equiv) in CH₂Cl₂. The iodide thus obtained was treated with *t*-BuLi (2.2 equiv) in ether¹⁰ followed by zincation with dry ZnBr₂. The Pd-catalyzed isoalkyl-alkenyl coupling of the in situ generated isoalkylzinc reagent with **4** in the presence of 1 mol % of Pd(PPh₃)₄ followed by desilylation with TBAF (1.2 equiv), provided **5** as a ≥98% pure compound in 74% yield from the two iodide precursors,¹¹ and its acetylation gave the desired yellow scale pheromone (**1**) of ≥98% isomeric purity in 92% yield. Thus, **1** was synthesized in 34% yield over six steps in the longest linear sequence from TBDPS-protected 3-buten-1-ol or in 54% yield over four steps from 3-methyl-1-butyne.

In summary, the following notable findings have been discovered in this work. (1) The hitherto essentially unexplored Pd-catalyzed allylzinc-alkenyl electrophile coupling has been investigated. The parent allylzinc bromide does not give the desired allyl-alkenyl coupling products under the conditions used. Fortunately, all of the other allylzinc derivatives with one or two substituents in the C2 or C3 position have smoothly reacted to give the desired allyl-alkenyl coupling products in good yields. As anticipated, however, this allyl-alkenyl coupling has proven to be more readily prone to regio- and/or stereoscrumbling than the previously developed alkenyl-allyl coupling.⁸ (2) The Pd-catalyzed allyl-alkenyl coupling proved to be exactly what was needed for converting the alkyne bromoboration product **3** into one of the key intermediates **4** for the synthesis of yellow scale pheromone (**1**), which proceeded in 77% yield without being accompanied by any isomerization or unwanted debromoboration of **3** to detectable extents. (3) The other key intermediate **2** was prepared in 52% yield as a 99% pure *R* isomer via the ZACA reaction of *t*-BuPh₂Si-protected homoallyl alcohol followed by enantiomeric purification by Amano PS lipase-catalyzed acetylation of the unwanted *S* isomer. (4) Iodination of **2** in 97% yield was followed by its in situ lithiation-zincation and Pd-catalyzed cross-coupling of the isoalkylzinc reagent thus generated with the alkenyl iodide **4** cleanly produced isomerically pure ($\geq 98\%$) **5** in 74% yield from the two iodide precursors. Acetylation of **5** produced the sex pheromone of yellow scale (**1**) of $\geq 98\%$ isomeric purity even before purification in 34% yield from *t*-BuPh₂Si-protected homoallyl alcohol in six steps or in 54% yield from 3-methyl-1-butyne in four steps.

The results presented herein provide yet another example demonstrating the synthetic utility of the ZACA-Pd catalyzed cross-coupling synergy,¹² which, in this case, is further promoted by alkyne haloboration.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


Acknowledgments

We thank the National Institutes of Health (GM 36792) and Purdue University for support of this research. We also thank Albemarle, Wako Chemical Co., and NICHIA Corp. for their support and Dr. Z. Huang of our laboratories for technical assistance.

References

1. For seminal contributions, see: (a) Negishi E, Baba S. *J. Chem. Soc., Chem. Commun* 1976;596:597.. (b) Baba S, Negishi E. *J. Am. Chem. Soc* 1976;98:6729–6731.. (c) Negishi E, Van Horn DE. *J. Am. Chem. Soc* 1977;99:3168–3170.. (d) Okukado N, Van Horn DE, Klima WL, Negishi E. *Tetrahedron Lett* 1978;1027:1030.. (e) Negishi E, Okukado N, King AO, Van Horn DE, Spiegel BI. *J. Am. Chem. Soc* 1978;100:2254–2256.. (f) Miyaura N, Yamada K, Suzuki A. *Tetrahedron Lett* 1979;3437:3440.
2. For recent reviews, see: (a) Negishi, E., editor. *Handbook of Organopalladium Chemistry for Organic Synthesis*. Wiley-Interscience; New York: 2002. p. 3279 Chapters III.2.6, III.2.9, III.2.11, III.2.15. (b) de Meijere, A.; Diederich, F., editors. *Metal-catalyzed Cross-Coupling Reactions*. Vol. 2004. Wiley-VCH; Weinheim: p. 916. (c) Negishi E, Hu Q, Huang Z, Qian M, Wang G. *Aldrichim. Acta* 2005;38:71–88.. (d) Negishi E, Huang Z, Wang G, Mohan S, Wang C, Hattori H. *Acc. Chem. Res.* in press
3. (a) Gieselmann MJ, Moreno DS, Fargerleund J, Tashiro H, Roelofs WL. *J. Chem. Ecol* 1979;5:25–31. (b) Anderson RJ, Henrick CA. *J. Chem. Ecol* 1979;5:773–779. (c) Suguro T, Roelofs WL, Mori K. *Agric. Biol. Chem* 1981;45:2509–2514. (d) Masuda S, Kuwahara S, Suguro T, Mori K. *Agric. Biol. Chem* 1981;45:2515–2520. (e) Mori K, Kuwahara S. *Tetrahedron* 1982;38:521–525. (f) Millar JG. *Tetrahedron Lett* 1989;30:4913–4914.
4. For a seminal paper on silylcupration, see: Fleming I, Newton TW, Roessler F. *J. Chem. Soc., Perkin Trans. 1* 1981;252:7–2532.



5. For seminal and pertinent papers on the ZACA reaction, see: (a) Kondakov D, Negishi E. J. Am. Chem. Soc 1995;117:10771–10772.. (b) Kondakov D, Negishi E. J. Am. Chem. Soc 1996;118:1577–1578.. (c) Huo S, Shi J, Negishi E. Angew. Chem., Int. Ed 2002;41:2141–2143.. (d) Huang Z, Tan Z, Novak T, Zhu G, Negishi E. Adv. Synth. Catal 2007;349:539–545.
6. Erker G, Aulbach M, Knickmeier M, Wingbermuhle D, Kruger C, Nolte M, Werner S. J. Am. Chem. Soc 1993;115:4590.
7. Satoh Y, Serizawa H, Miyaura N, Hara S, Suzuzki A. Tetrahedron Lett 1988;29:1811–1814. For a seminal report on alkyne haloboration, see: Lappert MF, Prokai B. J. Organomet. Chem 1964;1:384–400.
8. Matsushita H, Negishi E. J. Am. Chem. Soc 1981;103:2882–2884. For a review, see: Negishi E, Liu F. Chapter III.2.9 in ref 2a
9. (a) Shriner, RL.; Neumann, FW. Organic Synthesis. Vol. Vol. III. Wiley; New York: 1955. p. 73 (b) Negishi E, Matsushita H, Okukado N. Tetrahedron Lett 1981;22:2715–2718.
10. (a) Negishi E, Swanson DR, Rousset CJ. J. Org. Chem 1990;55:5406–5409. (b) Bailey WF, Punzalan ER. J. Org. Chem 1990;55:5404–5406.
11. For conceptually related applications of tandem Negishi coupling processes to the synthesis of allenynes and enediynes, see: (a) Wang KK, Wang Z. Tetrahedron Lett 1994;35:1829–1832.. (b) Wang KK, Wang Z, Tarli A, Gannett P. J. Am. Chem. Soc 1996;118:10783–10791.. (c) Tarli A, Wang KK. J. Org. Chem 1997;62:8841–8847.
12. (a) Negishi E, Tan Z, Liang B, Novak T. Proc. Nat. Acad. Sci. U.S.A 2004;101:5782–5787. (b) Tan Z, Negishi E. Angew. Chem., Int. Ed 2004;43:2911–2914. (c) Magnin-Lachaux M, Tan Z, Liang B, Negishi E. Org. Lett 2004;6:1425–1427. [PubMed: 15101758] (d) Novak T, Tan Z, Liang B, Negishi E. J. Am. Chem. Soc 2005;127:2838–2839. [PubMed: 15740104] (e) Liang B, Novak T, Tan Z, Negishi E. J. Am. Chem. Soc 2006;128:2770–2771. [PubMed: 16506734] (f) Zhu G, Negishi E. Org. Lett 2007;9:2771–2774. [PubMed: 17583343] (g) Zhu G, Liang B, Negishi E. Org. Lett 2008;10:1099–1101. [PubMed: 18275210]



**Scheme 1.**


Regio- and Stereoselective Synthesis of Yellow Scale Pheromone

Table 1

Scope of a Tandem Process Consisting of Alkyne Haloboration-Pd-Catalyzed Negishi Coupling with Allylzinc Bromide^a












5 Pd(PPh₃)₂Cl₂







^a All reactions were run with 1.0 equiv of BB_r3, 1.2 equiv of allylzinc bromide generated by treating the corresponding allylic bromides with Zn dust, 2.0 equiv of I₂, and 1.5 equiv of NaOAc.

^b All yields refer to isolated yields.

^c No allyl-alkenyl coupling product was observable by GC and ¹H NMR.