

Short report

Open Access

In vivo evidence that truncated *trkB.T1* participates in nociception

Cynthia L Renn¹, Carmen C Leitch¹ and Susan G Dorsey^{*1,2,3}

Address: ¹School of Nursing, University of Maryland, Baltimore, MD 21201, USA, ²Program in Neuroscience, University of Maryland, Baltimore, MD 21201, USA and ³Program in Oncology, University of Maryland, Baltimore, MD 21201, USA

Email: Cynthia L Renn - renn@son.umaryland.edu; Carmen C Leitch - cleit001@son.umaryland.edu; Susan G Dorsey* - sdorsey@son.umaryland.edu

* Corresponding author

Published: 29 October 2009

Received: 5 August 2009

Molecular Pain 2009, **5**:61 doi:10.1186/1744-8069-5-61

Accepted: 29 October 2009

This article is available from: <http://www.molecularpain.com/content/5/1/61>

© 2009 Renn et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Brain-Derived Neurotrophic Factor (BDNF) is a central nervous system modulator of nociception. In animal models of chronic pain, BDNF exerts its effects on nociceptive processing by binding to the full-length receptor tropomyosin-related kinase B (*trkB.FL*) and transducing intracellular signaling to produce nocifensive behaviors. In addition to *trkB.FL*, the *trkB* locus also produces a widely-expressed alternatively-spliced truncated isoform, *trkB.T1*. *TrkB.T1* binds BDNF with high affinity; however the unique 11 amino acid intracellular cytoplasmic tail lacks the kinase domain of *trkB.FL*. Recently, *trkB.T1* was shown to be specifically up-regulated in a model of HIV-associated neuropathic pain, potentially implicating *trkB.T1* as a modulator of nociception. Here, we report that *trkB.T1* mRNA and protein is up-regulated in the spinal dorsal horn at times following antiretroviral drug treatment and hind paw inflammation in which nocifensive behaviors develop. While genetic depletion of *trkB.T1* did not affect baseline mechanical and thermal thresholds, the absence of *trkB.T1* resulted in significant attenuation of inflammation- and antiretroviral-induced nocifensive behaviors. Our results suggest that *trkB.T1* up-regulation following antiretroviral treatment and tissue inflammation participates in the development and maintenance of nocifensive behavior and may represent a novel therapeutic target for pain treatment.

Findings

BDNF is a potent modulator of pain processing in the CNS [1,2]; for review see [3,4]. BDNF exerts effects on nociception by binding to *trkB.FL* and initiating intracellular signaling cascades that lead to transcriptional changes. Noxious stimulation increases BDNF and *trkB.FL* production in the spinal dorsal horn [5-9] and brainstem [10,11], leading to hyperalgesia and the formation of nocifensive behaviors. Interfering with *trkB.FL* signaling via *trkB.FL*-specific exogenous antibody administration [12] or by blocking receptor activation in the chemical-genetic transgenic *trkB^{FG16A}* mice [13] prevents the devel-

opment of tissue- and nerve injury-induced thermal and mechanical hypersensitivity [14].

In addition to *trkB.FL*, the *Ntrk2* (*trkB*) locus encodes for several alternatively-spliced isoforms of the receptor [15], including *trkB.T1*, the predominant isoform expressed in the adult mammalian nervous system. The extracellular domain of *trkB.T1* is identical to the full-length isoform, which enables high-affinity BDNF binding [15,16]. However, the 11 amino acid intracellular portion of *trkB.T1* lacks the kinase activation domain necessary to activate classical signal transduction pathways. Since *trkB.T1* het-

erodimerization with *trkB.FL* inhibits trans-autophosphorylation of the *trkB.FL* kinase domain, studies support a model in which *trkB.T1* functions to reduce BDNF signaling [17-22]. However, we [23,24] and others [25-31] have demonstrated evidence that *trkB.T1* may signal independently.

TrkB.T1 is up-regulated in pathological states including human Alzheimer's disease [32] and it is the mechanism underlying premature hippocampal cell death in a mouse model of Down Syndrome [23,33]. Moreover, altered *trkB.T1* expression can affect synaptic plasticity. Overexpression of *trkB.T1* in hippocampal neurons leads to the inhibition of synaptic potentiation [34] and overexpressing *trkB.T1* at neuromuscular synapses results in disassembly of acetylcholine receptor clustering [18]. Recently, *trkB.T1* specifically was shown to be significantly up-regulated in dorsal root ganglion neurons in a rodent model of HIV-associated neuropathic pain [35]. Thus, it is reasonable to hypothesize that *trkB.T1* up-regulation might have a specific role in pain processing.

First, we examined *trkB.T1* mRNA expression in the spinal dorsal horn of a mouse model of antiretroviral-associated neuropathic pain [36], a model in which significant mechanical hypersensitivity occurs one day after tail vein injection with stavudine (d4T) [37]. Since behavioral changes occur soon after treatment [37], we examined early time points and found that *trkB.T1* mRNA was significantly up-regulated at 3 hours after tail vein injection of d4T compared with mice injected with an equivalent volume of saline vehicle (Figure 1A). At later time points (6 h, 12 h, 18 h, 3 d) *trkB.T1* mRNA levels returned to saline control levels (data not shown). Next, we asked whether *trkB.T1* protein was regulated in the dorsal horn after treatment. In Figure 1B, we demonstrate that *trkB.T1* protein is significantly up-regulated one day after d4T injection compared with saline control-injected animals. By three days after d4T treatment, *trkB.T1* levels returned to saline control levels, a result consistent with our mRNA data. We next asked whether *trkB.T1* was up-regulated in the dorsal horn of mice with hind paw inflammation. As demonstrated in Figures 1C and 1D, both *trkB.T1* mRNA and protein are significantly up-regulated in CFA-treated mice compared with mice receiving a saline control-injected hind paw. These results raise the possibility that the *trkB.T1* receptor is involved in nociception in these two pain models.

If *trkB.T1* is up-regulated in the dorsal horn following noxious stimulation, then genetic depletion of *trkB.T1* [23] would be expected to mitigate nocifensive behaviors in these models. We first tested this hypothesis in the antiretroviral model (i.e. stavudine) of mechanical hypersensitivity [37]. There were no differences in mechanical

threshold (g) in naïve *trkB.T1* null mice at baseline (Figure 2A). Genetic deletion of *trkB.T1* provided significant attenuation of mechanical hypersensitivity throughout the testing period (Figure 2A). Similarly, in the well-established CFA model of inflammation although we found no difference in thermal paw withdrawal latency (s) in *trkB.T1* KO mice at baseline, the absence of *trkB.T1* in vivo significantly attenuated thermal hypersensitivity (Figure 2B). In Figure 2C, we show that there were no differences in the degree of hind paw thickness secondary to CFA injection. As this is an indirect measure of inflammation we conclude that there was no difference in the inflammatory response in the *trkB.T1* knockout mice. Next, we asked whether genetic deletion of *trkB.T1* would provide a benefit for mice treated with capsaicin, an agonist of TRPV1 receptors that are expressed in BDNF-immunoreactive nociceptors. Similar to results in our previous two models, the absence of *trkB.T1* provided significant attenuation of the latency to lick, number of licking bouts and duration of licking after capsaicin treatment. Thus, we conclude that the absence of *trkB.T1* provides significant protection against the development of thermal and mechanical hypersensitivity.

BDNF signaling modulates nociceptive processing across a variety of pre-clinical models of pain (for review, [38]). Most studies have focused on *trkB.FL* as the receptor responsible for BDNF-mediated nociception, since antibodies specific for *trkB.FL* [12] and abrogation of *trkB.FL* activation using the *trkB^{FG616A}* transgenic mouse [14] attenuate thermal and mechanical hypersensitivity. Few studies, however, have examined a specific role for *trkB.T1*. Yajima et al. [12] concluded that *trkB.T1* did not have a role in nociception since repeated intrathecal injections with anti-*trkB.T1* did not produce changes in nerve injury-induced thermal hyperalgesia. However, we (Dorsey and Renn, unpublished) and others (Dr. Rita Balice-Gordon, University of Pennsylvania, personal communication) have shown that the *trkB.T1* antibody used in the Yajima et al. [12] study cross-reacts with other epitopes *in vivo*. Moreover, as acknowledged in Wang et al. [14], indirect methods of targeting receptor activation, for example via exogenous antibody administration, can be inconclusive. In support of a role for *trkB.T1* in nociception, a recent microarray study in a model of HIV-associated neuropathic pain found that *trkB.T1*, but not *trkB.FL* or *trkB.T2* (a second truncated isoform), was up-regulated in dorsal root ganglion neurons [35].

Since *trkB.T1* and *trkB.FL* are presumably regulated by a common promoter, we also undertook a careful examination of spinal dorsal horn *trkB.FL* mRNA and protein regulation in the antiretroviral and inflammatory models in both wildtype and *trkB.T1* knockout animals. As demonstrated in Additional File 1, we detected no significant reg-




Figure 1 (see legend on next page)

Figure 1 (see previous page)

TrkB.T1 expression in the spinal dorsal horn of d4T and CFA-treated mice **A.** Quantification of trkB.T1 mRNA expression in the dorsal horn of wildtype mice 3 hours after receiving a saline tail injection (black bar; n = 6) or an injection of d4T [50 mg/kg] (blue bar; n = 6). **B.** TrkB.T1 protein expression in the dorsal horn in the antiretroviral model was assayed via western blot. The top panel shows a representative western blot of trkB.T1 following saline or d4T injection at 1d and 3d. Below the panel the results from saline treated (black bar; n = 6) and 1d (blue bar; n = 6) or 3d (red bar; n = 6) after d4T treatment are quantified. *indicates $p < 0.05$ by ANOVA with Tukey post-hoc testing. **C.** Quantification of trkB.T1 mRNA expression in the dorsal horn of wildtype mice 3 days after receiving a saline hind paw injection (black bar; n = 6) or an injection of CFA (blue bar; n = 6). **D.** TrkB.T1 protein expression in the dorsal horn in the CFA model was assayed via western blot. The top panel shows a representative western blot of trkB.T1 following a saline or CFA hind paw injection at 3d. Below the panel the results from saline treated (black bar; n = 6) and 3d CFA (blue bar; n = 6) are quantified. *indicates $p < 0.05$ by t-test.

ulation of trkB.FL mRNA (panels A-D) following antiretroviral or inflammatory treatment compared with vehicle controls in either genotype. Similar results were found for protein levels (panels E-H), although trkB.FL was down-regulated approximately 25% in wildtype CFA-treated animals. Thus, we conclude that the up-regulation of trkB.T1 following noxious stimulation is specific for this isoform.

Here, we demonstrate that trkB.T1 mRNA and protein are both significantly up-regulated in the spinal dorsal horn following systemic antiretroviral drug administration and hind paw inflammation (Figure 1). We show that *in vivo* deletion of trkB.T1 did not affect acute nociceptive processing, since baseline mechanical and thermal thresholds were not significantly different from wildtype strain-matched controls (Figure 2). However, following antiretroviral administration or hind paw inflammation via CFA injection or capsaicin treatment, animals lacking trkB.T1 were significantly protected against the development of mechanical and thermal hypersensitivity (Figure 2A, B, D). These results suggest that trkB.T1 participates in the development and maintenance of mechanical and thermal hypersensitivity, and that upregulation specifically contributes to the degree of thermal and mechanical hypersensitivity since the null mutant demonstrates attenuated hypersensitivity to our noxious stimulation paradigms.

The expression of trkB.FL and trkB.T1 is conserved across species and throughout evolution, and trkB isoform expression is regulated during development when the dominant trkB isoform expression in the brain switches from trkB.FL to trkB.T1 [39]. Receptor expression is cell specific and not all cells express both isoforms. For example, neurons express both isoforms [33] while astrocytes only express trkB.T1 [23,40]. The prevailing view has been that trkB.T1 expression leads to dominant negative inhibition of full-length trkB signaling and a reduction in the activation of downstream signaling molecules [17-23,33]. If trkB.T1 were functioning solely as a dominant negative inhibitor of signaling in the dorsal horn, the prediction

would be that deletion of trkB.T1 would result in more, not less pain. However, in the animal models of pain tested in this study, genetic depletion of trkB.T1 results in attenuation of pain. Several studies have provided evidence to suggest that trkB.T1 can signal independently, however the precise nature and functional relevance of this signal has been less clear. The available evidence regarding trkB.T1 signaling suggests that BDNF binding to trkB.T1 causes increased intracellular calcium release [28]. We have shown that neurons from a mouse model of neurodegeneration over-express trkB.T1, resulting in a significant increase in resting intracellular calcium; reducing trkB.T1 expression in those neurons reduced intracellular calcium to wildtype levels [23]. In a rodent model of antiretroviral-induced neuropathy, intrathecal injection of TMB-8, a drug that buffers calcium, provided near complete abrogation of mechanical hypersensitivity [36]. Thus, our results demonstrating attenuation in thermal and mechanical sensitivity in the absence of trkB.T1 could be because increased trkB.T1 expression induces increased intracellular calcium signaling which promotes hypersensitivity.

Our results support the idea that trkB signaling is complex, and that trkB.T1 participates in the development and maintenance of persistent pain. Genetic deletion of trkB.T1 provides significant protection from the development of thermal and mechanical hypersensitivity following noxious stimulation. Thus, we conclude that while previous studies have demonstrated a specific role for BDNF-mediated trkB.FL signaling in nociception, our results support a role for trkB.T1 in nociception that has not been previously described.

Materials and methods

Western blots

Procedures have been described elsewhere [23,33]. Primary antibodies included anti-trkB.T1 (Santa Cruz TrkB (C-13) sc-119), anti-trkB out (generous gift from David Kaplan, Hospital for Sick Kids, Toronto, Canada) and anti-actin (Sigma A2547). Secondary antibodies conju-




Figure 2 (see legend on next page)

Figure 2 (see previous page)

Nocifensive behavior in *trkB.T1* wildtype and knockout mice. **A.** Mice were baseline tested for mechanical threshold and then randomized within genotype (wildtype, knockout; n = 12 per genotype) to receive a saline (n = 6 per genotype) or d4T (n = 6 per genotype) tail vein injection. Following tail vein injection, mice were tested at 1, 7, 14, 21 and 28d for the presence of mechanical sensitivity. * indicates T1 WT (blue circles) and T1 KO (green triangles) significantly different from saline controls; p < 0.05 by Repeated Measures ANOVA with Tukey post-hoc testing. **B.** Mice were baseline tested for thermal threshold and then randomized within genotype (wildtype, knockout; n = 12 per genotype); to receive saline (n = 6 per genotype) or CFA (n = 6 per genotype) hind paw injection. Following hind paw injection, mice were testing for thermal sensitivity at 3 h and 1, 3, 7, 14, 21, 28 and 35d.* indicates T1 WT (blue circles) and T1 KO (green triangles) significantly different than saline controls; p < 0.05 by Repeated Measures ANOVA with Tukey post-hoc testing. **C.** The hind paws from mice in each group tested in B. were measured using calipers and the thickness recorded in mm. **D.** Latency to licking, number of licking bouts and duration of licking following capsaicin administration were recorded and the mean and s.e.m. presented for each genotype (n = 6). *indicates p < 0.05 by t-test.

gated to peroxidase included anti-mouse IgG and anti-rabbit IgG (Amersham Pharmacia Biotech).

qPCR

Methods for RNA extraction and qPCR techniques have been previously described (Dorsey et al., 2009). Primers specific for *trkB.T1* included forward: 5'-TGG TGA TGT TGC TCC TGC TCA AGT-3'; reverse: 5'-CCC ATC CAG TGG GAT CTT ATG AAA C-3'. Primers specific for *trkB.FL* included forward: 5'-ACT TTG GCA TCA CCA ACA GTC AGC-3'; reverse: 5'-AGG TTG TAG CAC TCG GCA AGG AAA-3'. The β -actin gene was used as the reference gene with specific primers: forward: 5'-TGT GGT GCC AGA TCT TCT CCA TGT-3'; reverse: 5'-TGT GGT GCC AGA TCT TCT CCA TCT-3' (Integrated DNA Technologies).

Generation of *trkB.T1* wildtype and knockout mice

Mice were generated as previously described [23]. Wildtype and knockout experimental animals were the progeny of heterozygous *trkB.T1* matings from animals at N13 generation.

Nocifensive behavior

All experiments were approved by the University of Maryland Baltimore Institutional Animal Care and Use Committee and were performed following the guidelines of the International Association for the Study of Pain. All experiments were performed with the tester blind to the genotype of the mice and the treatment that they received. All mice tested were adult male mice (2-6 months of age weighing approximately 30 g. Details regarding the production of the inflammatory model using Complete Freund's adjuvant (CFA) can be found in Renn et al. [11] and the antiretroviral model methods are detailed in Dorsey et al. [37]. For capsaicin experiments, capsaicin (Sigma, St. Louis, MO) was dissolved in 100% dimethylsulfoxide (DMSO) followed by dilution with 0.9% saline to a concentration of 0.08 $\mu\text{g}/\mu\text{l}$. Using a 50 μl Hamilton syringe with a 1/2" 30-gauge needle attached, each mouse was given a 20 μl subcutaneous injection of capsaicin (1.6 $\mu\text{g}/$

20 μl) into the plantar surface of the left hind paw. The mice were placed in individual Plexiglas cubicles and allowed to acclimate for approximately one hour before the injection of capsaicin. Licking induced by the capsaicin injection was used to indicate a nocifensive response. Each mouse was examined for a period of 5 minutes following the capsaicin injection by a blinded observer. The time (in seconds) to onset of licking, total accumulated time (in seconds) spent licking and the number of bouts of licking the capsaicin-injected paw was measured. Thermal Hyperalgesia: The Paw Thermal Stimulator (PTS, UARDG, Department of Anesthesiology, University of California, San Diego; attn George Ozaki) was used to assess thermal sensitivity. Mice were placed in individual Plexiglas cubicles on a glass surface maintained at a constant temperature of $30^{\circ}\text{C} \pm 0.1^{\circ}\text{C}$, and allowed to acclimate for approximately one hour prior to testing. Paw withdrawal latencies were defined as the time in seconds required from the onset of the stimulus to the point of a brisk withdrawal of the hind paw from the stimulus. An automatic 20-second cut-off was used to avoid tissue injury from prolonged exposure to the thermal stimulus. Each hind paw was tested three times and the response latency was the average of the three tests. Approximately 10 minutes elapsed between repeat test exposures of each hind paw to avoid tissue injury from the repeated applications of the thermal stimulus. Mechanical Allodynia: The nocifensive behavior of paw withdrawal from a mechanical stimulus was used to assess the development of mechanical allodynia. The mice were placed in individual Plexiglas cubicles on a wire mesh platform, and allowed to acclimate for approximately one hour. A series of von Frey filaments (Touch Test Sensory Evaluator Kit, myNeuroLab.com, St. Louis, MO), with bending forces that ranged from 0.04 g to 1.40 g, was used to deliver the mechanical stimuli. Each filament was applied to the left hind paw until the filament just bent and was held in place for 5 seconds or until the mouse withdrew his paw. Each filament was tested 5 times on the left hind paw, with a period of 3-5 minutes elapsing between trials. A

positive response to the stimulus was defined as a brisk withdrawal, with or without shaking or licking, of the hind paw during or immediately upon removal of the filament application. Using a procedure modified after Ren [41], the mechanical stimuli were applied to the plantar surface of the left hind paw, starting with the 0.4 g filament. If the 0.4 g filament elicited 3 positive responses out of 5 trials, then the mouse was tested moving downward through the series to the 0.04 g filament and the number of withdrawals was recorded for each filament. If the 0.4 g filament did not elicit 3 positive responses, then the mouse was tested moving upward through the series to the 1.4 g filament and the number of withdrawals was recorded for each filament. Threshold was defined as the filament with the lowest bending force that elicited at least 3 positive responses out of 5 trials.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

CLR assisted with the design of experiments, conducted the behavioral experiments and assisted with data analysis; CCL conducted the qPCR and western blot experiments; SGD designed and supervised the experiments, analyzed the data and wrote the manuscript. All authors have read and approved the final manuscript.

Additional material

Additional file 1

trkB.FL mRNA and protein expression in the antiretroviral and inflammatory models in trkB.T1 WT and KO animals. A and B. Quantification of trkB.FL mRNA expression in the dorsal horn of trkB.T1 wildtype mice (A) and trkB.T1 knockout mice (B) after receiving a saline tail injection (black bar; n = 3) or 1d (blue bar; n = 3) or 3d (red bar; n = 3) after an injection of d4T [50 mg/kg]. C and D. Quantification of trkB.FL mRNA expression in the dorsal horn of trkB.T1 wildtype mice (C) and trkB.T1 knockout mice (D) after receiving a saline hind paw injection (black bar; n = 3) or a hind paw injection of CFA (blue bar; n = 3). E and F. TrkB.FL protein expression in the dorsal horn of trkB.T1 wildtype (E) and trkB.T1 knockout (F) mice in the antiretroviral model was assayed via western blot. The top panels shows a representative western blot of trkB.FL following saline or d4T injection at 1d and 3d. Blots were stripped and re-probed with anti-actin to control for protein loading (shown in lower panel). Below the panel the results from saline treated (black bar; n = 3) and 1d (blue bar; n = 3) or 3d (red bar; n = 3) after d4T treatment are quantified. G and H. E and F. TrkB.FL protein expression in the dorsal horn in trkB.T1 wildtype (G) and trkB.T1 knockout (H) mice in the CFA model was assayed via western blot. The top panel shows a representative western blot of trkB.FL following a saline or CFA hind paw injection at 3d. Blots were stripped and re-probed with anti-actin to control for protein loading (shown in lower panel). Below the panel the results from saline treated (black bar; n = 3) and 3d CFA (blue bar; n = 3) are quantified.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1744-8069-5-61-S1.pdf>]

Acknowledgements

This work was supported by a grant from the National Institutes of Health to S. G. D. (NR009672).

References

- Thompson SW, Bennett DL, Kerr BJ, Bradbury EF, McMahon SB: **Brain-derived neurotrophic factor is an endogenous modulator of nociceptive responses in the spinal cord.** *Proc Natl Acad Sci USA* 1999, **96**:7714-7718.
- Miletic G, Miletic V: **Increases in the concentration of brain derived neurotrophic factor in the lumbar spinal dorsal horn are associated with pain behavior following chronic constriction injury in rats.** *Neurosci Lett* 2002, **319**:137-140.
- McMahon SB, Cafferty WB: **Neurotrophic influences on neuropathic pain.** *Novartis Found Symp* 2004, **261**:68-92.
- Pezet S, McMahon SB: **Neurotrophins: Mediators and modulators of pain.** *Annu Rev Neurosci* 2006, **29**:507-538.
- Michael GJ, Averill S, Shortland PJ, Yan Q, Priestley JV: **Axotomy results in major changes in BDNF expression by dorsal root ganglion cells: BDNF expression in large trkB and trkC cells, in pericellular baskets, and in projections to deep dorsal horn and dorsal column nuclei.** *Eur J Neurosci* 1999, **11**:3539-3551.
- Ha SO, Kim JK, Hong HS, Kim DS, Cho HJ: **Expression of brain-derived neurotrophic factor in rat dorsal root ganglia, spinal cord and gracile nuclei in experimental models of neuropathic pain.** *Neuroscience* 2001, **107**:301-309.
- Pezet S, Malcangio M, Lever IJ, Perkinson MS, Thompson SW, Williams RJ, McMahon SB: **Noxious stimulation induces trkB receptor and downstream ERK phosphorylation in spinal dorsal horn.** *Mol Cell Neurosci* 2002, **21**:684-695.
- Coull JA, Beggs S, Boudreau D, Boivin D, Tsuda M, Inoue K, Gravel C, Salter MW, DeKoningk Y: **BDNF from microglia causes the shift in neuronal anion gradient underlying neuropathic pain.** *Nature* 2005, **438**:1017-1021.
- Li L, Xian CJ, Zhong JH, Zhou XF: **Upregulation of brain-derived neurotrophic factor in the sensory pathway by selective motor nerve injury in adult rats.** *Neurotox Res* 2006, **9**:269-283.
- Guo W, Robbins MT, Wei F, Zhou S, Dubner R, Ren K: **Supraspinal brain-derived neurotrophic factor signaling: A novel mechanism for descending pain facilitation.** *J Neurosci* 2006, **26**:126-137.
- Renn CL, Lin L, Thomas S, Dorsey SG: **Full-length tropomyosin-related kinase B expression in the brainstem in response to persistent inflammatory pain.** *NeuroReport* 2006, **17**:1175-1179.
- Yajima Y, Narita M, Narita M, Matsumoto N, Suzuki T: **Involvement of a spinal brain-derived neurotrophic factor/full-length trkB pathway in the development of nerve injury-induced thermal hyperalgesia in mice.** *Brain Res* 2002, **958**:338-346.
- Chen X, Ye H, Kuruvilla R, Ramanan N, Scangos KW, Zhang C, Johnson NM, England PM, Shokat KM, Ginty DD: **A chemical-genetic approach to studying neurotrophin signaling.** *Neuron* 2005, **46**:13-21.
- Wang X, Ratnam J, Zou B, England PM, Basbaum AI: **TrkB signaling is required for both the induction and maintenance of tissue and nerve injury-induced persistent pain.** *J Neurosci* 2009, **29**:5508-5515.
- Middlemas DS, Lindberg RA, Hunter T: **TrkB, a neural receptor protein-tyrosine kinase: Evidence for a full-length and two truncated receptors.** *Mol Cell Biol* 1991, **11**:143-153.
- Biffo S, Offenhauser N, Carter BD, Barde YA: **Selective binding and internalization by truncated receptors restrict the availability of BDNF during development.** *Development* 1995, **121**:2462-2470.
- Eide FF, Vinig ER, Eide BL, Zang K, Wang XY, Reichardt LF: **Naturally occurring truncated trkB receptors have dominant inhibitory effects on brain-derived neurotrophic factor signaling.** *J Neurosci* 1996, **16**:3123-3129.
- Gonzalez M, Ruggiero FP, Chang Q, Shi YJ, Rich MM, Kraner S, Balice-Gordon RJ: **Disruption of trkB-mediated signaling induces disassembly of postsynaptic receptor clusters at neuromuscular junctions.** *Neuron* 1999, **24**:567-583.
- Saarelainen T, Pussinen R, Koponen E, Alhonen L, Wong G, Sirvio J, Castren E: **Transgenic mice overexpressing truncated trkB neurotrophin receptors in neurons have impaired long-term**

- spatial memory but normal hippocampal LTP.** *Synapse* 2000, **38**:102-104.
20. Haapasalo A, Koponen E, Hoppe E, Wong G, Castren E: **Truncated *trkB.T1* is a dominant negative inhibitor of *trkB.TK+*-mediated cell survival.** *Biophys Res Commun* 2001, **280**:1352-1358.
 21. Lahtinen S, Pitkanen A, Saarelainen T, Nissinen J, Koponen E, Castren E: **Decreased BDNF signaling in transgenic mice reduces epileptogenesis.** *Eur J Neurosci* 2002, **15**:721-734.
 22. Luikart BW, Nef S, Shipman T, Parada LF: **In vivo role of truncated *trkB* receptors during sensory ganglion neurogenesis.** *Neuroscience* 2003, **117**:847-858.
 23. Dorsey SG, Renn CL, Carim-Todd L, Barrick CA, Bambrick L, Krueger BK, Ward CW, Tessarollo L: **In vivo restoration of physiological levels of truncated *trkB.T1* receptor rescues neuronal cell death in a trisomic mouse model.** *Neuron* 2006, **51**:21-18.
 24. Carim-Todd L, Bath KG, Fulgenzi G, Yanpallewar S, Jing D, Barrick CA, Becker J, Buckley H, Dorsey SG, Lee FS, Tessarollo L: **Endogenous truncated *trkB.T1* receptor regulates neuronal complexity and *trkB* kinase receptor function in vivo.** *J Neurosci* 2009, **29**:678-685.
 25. Baxter GT, Radeke MJ, Kuo RC, Makrides V, Hinkle B, Hoang R, Medina-Selby A, Coit D, Valenzuela P, Feinstein SC: **Signal transduction mediated by the truncated *trkB* receptor isoforms, *trkB.T1* and *trkB.T2*.** *J Neurosci* 1997, **17**:2683-2690.
 26. Haapasalo A, Saarelainen T, Moshnyakov M, Arumae U, Kiema TR, Saarna M, Wong G, Castren E: **Expression of the naturally occurring truncated *trkB* neurotrophin receptor induces outgrowth of filopodia and processes in neuroblastoma cells.** *Oncogene* 1999, **18**:1285-1296.
 27. Yacoubian TA, Lo DC: **Truncated and full-length *trkB* receptors regulate distinct modes of dendritic growth.** *Nat Neurosci* 2000, **3**:342-349.
 28. Rose CR, Blum R, Pichler B, Lepier A, Kafitz KW, Konnerth A: **Truncated *trkB-T1* mediates neurotrophin-evoked calcium signaling in glia cells.** *Nature* 2003, **426**:74-78.
 29. Hartmann M, Brigadski T, Erdmann KS, Holtmann B, Sendtner M, Narz F, Lessmann V: **Truncated *trkB* receptor-induced outgrowth of dendritic filopodia involves the p75 neurotrophin receptor.** *J Cell Sci* 2004, **117**:5803-5814.
 30. Ohira K, Kumanoogoh H, Sahara Y, Homma KJ, Hirai H, Nakamura S, Hayashi M: **A truncated tropomyosin-related kinase B receptor, *T1*, regulates glial cell morphology via Rho GDP dissociation inhibitor 1.** *J Neurosci* 2005, **25**:1343-1353.
 31. Ohira K, Homma KJ, Hirai H, Nakamura S, Hayashi M: ***TrkB-T1* regulates the RhoA signaling and actin cytoskeleton in glioma cells.** *Biochem Biophys Res Commun* 2006, **342**:867-874.
 32. Ferrer I, Marin C, Rey MJ, Ribalta T, Goutan E, et al.: **BDNF and full-length and truncated *trkB* expression in Alzheimer disease: Implications in therapeutic strategies.** *J Neuropathol Exp Neurol* 1999, **58**:729-739.
 33. Dorsey SG, Bambrick LL, Balice-Gordon RJ, Krueger BK: **Failure of brain-derived neurotrophic factor-dependent neuron survival in mouse trisomy 16.** *J Neurosci* 2002, **22**:2571-2578.
 34. Li YX, Xu Y, Ju D, Lester HA, Davidson N, Schuman EM: **Expression of a dominant negative *trkB* receptor, *T1*, reveals a requirement for presynaptic signaling in BDNF-induced synaptic potentiation in cultured hippocampal neurons.** *Proc Natl Acad Sci USA* 1998, **95**:10884-10889.
 35. Maratou K, Wallace VC, Hasnie FS, Okuse K, Hosseini R, et al.: **Comparison of dorsal root ganglion gene expression in rat models of traumatic and HIV-associated neuropathic pain.** *Eur J Pain* 2009, **13**:87-98.
 36. Joseph EK, Chen X, Khasar SG, Levine JD: **Novel mechanism of enhanced nociception in a model of AIDS-therapy-induced painful peripheral neuropathy in the rat.** *Pain* 2004, **107**:147-158.
 37. Dorsey SG, Leitch CC, Renn CL, Lessans S, Smith BA, Wang XM, Dionne RA: **Genome-wide screen identifies drug-induced regulation of the gene giant axonal neuropathy (*Gan*) in a mouse model of antiretroviral-induced painful peripheral neuropathy.** *Biol Res Nurs* 2009, **11**(1):7-16.
 38. Merighi A, Salio C, Ghirri A, Lossi L, Ferrini F, Betelli C, Bardoni R: **BDNF as a pain modulator.** *Prog Neurobiol* 2008, **85**:297-317.
 39. Klein R, Smeyne RJ, Wurst W, Long LK, Auerbach BA, Joyner AL, Barbacid M: **Targeted disruption of the *trkB* neurotrophin receptor gene results in nervous system lesions and neonatal death.** *Cell* 1993, **75**:113-122.
 40. Frisen J, Verge VM, Fried K, Risling M, Persson H, Trotter J, Hokfelt T, Lindholm D: **Characterization of glial *trkB* receptors: Differential response to injury in the central and peripheral nervous systems.** *Proc Natl Acad Sci USA* 1993, **90**:4971-4975.
 41. Ren K: **An improved method for assessing mechanical allodynia in the rat.** *Physiol Behav* 1999, **67**:711-716.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

